



Publication Year	2015
Acceptance in OA @INAF	2020-03-18T09:51:10Z
Title	The physics inside the scaling relations for X-ray galaxy clusters: gas clumpiness, gas mass fraction and slope of the pressure profile
Authors	ETTORI, STEFANO
DOI	10.1093/mnras/stu2292
Handle	http://hdl.handle.net/20.500.12386/23344
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
Number	446

The physics inside the scaling relations for X-ray galaxy clusters: gas clumpiness, gas mass fraction and slope of the pressure profile

S. Ettori^{1,2}★

¹*INAF, Osservatorio Astronomico di Bologna, via Ranzani 1, I-40127 Bologna, Italy*

²*INFN, Sezione di Bologna, viale Berti Pichat 6/2, I-40127 Bologna, Italy*

Accepted 2014 October 29. Received 2014 October 14; in original form 2014 May 14

ABSTRACT

In galaxy clusters, the relations between observables in X-ray and millimetre wave bands and the total mass have normalizations, slopes and redshift evolutions that are simple to estimate in a self-similar scenario. We study these scaling relations and show that they can be efficiently expressed, in a more coherent picture, by fixing the normalizations and slopes to the self-similar predictions, and advocating, as responsible of the observed deviations, only three physical mass-dependent quantities: the gas clumpiness C , the gas mass fraction f_g and the logarithmic slope of the thermal pressure profile β_P . We use samples of the observed gas masses, temperature, luminosities and Compton parameters in local clusters to constrain normalization and mass dependence of these three physical quantities, and measure $C^{0.5}f_g = 0.110(\pm 0.002 \pm 0.002)(E_z M/5 \times 10^{14} M_\odot)^{0.198(\pm 0.025 \pm 0.04)}$ and $\beta_P = -\ln P/\ln r = 3.14(\pm 0.04 \pm 0.02)(E_z M/5 \times 10^{14} M_\odot)^{0.071(\pm 0.012 \pm 0.004)}$, where both a statistical and systematic error (the latter mainly due to the cross-calibration uncertainties affecting the *Chandra* and *XMM-Newton* results used in the present analysis) are quoted. The degeneracy between C and f_g is broken by using the estimates of the Compton parameters. Together with the self-similar predictions, these estimates on C , f_g and β_P define an intercorrelated internally consistent set of scaling relations that reproduces the mass estimates with the lowest residuals.

Key words: galaxies: clusters: general – cosmology: miscellaneous – X-rays: galaxies: clusters.

1 INTRODUCTION

To use galaxy clusters as probes of the background Universe in which they form and evolve is essential to link some of their observed properties in the electromagnetic spectrum to their gravitational potential (see e.g. Allen, Evrard & Mantz 2011; Kravtsov & Borgani 2012). Many proxies at different wavelengths, from radio to X-ray band, are nowadays available and robustly determined. Some attempts are already started to combine few of these proxies to improve the constraints on the inferred mass (e.g. Okabe et al. 2010; Stanek et al. 2010; Ettori et al. 2012; Ettori 2013; Evrard et al. 2014; Maughan 2014; Rozo et al. 2014).

In this work, we focus on the intracluster medium (ICM), the hot fully ionized optically thin plasma that collapses into the cluster gravitational potential. The physical processes occurring in the ICM can be mapped both with the X-rays, produced via bremsstrahlung radiation (e.g. Böhringer & Werner 2010) and through the Sunyaev–

Zeldovich (hereafter SZ) effect, that traces the Compton scattering of the photons of the cosmic microwave background on the electrons of the same plasma (Sunyaev & Zeldovich 1980).

In particular, we consider the scaling relations between cluster masses and the X-ray/SZ observables (see Giodini et al. 2013, for a recent review on this topic). We obtain, first, the analytic expressions that relate gas mass, temperature, luminosity and Compton parameter to the total mass and, then, we show that these relations, with the normalizations and slopes fixed to the analytic values, can be used more efficiently to estimate the total mass, once a set of three physically motivated quantities are defined also in their mass dependence.

The paper is organized as follows. In Section 2, we introduce the scaling relations considered for our analysis, providing a numerical value for the normalization that depends just on three unknown quantities, i.e. the average gas clumpiness, the cluster gas mass fraction and the slope of the gas pressure profile. In Section 3, we describe how we can calibrate the investigated scaling relations by using the largest sample available of hydrostatic mass measurements. In Section 4, we summarize our main findings. Hereafter,

★E-mail: stefano.ettori@oabo.inaf.it

all the physical quantities considered refer to the cosmological parameters $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $\Omega_m = 1 - \Omega_\Lambda = 0.3$, unless stated otherwise.

2 THE X-RAY AND SZ SCALING LAWS FOR THE TOTAL MASS

For a galaxy cluster in hydrostatic equilibrium, the radial profile of the total mass is described by the equation (e.g. Ettori et al. 2013)

$$M(< R) \equiv M = -\frac{R T(R) \frac{d \ln P}{d \ln r}}{\mu m_p G} = \frac{R T f_T \beta_P}{\mu m_p G}, \quad (1)$$

where $\beta_P = -d \ln P / d \ln r > 0$ is the opposite of the logarithmic slope of the gas pressure profile, and $f_T = T(R)/T$ is defined as the ratio between the 3D value of the gas temperature at the radius R and the mean spectroscopic estimate T , which will appear in the scaling relations.

Studies of the properties of the self-similar scaling scenario have shown to be more convenient to refer to cluster's regions defined with respect to a fixed overdensity when haloes with different masses and redshifts are considered (e.g. Böhringer, Dolag & Chon 2012). In our analysis, we consider physical quantities estimated within a radius R_Δ , which defines a spherical region where the mean mass overdensity Δ is evaluated with respect to the critical density of the Universe at the cluster's redshift z , $\rho_{c,z} = 3H_z^2/(8\pi G)$: $\Delta = 3M/(4\pi\rho_{c,z}R_\Delta^3) = 2GM/(H_z^2R_\Delta^3)$, where the Hubble constant $H_z = H_0E_z$ includes the factor describing its cosmic evolution $E_z = [\Omega_m(1+z)^3 + 1 - \Omega_m]^{1/2}$ for a flat cosmology with matter density parameter Ω_m .

We assume

(i) a gas mass fraction $f_g = M_g/M$;

(ii) that the X-ray emission is mostly due to bremsstrahlung processes so that the bolometric luminosity $L \equiv L_{\text{bol}} = \int n_e n_p \Lambda(T) dV = f_L M_g^2 c_f / (\mu_e^2 m_{\text{amu}}^2 V)$, where $V = 4/3\pi R_\Delta^3$ is the cluster volume; $f_L = \int n_e^2 dV / (\int n_e dV)^2 V$ is the correction needed to consider the gas mass ($\int n_e dV$) instead of the emission integral ($\int n_e^2 dV$) for the scaling purpose and is equal to 1.80 for a gas density distribution described by a β -model with $\beta = 0.65$ and $R_{500} = 5 \times$ the core radius, that are the median values of the estimated parameters in the sample of the brightest 45 nearby galaxy clusters in Mohr, Mathiesen & Evrard (1999);¹ the cooling function c_f is equal to $c_{f,0} \times T_{\text{keV}}^{0.5} \text{ erg s}^{-1} \text{ cm}^3$, with $c_{f,0} = 0.85 \times 10^{-23} c_{\text{pe}}$ (this value is completely consistent with, e.g., Sutherland & Dopita 1993 as tabulated in Tozzi & Norman 2001);² a conversion factor from protons to electrons $c_{\text{pe}} = 1.1995$, an electronic weight $\mu_e = 1.1738$ and an atomic mass $m_{\text{amu}} = 1.66 \times 10^{-24} \text{ g}$ are used;

¹ Using the extremes of the inter-quartile ranges of the estimated values of β and R_{500} in the Mohr et al. sample, we estimate the variations on f_L between -15 and $+29$ per cent, and, through the dependence to the power of $-3/4$, on the quoted normalization of the M - L relation between -17 and $+13$ per cent.

² The normalization of the cooling function is estimated by fitting a function $cT^{0.5}$ to the values of the cooling function evaluated with the thermal model `apec` in `XSPEC` (Arnaud 1996) where a metallicity of 0.3 times the solar abundance as tabulated in Anders & Grevesse (1989) and a set of temperature between 2 and 12 keV are considered; adopting a metallicity of 0.1 decreases the normalization of c_f by 5 per cent; the difference is -6 per cent when a metallicity of 0.3 and the more recent table of solar abundance from Asplund et al. (2009) are considered.

(iii) that the millimetre wave emission is due to the SZ effect which is proportional to the integrated pressure of the X-ray emitting plasma along the line of sight and is described from the integrated Compton parameter $Y_{\text{SZ}} D_\Delta^2 = (\sigma_T / m_e c^2) \int P dV$, where D_Δ is the angular distance to the cluster, $\sigma_T = 8\pi/3(e^2/m_e c^2)^2 = 6.65 \times 10^{-25} \text{ cm}^2$ is the Thomson cross-section, m_e and e are the electron rest mass and charge, respectively, c is the speed of light and $P = n_e T$ is the electron pressure profile.

We can write the following scaling laws with their calculated normalization

$$\begin{aligned} & \frac{F_z M}{5 \times 10^{14} M_\odot} \\ &= 1.0 \left(\frac{C^{0.5} f_g}{0.1} \right)^{-1} \frac{F_z M_g}{5 \times 10^{13} M_\odot} \\ &= 0.832 \left(\frac{\beta_P}{3} \right)^{3/2} \left(\frac{kT}{5 \text{ keV}} \right)^{3/2} \\ &= 0.962 \left(\frac{\beta_P}{3} \right)^{3/8} \left(\frac{C^{0.5} f_g}{0.1} \right)^{-3/2} \left(\frac{F_z^{-1} L}{5 \times 10^{44} \text{ erg s}^{-1}} \right)^{3/4} \\ &= 1.748 \left(\frac{\beta_P}{3} \right)^{3/5} \left(\frac{f_g}{0.1} \right)^{-3/5} \left(\frac{F_z Y_{\text{SZ}} D_\Delta^2}{10^{-4} \text{ Mpc}^2} \right)^{3/5}. \end{aligned} \quad (2)$$

Here, we define $F_z = E_z \times (\Delta/500)^{0.5}$ (see e.g. Ettori et al. 2004); the clumpiness in the gas density $C = \langle n_g^2 \rangle / \langle n_g \rangle^2$ that affects the measurement of the gas density as obtained from the deprojection of the X-ray data produced from free-free emission, but not from SZ signal due to inverse Compton (see e.g. Eckert et al. 2013a,b; Roncarelli et al. 2013); the mean atomic weight $\mu = 0.61$. As reference values, we adopt an overdensity of $\Delta = 500$ (and therefore $F_z = E_z$), for which $f_T \approx 0.67$ (e.g. Vikhlinin et al. 2006; Baldi et al. 2012); a gas fraction of 0.1 (see e.g. Ettori et al. 2009; Mantz et al. 2014); a logarithmic slope of the gas pressure profile at R_{500} of -3 , which is consistent with the values in the range $(-3.2, -2.8)$ of the profiles adopted in Arnaud et al. (2010) and in the papers of the Planck Intermediate Results V (2013). We refer to the appendix for further details on how the normalizations are estimated (Appendix A) and to the extension of the M - L relation to no-bolometric energy bands (Appendix B).

Following Ettori (2013, hereafter E13), where a generalized form for the scaling laws has been presented, a concise form of all the set of the above equations can be written as

$$F_z M \sim \beta_P^\theta f_g^{-\phi} (F_z^{-1} L)^\alpha (F_z M_g)^\beta T^\gamma \quad (3)$$

where the relations

$$4\alpha + 3\beta + 2\gamma = 3$$

$$\theta = \alpha/2 + \gamma$$

$$\phi = 2\alpha + \beta \quad (4)$$

among the exponents hold in a self-similar scenario (e.g. the M - T relation is recovered by imposing the absence of any dependence on M_g and L , i.e. $\alpha = \beta = 0$; then $\gamma = 3/2$, $\theta = \gamma = 3/2$ and $\phi = 0$), and Y_{SZ} is here represented as the product of gas mass and temperature.




Figure 1. Left: comparison between our collection of hydrostatic masses (S09 = Sun et al. 2009; P09 = Pratt et al. 2009; R11 = Reichert et al. 2011; M13 = Mahdavi et al. 2013; M14 = Maughan 2014) and M_{YX} as estimated in Planck Collaboration XXIX (2014). Right: median values (with the lower and upper quartile) of the ratios M/M_{YX} for the whole sample (black), in four different mass bins (green) and for the different considered data sets.

3 THE CALIBRATION OF THE SCALING RELATIONS

To check the consistency between the normalizations in equation (2) and the observed distributions, we consider X-ray mass estimates obtained through the application of the equation of the hydrostatic equilibrium under the assumptions that any gas velocity is zero and that the ICM is distributed in a spherically symmetric way into the cluster gravitational potential (see e.g. Ettori et al. 2013). We start with the sample described in E13 [120 entries, 113 of which are unique hydrostatic mass measurements collected from the public catalogues in Sun et al. (2009), Pratt et al. (2009),³ Mahdavi et al. (2013) and Maughan (2014)] and add the 110 (out of 232) objects present in the Reichert et al. (2011) sample and not considered in E13, for a total number of 213 galaxy clusters with reliable hydrostatic masses estimated at $\Delta = 500$. Using this sample, we investigate, as described below, the normalization and slope of the M - T relation (213 objects), M - L relation (199 objects) and M - M_{gas} relation (113 objects). Then, we consider the Planck catalogue (file *COM_PCCS_SZ-validation_R1.13.fits* available at http://www.sciops.esa.int/index.php?page=Planck_Legacy_Archive&project=planck; see Planck Collaboration XXIX 2014) with 1227 entries, 455 of which with estimated redshift and $Y_{\text{SZ}} \equiv Y_{500, \text{PSX}} > 0$. We obtain that 94 are the systems in common between the 213 galaxy cluster with hydrostatic masses and the 455 Planck clusters. In the published catalogue, also estimates of the mass, M_{YX} , obtained through the $Y_X = M_g T$ parameter (e.g. Kravtsov, Vikhlinin & Nagai 2006; Arnaud et al. 2010) are provided. For the 94 objects

³ The masses derived in Pratt et al. (2009) for the objects in the REXCESS sample are not obtained from the equation of the hydrostatic equilibrium, but are estimated from the $Y_X (= M_g T)$ - M relation as calibrated in Arnaud, Pointecouteau & Pratt (2007). We consider them in our sample for the wealth of information associated with the REXCESS catalogue.

in common, we calculate the ratio between the collected values of the hydrostatic mass and M_{YX} . We obtain overall perfect agreement (median value: 0.99). On the other hand, we also notice a clear bias depending on the total hydrostatic mass, with systems at lower ($< 3 \times 10^{14} M_{\odot}$) and higher ($> 7 \times 10^{14} M_{\odot}$) masses showing the highest deviations (median values of 1.37 and 0.74, respectively; see Fig. 1), indicating that the collected hydrostatic estimates over (under) predict the high (low) values of M_{YX} .

The main properties of the sample here analysed are listed in Table 1.

We fit these quantities using the linear function

$$\mathcal{Y} = n + a\mathcal{X} \quad (5)$$

and minimizing the merit function

$$\chi^2 = \sum_{i=1}^N \frac{(\mathcal{Y}_i - n - a\mathcal{X}_i)^2}{\epsilon_i^2} \quad (6)$$

$$\epsilon_i^2 = \epsilon_{\mathcal{Y},i}^2 + a^2 \epsilon_{\mathcal{X},i}^2 - 2a\rho \epsilon_{\mathcal{Y},i} \epsilon_{\mathcal{X},i},$$

where $\mathcal{Y} = \log\left(\frac{F_z M}{5 \times 10^{14} M_{\odot}}\right)$, $\mathcal{X} = \log(X)$ and X is equal to $\frac{F_z M_g}{5 \times 10^{13} M_{\odot}}$, $\frac{T}{5 \text{ keV}}$, $\frac{F_z^{-1} L_{\text{bol}}}{5 \times 10^{44} \text{ erg s}^{-1}}$, $\frac{F_z Y_{\text{SZ}} D_A^2}{10^{-4} \text{ Mpc}^2}$; ‘log’ indicates the base-10 logarithm; the associated errors $\epsilon_{\mathcal{Y}}$ and $\epsilon_{\mathcal{X}}$ are obtained through the propagation of the measured uncertainties; N is the number of data points and $D = N - p$ are the degrees of freedom given a number p of fitted parameters (either 2 – slope and normalization – or the normalization only); ρ is the Pearson’s correlation coefficient among the variables \mathcal{Y} and \mathcal{X} . An intrinsic scatter is estimated by adding it in quadrature to ϵ_i and re-iterating the fitting procedure until a reduced χ^2 of 1 is obtained. The relative error on it is obtained as discussed in E13. The fit is performed using the IDL routine MPFIT (Markwardt 2008).

Table 1. Properties of the multiwavelength samples considered in the present analysis. In the columns M and X , the median value, the range covered (in parentheses) and the relative error of the mass and the investigate observable, respectively, are quoted. The units for the observables X are $10^{13} M_\odot$ for M_g , keV for T , 10^{44} erg s $^{-1}$ for L and $D_A^2 10^4$ Mpc $^{-2}$ for Y_{SZ} .

Sample	N	z	M ($10^{14} M_\odot$)	X
All M	213	0.226 (0.012–1.390)	3.51 (0.15–22.80); 0.19	–
$M-M_g$	109	0.141 (0.012–0.550)	3.15 (0.15–14.52); 0.19	3.91 (0.08–26.7); 0.09
$M-T$	213	0.226 (0.012–1.390)	3.51 (0.15–22.80); 0.19	5.30 (0.81–12.5); 0.07
$M-L$	199	0.231 (0.012–1.390)	3.52 (0.15–22.80); 0.20	6.80 (0.02–118.2); 0.04
$M-Y_{SZ}$	94	0.176 (0.048–0.548)	5.99 (0.98–14.52); 0.19	0.66 (0.06–3.8); 0.14

Although we provide all the calculations needed to investigate the evolution with redshift of the scaling relations, we prefer not to study it in the present work because of the heterogeneous origin of the considered data set that, without a proper weight provided from a redshift-dependent selection function, could affect any conclusion on the redshift evolution.

3.1 A mass-dependent deviation from self-similarity

Since the first evidence of the deviations of the observed slopes of the X-ray scaling laws from the self-similar expectations, it has been suggested that a possible solution to reconcile the predicted and observed values can be obtained by assuming that at least one of the physical quantities (like, e.g., the gas mass fraction) appearing in the derivation of the scaling law has a not-negligible mass dependence (see e.g. Arnaud & Evrard 1999; Pratt et al. 2009). In this section, we investigate how we can constrain the mass dependence of the set of the physical quantities we need for a complete description of the scaling relations, by imposing that this mass dependence is fully responsible for any observed deviation from the self-similar prediction.

In general, we can write the scaling relations here investigated between the mass M and an observable X as $M = \mathcal{N}^c X^a E_z^b$. Note that, for sake of completeness, we are considering a normalization with its own exponent (to treat the case of, e.g., β_P and f_g in equation 2) and a term for the redshift evolution proportional to E_z . In the case that the normalization \mathcal{N} depends upon the mass M , $\mathcal{N} = \mathcal{N}_0 M^m$, the scaling relations are then modified accordingly:

$$M = \mathcal{N}_0^{c/(1-cm)} X^{a/(1-cm)} E_z^{b/(1-cm)} = \mathcal{N}_{\text{obs}} X^{a_{\text{obs}}} E_z^{b_{\text{obs}}}, \quad (7)$$

where the subscript *obs* in the last member refers to the values measured by a best-fitting procedure leaving normalization (\mathcal{N}_{obs}), slope (a_{obs}) and redshift evolution (b_{obs}) free to vary. By imposing $c = a = a_{\text{exp}}$, where a_{exp} is the nominal exponent predicted in the self-similar scenario, and equating the second and third members of equation (7), we obtain that the ‘intrinsic’, mass-corrected scaling relations can be recovered by estimating

$$\begin{aligned} m &= 1/a_{\text{exp}} - 1/a_{\text{obs}} \\ \mathcal{N}^c &= \mathcal{N}_{\text{obs}}^{1-cm} = \mathcal{N}_{\text{obs}}^{a_{\text{exp}}/a_{\text{obs}}} \\ b &= b_{\text{obs}}(1 - cm) = b_{\text{obs}} a_{\text{exp}}/a_{\text{obs}}. \end{aligned} \quad (8)$$

Following the expression of equation (2), where all the scaling relations can be written as a function of $(E_z M)$, we assume a similar dependence on the mass of the three unknown parameters (i.e. the logarithmic slope of the pressure β_P , the gas mass fraction f_g and

the gas clumpiness C):

$$\begin{aligned} \beta_P &= \beta_{P,0} (E_z M)^{m_1} \\ f_g &= f_{g,0} (E_z M)^{m_2} \\ C &= C_0 (E_z M)^{m_3}. \end{aligned} \quad (9)$$

Using equation (9), together with equations (3) and (7), we can then write a general expression in the form

$$(E_z M)^{1-\theta m_1 + \phi m_2 + \phi m_3/2} \sim \beta_{P,0}^\theta C_0^{-\phi/2} f_{g,0}^{-\phi} (E_z^{-1} L)^\alpha (E_z M_g)^\beta T^\gamma, \quad (10)$$

which can be resolved in each of the scaling laws considered here (see equations 2 and 4) as

$$\begin{aligned} E_z M &\sim (C_0^{0.5} f_{g,0})^{-1/(1-m)} (E_z M_g)^{1/(1-m)}; \quad m = -m_2 - m_3/2 \\ &\sim \beta_{P,0}^{3/2/(1-3/2m)} (kT)^{3/2/(1-3/2m)}; \quad m = m_1 \\ &\sim \beta_{P,0}^{3/8/(1-3/4m)} f_{g,0}^{-3/2/(1-3/4m)} \\ &\quad (E_z^{-1} L)^{3/4/(1-3/4m)}; \quad m = m_1/2 - 2m_2 - m_3 \\ &\sim \beta_{P,0}^{3/5/(1-3/5m)} f_{g,0}^{-3/5/(1-3/5m)} \\ &\quad (E_z Y_{SZ} D_A^2)^{3/5/(1-3/5m)}; \quad m = m_1 - m_2. \end{aligned} \quad (11)$$

Here, the symbol ‘ \sim ’ is used to replace all the factors and pivot values shown in equation (2). These equations show explicitly the quantities that can be constrained by fitting a linear function, with normalization and slope as free parameters, to the logarithmic values of the mass and of the observables. For example, by fitting the $M-T$ relation, one can directly estimate $m = m_1$ from the best-fitting value of the slope and $\beta_{P,0}$ from the best-fitting value of the normalization. In the following subsection, we show how we can constrain the parameters of our interest, defined in equation (9), by combining the results obtained from the linear fit of the scaling relations and quoted in Table 2.

Once the dependence on the mass is assessed, we can fit the scaling relation by fixing the expected slope a_{exp} and propagating the correction to the total mass:

$$(1 - \theta m_1 + \phi m_2 + \phi m_3/2) \log(E_z M) = \bar{n} + a_{\text{exp}} \log(X). \quad (12)$$

The normalization \bar{n} is the only free parameter and is used to calibrate finally the gas mass fraction, $f_{g,0}$, the gas clumpiness, C_0 , and the logarithmic slope of the gas pressure, $\beta_{P,0}$.

Table 2. Best-fitting results for the scaling relations investigated. N is the number of fitted data; ρ is the Pearson’s correlation coefficient; n and a refer to equation (5); χ_r^2 is the reduced χ^2 ; σ_i is the intrinsic scatter in $\log M$ at given observable \mathcal{X} ; m and \mathcal{N}^c are described in equation (8) and are obtained from the fits with normalization n and slope a as free parameters. For each scaling relation, we provide the best-fitting results obtained with (1) equation (5) with the slope fixed to the self-similar value and the sub-sample of ‘local’ (i.e. $z < 0.15$) and ‘massive’ (i.e. $M_{500} > 3 \times 10^{14} M_\odot$) objects; (2) as for (1), but leaving the slope free to vary; (3) equation (12); (4–6) as for (1–3), but for the subsample of the all ‘local’ clusters. In particular, the third row of the ‘local’ samples [i.e. fit (3) and (6)] refers to the best-fitting results for a fixed slope and including the mass dependence described in equation (12). Finally, the fit (7) is obtained by equation (12) with the normalization fixed after the calibration of the physical quantities described in equation (9) (see the instructions at the end of Section 3.2). In Fig. C1, we show the samples and the best-fitting lines that represent the results for the fits (4–7).

Sample	N	ρ	$10^n = \mathcal{N}_{\text{obs}}$	a	χ_r^2	σ_i	m	\mathcal{N}^c
$M-M_g$								
(1) Local massive	16	0.94	0.883 ± 0.012	1.000	6.9	0.053 ± 0.013	–	–
(2)			0.923 ± 0.033	0.751 ± 0.067	2.3	0.028 ± 0.023	-0.332 ± 0.119	0.899 ± 0.112
(3) (equation 12)			0.912 ± 0.017	1.000	1.4	0.019 ± 0.016	–	–
(4) Local all	59	0.97	1.065 ± 0.005	1.000	44.4	0.096 ± 0.010	–	–
(5)			0.848 ± 0.025	0.835 ± 0.017	7.8	0.048 ± 0.007	-0.198 ± 0.025	0.821 ± 0.031
(6) (equation 12)			0.912 ± 0.011	1.000	1.0	0.000 ± 0.003	–	–
(7)			0.912	1.000	1.0	0.000 ± 0.004	–	–
$M-T$								
(1) Local massive	29	0.90	0.927 ± 0.017	1.500	1.7	0.043 ± 0.016	–	–
(2)			0.947 ± 0.034	1.390 ± 0.119	1.6	0.044 ± 0.018	-0.053 ± 0.062	0.943 ± 0.094
(3) (equation 12)			0.890 ± 0.017	1.500	1.9	0.042 ± 0.014	–	–
(4) Local all	73	0.97	0.780 ± 0.008	1.500	4.9	0.073 ± 0.009	–	–
(5)			0.873 ± 0.021	1.679 ± 0.033	3.6	0.055 ± 0.008	0.071 ± 0.012	0.886 ± 0.027
(6) (equation 12)			0.881 ± 0.014	1.500	1.0	0.007 ± 0.011	–	–
(7)			0.890	1.500	1.1	0.009 ± 0.011	–	–
$M-L$								
(1) Local massive	22	0.87	0.830 ± 0.022	0.750	2.2	0.060 ± 0.019	–	–
(2)			0.855 ± 0.054	0.679 ± 0.088	2.2	0.061 ± 0.021	-0.138 ± 0.190	0.841 ± 0.131
(3) (equation 12)			0.909 ± 0.035	0.750	1.6	0.070 ± 0.031	–	–
(4) Local all	60	0.90	1.118 ± 0.017	0.750	11.1	0.158 ± 0.017	–	–
(5)			0.821 ± 0.046	0.609 ± 0.028	12.6	0.103 ± 0.017	-0.309 ± 0.076	0.784 ± 0.063
(6) (equation 12)			0.837 ± 0.023	0.750	2.5	0.122 ± 0.020	–	–
(7)			0.852	0.750	2.6	0.124 ± 0.020	–	–
$M-Y_{\text{SZ}}$								
(1) Local massive	27	0.79	1.960 ± 0.038	0.600	4.0	0.089 ± 0.017	–	–
(2)			1.923 ± 0.169	0.578 ± 0.093	4.1	0.091 ± 0.018	-0.063 ± 0.277	1.971 ± 0.371
(3) (equation 12)			2.113 ± 0.052	0.600	2.5	0.080 ± 0.021	–	–
(4) Local all	36	0.90	1.753 ± 0.024	0.600	9.6	0.110 ± 0.016	–	–
(5)			2.165 ± 0.369	0.774 ± 0.127	9.2	0.098 ± 0.032	0.374 ± 0.212	1.820 ± 0.387
(6) (equation 12)			2.083 ± 0.044	0.600	2.9	0.101 ± 0.024	–	–
(7)			2.113	0.600	2.9	0.103 ± 0.024	–	–

3.2 The best-fitting constraints

In an ideal case, where the samples analysed have a well-known selection function, a direct constraint on the dependence of the scaling relations on the mass (and the redshift) could be obtained by applying the equations listed above. In particular, from equation (11), one can recover

- (i) $\beta_{P,0}$ and m_1 from the best-fitting normalization and slope of the observed $M-T$ relation;
- (ii) using (i), $f_{g,0}$ and m_2 from the best-fitting normalization and slope of the observed $M-Y_{\text{SZ}}$ relation;
- (iii) using (ii), C_0 and m_3 from the best-fitting normalization and slope of the observed $M-M_g$ (or $M-L$) relation.

However, considering that (a) our data sets have been collected from the literature (see discussion in E13) and, thus, cannot be treated as a statistically well-defined sample, and (b) a different definition of R_{500} as recovered from, e.g., hydrostatic masses and

M_{Y_X} (see, for instance, the mass-dependent bias shown in Fig. 1) affects the reconstructed Planck Y_{SZ} signal, we decide to proceed differently. First, we decide not to use the $M-Y_{\text{SZ}}$ relation to calibrate the gas mass fraction. This implies that we have to deal with a degeneracy between the mass dependence on f_g and C . Therefore, we fix $m_3 = 0$, assuming that the gas clumpiness does not have any significant dependence on the cluster mass (see e.g. Nagai & Lau 2011 and Roncarelli et al. 2013, where a marginal mass dependence for simulated systems appears at radii beyond R_{200} , but it is almost negligible at R_{500}). Secondly, we use the whole sample of local systems (‘local all’ sample in Table 2) to quantify the mass dependence in equation (9). To do that, we use equation (8) and compute the corrected values of the normalization \mathcal{N}^c from the observed best-fitting parameters. The best-fitting values of m are quoted in Table 2 and imply that $m_1 = 0.071 \pm 0.012$ and $m_2 = 0.198 \pm 0.025$. Thirdly, to constrain the normalizations $\beta_{P,0}$ and $C^{0.5} f_{g,0}$, we analyse the sub-samples of the nearby ($z < 0.15$), massive ($M_{500} > 3 \times 10^{14} M_\odot$) galaxy clusters (‘local massive’ sample in Table 2). Doing that, we




Figure 2. From left top to right bottom, we show the process of the calibration of the three physical quantities we require to define the normalization of the scaling relations. Left top: ratios between the observed and expected normalizations of equation (2). Filled points refer to the best-fitting results measured for nearby massive systems ($M_{500} > 3 \times 10^{14} M_{\odot}$ and $z < 0.15$). For sake of completeness, we also show (but do not use in our calculations) the ratios measured for clusters at $z > 0.15$ and divided into two redshift bins (biggest points: subsample including the half of the most massive ones). Right top: calibration of $C^{0.5} f_{\text{gas}}$ and β_{P} using only the nearby massive systems (filled dots in the left-hand panel). The red circle represents the reference values of $(\beta_{\text{P}}, f_{\text{g}}) = (3, 0.1)$ with a relative uncertainty of 10 per cent. Left bottom: combination of the constraints from the normalizations of the M – M_{g} , M – L and M – Y_{SZ} relations in the gas mass fraction–gas clumpiness plane. The label ‘ (M_{Y_X}) ’ indicates the constraint on f_{g} obtained from the M_{Y_X} – Y_{SZ} relation (see Section 3.2). Right bottom: as in the upper-left panel, after correcting for the best-fitting values of $(\beta_{\text{P}}, f_{\text{g}})$ in equation (13) as discussed in Sections 3.1 and 3.2.

minimize the effect of a mass and redshift dependence on these values, and avoid any significant Malmquist bias due to the fact that the average luminosity of selected clusters is higher than that in the parent population in a flux limited sample (e.g. Stanek et al. 2006; Pratt et al. 2009). Proceeding in this way, we constrain $\beta_{\text{P},0}$ and $C^{0.5} f_{\text{g},0}$ from the best-fitting normalization of the observed M – T and M – M_{g} relation, respectively, and obtain $C_0^{0.5} f_{\text{g},0} = 0.110(\pm 0.002)$ and

$\beta_{\text{P},0} = 3.14(\pm 0.04)$, respectively, at $\Delta = 500$ (see the top-right panel of Fig. 2).

For sake of completeness, we show in Fig. 2 also the ratios between the estimated normalization and the expected value obtained in two redshift bins (defined with respect to the median value in the interval $0.15 - \max(z)$) and in two mass bins (build accordingly to the median value in each redshift bin). These ratios indicate that

our procedure is already capable of reproducing reasonably well the scaling relations for systems in the low-mass and/or high-redshift regime. On the other hand, a proper treatment of these cases requires the adoption of the selection function used to define our sample. This treatment is beyond the purpose of the present work and can be avoided just considering local, and massive, objects.

Considering now the M - Y_{SZ} relation, where the normalization is independent from the clumpiness, we can break the degeneracy between C and f_g (see the bottom-left panel of Fig. 2) and obtain $C_0 = 2.07 (\pm 0.02)$ and $f_{g,0} = 0.076 (\pm 0.003)$.

To summarize, we calibrate the new formalism in the following way:

(i) using the ‘local all’ sample, we quantify the mass dependence m_1 and m_2 (m_3 is fixed equal to 0) using equation (11) (see the fit labelled (5) in Table 2);

(ii) we estimate $\beta_{p,0}$ and $C_0^{0.5} f_{g,0}$ in the ‘local massive’ samples through the M - T and M - M_g relation, respectively, by equation (12) (see the fit labelled (3) in Table 2);

(iii) the degeneracy between C_0 and $f_{g,0}$ is broken with the M - Y_{SZ} relation for the ‘local massive’ systems.

All the quoted errors are at 1σ level and originate from the statistical uncertainties only. When we take into account the uncertainties related to the cross-calibration between *Chandra* and *XMM-Newton* on the gas temperature, gas mass, gas luminosity and hydrostatic mass as discussed, e.g., in Maughan (2014) and Mahdavi et al. (2013; also private communication), systematic errors of ± 0.002 and ± 0.02 affect the normalization of $C_0^{0.5} f_g$ and β_p , respectively, whereas the error associated with the slope of the mass dependence is about ± 0.04 and ± 0.004 , respectively.

Once we have constrained the normalizations and mass dependence of the quantities in equation (9), we re-estimate the ratios between the normalizations of the scaling relations and the predicted values. As shown in Fig. 2 (panel at the bottom right), we obtain a match of the order of few per cent for all the set of scaling laws investigated. The fit labelled (7) in Table 2 indicates the results obtained by fixing both the slope (to the self-similar expectation) and the normalization (after the calibration described above) of the scaling relations. Both the reduced χ^2 and the intrinsic scatter are lower than in the scaling laws where normalizations and slopes are used as free parameters.

3.3 Comparison with previous work

The constraint on the value of $C_0^{0.5} f_g$ is perfectly consistent with the results on the gas mass fraction obtained from recent work on both X-ray observations and the most recent hydrodynamical numerical simulations. By combining observational constraints from Vikhlinin et al. (2006), Arnaud et al. (2007) and Sun et al. (2009), Pratt et al. (2009) quote a gas mass fraction at $\Delta = 500$ of $0.113 (\pm 0.005) (M/5 \times 10^{14} M_\odot)^{0.21 (\pm 0.03)}$. Planelles et al. (2013), using a set of cosmological smoothed particle hydrodynamical simulations of massive ($M_{500} > 2.8 \times 10^{14} M_\odot$) galaxy clusters, measure, in the redshift range 0–1, a mean gas mass fraction in the range between 0.105 (for simulations including radiative cooling, star formation and feedback from supernovae) and 0.140 (for the non-radiative set), with an average value of 0.117 (and an rms of 0.008) for the objects simulated also accounting for the effect of feedback from active galactic nuclei. This would require $C \approx 1$, implying that the considered Y_{SZ} signal is biased high by about $(0.110/0.076) \sim 45$ per cent at given mass. This amount is difficult to explain with some selection effect, also considering that 22 (out




Figure 3. Constraints on the logarithmic slope of the pressure profile as a function of M_{500} . The dotted lines show the 1σ uncertainty associated with the best-fitting result (dashed line; see equation 13). The points refer to the best-fitting models adopted in Planck Intermediate Results V (2013) and Arnaud et al. (2010) as labelled.

of 27) of the systems included in the ‘local massive’ sample have a signal-to-noise ratio related to the SZ detection in correspondence of the X-ray position larger than 7 (all the local, massive objects have a signal-to-noise ratio in the range 5.7–26.5, with a median value of 10.3), making them less prone to any Malmquist-like bias propagated through the sample selection (see e.g. discussion in section 7.5.2 of Planck Collaboration XXIX 2014). On the other hand, if we replace the hydrostatic masses with the values M_{Y_x} estimated through the Y_x parameter (see a discussion on the comparison between them in Section 3 and Fig. 1) and fit equation (12), we measure $\mathcal{N}_{\text{obs}} = 1.670 (\pm 0.008)$ that implies a gas mass fraction of $0.113 (\pm 0.002)$ and, combined with the result on the M - M_g relation, a gas clumpiness slightly lower than the physically motivated lower bound of 1 ($C \sim 0.95$). We conclude that, for the available data set, some tension between hydrostatic M and M_{Y_x} is present (see, for instance, the mass-dependent bias shown in Fig. 1) that does not permit to break univocally the degeneracy between f_g and C . We recognize also that more work on this topic, with a more extended and detailed comparison between hydrostatic masses and integrated Compton parameters, is needed, but beyond the purpose of the present study.

The mass dependence of the gas pressure profile (see results for the M - T relation for the ‘local all’ sample in Table 2) is not in contrast with the present observational constraints (e.g. Arnaud et al. 2010; Sun et al. 2011; Planck Intermediate Results V 2013). In Fig. 3, we show our best-fitting constraints compared to the predictions from the best-fitting values of the universal model presented in Arnaud et al. (2010) and in Planck Intermediate Results V (2013). This universal model is obtained by combining observational data based on *XMM-Newton* observations in the radial range 0.03 – $1 R_{500}$ with hydrosimulation results out to $4 R_{500}$ and using a functional form as originally proposed in Nagai, Kravtsov & Vikhlinin (2007) to fit the combined re-scaled profile. Our result on the logarithmic slope of the pressure profile at R_{500} , β_p , shows a steeper mass

dependence, with values that lie between 2.8 at $\sim 10^{14} M_{\odot}$, preferred also from the Planck Collaboration best-fitting parameters, and 3.3 at $\sim 10^{15} M_{\odot}$, more in agreement with the Arnaud et al. profile.

4 SUMMARY AND DISCUSSION

In the present work, we estimate the predicted values of the normalization and slope of the scaling relations holding between the hydrostatic mass and (i) the gas mass, (ii) the gas temperature, (iii) the X-ray bolometric luminosity and (iv) the integrated Compton parameter. We show in details how these normalizations depend upon the gas density clumpiness C , the gas mass fraction f_g and the logarithmic slope of the thermal pressure profile β_p . We argue that the deviations of the observed slopes from the self-similar expectations can be fully explained with a mass dependence of the gas mass fraction and the logarithmic slope of the thermal pressure profile.

Relying on the availability of large data base of measured hydrostatic masses and observables in X-ray and millimetre wave bands, we constrain at high significance the normalization and mass dependence of the gas mass fraction and the logarithmic slope of the thermal pressure profile, putting also some limits on the level of gas clumpiness requested to accommodate in a self-consistent scenario all the set of the scaling relations. We conclude the following.

(i) The three astrophysical quantities (i.e. gas clumpiness, gas mass fraction and slope of the pressure profile) advocated to explain consistently the predicted M - $\{M_g, T, L, Y_{SZ}\}$ relations are sufficient to define the observed normalization and slope of these scaling laws.

(ii) Using nearby ($z < 0.15$), massive ($M_{500} > 3 \times 10^{14} M_{\odot}$) galaxy clusters, the M - M_g requires ($C^{0.5} f_g$) = 0.110 (± 0.002). Using the further constraint obtained from the clumpiness-free normalization of the M - Y_{SZ} relation, we obtain that, within R_{500} , the gas clumpiness is 2.07 (± 0.02) and the gas mass fraction is 0.076 (± 0.003) (see Fig. 2).

(iii) We note, however, that being the constraint on ($C^{0.5} f_g$) well in agreement with results from, e.g., recent hydrodynamical simulations on the cluster gas mass fraction at $\Delta = 500$ (e.g. Planelles et al. 2013), it would suggest that $C \approx 1$ and that the considered Y_{SZ} signal is biased high by $(0.110/0.076) \sim 45$ per cent at given mass.

(iv) Considering that most of the galaxy clusters included in the ‘local massive’ sample have a signal-to-noise ratio related to the SZ detection in correspondence of the X-ray position larger than 7, we exclude any significant Malmquist-like bias affecting the analysed sample.

(v) On the other hand, if we replace the hydrostatic masses with the values obtained from the Planck Collaboration through the Y_X parameter and carry on the same analysis, we obtain indeed that $C \sim 1$. However, this result highlights a tension between the measurements of M and M_{Y_X} for the same objects, with the hydrostatic estimates that over (under) predict the high (low) values of M_{Y_X} by about 30 per cent (see Fig. 1).

(vi) Using the same sample of local and massive galaxy clusters and the M - T relation, we constrain $\beta_p = -\ln P / \ln r = 3.14 (\pm 0.04)$.

(vii) We quantify the dependence upon the mass of the two adopted quantities (the clumpiness is assumed to be independent from the mass, i.e. $m_3 = 0$) through the best-fitting parameters of equation (11), and obtain $f_g \sim M^{0.20 \pm 0.02}$ and $\beta_p \sim M^{0.07 \pm 0.01}$; while the former is in good agreement both with other observational results and profiles predicted from hydrodynamical simulations, the latter one shows agreement with the Planck Intermediate Results

V (2013) best-fitting parameters at lower ($\sim 10^{14} M_{\odot}$) masses and with the Arnaud et al. (2010) profile at higher ($\sim 10^{15} M_{\odot}$) masses, requiring a steeper mass dependence.

(viii) By adjusting for the mass dependence of f_g and β_p , we demonstrate (see results labelled with ‘equation (12)’ and fit (7) in Table 2) that the scaling relations with a slope fixed to the expected value in the self-similar scenario provide best-fitting results with a reduced χ^2 and an intrinsic scatter comparable to the results obtained leaving the slope free to vary.

Therefore, we conclude that the scaling relations based on X-ray/SZ quantities have a simple and predictable behaviour that can be fully described at $\Delta = 500$ by the equations (2) and (9) (or their formal extension in equation 10), where

$$C^{0.5} f_g = 0.110(\pm 0.002) \left(\frac{E_z M}{5 \times 10^{14} M_{\odot}} \right)^{0.198(\pm 0.025)}$$

$$\beta_p = - \frac{d \ln P}{d \ln r} = 3.14(\pm 0.04) \left(\frac{E_z M}{5 \times 10^{14} M_{\odot}} \right)^{0.071(\pm 0.012)}. \quad (13)$$

The quoted uncertainties are statistical only and are the products of the propagation of the relative error available to the estimates of the hydrostatic masses, gas masses, temperature and luminosity and the size of the cluster sample analyzed. When the uncertainties related to the cross-calibration between *Chandra* and *XMM-Newton* on the gas temperature and hydrostatic mass are taken into account as discussed, e.g., in Maughan (2014) and Mahdavi et al. (2013), systematic errors of the order of (i) ± 0.002 and ± 0.04 and (ii) ± 0.02 and ± 0.004 affect the normalization and the slope of the mass dependence of $C^{0.5} f_g$ and β_p , respectively.

Inserting these values into equation (12), the gravitating mass can be recovered with, for instance, a lower intrinsic scatter associated with it than the one measured by using the standard relations with normalization and slope free to vary.

For the set of the four relations here investigated, these results provide a significant simplification in terms of number of free parameters to be constrained: routinely, a slope and a normalization have to be estimated (for a total of eight free parameters), whereas in our new framework, one needs only to limit the normalization of C , f_g and β_p and the mass dependence of the latter two, for a total of five free parameters. This evidence can also be formalized by the estimates of the Akaike information criterion (AIC; Akaike 1974), or equivalent information criteria (see e.g. Liddle 2007). All our models that adopt the self-similar scaling laws with the mass-dependent physical quantities perform significantly better (from a statistical point of view) than the power-law fits where normalization and slope are left free to vary (apart from the M - T relation of the sample ‘local massive’, where AICs are comparable) with an evidence ratio $e^{0.5\Delta}$, with Δ being the difference between the AIC estimated for ‘free parameters’ model and the one for the modified scaling relations, larger than 600 (compare, e.g., fits labelled (3) and (6) with the ones labelled (2) and (5) in Table 2). When the four scaling relations are considered together, we obtain a cumulative χ^2 of 1743 and 379 with 228 data points for the ‘local all’ sample (222, 173 and 94, respectively, for the ‘local massive’ one) for the set of the scaling laws with eight (all the normalizations and slopes) and five free parameters, respectively, implying ‘decisive’ evidence (according to the Jeffreys’ scale in Kass & Raftery 1995) in favour of our alternative scenario.

We also note that the formalism described in Section 3.1 (e.g. equation 10) is ready to accommodate the redshift evolution of the scaling relations through the assumed expressions in equation (9).

As we present in Fig. 2, preliminary plots that do not consider any selection function show encouraging agreements between the observed distributions and the expected ones. More dedicated work to characterize properly the studied samples both as a function of mass and redshift (for instance, to measure the relative weight of low-mass and high-redshift systems in the fit of the scaling relations) is however needed.

The result of this study opens a very promising prospective to have a full set of intercorrelated and internally consistent scaling relations that rely on the ones predicted from the self-similar scenario with an extension depending on well-identified astrophysical properties that can be investigated independently (like, e.g., the mass dependence of the thermal pressure profile or of the gas mass fraction).

ACKNOWLEDGEMENTS

We thank the anonymous referee for helpful comments that improved the presentation of the work. We also thank Mauro Sereno, Marco De Petris and Dominique Eckert for discussion and comments on the manuscript. We acknowledge the financial contribution from contracts ASI-INAF I/009/10/0 and PRIN-INAF 2012.

REFERENCES

Akaike H., 1974, *IEEE Trans. Autom. Control*, 19, 716
 Allen S. W., Evrard A. E., Mantz A. B., 2011, *ARA&A*, 49, 409
 Anders E., Grevesse N., 1989, *Geochim. Cosmochim. Acta*, 53, 197
 Arnaud K. A., 1996, in Jacoby G., Barnes J., eds, *ASP Conf. Ser. Vol. 101, Astronomical Data Analysis Software and Systems V*. Astron. Soc. Pac., San Francisco, p. 17
 Arnaud M., Evrard A. E., 1999, *MNRAS*, 305, 631
 Arnaud M., Pointecouteau E., Pratt G. W., 2007, *A&A*, 474, L37
 Arnaud M., Pratt G. W., Piffaretti R., Böhringer H., Croston J. H., Pointecouteau E., 2010, *A&A*, 517, 92
 Asplund M., Grevesse N., Sauval A. J., Scott P., 2009, *ARA&A*, 47, 481
 Baldi A., Ettori S., Molendi S., Gastaldello F., 2012, *A&A*, 545, 41
 Böhringer H., Werner N., 2010, *A&AR*, 18, 127
 Böhringer H., Dolag K., Chon G., 2012, *A&A*, 539, A120
 Eckert D., Molendi S., Vazza F., Ettori S., Paltani S., 2013a, *A&A*, 551, 22
 Eckert D., Ettori S., Molendi S., Vazza F., Paltani S., 2013b, *A&A*, 551, 23
 Ettori S., 2013, *MNRAS*, 435, 1265 (E13)
 Ettori S. et al., 2004, *MNRAS*, 354, 111
 Ettori S., Morandi A., Tozzi P., Balestra I., Borgani S., Rosati P., Lovisari L., Terenziani F., 2009, *A&A*, 501, 61
 Ettori S., Rasia E., Fabjan D., Borgani S., Dolag K., 2012, *MNRAS*, 420, 2058
 Ettori S., Donnarumma A., Pointecouteau E., Reiprich T. H., Giodini S., Lovisari L., Schmidt R. W., 2013, *Space Sci. Rev.*, 177, 119
 Evrard A. E., Arnault P., Huterer D., Farahi A., 2014, *MNRAS*, 441, 3562
 Giodini S., Lovisari L., Pointecouteau E., Ettori S., Reiprich T. H., Hoekstra H., 2013, *Space Sci. Rev.*, 177, 247
 Kass R. E., Raftery A. E., 1995, *J. Am. Stat. Assoc.*, 430, 791
 Kravtsov A. V., Borgani S., 2012, *ARA&A*, 50, 353
 Kravtsov A. V., Vikhlinin A., Nagai D., 2006, *ApJ*, 650, 128
 Liddle A. R., 2007, *MNRAS*, 377, L74
 Mahdavi A., Hoekstra H., Babul A., Bildfell C., Jeltema T., Henry J. P., 2013, *ApJ*, 767, 116
 Mantz A. B., Allen S. W., Morris R. G., Rapetti D. A., Applegate D. E., Kelly P. L., von der Linden A., Schmidt R. W., 2014, *MNRAS*, 440, 2077
 Markwardt C. B., 2008, in Bohlender D., Dowler P., Durand D., eds, *ASP Conf. Ser. Vol. 411, Astronomical Data Analysis Software and Systems XVIII*. Astron. Soc. Pac., San Francisco, p. 251
 Maughan B. J., 2014, *MNRAS*, 437, 1171

Mohr J. J., Mathiesen B., Evrard A. E., 1999, *ApJ*, 517, 627
 Nagai D., Lau E. T., 2011, *ApJ*, 731, L10
 Nagai D., Kravtsov A. V., Vikhlinin A., 2007, *ApJ*, 668, 1
 Okabe N., Zhang Y.-Y., Finoguenov A., Takada M., Smith G. P., Umetsu K., Futamase T., 2010, *ApJ*, 721, 875
 Planck Collaboration XXIX, 2014, *A&A*, 571, A29
 Planck Intermediate Results V, 2013, *A&A*, 550, 131
 Planelles S., Borgani S., Dolag K., Ettori S., Fabjan D., Murante G., Tornatore L., 2013, *MNRAS*, 431, 1487
 Pratt G. W., Croston J. H., Arnaud M., Böhringer H., 2009, *A&A*, 498, 361
 Reichert A., Böhringer H., Fassbender R., Mühlegger M., 2011, *A&A*, 535, A4
 Roncarelli M., Ettori S., Borgani S., Dolag K., Fabjan D., Moscardini L., 2013, *MNRAS*, 432, 3030
 Rozo E., Bartlett J. G., Evrard A. E., Rykoff E. S., 2014, *MNRAS*, 438, 78
 Stanek R., Evrard A. E., Böhringer H., Schuecker P., Nord B., 2006, *ApJ*, 648, 956
 Stanek R., Rasia E., Evrard A. E., Pearce F., Gazzola L., 2010, *ApJ*, 715, 1508
 Sun M., Voit G. M., Donahue M., Jones C., Forman W., 2009, *ApJ*, 693, 1142
 Sun M., Sehgal N., Voit G. M., Donahue M., Jones C., Forman W., Vikhlinin A., Sarazin C., 2011, *ApJ*, 727, L49
 Sunyaev R. A., Zeldovich I. B., 1980, *ARA&A*, 18, 537
 Sutherland R. S., Dopita M. A., 1993, *ApJS*, 88, 235
 Tozzi P., Norman C., 2001, *ApJ*, 546, 63
 Vikhlinin A., Kravtsov A., Forman W., Jones C., Markevitch M., Murray S. S., Van Speybroeck L., 2006, *ApJ*, 640, 691

APPENDIX A: NUMERICAL ESTIMATES OF THE NORMALIZATION

For sake of completeness, we provide here the details on how the numbers of equation (2) are obtained. Let us define

$$k_0 = \frac{4}{3} \pi \Delta \rho_{c,0} = \frac{\Delta H_0}{2G} = 1.928 \times 10^{-26} \frac{\Delta}{500} \text{ g s cm}^{-3}$$

$$k_1 = \mu m_{\text{amu}} G k_0^{1/3} = 1.820 \times 10^{-40} \text{ g}^{1/3} \text{ s}^{-5/3} \text{ cm}^2. \quad (\text{A1})$$

Then, the normalization for the $M-M_g$, $M-T$, $M-L$ and $M-Y_{\text{SZ}}$ relations can be estimated as

$$n_{MM_g} = (C_0^{0.5} f_{g,0})^{-1} \frac{M_{g,0}}{M_0}$$

$$n_{MT} = \left(\frac{f_T}{k_1} \right)^{3/2} \beta_{P,0}^{3/2} \frac{T_0^{3/2}}{M_0}$$

$$n_{ML} = \left(\frac{4/3 \pi \mu_e^2 m_{\text{amu}}^2 f_T^{0.5}}{f_L c_{t,0} k_0 k_1^{0.5}} \right)^{3/4} \beta_{P,0}^{3/8} (C_0^{0.5} f_{g,0})^{-3/2} \frac{L_0^{3/4}}{M_0}$$

$$n_{MY} = \left(\frac{m_e c^2 \mu_e m_{\text{amu}} f_T}{k_1 \sigma_T} \right)^{3/5} \beta_{P,0}^{3/5} f_{g,0}^{-3/5} \frac{Y_0^{3/5}}{M_0}, \quad (\text{A2})$$

where M_0 , $M_{g,0}$, T_0 , L_0 and Y_0 are the pivot values in cgs unit and are equal to $5 \times 10^{14} M_\odot$, $5 \times 10^{13} M_\odot$, 5 keV , $5 \times 10^{44} \text{ erg s}^{-1}$ and 10^{-4} Mpc^2 , respectively, in the present work.

APPENDIX B: ENERGY BAND DEPENDENCE OF THE $M-L$ RELATION

The gas luminosity considered in our analysis is the X-ray bolometric one, i.e. it has been evaluated in the energy band 0.01–100 keV. We indicate here how the $M-L$ relation is modified once the luminosity is estimated in different energy bands. In these cases, the




Figure C1. These plots show the samples and the best-fitting results, with the associated residuals χ , described from lines (4)–(7) in Table 2. Note that the best-fitting lines labelled (6) and (7) have been corrected by the factor $(1 - \theta m_1 + \phi m_2 + \phi m_3/2)$ in equation (12) for the sake of representation. The sum of the squared residuals χ provides the quoted total χ^2 .

cooling function c_f will not show a dependence upon the temperature to the power of $1/2$. By approximating the cooling function as a power law of the temperature, we can write $c_f = c_{f,0} \times T^\tau$ and

$$\frac{F_z M}{5 \times 10^{14} M_\odot} = n_{MLE} \left(\frac{\beta_P}{3} \right)^{\tau/(1+2\tau/3)} \left(\frac{C^{0.5} f_g}{0.1} \right)^{-2/(1+2\tau/3)} \left(\frac{F_z^{-1} L}{5 \times 10^{44} \text{ erg s}^{-1}} \right)^{1/(1+2\tau/3)},$$

$$n_{MLE} = \left(\frac{4/3\pi \mu_e^2 m_{\text{amu}}^2 f_T^\tau}{f_L c_{f,0} k_0 k_1^\tau} \right)^{1/(1+2\tau/3)}. \quad (\text{B1})$$

We quote here $c_{f,0}$, τ and the modified M – L relation for the most commonly used energy bands.

(i) (0.1–2.4) keV: formally, the best-fitting values with a power law of the cooling function in the range 2–12 keV are $c_f = 1.12 \times 10^{-23} c_{pe} T_{\text{keV}}^{-0.11} \text{ erg s}^{-1} \text{ cm}^3$. Adopting an exponent $\tau = 0$, $c_{f,0} = 0.91 \times 10^{-23} c_{pe}$ and the M – L relation can be written as

$$\frac{F_z M}{5 \times 10^{14} M_\odot} = 2.110 \left(\frac{C^{0.5} f_g}{0.1} \right)^{-2} \left(\frac{F_z^{-1} L}{5 \times 10^{44} \text{ erg s}^{-1}} \right). \quad (\text{B2})$$

(ii) (0.5–2 keV): as above, $c_f = 0.68 \times 10^{-23} c_{pe} T_{\text{keV}}^{-0.10} \text{ erg s}^{-1} \text{ cm}^3$. With a null dependence upon

the temperature, $c_f = c_{f,0} = 0.56 \times 10^{-23} c_{\text{pe}}$ and the M - L relation can be written as

$$\frac{F_z M}{5 \times 10^{14} M_\odot} = 3.400 \left(\frac{C^{0.5} f_g}{0.1} \right)^{-2} \left(\frac{F_z^{-1} L}{5 \times 10^{44} \text{ erg s}^{-1}} \right). \quad (\text{B3})$$

(iii) (2–10 keV): in this case, $c_f = 0.38 \times 10^{-23} c_{\text{pe}} T_{\text{keV}}^{0.5} \text{ erg s}^{-1} \text{ cm}^3$. Then,

$$\frac{F_z M}{5 \times 10^{14} M_\odot} = 1.749 \left(\frac{\beta_p}{3} \right)^{3/8} \left(\frac{C^{0.5} f_g}{0.1} \right)^{-3/2} \left(\frac{F_z^{-1} L}{5 \times 10^{44} \text{ erg s}^{-1}} \right)^{3/4}. \quad (\text{B4})$$

APPENDIX C: PLOTS OF THE INVESTIGATED SCALING RELATIONS

We show in Fig. C1 the plots, with the best-fitting lines and the corresponding residuals χ_i of equation (6), of the samples described in Table 2. The normalizations of these best-fitting scaling relations are shown in Fig. 2.

This paper has been typeset from a \LaTeX file prepared by the author.