



Publication Year	2015
Acceptance in OA @INAF	2020-03-25T10:14:42Z
Title	Spectroscopic evidence of multiple populations in globular clusters
Authors	GRATTON, Raffaele; LUCATELLO, Sara; CARRETTA, Eugenio; BRAGAGLIA, Angela
DOI	10.1017/S1743921314005547
Handle	http://hdl.handle.net/20.500.12386/23535
Series	PROCEEDINGS OF THE INTERNATIONAL ASTRONOMICAL UNION
Number	vol. 10, H16 (Highlights of Astronomy)

Spectroscopic evidence of multiple populations in globular clusters

R. Gratton¹, S. Lucatello¹, E. Carretta² and A. Bragaglia²

¹INAF-Osservatorio Astronomico di Padova

²INAF-Osservatorio Astronomico di Bologna

Abstract. We review spectroscopic evidence of multiple stellar populations in globular clusters. First, we lay down the basic data: the C-N, Na-O, Mg-Al anti-correlations among red giants and main sequence stars, and discuss how they appear to be general properties of globular clusters, in spite of cluster-to-cluster differences. We will then describe what is currently known about He from spectroscopy. We will then present the implications and current observations for the interpretation of the horizontal branches, showing that the multiple population phenomenon is strongly related to the distribution of stars along them. We will briefly mention the spectroscopic evidence related to some less understood cases, like the clusters with multiple subgiant branches. Finally, we summarize the relation between multiple populations and general properties for globular clusters, and their implications for the formation scenario.

Keywords. (Galaxy:) globular clusters: general, stars: abundances, Galaxy: halo

The last ten years have seen a revolution in our concept of globular clusters (GCs) thanks to data coming from new efficient multi-object spectrographs on 8-m telescopes (e.g. Carretta *et al.* 2009a, b) and the exquisite photometric precision of HST (e.g. Bedin *et al.* 2004). These studies clarified the meaning of large star-to-star abundance variations for several light elements known to be present in GCs for almost forty years (see Gratton *et al.* 2004 and 2012 for recent reviews) and revealed multiple sequences on the red giant branch (RGB), subgiant branch (SGB), and main sequence (MS) in several GCs. Regarded for a long time as intriguing abundance anomalies restricted to some cluster stars, the peculiar chemical patterns only recently were explicitly understood as a universal phenomenon in GCs related to their very same nature/origin (Carretta 2006; Carretta *et al.* 2006, 2009a, b, 2010). The observational pattern is quite well assessed (see the reviews by Gratton *et al.* 2004, 2012, and Piotto 2009), thanks to several important milestones:

(i) Abundance variations of the heavier species (O-Na and Mg-Al anti-correlations) are restricted to the denser cluster environment. Evolutionary changes along the RGB additionally affect other light elements (Li, C, and N: Charbonnel *et al.* 1998; Gratton *et al.* 2000).

(ii) This pattern is established in p -capture reactions of the CNO, NeNa and MgAl chains in high temperature H-burning (Denisenkov & Denisenkova 1989).

(iii) The variations are found also among unevolved stars currently on the MS of GCs (Briley *et al.* 1996; Gratton *et al.* 2001). This calls for a class of now-extinct stars, more massive than the low-mass ones presently evolving in GCs, as the site for the nucleosynthesis.

(iv) The observed variations in the abundances of p -capture elements are connected to that of He, i.e., stars that are Na-rich and O-poor are also He-rich, although the relation between these abundance variations may be complicated. Multiple MSs, attributed to populations with different He fraction Y , have been found in some GCs (Bedin *et al.*

2004, Piotto *et al.* 2007). There are indications that Y differences can be traced also on the RGB (Carretta *et al.* 2007, Bragaglia *et al.* 2010, Pasquini *et al.* 2011), and contribute to explaining the horizontal branch (HB) second parameter problem (Gratton *et al.* 2010).

In summary, in contrast with the common assumption, GCs are not simple stellar populations: they harbour various stellar generations, distinct by their chemistry. These populations may be separated using the patterns of anti-correlated Na-O, Mg-Al or photometric methods. They provide insights into the early phases of GCs, since the time scale for the release of matter processed by H-burning at high temperatures is less than 10^8 year. To decipher the relevant information, large and homogeneous datasets are needed, like the one recently gathered by Carretta *et al.* (2009a, 2009b), from which Carretta *et al.* (2010) were able to place the properties of different stellar generations in GCs and global cluster parameters into a general framework for the formation of GCs. The Na-O anti-correlation may be used to separate GCs (hosting multiple stellar populations) from open clusters (OCs, where a single stellar population is present), and first and second generation (FG and SG) stars within each GC. The emerging scenario for the formation of GCs includes various phases: a large episode of star formation (the FG) and then the formation of the current GC (mainly made of SG stars) within a cooling flow formed by the slow winds of a fraction of this FG (D'Ercole *et al.* 2008). Some stars of the FG remain in the cluster. Since the ejecta of only a small fraction of the FG stars are used to form the SG, a large fraction of the FG must have been lost by the proto-GCs and may represent the main component of halo field stars (see also Martell & Grebel 2010; Vesperini *et al.* 2010; Schaerer & Charbonnel 2011). This links directly the formation of GC to that of galaxies.

References

- Bedin, L. R., *et al.*, 2004, *ApJ*, 605, L125
Bragaglia, A., *et al.*, 2010, *A&A*, 519, 60
Briley, M., *et al.*, 1996, *Nature*, 383, 604
Charbonnel, C., *et al.*, 1998, *A&A*, 332, 204
Carretta E., 2006, *AJ*, 131, 1766
Carretta, E., *et al.*, 2006, *A&A*, 450, 523
Carretta, E., *et al.*, 2007, *A&A*, 464, 927
Carretta, E., *et al.*, 2009a, *A&A*, 505, 117
Carretta, E., *et al.*, 2009b, *A&A*, 505, 139
Carretta, E., *et al.* 2010, *A&A*, 516, 55
Denisenkov, P. A. & Denisenkova, S. N. 1989, *A.Tsir.*, 1538, 11
D'Ercole, A., *et al.* 2008, *MNRAS*, 391, 825
Gratton, R. G., *et al.*, 2000, *A&A*, 358, 671
Gratton, R. G., *et al.*, 2001, *A&A*, 369, 87
Gratton, R. G., *et al.*, 2004, *ARA&A*, 42, 38
Gratton, R. G., *et al.*, 2010, *A&A*, 517, 8
Gratton, R. G., *et al.*, 2012, *A&ARv*, 20, 50
Martell, S. L. & Grebel, E. K. 2010, *A&A*, 519, 14
Pasquini, L., *et al.*, 2011, *A&A*, 531, 35
Piotto, G. 2009, in *IAUS* 258, 233
Piotto, G., *et al.*, 2007, *ApJ*, 661, L53
Schaerer, D. & Charbonnel, C., 2011, *MNRAS* 413, 2297
Vesperini, E., *et al.*, 2010, *ApJ*, 718, L112