

Publication Year	2015
Acceptance in OA@INAF	2020-03-31T14:04:19Z
Title	Molecular depletion times and the CO-to-H <sub>2</sub> conversion factor in metal-poor galaxies
Authors	HUNT, Leslie Kipp; García-Burillo, S.; CASASOLA, VIVIANA; Caselli, P.; Combes, F.; et al.
DOI	10.1051/0004-6361/201526553
Handle	http://hdl.handle.net/20.500.12386/23766
Journal	ASTRONOMY & ASTROPHYSICS
Number	583

# Molecular depletion times and the CO-to-H<sub>2</sub> conversion factor in metal-poor galaxies\*

L. K. Hunt<sup>1</sup>, S. García-Burillo<sup>2</sup>, V. Casasola<sup>1</sup>, P. Caselli<sup>3</sup>, F. Combes<sup>4</sup>, C. Henkel<sup>5,6</sup>, A. Lundgren<sup>7</sup>, R. Maiolino<sup>8</sup>, K. M. Menten<sup>5</sup>, L. Testi<sup>9</sup>, and A. Weiss<sup>5</sup>

- <sup>1</sup> INAF–Osservatorio Astrofisico di Arcetri, Largo E. Fermi, 5, 50125 Firenze, Italy e-mail: hunt@arcetri.astro.it
- <sup>2</sup> Observatorio Astronómico Nacional (OAN) Observatorio de Madrid, Alfonso XII, 3, 28014 Madrid, Spain
- <sup>3</sup> Max-Planck-Institut für extraterrestrische Physik, Giessenbachstrasse 1, 85748 Garching, Germany
- <sup>4</sup> Observatoire de Paris, LERMA, Collège de France, CNRS, PSL, Sorbonne University UPMC, 75014 Paris, France
- <sup>5</sup> Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany
- <sup>6</sup> Astronomy Department, King Abdulaziz University, PO Box 80203 Jeddah, Saudia Arabia
- <sup>7</sup> ALMA JAO, Alonso de Cordova 3107, Vitacura, Casilla 19001, Santiago, Chile
- <sup>8</sup> Cavendish Laboratory, University of Cambridge, 19 J.J. Thomson Avenue, Cambridge CB3 0HE, UK
- <sup>9</sup> ESO, Karl Schwarzschild str. 2, 85748 Garching bei München, Germany

Received 18 May 2015 / Accepted 14 September 2015

# ABSTRACT

Tracing molecular hydrogen content with carbon monoxide in low-metallicity galaxies has been exceedingly difficult. Here we present a new effort, with IRAM 30-m observations of <sup>12</sup>CO(1–0) of a sample of 8 dwarf galaxies having oxygen abundances ranging from  $12 + \log(O/H) \sim 7.7$  to 8.4. CO emission is detected in all galaxies, including the most metal-poor galaxy of our sample  $(0.1 Z_{\odot})$ ; to our knowledge this is the largest number of <sup>12</sup>CO(1–0) detections ever reported for galaxies with  $12 + \log(O/H) \leq 8 (0.2 Z_{\odot})$  outside the Local Group. We calculate stellar masses,  $M_{star}$ , and star-formation rates (SFRs), and analyze our results in conjunction with galaxy samples from the literature. Extending previous results for a correlation of the molecular gas depletion time,  $\tau_{dep}$ , with  $M_{star}$ and specific SFR (sSFR), we find a variation in  $\tau_{dep}$  of a factor of 200 or more (from  $\leq 50$  Myr to  $\sim 10$  Gyr) over a spread of  $\sim 10^3$ in sSFR and  $M_{star}$ . We exploit the variation of  $\tau_{dep}$  to constrain the CO-to-H<sub>2</sub> mass conversion factor  $\alpha_{CO}$  at low metallicity, and assuming a power-law variation find  $\alpha_{CO} \propto Z/Z_{\odot}^{-2}$ , similar to results based on dust continuum measurements compared with gas mass. By including H I measurements, we show that the fraction of total gas mass relative to the baryonic mass is higher in galaxies that are metal poor, of low mass, and with high sSFR. Finally, comparisons of the data with star-formation models of the molecular gas phases show that the models are generally quite successful, but at metallicities  $Z/Z_{\odot} \leq 0.2$ , there are some discrepancies.

Key words. galaxies: dwarf – galaxies: starburst – galaxies: star formation – galaxies: ISM – ISM: general

#### 1. Introduction

Dense clumps in cool molecular clouds are thought to be the "cradles" of star formation, but finding molecules at low metallicities has been exceedingly difficult. In dwarf galaxies with nebular oxygen abundances  $\leq 0.3 Z_{\odot} (12 + \log(O/H) \leq 8.2)^1$ , despite vigorous on-going star formation, there seems to be very little CO (Sage et al. 1992; Taylor et al. 1998; Gondhalekar et al. 1998; Barone et al. 2000; Leroy et al. 2005, 2007; Buyle et al. 2006; Schruba et al. 2012; Cormier et al. 2014). Intriguingly, this is the same metallicity "threshold" below which little or no polycyclic aromatic hydrocarbon (PAH) emission is detected in metal-poor galaxies (Engelbracht et al. 2005; Wu et al. 2006; Madden et al. 2006; Hunt et al. 2010).

The reason for this deficit in CO at low metallicities is not yet completely understood, although there is strong evidence for "CO-dark" molecular gas in the interstellar medium (ISM; e.g., Wolfire et al. 2010; Leroy et al. 2011; Planck Collaboration XIX 2011; Paradis et al. 2012; Lebouteiller et al. 2012; Pineda et al. 2014). Such a component would result from a diffuse or metalpoor ISM in which CO suffers from enhanced photodissociation relative to H<sub>2</sub> (e.g., Maloney & Black 1988; Bolatto et al. 1999; Glover & Mac Low 2011; Narayanan et al. 2012). H<sub>2</sub> can effectively avoid the photodissociation effects of UV radiation through self-shielding, even at relatively low gas surface densities,  $\Sigma_{ISM}$ . On the other hand, CO needs higher  $\Sigma_{ISM}$  to selfshield, thus implying that the CO-to-H<sub>2</sub> ratio should vary even within an individual molecular cloud (e.g., Röllig et al. 2006; Leroy et al. 2013a; Hughes et al. 2013). Under conditions of low  $\Sigma_{\rm ISM}$  and low dust extinction  $A_V$ , most of the carbon would be in other forms such as CI or CII, rather than in CO.

Nevertheless, even under extreme conditions in a metalpoor ISM, it is likely that stars form in cool, dense molecular clouds (e.g., Krumholz et al. 2011). However, H<sub>2</sub>, the dominant

<sup>\*</sup> Based on observations carried out with the IRAM 30 m; IRAM is supported by the INSU/CNRS (France), MPG (Germany), and IGN (Spain).

<sup>&</sup>lt;sup>1</sup> Throughout this work, we will rely on the Asplund et al. (2009) solar abundance calibration of  $12 + \log(O/H) = 8.69$ .

#### Table 1. Observing targets.

Name	Other name	Redshift <sup>a</sup>	Systemic velocity <sup>a</sup> (km s <sup>-1</sup> )	Distance <sup><i>a</i></sup> (Mpc)	Distance <sup>a</sup> method	$12 + \log(O/H)^b$	Dimensions <sup>a</sup> (arcsec <sup>2</sup> )	$\frac{SFR^c}{(M_{\odot}{\rm yr}^{-1})}$	$Log^c$ $M_{star}$ $(M_{\odot})$	$Log^d$ $M_{\rm HI}$ $(M_{\odot})$	$Log(Total)^e$ $M_{\rm HI}$ $(M_{\odot})$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
CGCG 007-025		0.0048	1449	24.5	CMB	7.74	$27.0 \times 16.2$	0.23	8.36	8.90	8.90
II Zw 40	UGCA116	0.0026	789	11.7	CMB	8.12	$33.6 \times 13.2$	1.75	8.25	7.54	8.77
Mrk 996		0.0054	1622	18.1	CMB	8.00	$36 \times 30$	0.16	9.18	8.00	8.00
NGC 1140	Mrk 1063	0.0050	1501	19.7	TF	8.18	$102 \times 54$	0.82	9.76	8.56	9.51
NGC 1156		0.0013	375	8.1	Stars	8.23	$198 \times 150$	0.23	9.55	8.66	9.04
NGC 3353	Haro 3,	0.0031	944	18.1	TF	8.37	$72.0 \times 49.8$	1.28	9.54	8.87	8.87
	Mrk 35										
NGC 7077	Mrk 900	0.0038	1152	17.2	TF	8.03	$48 \times 42$	0.12	8.98	8.15	8.15
SBS 0335-052E		0.0135	4043	53.6	CMB	7.25	$13.8 \times 12.0$	1.30	8.51	7.81	8.61
UM 448	Arp 161,	0.0186	5564	81.2	CMB	8.04	$24 \times 24$	11.11	10.91	9.84	9.84
	UGC 6665										
UM 462	Mrk 1307 UGC 6850	0.0035	1057	19.5	CMB	7.97	$36 \times 30$	0.29	8.56	8.30	8.72

**Notes.** <sup>(a)</sup> NASA/IPAC Extragalactic Database (NED, http://nedwww.ipac.caltech.edu/): CMB corresponds to Cosmic Microwave Background and TF to Tully-Fisher; velocities are heliocentric; dimensions are optical major and minor diameters. <sup>(b)</sup> Direct-temperature metallicities for CGCG 007–025, NGC 3353, UM 448, UM 462, SBS 0335–052E from Izotov et al. (2007); II Zw 40 (Thuan & Izotov 2005); Mrk 996 (Thuan et al. 1996); NGC 1140 (Izotov & Thuan 2004); NGC 1156 (Vigroux et al. 1987); NGC 7077 (van Zee & Haynes 2006). <sup>(c)</sup> These quantities are newly derived, as described in Sect. 2.3. <sup>(d)</sup> When these values differ from those in Col. (12), they are calculated from the mean H<sub>I</sub> surface density integrated over the optical size of the galaxy. <sup>(e)</sup> The H<sub>I</sub> measurements are taken from van Zee (2001, CGCG 007–025, NGC 7077), van Zee et al. (1998, II Zw 40, UM 462), Thuan et al. (1999, Mrk 996), Hunter et al. (1994a, NGC 1140), Swaters et al. (2002, NGC 1156), Hunter et al. (1982, NGC 3353), Ekta et al. (2009, SBS 0335–052E), Paturel et al. (2003, UM 448).

molecular component of giant molecular clouds, is not directly observable, but rather inferred from line emission of rotational transitions of <sup>12</sup>CO(1–0), unfortunately deficient at low metallicity. Constraining the properties of metal-poor molecular clouds observationally is difficult because the factor,  $\alpha_{CO}$ , relating CO luminosity  $L'_{CO}$  to H<sub>2</sub> mass  $M_{H2}$  can vary dramatically, through the presence of CO-dark gas.  $\alpha_{CO}$  can be more than 50 times higher than the Galactic value in metal-poor environments and in regions of low  $A_V$  and low  $\Sigma_{ISM}$  (Israel 1997; Leroy et al. 2009b, 2011; Pineda et al. 2010; Bolatto et al. 2011; Schruba et al. 2012; Sandstrom et al. 2013). Although the relation between CO and H<sub>2</sub> has been the subject of much debate (see Bolatto et al. 2013, for a recent review), it is clear that at some low metallicity, probably  $\leq 0.05-0.1 Z_{\odot}$ , CO ceases to become a useful tracer of total H<sub>2</sub> mass.

In this paper, we begin to explore the limits to which CO can trace H<sub>2</sub> in metal-poor star-forming galaxies. We first describe the targets and ancillary data in Sect. 2, and the <sup>12</sup>CO(1–0) observations and data analysis in Sect. 3. In Sect. 4, we compare our results with previous work, and assess the dependence of CO luminosity on star-formation rate (SFR). Section 5 describes constraints on  $\alpha_{CO}$  and  $\tau_{dep}$ , the molecular gas depletion time, i.e., the time required to consume the H<sub>2</sub> gas reservoir. We adopt a variation of the method proposed by Saintonge et al. (2011b), extending it to low metallicity in order to separate the effects of  $\tau_{dep}$  variations from variations of  $\alpha_{CO}$ . In Sect. 6, we analyze the H<sub>2</sub> content of our sample and compare it with H<sub>1</sub> for a compilation of galaxies taken from the literature. We discuss the implications of our results in Sect. 7, and, in Sect. 8, summarize our results and present our conclusions.

#### 2. The targets and the ancillary data

The observing sample was defined mainly on the basis of detected  $H_2$  emission, either in the ro-vibrational transitions

at  $2\mu$ m, or the purely rotational transitions in the mid-infrared (e.g., *Spitzer*/IRS); however, we also targeted some galaxies that had already been observed in CO (see Sect. 3.4). Table 1 lists the sample galaxies (ordered alphabetically by name), together with relevant observational parameters; Figs. 1 and 2 show thumbprint images.

#### 2.1. Individual source description

The targets have several characteristics in common, in addition to their low metallicities. All of them, with the exception of NGC 7077 and SBS 0335–052E, are sufficiently bright in the infrared (IR) to have been identified in the (faint) IRAS pointsource catalogue. In fact, II Zw 40 was identified as a 60- $\mu$ m peaker source by Vader et al. (1993). Except for NGC 7077, all galaxies host conspicuous Wolf-Rayet populations (Schaerer et al. 1999; Guseva et al. 2000; Zhang et al. 2007); they are all also rich in atomic gas (see Sect. 6). This section describes particular features of the individual sources (in alphabetical order by name as in Table 1).

#### \* CGCG 007-025

This galaxy is the most metal-poor  $(12 + \log(O/H) = 7.7, ~0.1 Z_{\odot})$  <sup>12</sup>CO(1–0) detection of our sample (see Sect. 4). Although it appears in an isolated galaxy catalogue (van Zee 2001), it has a companion, a dwarf irregular galaxy at 8.3 kpc projected separation (84 km s<sup>-1</sup> velocity separation) which may have triggered the current star-formation episode (van Zee 2001). It is also found to be a member of a compact group by Focardi & Kelm (2002). CGCG 007–025 has the bluest U - B color of any dwarf galaxy in the van Zee (2001) sample, with a very clumpy morphology as seen in Fig. 1. The radiation field is sufficiently hard to excite [FeV], a line that requires ionizing radiation with photon energies exceeding 4 Ryd (54.4 eV) (Thuan & Izotov 2005). Although a population of Wolf-Rayet stars has

L. K. Hunt et al.: Molecular depletion times at low metallicity



**Fig. 1.** *Top row, from left to right* in alphabetical order of name: SDSS composite (*ugriz*) images of CGCG 007–025, Haro 3 (NGC 3353), II Zw 40, Mrk 996. *Bottom* row, *leftmost panel*: NGC 1156 (image taken from DSS2, reprocessed); then *from left to right* SDSS images of NGC 7077, UM 448, UM 462. Images are oriented with North up, East left, and are 50" on a side, except for NGC 1156 which is ~10'. The color of II Zw 40 is reddened by >2 mag  $A_V$  from the Galaxy. NGC 1140 and SBS 0335–052E, both of which harbor SSCs and whose images are not from SDSS, are shown in Fig. 2.



**Fig. 2.** NGC 1140 and SBS 0335–052E taken from composite HST images; both galaxies host super-star clusters (Hunter et al. 1994b; Reines et al. 2008). The image of NGC 1140 is  $\sim$ 90" on a side, and SBS 0335–052E  $\sim$ 5", and orientation is North up, East left.

been detected in this galaxy (Zhang et al. 2007), these are insufficient to power the high-excitation emission lines, thought instead to be due to fast radiative shocks (Thuan & Izotov 2005). Hunt et al. (2010) found evidence for rotational H<sub>2</sub> emission, and despite the low metallicity of CGCG 007–025, also 7.7  $\mu$ m and 11.3  $\mu$ m emission from PAHs. Judging from the relatively high density of its ionized gas (~300 cm<sup>-3</sup>, Izotov et al. 2007), CGCG 007–025 is an example of an "active" starburst, forming stars in compact dense regions (Hirashita & Hunt 2004).

# \* II Zw 40 (UGCA 116)

Although we did not observe this source in  ${}^{12}CO(1-0)$ , it is included in our CO survey of higher-*J* lines (Hunt et al., in prep.); thus we consider it in our sample here. From its discovery, together with IZw 18, as an extragalactic HII region (Sargent & Searle 1970), IIZw 40 has been the subject of much scrutiny.

Its two-tailed optical morphology and kinematics suggest that it is the result of a collision between two gas-rich dwarf galaxies which triggered the current starburst (Baldwin et al. 1982; van Zee et al. 1998). Near-infrared recombination lines show that more than half the  $2 \mu m$  flux comes from free-free emission, and warm dust contributes at least ~25% to the 3–4  $\mu$ m fluxes (Joy & Lester 1988). Emission from the ro-vibrational transitions of H<sub>2</sub> at  $2\mu$ m was discovered by Vanzi et al. (1996), and later found to be in a cloud distinct from the primary HII region (Vanzi et al. 2008). Visual extinction  $A_V$  is spatially variable in II Zw 40 and can be as high as  $\gtrsim 10$  mag in the nucleus (Vanzi et al. 2008). This galaxy is undergoing a massive starburst in a region of  $\sim$ 150 pc in diameter powered by  $\sim$ 5000–10000 OV stars (Beck et al. 2002; Kepley et al. 2014). These massive stars are driving copious stellar winds which alter the observed kinematics (Bordalo et al. 2009), and also produce high-ionization [FeV] (Thuan & Izotov 2005). Like CGCG 007-025, the high density derived from optical spectra (~500-1400 cm<sup>-3</sup>, Bordalo et al. 2009) and the small dimensions of the star-formation episode imply that it is an "active" starburst.

# \* Mrk 996

Mrk 996 is an unusual blue compact dwarf galaxy, with smooth early-type morphology and a system of globular clusters (Thuan et al. 1996). Virtually all its star formation occurs in the central region of the galaxy ( $\sim$ 315 pc in radius). Because of its high central density ( $\sim$ 10<sup>6</sup> cm<sup>-3</sup>) and broad emission lines ( $\sim$ 900 km s<sup>-1</sup>, Thuan et al. 1996), it was thought that there could be an accreting intermediate-mass black hole; however X-ray observations revealed no compact point source (Georgakakis et al. 2011). A high ionization [OIV] line is detected in Mrk 996 (Thuan et al. 2008), that, like [FeV], needs  $\sim$ 4 Ryd to excite it. Thus, fast radiative shocks are also present in this galaxy, as its emission-line properties are inconsistent with any other source of ionization (Georgakakis et al. 2011). The lowest rotational transition of  $H_2$  (~28  $\mu$ m) is found in Mrk 996, the only one in the sample of Hunt et al. (2010) in which it was detected.

#### \* NGC 1140 (Mrk 1063)

NGC 1140 is a barred dwarf irregular galaxy with peculiar morphology and a central starburst. The H $\alpha$  luminosity of its central HII complex is 40 times higher than that of the giant HII region 30 Doradus in the Large Magellanic Cloud. Within this complex, roughly 500 pc in diameter, there are 6–7 blue super-star clusters (SSCs; Hunter et al. 1994b; de Grijs et al. 2004), the most luminous of these containing almost 6000 O stars (Moll et al. 2007). Although there is no galactic-scale outflow in this galaxy, powerful shocks from the massive SSCs are disrupting the ISM on scales up to 1–2 kpc radius (Westmoquette et al. 2010). Kinematic and other evidence suggests that the formation of the SSCs and the larger-scale starburst have been triggered by an interaction or merger event with a low-luminosity late-type spiral about 1 Gyr ago (Hunter et al. 1994a,b; Westmoquette et al. 2010).

#### \* NGC 1156

A typical barred Magellanic irregular about 25% brighter in the *B* band than the Large Magellanic Cloud, NGC 1156 is the most quiescent galaxy in our sample. It has been described as the least disturbed galaxy in the Local Universe, with no significant galaxy within 700 kpc (Karachentsev et al. 1996). More recent H<sub>I</sub> observations revealed a single dwarf galaxy at a distance of 80 kpc (Minchin et al. 2010) which however is too faint to provide any measurable perturbation. The star formation in NGC 1156 is spread out in HII regions spanning the entire bar+disk, and the kinematics of the ionized (and atomic) gas and stars is consistent with circular rotation about a common axis (Hunter et al. 2002).

# \* NGC 3353 (Haro 3, Mrk 35)

NGC 3353, the most metal-rich galaxy in our sample  $(12 + \log(O/H) \sim 8.4$ , see Table 1), is a dwarf irregular hosting a galaxy-wide starburst. However, high-resolution radio images show that most of the starburst activity is concentrated within a single knot to the northwest of the galaxy nucleus (Johnson et al. 2004). The star clusters in this knot are quite young (1–4 Myr) and massive (~10<sup>6</sup>  $M_{\odot}$ ) and reside in a region with  $A_V \sim 2-8$  mag, implying that much of the current star formation is currently enshrouded in dust (Johnson et al. 2004). In fact, optical measures of the extinction are smaller than those measured with the radio and near-infrared observations (Steel et al. 1996; Cairós et al. 2007). Detailed photoionization models of the infrared fine-structure lines suggest that there are at least two emission zones in this galaxy: one with low extinction, visible in the optical, and another, optically invisible with much higher extinction (Hunt et al. 2006). Rotational H<sub>2</sub> has been detected in NGC 3353, and the detection of the high-excitation [OIV] line  $(\gtrsim 4 \text{ Ryd})$  implies that radiative shocks are also present (Hunt et al. 2010). Like other galaxies in this sample, kinematic evidence suggests that a merger is responsible for triggering the starburst in NGC 3353 (Cairós et al. 2007).

# \* NGC 7077 (Mrk 900)

NGC 7077 is classified as an early-type galaxy, E/S0, but with peculiar morphology, and relatively blue colors (Gil de Paz et al. 2003; Micheva et al. 2013). It is an isolated galaxy, part of the van Zee (2000) sample, with red colors in its outer regions (see also Fig. 1). These red colors are in contrast with the strong (blue) nuclear starburst which has an intensity comparable to that

of 30 Doradus (van Zee 2000). Perhaps the most salient feature of NGC 7077 is its extended ultraviolet disk, suggesting that the disk of this galaxy is in an active growth phase (Moffett et al. 2012).

#### \* SBS 0335-052E

Like, II Zw 40, we did not observe this source in  ${}^{12}CO(1-0)$ , but it is included in our CO survey of higher-J lines (Hunt et al., in prep.); thus we consider it here. This is the most metal-poor galaxy of our sample  $(12 + \log(O/H) \sim 7.2)$ , with a neighbor at ~22 kpc distance (SBS 0335-052W, Pustilnik et al. 2001). This companion, embedded in a common HI cloud, is the most metal-poor star-forming galaxy known in the Local Universe  $(12 + \log(O/H) \sim 7.1$ , Lipovetsky et al. 1999; Izotov et al. 2005). High-resolution optical and near-infrared imaging shows that the star formation activity in SBS 0335-052E is confined to six massive SSCs, with the two youngest ones ( $\leq 3$  Myr) each comprising ~5000 O7.5 V stars (Reines et al. 2008). Despite its extremely low abundance, SBS 0335-052E is known to have extremely high extinction ( $A_V \gtrsim 15$  mag, Plante & Sauvage 2002; Houck et al. 2004), and  $\gtrsim 10^4 M_{\odot}$  of dust surrounding only the two youngest SSCs (Hunt et al. 2001, 2014). Warm H<sub>2</sub> is clearly present in this galaxy, with strong emission from ro-vibrational transitions at  $2 \mu m$  (Vanzi et al. 2000, 2011). The cm-regime radio spectrum is highly self-absorbed, implying ionized-gas densities of  $\gtrsim 3000 \text{ cm}^{-3}$  (Hunt et al. 2004; Johnson et al. 2009). SBS 0335-052E is the proto-typical "active" starburst with its star-formation activity occurring in compact (unresolved at HST resolution), dense regions (Hirashita & Hunt 2004).

#### \* UM 448 (Arp 161, UGC 6665)

UM 448 is a merger remnant with residual tidal tails, and so IR luminous ( $L_{IR} = 6.3 \times 10^{10} L_{\odot}$ ) that it would almost be classified as a Luminous IR Galaxy (LIRG, Dopita et al. 2002). Despite its low metal abundance ( $12 + \log(O/H) \sim 8.0, 0.2 Z_{\odot}$ , see Table 1), this galaxy shows clear PAH emission (Galliano et al. 2008). This is perhaps because of a metallicity gradient which reaches a peak value of  $12 + \log(O/H) \sim 8.4$  toward the northwest, a region of lower temperature (James et al. 2013). There is evidence for age gradients in the stellar populations and complex ionized emission-line kinematics, both of which probably reflect the merger nature of UM 448 (James et al. 2013).

#### ★ UM 462 (Mrk 1307, UGC 6850)

Like many of the galaxies in our sample, UM 462 is possibly tidally interacting with a relatively close neighbor, UM 461 (Taylor et al. 1995), although van Zee et al. (1998) concluded that the H<sub>I</sub> kinematics are inconsistent with this. Most of the star formation in UM 462 occurs in two main knots of star formation, although high-resolution observations resolve at least four starbursting regions with disrupted kinematics (Vanzi et al. 2002; James et al. 2010). The Wolf-Rayet content of this galaxy has been disputed by James et al. (2010) who do not find evidence to support a significant Wolf-Rayet classification. Vanzi et al. (2002) found ro-vibrational H<sub>2</sub> emission in this galaxy, but no evidence for [FeII], implying that the star-formation episodes must be younger than ~10 Myr. The knots of star formation are so compact and luminous as to be classified as SSCs (Vanzi 2003).

#### 2.2. An additional source, I Zw 18

Although IZw 18 was not included in our sample, it was observed, although not detected, in the  ${}^{12}CO(1-0)$  transition by Leroy et al. (2007). Because of its extreme low metallicity, similar to SBS 0335–052E, we have incorporated this galaxy in our analysis (see also Schruba et al. 2012). We have adopted the SFR from Hunt et al. (2005), and calculated the stellar mass,  $M_{\text{star}}$ , as described below (see also Fumagalli et al. 2010).

Like II Zw 40, I Zw 18 was discovered as an extragalactic HII region by Sargent & Searle (1970). Together with the SBS 0335-052E and SBS 0035-052W pair, it is the most metal-poor star-forming galaxy in the Local Universe with  $12 + \log(O/H) = 7.17 - 7.22$  (Thuan & Izotov 2005). Most, if not all, of the current star formation occurs in two main clusters, although the "C component" (or Zwicky's flare) is also forming stars (Izotov et al. 2001). However, the two clusters are not as extreme as those in SBS 0335-052E (and other galaxies in our sample), being somewhat less luminous than 30 Doradus (Hunter & Thronson 1995). Like SBS 0335-052E, around the main star clusters there is a complex structure of filaments and arcs from ionized gas, and the complex HI morphology and kinematics suggest that the current star-formation episode was triggered by an interaction with the C component (Lelli et al. 2012). Unlike SBS 0335-052E, however, there is no evidence for a dense gas component in IZw 18; the radio continuum spectrum follows the trend expected for optically thin free-free emission (Hunt et al. 2005) with an ionized-gas density of  $\leq 100 \text{ cm}^{-3}$ , similar to that found from optical spectra (Izotov et al. 2001). Despite their similarly low metal abundance, the dust masses, dust-to-gas, and dust-to-stars mass ratios in IZw 18 are more than 100 times lower than in SBS 0335-052E (Hunt et al. 2014), implying that the density of the cool gas in the ISM may play an important role in dust formation (Schneider et al. 2015).

#### 2.3. SFRs and stellar masses

We have calculated the SFRs for our targets according to Calzetti et al. (2010) using H $\alpha$  and 24  $\mu$ m luminosities (their Eq. (17)). This approach obviates the need for uncertain extinction corrections, and has the added advantage of accounting for the general behavior of dust opacity in galaxies. Many of the H $\alpha$  total fluxes come from Gil de Paz et al. (2003, Haro 3, II Zw 40, Mrk 996, NGC 7077), but others come from van Zee (2000, CGCG 007–025), Kennicutt et al. (2009, NGC 1140, NGC 1156), Dopita et al. (2002, UM 448), and James et al. (2010, UM 462). The 24  $\mu$ m fluxes are taken from Engelbracht et al. (2008), Temi et al. (2009, NGC 7077), or Hunt et al. (in prep.). For SBS 0335–052E and IZw 18, we take the SFR measured by high-frequency radio free-free emission (Johnson et al. 2009; Hunt et al. 2005).

Following Grossi et al. (2015), stellar masses,  $M_{\text{star}}$ , were estimated according to the recipe given by Wen et al. (2013) based on WISE W1 (3.4  $\mu$ m) luminosities. This approach exploits the relatively constant mass-to-light ratios of stellar populations, independently of metallicity and age, at near-infrared wavelengths (Norris et al. 2014; McGaugh & Schombert 2014). However, when W1 photometry was not available, we used IRAC 3.6  $\mu$ m photometry instead. In fact, the two bands are very similar; using data from Brown et al. (2014), Grossi et al. (2015) find for spirals a mean flux ratio  $F_{3.4}/F_{3.6} = 1.02 \pm 0.035$ . Including also the data for dwarf irregulars from Brown et al. (2014) gives a mean flux ratio of the W1 and IRAC 3.6  $\mu$ m bands is unity, with 5–6% scatter for galaxies like our targets.

We used the HII-galaxy formulation by Wen et al. (2013), rather than what they found for their full sample; the HII galaxies

have the lowest mass-to-light ratios in their compilation, corresponding roughly to the bluest regions of the galaxies studied by Zibetti et al. (2009). To better take into account the weak trends with abundance found by Wen et al. (2013), we also applied an approximate correction for low metallicity (by multiplying the mass-to-light ratio by 0.8: Wen et al. 2013, see their Fig. 17).

However, before applying the HII-galaxy relation by Wen et al. (2013), we first subtracted nebular emission and emission from hot dust. Such contamination can be very important in the near-infrared and can contribute 50% or more to the observed flux at these wavelengths (Hunt et al. 2001, 2002, 2012; Smith & Hancock 2009). The ionized gas continuum contribution to the  $3.4-3.6\,\mu\text{m}$  flux was estimated from the SFR using the emission coefficients from Osterbrock & Ferland (2006). The subtraction of hot dust is based on the assumption that H-band emission is entirely stellar. Because H-band photometry is available for most of our sample, we used the data from Brown et al. (2014) to estimate the maximum possible IRAC  $3.6 \,\mu\text{m}/$ H-band ratio in galaxies similar to our targets; 95% of the spiral/dwarf irregular galaxies have a flux ratio ≤2.4. This corresponds to a (Vega-based) [H-3.6] color of ~0.8, consistent with what is found for the pure stellar component in star-forming galaxies (Hunt et al. 2002). After subtraction of the nebular component, any excess over this ratio was attributed to hot dust and subtracted. We found that hot dust can contribute between 10% and 50% in the 3.4–3.6  $\mu$ m bands (most significantly for II Zw 40, SBS 0335-052E).

We compared the stellar masses obtained with the formulation of Wen et al. (2013) to those calculated according to Lee et al. (2006) based on IRAC 4.5  $\mu$ m luminosities (used by Hunt et al. 2012), and find they are consistent to within 0.2 dex. Moreover, for the galaxies having both W1 and IRAC data, the stellar masses obtained from 3.6  $\mu$ m luminosities are within ~5% of those from W1 as expected. The Kroupa-like broken powerlaw initial mass function (IMF) used by Calzetti et al. (2010) and consequently for our calculation of SFRs is consistent with the IMF for  $M_{\text{star}}$  by Wen et al. (2013).

#### 2.4. HI measurements

All the sample galaxies have been observed in H<sub>I</sub>, half of them with interferometric maps. For the galaxies with resolved observations, for comparison with CO, we have attempted to compute the H<sub>I</sub> mass,  $M_{\rm HI}$ , by integrating the mean H<sub>I</sub> surface density within only the optical radius. Because virtually all metal-poor dwarf galaxies show H<sub>I</sub> emission well outside their optical radius (e.g., Hunter et al. 1994a; van Zee et al. 1998; van Zee 2000; Ekta et al. 2009; Lelli et al. 2012, 2014), these values of  $M_{\rm HI}$  are typically a factor of 2–10 lower than the total H<sub>I</sub> mass, integrated over the entire H<sub>I</sub> extent. Table 1 reports both values of  $M_{\rm HI}$  when resolved H<sub>I</sub> maps are available.

#### 2.5. Comparison samples

To expand the parameter space for our analysis, we also considered another dwarf sample with CO detections (Schruba et al. 2012), as well as normal star-forming galaxies, LIRGs, and Ultra-LIRGs (ULIRGs, Gao & Solomon 2004; Graciá-Carpio et al. 2008; García-Burillo et al. 2012). For the latter galaxies, we converted their published IR luminosity to SFR according to Kennicutt (1998), after applying the correction to a Kroupa (2001) IMF for consistency with our SFRs and those of Schruba et al. (2012). For the Schruba et al. CO(2–1) observations, we applied the factor they recommend to convert  ${}^{12}$ CO(2–1) to  ${}^{12}$ CO(1–0):  $I_{CO2-1}/I_{CO1-0} = 0.7$ . Stellar masses for the Gao & Solomon (2004) and García-Burillo et al. (2012) samples were taken from Skibba et al. (2011), and for the galaxies from Schruba et al. (2012), masses were adopted either from Hunt et al. (2012) or Lee et al. (2006). These latter estimates are based on IRAC photometry similar to the method used here, as described in Sect. 2.3.

We incorporated also the galaxies from the CO(2-1) APEX Low-redshift Legacy Survey of MOlecular Gas (ALLSMOG, Bothwell et al. 2014). This sample is essentially an extension to lower stellar masses (8.5 <  $\log(M_{\text{star}}/M_{\odot}) \le 10$ ) of the COLDGASS galaxies studied by Saintonge et al. (2011a,b). For consistency with Schruba et al. (2012), to convert <sup>12</sup>CO(2-1) to <sup>12</sup>CO(1–0), we again use  $I_{CO2-1}/I_{CO1-0} = 0.7$ , rather than  $I_{\text{CO2-1}}/I_{\text{CO1-0}} = 1$  as in Bothwell et al. (2014). Because of the ~0.4 dex offset between the ALLSMOG metallicity calibration and the mainly direct-temperature<sup>2</sup> determinations used here (e.g., Hunt et al. 2012; Andrews & Martini 2013), the oxygen metallicities given by Bothwell et al. (2014) were converted to the Pettini & Pagel (2004) calibration according to Kewley & Ellison  $(2008)^3$ . Finally, to extend the metallicity range to extremely metal-poor galaxies (XMPs), we incorporate the sample from Filho et al. (2013) which, however, have no CO measurements; we include them to better investigate gas content at low metallicities.

Both the ALLSMOG and the Filho et al. (2013) galaxies have stellar masses and SFRs taken from the MPA-JHU catalogue (Kauffmann et al. 2003; Brinchmann et al. 2004), the same basis for calibration used by Wen et al. (2013). However, the automatic algorithm used by the SDSS does not always provide the correct global photometry for galaxies whose surface brightness is not uniformly distributed (e.g., in clumps or knots). This should not be a problem for the ALLSMOG galaxies, which tend to be extended spirals, but could be a problem for the comparison with the XMP sample by Filho et al. (2013). In fact, Filho et al. (2013) found that comparing mass-to-light ratios based on optical colors gave as much as an order of magnitude  $M_{\text{star}}$  excess over the SDSS estimates; the discrepancies tend to be largest for galaxies with either cometary or two-/multiple-knot morphology. To check for systematic differences between the nearinfrared method used here (IRAC, WISE) and in Hunt et al. (2012; see also Madden et al. 2013) and the optically-determined (MPA-JHU) SDSS values, we searched the MPA-JHU database and found 17 low-metallicity galaxies for which we had calculated stellar masses in Hunt et al. (2012) using  $4.5 \,\mu$ m luminosities. Although the contributions of the nebular continuum and hot dust emission have been subtracted, the values of  $M_{\text{star}}$  using IRAC/WISE photometry are, on average, 17 times larger than those with SDSS with a scatter of  $\sim 0.4$  dex. We investigated the morphology of the galaxies in this comparison and find that they tend to be very clumpy, with either multiple knots or cometary features. Comparing the SDSS photometry with the broad-band JHK photometry for these galaxies gives NIR-optical colors that are extremely red, unrealistically so for these blue galaxies; the implication is that the optical fluxes are being underestimated. We thus ascribe this difference to the difficulty of the

 $^2$  The electron temperature derived from the ratio of the [OIII]  $\lambda$ 4363 auroral line to lower-excitation lines ([OIII]  $\lambda$ 4959, 5007) is converted to a metallicity after correcting for unseen phases of ionization (e.g., Osterbrock & Ferland 2006).

SDSS automated photometric algorithms when confronted with multiple clumps. As a consequence, in our comparisons with the XMP sample, these uncertainties should be kept in mind.

# 3. The CO observations

We have observed  ${}^{12}CO(1-0)$  in our sample galaxies over a three-year period from 2008 to 2010 with the IRAM 30-m telescope (Pico Veleta, Spain).

#### 3.1. IRAM observations

In September, 2009, and September, 2010, we observed six galaxies<sup>4</sup> in <sup>12</sup>CO(1–0) with the Eight Mixer Receiver (EMIR) using the Wideband Line Multiple Autocorrelator (WILMA) backend. Observations of <sup>12</sup>CO(1–0) for NGC 1140 and NGC 1156 were performed with the AB receivers in November, 2008, using the 1 and 4 MHz backends at 3 and 1.3 mm (88–115 GHz, 219–229 GHz, respectively). The precipitable water vapor was  $\leq 4$  mm during the 2008 run, giving system temperatures of  $T_{sys}$  of  $\leq 370$  K at 3 mm. In 2009, the conditions were worse ( $\leq 11$  mm), but the EMIR receivers had comparable  $T_{sys}$ ,  $\sim 335$  K at 3 mm; in 2010, our most extensive run,  $T_{sys}$  varied from  $\sim 230$  K to  $\geq 400$  K, with precipitable water vapor ranging from  $\sim 4$  mm to 10 mm. In all runs, we used wobbler switching with a throw of 90″. Intensity calibration was performed with two absorbers at different temperatures.

Pointing was checked at regular intervals (every 1.5 h) on nearby planets or bright quasars (e.g., 0234+285), and focus tests were carried out every 4 h during the night and every 3 h during the day. Pointing errors never exceeded  $\sim 3''$ , and were usually  $\leq 2''$ . Given the beam size (half-power beam width) of  $\sim 22''$ , this was more than sufficient.

#### 3.2. Data reduction

We adopted the GILDAS/CLASS data reduction package to obtain mean spectra<sup>5</sup>. To remove the continuum from each spectrum, a polynomial baseline was fitted to the line-free regions of each scan (defined a priori) and subtracted; the scans were thereafter smoothed and averaged, and a constant baseline subtracted. We measured the peak intensities, central velocities, full width half-maximum (FWHM) and velocity-integrated fluxes of the detected lines by fitting Gaussian profiles to the data. In most cases, there is significant velocity structure in the emission, so we fit the line profiles to multiple Gaussians. In these cases we use the sum of the integrated line intensities from the multiple Gaussian fits in the analysis (see Table 2).

Antenna temperatures  $(T_A^*)$  have been converted to mainbeam temperatures  $(T_{mb})$  by dividing the antenna temperatures by  $\eta \equiv B_{eff}/F_{eff}$ , where  $B_{eff}$  (=0.78) and  $F_{eff}$  (=0.94) are the beam and forward hemisphere efficiencies, respectively (i.e.,  $T_{mb} = T_A^*/\eta$ ). To convert the measured <sup>12</sup>CO(1–0) brightness temperatures,  $[T_{mb}$  (K)] to fluxes [S (Jy)], we used the standard IRAM 30-m conversion factor of 3.906 Jy K<sup>-1</sup>. The <sup>12</sup>CO(1–0) line profiles are shown in Fig. 3, together with the Gaussian fits obtained using the GILDAS/CLASS package as described above and reported in Table 2. Although a few of the individual velocity components are too narrow to be significant (e.g., NGC 1140,

<sup>&</sup>lt;sup>3</sup> These metallicites are shown in Figs. 5, 6, 8, 9, 11, 13, and 14.

 $<sup>^4\,</sup>$  II Zw 40 and SBS 0335–052E were observed only in the higher-J lines with APEX.

http://www.iram.fr/IRAMFR/GILDAS

#### Table 2. <sup>12</sup>CO(1–0) emission line parameters.

Name	Telescope time (min)	Component	Channel ( km s <sup>-1</sup> )	T <sub>rms</sub> (mb) <sup>a</sup> (K)	Offset ( km s <sup>-1</sup> )	<i>FWHM</i> ( km s <sup>-1</sup> )	Integrated intensity <sup>b</sup> $(K  km  s^{-1})$
CGCG007-025	524	Total Total	10.5 15.7	0.0018 0.0016	-39 -39	115 105	0.286 (0.095) 0.285 (0.102)
II $\mathbb{Z}$ w 40 <sup>c</sup>	-						
Mrk 996	573	1 2 Total	5.2 5.2 5.2	0.0019	$-1 \\ 39 \\ 0$	28 10 -	0.18 (0.03) 0.05 (0.02) 0.23 (0.04)
NGC 1140	831	1 2 3 4 Total Total <sup>d</sup>	5.2 5.2 5.2 5.2 5.2 5.2 5.2	0.0016	-17 3 30 64 0	26 8 38 7 -	$\begin{array}{c} 0.32 \ (0.05) \\ 0.07 \ (0.04) \\ 0.18 \ (0.05) \\ 0.04 \ (0.02) \\ 0.61 \ (0.08) \\ 1.00 \ (0.13) \end{array}$
NGC 1156	47	1 2 3 Total Total <sup>d</sup>	10.4 10.4 10.4 10.4 10.4	0.0076 0.0076	$-84 \\ -35 \\ 44 \\ 0 \\ 0$	13 100 42 -	$\begin{array}{c} 0.36 \ (0.09) \\ 0.89 \ (0.09) \\ 0.48 \ (0.09) \\ 1.25 \ (0.16) \\ 4.04 \ (0.52) \end{array}$
NGC 3353 [0", 0"]	79	1 2 Total	5.2 5.2 5.2	0.0056	-31 21 0	60 16 -	1.83 (0.17) 0.34 (0.11) 2.17 (0.20)
NGC 3353 [0", +10"]	77	lotal <sup>a</sup> 1 2 Total	5.2 5.2 5.2 5.2	0.0045	-35 19	42 22	3.07 (0.29) 1.38 (0.12) 0.43 (0.12) 1.81 (0.17)
NGC 3353 [-6", -9"]	80	1 2 3 Total	5.2 5.2 5.2 5.2 5.2	0.0046	-17 25 98 0	71 36 31 -	$\begin{array}{c} 1.81 (0.17) \\ 0.84 (0.20) \\ 0.62 (0.18) \\ 0.30 (0.08) \\ 1.46 (0.27) \end{array}$
NGC 7077	585	1 2 Total	5.2 5.2 5.2	0.0017	$-5 \\ 35 \\ 0$	20 30	0.25 (0.03) 0.20 (0.04) 0.45 (0.05)
$SBS0335-052E^e$	_						
UM 448	454	1 2 3 Total	5.3 5.3 5.3 5.3	0.0017	-115 -58 1 0	20 63 44 -	0.11 (0.04) 0.42 (0.16) 0.52 (0.14) 1.04 (0.22)
UM 462	119	Total Total	10.4 15.7	0.0034 0.0032	-9 -8	121 116	0.824 (0.173) 0.749 (0.207)

**Notes.** <sup>(a)</sup> This and all temperature units in this table are main-beam,  $T_{mb}$ . <sup>(b)</sup> Values in parentheses are the uncertainties. The total line flux is the sum of the individual components, and the total uncertainty is calculated by adding in quadrature the individual uncertainties. <sup>(c)</sup> As discussed in the text, we did not observe this galaxy in <sup>12</sup>CO(1–0), but rather relied on the data reported by Taylor et al. (1998). <sup>(d)</sup> With aperture corrections as described in the text. <sup>(e)</sup> We did not observe this galaxy in <sup>12</sup>CO(1–0), but rather adopted the <sup>12</sup>CO(3–2) data reported by Hunt et al. (2014), converted to CO(1–0) assuming a CO(3–2)/CO(1–0) flux ratio of 0.6.

NGC 1156), we have checked that the total intensities are robust to the details of the fitting.

#### 3.3. Aperture corrections

The IRAM 30 m has a beam with an FWHM of  $\sim 22$  " at 3 mm. Consequently, the largest galaxies in our sample (NGC 1140,

NGC 1156, NGC 3353, see Table 1) may be missing flux. We have attempted to correct for this by adopting the approach of Saintonge et al. (2011a) who calibrated aperture corrections using the set of nearby galaxies observed by Kuno et al. (2007). We applied Eq. (2) by Saintonge et al. (2011a) based on optical size, but used the geometric means of the major and minor axes given in Table 1. These corrections amount to an increase







**Fig. 3.** <sup>12</sup>CO(1–0) average spectra for the observed galaxies. *Top row, from left to right* in alphabetical order of name: CGCG 007–025, Mrk 996, and NGC 1140. *Second row*: NGC 1156, NGC 3353 (Haro 3) two pointings ([0, 0], [0, +10"]. *Third row*: NGC 3353 third pointing [–6", –9"]), NGC 7077 (Mrk 900), UM 448. *Bottom row*: UM 462. The baselines are shown as a horizontal solid line, together with the multiple-component Gaussian fits as described in Sect. 3.2. The vertical axes are in  $T_A^*$  units, while in Table 2 the units are  $T_{\rm mb}$ , converted from  $T_A^*$  as described in the text. Velocity channels are as reported in Table 2.

in line intensities by a factor of 1.64 for NGC 1140 and 3.23 for NGC 1156.

For NGC 3353 (Haro 3), we have additional pointings spaced by roughly 0.5 beams, hence not independent. Where necessary, Saintonge et al. (2011a) used spacings of 0.75 beams along the major axis, and found a mean ratio between the offset flux and the central flux of 0.33. The corresponding ratio for the [-6", -9"] pointing of NGC 3353 (along the southern major axis) is 0.53, larger than the offset corresponding to the wider spacing by Saintonge et al. (2011a), but consistent with what could be expected for the closer spacing of our offset. Hence, for NGC 3353, we adopted Eq. (3) of Saintonge et al. (2011a) with their mean ratio between offset and central intensities (0.33); this correction gives a multiplicative factor of 1.41. Table 2 also reports the  ${}^{12}CO(1-0)$  intensities corresponding to the aperture corrections for these galaxies which we will adopt throughout the paper.

#### 3.4. Comparison with previous CO observations

Because some of the galaxies in our sample have previous CO observations, here we compare our results with those. Sage et al. (1992) observed NGC 3353, NGC 7077, UM 448, and UM 462, and securely detected only the former two. The results for NGC 3353, obtained with the IRAM 30-m, are in good agreement with our central pointing (they find an integrated CO intensity of  $1.98 \pm 0.28$  K km s<sup>-1</sup>), although we find ~70% of their CO emission for NGC 7077 (they find  $0.68 \pm 0.1$  K km s<sup>-1</sup>).

A114, page 8 of 17

Our detections of UM 448 and UM 462 are consistent with their marginal detections obtained with the NRAO 12-m.

UM 462 was also observed by Brinks & Taylor (1997) and Gondhalekar et al. (1998), neither of whom detected the galaxy; their upper limits (<0.28 K km s<sup>-1</sup> and <0.5 K km s<sup>-1</sup>, respectively) are somewhat discrepant with our detection (0.7–0.8 K km s<sup>-1</sup>), as well as with the marginal detection of Sage et al. (1992,  $\sim$ 0.7 K km s<sup>-1</sup>).

NGC 1140 was observed previously by Albrecht et al. (2004) and NGC 1156 by Leroy et al. (2005). The two previous observations are reasonably consistent with our results.

## 4. CO luminosity and SFR

Our new IRAM observations have successfully detected  ${}^{12}CO(1-0)$  down to a metallicity of  $12 + \log(O/H) = 7.74$  (CGCG 007–025, signal-to-noise ratio 3.0, see Table 2), which is the lowest metallicity at which CO has been (tentatively) detected (see WLM:  $12 + \log(O/H) = 7.8$ , Elmegreen et al. 2013). To our knowledge, Table 2 gives the largest number of  ${}^{12}CO(1-0)$  detections ever reported for  $12 + \log(O/H) \leq 8$  (0.2  $Z_{\odot}$ ) outside the Local Group.

In the following, we compare the integrated CO luminosity,  $L'_{CO}$ , with SFR as calculated in Sect. 2.3.  $L'_{CO}$  was derived according to the formulation of Solomon & Vanden Bout (2005), using the appropriate Jy-to-K conversion (see Sect. 3 for the IRAM value). The result is shown in Fig. 4.

Not surprisingly, most of the metal-rich star-forming galaxies in the comparison samples follow the linear correlation



**Fig. 4.** SFR vs.  $L'_{CO}$  for our sample and additional galaxies from the literature. As discussed in Sect. 2.2, we include IZw 18 as one of our sample galaxies (see also Schruba et al. 2012). The open triangles for Schruba et al. (2012) correspond to their compilation of literature data and the filled ones to their metal-poor dwarf observations. CO non-detections are shown by leftward arrows. The rightmost regression line corresponds to the linear correlation found by Gao & Solomon (2004), and the leftmost one to our new determination, also compatible with a slope of unity (see text). The horizontal dotted line gives the IR luminosity limit above which the linear LIR- $L'_{CO}$  correlation of Gao & Solomon (2004) no longer holds.

between SFR and  $L'_{CO}$ , as formulated by Gao & Solomon (2004), shown as the rightmost dashed line (corrected to the Kroupa IMF used here for SFR, see Sect. 2); the horizontal dotted line in Fig. 4 gives the IR luminosity limit above which the linear LIR- $L'_{CO}$  correlation of Gao & Solomon (2004) no longer holds. The metal-poor dwarf galaxies in our sample show a similar correlation but offset to lower  $L'_{CO}$  by a factor of ~30:

$$\log(SFR) = \log(L'_{CO}) - (6.92 \pm 0.07).$$

The best-fitting slope obtained with a robust fitting technique (Li 1985; Fox 1997) is unity within the errors  $(1.05 \pm 0.15)$ ; thus, for consistency with Gao & Solomon (2004), we fixed a unit slope and derived the corresponding intercept as given above and shown in Fig. 4. Although we did not use the data of Schruba et al. (2012) to establish the best-fit regression, most of those galaxies also follow the trend for our sample (see Fig. 4). The exceptions are the two most metal-poor galaxies, SBS 0335–052E and IZw 18.

Judging from our sample alone, the offset between SFRs and high  $L'_{CO}$  and low  $L'_{CO}$  is apparently independent of metallicity, at least to the lowest oxygen abundance of our sample,  $12 + \log(O/H) \sim 7.7$ . However, CGCG 007–025 at this metallicity is only tentatively detected, and our other detected galaxies have a relatively narrow spread in metallicities,  $12 + \log(O/H) \sim 8$ . If we include in the regression the Schruba et al. (2012) compilation and their five  ${}^{12}CO(2-1)$  detections (converted to  ${}^{12}CO(1-0)$  as described above), we find a significant correlation between the SFR/ $L'_{CO}$  ratio and  $12 + \log(O/H)$ , as shown in Fig. 5 (Pearson parametric correlation coefficient, -0.88, >99.999% significance level). Two galaxies are



**Fig. 5.** SFR/ $L'_{CO}$  vs. 12 + log(O/H) for our sample and additional galaxies from the literature. As discussed in Sect. 2.2, we include IZw 18 as one of our sources (see also Schruba et al. 2012). The open triangles for Schruba et al. (2012) correspond to their compilation of literature data and the filled ones to their metal-poor dwarf observations. CO non-detections are shown by upward arrows. The two horizontal dashed lines correspond to the regressions with unit slope shown in Fig. 4, and the dotted one to a robust regression as described in the text: log(*SFR*/ $L'_{CO}$ ) = (-2.25 ± 0.15) [12 + log(O/H)] + (11.31 ± 1.3).

discrepant with this metallicity trend, CGCG 007–025 and SBS 0335–052E. SBS 0335–052E was not directly observed in <sup>12</sup>CO(1–0), so there may be some problems with the multiplicative factor relating CO(3–2) to CO(1–0) (see Hunt et al. 2014). CGCG 007–025, on the other hand, is directly observed in <sup>12</sup>CO(1–0) but the detection is only tentative. More CO(1–0) observations at metallicities below 12 + log(O/H) = 8 are needed to verify this in <a href="#sonsistency">sonsistency</a>, especially because of the expectation that for  $Z/Z_{\odot} \leq 0.1$ , CO may cease to be a viable tracer of H<sub>2</sub> (Bolatto et al. 2013). We discuss the implications of the relation between SFR/ $L'_{CO}$  and abundance in the following section.

# 5. Molecular depletion times, the CO-to-H<sub>2</sub> conversion factor, and specific SFR

Although we have shown that  $L'_{CO}$  is correlated with SFR even at low metallicity, the main parameter of interest is H<sub>2</sub> mass,  $M_{H2}$ . In fact, the study of low-metallicity star formation and H<sub>2</sub> content is hampered by the unknown factor,  $\alpha_{CO}$ , used to convert observed CO intensities to  $M_{H2}$ . Observationally,  $M_{H2}$ ,  $\alpha_{CO}$ , and the time required to consume the molecular gas reservoir,  $\tau_{dep}$ , are intimately related and difficult to disentangle. By definition,  $\tau_{dep}$  depends inversely on the star-formation rate  $\tau_{dep} \equiv M_{H2}/SFR = L'_{CO} \alpha_{CO}/SFR$ , but the potential simplicity of predicting  $M_{H2}$  from SFR (or vice versa) through  $\tau_{dep}$  is compromised by the necessary CO-to-H<sub>2</sub> factor. The observed (or directly derivable) quantities are  $L'_{CO}$  and SFR; thus only the ratio,  $\tau_{dep}/\alpha_{CO}$ , is constrained.

Much effort has been devoted to finding the dependence of  $\alpha_{CO}$  on metal abundance (e.g., Wilson 1995; Israel 1997; Rosolowsky et al. 2003; Bolatto et al. 2008, 2013; Leroy et al. 2009b, 2011; Genzel et al. 2012; Schruba et al. 2012).



**Fig. 6.** Specific SFR vs.  $12 + \log(O/H)$  for our sample and including data from the literature (Hunt et al. 2012; Schruba et al. 2012; Filho et al. 2013; Bothwell et al. 2014). The metallicities for the ALLSMOG (Bothwell et al. 2014) sample have been modified as discussed in the text. As in Fig. 4, the open triangles for Schruba et al. (2012) correspond to their compilation of data from the literature and the filled ones to their metal-poor dwarf observations. The regression line is a robust best-fit of the Local Volume Legacy (LVL, Lee et al. 2009) and the dwarf irregulars from Lee et al. (2006). The various colors for the LVL give the *Hubble* type (early types red, magenta; late types blue, cyan).

High-resolution cloud virial mass measurements tend to find little or no dependence on metallicity (Wilson 1995; Rosolowsky et al. 2003; Bolatto et al. 2008). However, because CO is found only in dense regions, rather than in diffuse gas,  $\alpha_{CO}$  depends on the physical scales probed by the observations, especially at low metallicity (Rubio et al. 1993). Given the relatively constant  $\tau_{dep}$  in nearby star-forming spirals (e.g., Bigiel et al. 2008, 2011; Leroy et al. 2008), some groups have assumed a constant  $\tau_{dep}$  to infer  $\alpha_{CO}$  in metal-poor galaxies. Such an assumption tends to give a steep dependence of  $\alpha_{\rm CO}$  on O/H:  $\alpha_{\rm CO}^{-1} \propto (Z/Z_{\odot})^{\sim 2-3}$ (Genzel et al. 2012; Schruba et al. 2012). An alternative approach relies on IR observations to derive  $M_{\rm H2}$  and infer  $\alpha_{\rm CO}$ by estimating the dust-to-gas ratio and assuming it is the same in adjacent regions with/without H2. This method has the considerable advantage of being independent of CO observations altogether, but requires comparable quantities of both  $H_1$  and  $H_2$ , and relies on the assumption that emissivity properties of dust grains are the same in dense and diffuse environments. Recent work using this method, introduced by Israel (1997), also gives super-linear slopes, although somewhat shallower (and with large scatter) than those obtained with the assumption of con-stant  $\tau_{dep}$ :  $\alpha_{CO}^{-1} \propto (Z/Z_{\odot})^{\sim 1.6-2}$  (Gratier et al. 2010; Leroy et al. 2011; Bolatto et al. 2011).

Here we present a different approach to constrain  $\alpha_{CO}$  at low metallicity. We exploit the results by Saintonge et al. (2011b; see also Huang & Kauffmann 2014) who find a significant inverse correlation of  $\tau_{dep}$  with sSFR: galaxies with higher sSFR also have shorter molecular gas depletion times. However, sSFR tends to also be inversely correlated with metallicity (e.g., Salim et al. 2014), as shown in Fig. 6 (adapted from Hunt et al. 2012). Thus, we must find a way to disentangle the effects of sSFR on  $\tau_{dep}$  from those of metallicity.



**Fig. 7.**  $L'_{CO}$ /SFR vs. sSFR for our previous IRAM sample and including data from the literature (COLDGASS: Huang & Kauffmann 2014; Gao & Solomon 2004; García-Burillo et al. 2012; Schruba et al. 2012; Hunt et al. 2014). Our best-fit of the COLDGASS+Gao&Solomon+Garcia-Burillo samples is shown (which is consistent with the regression given by Saintonge et al. 2011b, for only the COLDGASS galaxies).

#### 5.1. The CO-to-H<sub>2</sub> conversion factor

Saintonge et al. (2011b) found that star-forming galaxies of lower mass have much shorter  $\tau_{dep}$  than more massive ones, while the depletion time for atomic gas remains relatively constant. They also found that  $\tau_{dep}$  is strongly correlated with sSFR, with a single relation being able to fit both "normal" ("mainsequence", e.g., Salim et al. 2007) galaxies and more extreme starbursts including LIRGs and ULIRGs. Their relation is based on a single value of  $\alpha_{CO}$ , appropriate for the properties of their sample, selected by stellar mass.

Here we explore the hypothesis that the Saintonge et al. correlation between  $\tau_{dep}$  and sSFR extends to the lower-mass (and higher sSFR) galaxies in our sample. However, for a better comparison with our SFR estimates, we use the COLDGASS data by Huang & Kauffmann (2014) who re-evaluated the SFRs within the IRAM beam using additional mid-infrared and UV data for COLDGASS galaxies. Because there is some evidence that the SFRs from fitting of spectral-energy distributions such as done by Saintonge et al. (2011b) are overestimated relative to those from H $\alpha$ +24  $\mu$ m or FUV+24  $\mu$ m (e.g., Leroy et al. 2013a; Huang & Kauffmann 2014), the comparison with the COLDGASS sample should be more consistent using the SFRs by Huang & Kauffmann (2014)<sup>6</sup>. In addition to the COLDGASS data, we also consider other data from the literature where both CO measurements and stellar masses are available (see Sect. 2.5). The dashed line in Fig. 7 indicates our fit with a robust linear regression to COLDGASS data (using the SFRs and  $M_{\text{star}}$  within a 22" aperture by Huang & Kauffmann 2014), together with the samples of Gao & Solomon (2004) and García-Burillo et al. (2012):

<sup>&</sup>lt;sup>6</sup> We have performed the entire analysis with both datasets and confirm an overestimate of ~0.2 dex in the SFRs measured by Saintonge et al. (2011b) relative to Huang & Kauffmann (2014). This change affects only slightly the metallicity dependence of  $\alpha_{CO}$ , steepening the powerlaw index by 0.08 with respect to the Huang & Kauffmann (2014) data.



**Fig. 8.** Residuals from the regression shown in Fig. 7 vs. sSFR (*left panel*) and vs.  $12 + \log(O/H)$  (*right*). We include here data for our sample and literature data (COLDGASS: Huang & Kauffmann 2014; Gao & Solomon 2004; García-Burillo et al. 2012; Schruba et al. 2012; Bothwell et al. 2014). The horizontal dashed line in the *left panel* shows 0 to aid the eye, while the regression in the *right panel* is the robust fit to the significant trend of the residuals of the  $L'_{CO}$ /SFR fit with metallicity  $\propto (Z/Z_{\odot})^{1.96}$ .



**Fig. 9.**  $\tau_{dep}$  vs. sSFR (*left panel*) and vs. 12 + log(O/H) (*right*) for our sample and including data from the literature (COLDGASS: Huang & Kauffmann 2014; Gao & Solomon 2004; García-Burillo et al. 2012; Schruba et al. 2012; Bothwell et al. 2014). As in previous figures, the open triangles for Schruba et al. (2012) correspond to their compilation of data from the literature and the filled ones to their metal-poor dwarf observations. The regression line is our fit to the Schruba et al. (2012) and our sample as described in the text.

 $log(L'_{CO}/SFR) = (-0.38 \pm 0.03) log(sSFR) + (4.75 \pm 0.31)$ . The new fit is consistent with that found by Huang & Kauffmann (2014) for only the COLDGASS galaxies. However, both the dwarf galaxies in our sample and those in Schruba et al. (2012) show large residuals relative to the general trend.

The residuals to this fit are plotted vs. sSFR and O/H in Fig. 8. Because  $\tau_{dep} = \alpha_{CO} L'_{CO}/SFR$ , the residuals from the correlation of  $L'_{CO}/SFR$  on sSFR should eliminate the dependence on  $\tau_{dep}$ :  $[L'_{CO}/SFR]$ (observed)/ $\tau_{dep}$ (fit) =  $[\alpha_{CO}(Z)/\alpha_{CO}(Z_{\odot})]^{-1}$ . Thus, these residuals should correspond directly to the inverse of  $\alpha_{CO}$ , relative to  $Z_{\odot}$ . The left panel of Fig. 8 demonstrates that there is no additional correlation with sSFR not accounted for by the COLDGASS relation; however, the right panel shows that there is a clear correlation with O/H. The correlation of the residuals with metallicity is able to almost completely rectify the deviation of the low-metallicity galaxies by almost two orders

of magnitude from the COLDGASS galaxies. We find a robust fitted slope of  $\alpha_{\rm CO}^{-1} \propto (Z/Z_{\odot})^{1.96\pm0.16}$ , more shallow than found with the assumption of constant  $\tau_{\rm dep}$  (Schruba et al. 2012; Genzel et al. 2012), but consistent with the  $\alpha_{\rm CO}$  metallicity dependence found by using dust continuum measurements (Leroy et al. 2011; Bolatto et al. 2011).

For the calculation of  $M_{\rm H2}$ , we adopt a solar value  $\alpha_{\rm CO} = 3.2 \, M_{\odot}$  (K km s<sup>-1</sup> pc<sup>2</sup>)<sup>-1</sup>, not including a factor of 1.36 for helium (see also Saintonge et al. 2011b). For our galaxies and those from Schruba et al. (2012) and Bothwell et al. (2014) having metallicities  $Z/Z_{\odot} < 1$  (12 + log(O/H) < 8.69), we applied  $\alpha_{\rm CO} \propto (Z/Z_{\odot})^{-1.96}$ ; for metallicities  $Z/Z_{\odot} \ge 1$ , we used the constant  $\alpha_{\rm CO}$  as given above. Figure 9 shows the result of converting our  $L'_{\rm CO}$  observations to  $M_{\rm H2}$ . Although there is some scatter, the trend of  $\tau_{\rm dep}$  with sSFR established by Saintonge et al. (2011b), Huang & Kauffmann (2014) seems to extend to the higher sSFRs



**Fig. 10.**  $\tau_{dep}$  vs.  $M_{star}$  for our sample and including data from the literature (Huang & Kauffmann 2014; Schruba et al. 2012; Bothwell et al. 2014). As in previous figures, the open triangles for Schruba et al. (2012) correspond to their compilation of data from the literature and the filled ones to their metal-poor dwarf observations. Our robust best-fit regression for these samples is shown as a dashed line; the dotted line corresponds to the slope for the COLDGASS galaxies alone (Huang & Kauffmann 2014).

and shorter  $\tau_{dep}$  of these dwarf galaxies. A steeper dependence of  $\alpha_{CO}$  on metallicity would be inconsistent with this trend, but more data at  $12 + \log(O/H) \leq 8$  are needed to better assess any systematics.

In any case, in agreement with previous work (e.g., Saintonge et al. 2011b; Bothwell et al. 2014; Huang & Kauffmann 2014), our analysis shows a strong variation of the molecular gas depletion times, with values ranging from ~10 Gyr for the COLDGASS galaxies to ~70 Myr for II Zw 40 and even smaller,  $\leq 10$  Myr for SBS 0335–052E, NGC 5253 and NGC 1569 (from Schruba et al. 2012). This variation is correlated with a range in sSFRs of almost three orders of magnitude.

The right panel of Fig. 9 suggests that  $\tau_{dep}$  depends weakly on metallicity, even after the  $\alpha_{CO}$  dependence has been accounted for. The regression line is the robust fit to our galaxies and those from Schruba et al. (2012); the correlation is significant (Pearson coefficient of 0.38, significance >99.9%), with a slope of 0.64 ± 0.11. A similar (weak) effect was also found by Leroy et al. (2013a), but over a narrower range in metallicities and  $\tau_{dep}$ . Because  $\tau_{dep}$  and sSFR are correlated, and sSFR and O/H are also correlated (see Fig. 6), such a result is perhaps not surprising; it could be the result of the secondary dependence on sSFR or on  $M_{star}$ . In Sect. 7, we discuss forms of  $\alpha_{CO}$  and its metallicity dependence (e.g., Wolfire et al. 2010; Krumholz et al. 2011).

Because O/H and  $M_{\text{star}}$  are correlated through the massmetallicity relation, in Fig. 10, we have investigated the possibility that  $\tau_{\text{dep}}$  depends on  $M_{\text{star}}$ . This correlation is also formally significant, even more so than the one with O/H (Pearson coefficient of 0.56, significance >99.999%). The slope of 0.46 ± 0.03 (shown by a dashed line in Fig. 10) is similar to that (0.41±0.05) found for the COLDGASS galaxies alone. Our results suggest that low-mass metal-poor galaxies tend to have high sSFR (Fig. 6) and shorter H<sub>2</sub> depletion times (Figs. 9, 10), and the two properties are correlated. This result is not altogether new, although we have shown that it extends to much lower stellar masses and higher sSFRs than the COLDGASS sample.

#### 6. Atomic and molecular gas content

The implication is that for low-mass, metal-poor galaxies, with high sSFR,  $\tau_{dep}$  is systematically shorter than for more massive, metal-rich systems. Because our sample combined with the Schruba et al. (2012) and the Filho et al. (2013) compilation spans a wide range of O/H and sSFR, with <sup>12</sup>CO(1–0) detections down to  $Z/Z_{\odot} \sim 0.1$ , it is well suited to assess the dependence on these parameters of molecular and total gas (H<sub>1</sub>+H<sub>2</sub>) content, relative to stellar mass and total baryonic mass. The H<sub>I</sub> masses,  $M_{\rm HI}$ , for our sample are given in Table 1 (see also Sect. 2.4) and we have taken  $M_{\rm HI}$  for the Schruba et al. (2012) galaxies from Walter et al. (THINGS, 2008) when available, or otherwise from Kennicutt et al. (2003).

We first investigate any trends of  $M_{\rm H2}$ -to-total gas mass fraction in Fig. 11, both as a function of sSFR (left panel) and 12 + log(O/H) (right). We use total H<sub>I</sub> mass for this comparison, although it is almost certainly more extended than the H<sub>2</sub> (see Table 1), which usually resides within the optical confines of the galaxy (Regan et al. 2001; Leroy et al. 2009a). The combined samples suggest a weak trend for H<sub>2</sub> fractions to decrease at low sSFR, but there is no trend with O/H. A better assessment of both trends would require more CO data from high-sSFR and metal-poor galaxies (and resolved H<sub>I</sub> measurements).

Figure 12 shows the total gas  $M_{\text{gas}} \equiv M_{\text{H2}} + M_{\text{HI}}$  fraction relative to the total baryonic mass  $M_{\text{baryonic}} \equiv M_{\text{gas}} + M_{\text{star}}$ ) as a function of stellar mass. As in Fig. 11, we use the total H I mass for our targets (Col. 12 in Table 1). By including also the data from Schruba et al. (2012), Filho et al. (2013), Bothwell et al. (2014), Huang & Kauffmann (2014), we find there is a significant correlation between gas mass fraction and  $M_{\text{star}}$ , as shown by the robust regression line. This is mainly due to the spread in  $M_{\text{star}}$  of the two most extreme samples, COLDGASS and Filho et al. (2013). At low gas-mass fractions, this may also be partially induced by the co-dependence of the two axes, because of the presence of  $M_{\text{star}}$ . Nevertheless, the trend is roughly continuous over a factor of 10<sup>5</sup> of  $M_{\text{star}}$ , from ~10<sup>6</sup> to ~10<sup>11</sup>  $M_{\odot}$ .

Two of our sample galaxies are clear outliers, Mrk 996 and NGC 7077, both of which have a smooth elliptical-type morphology despite their low metallicity. Thus they may have already exhausted much of their H<sub>1</sub>, since the H<sub>2</sub>-to-total gas mass fractions are relatively high, ~40%. These two galaxies could be in the process of quenching their star formation, unless more gas is accreted from the outside.

We examine in Fig. 13 the trend of total gas mass relative to  $M_{\text{baryonic}}$  vs. sSFR (left panel) and  $12 + \log(O/H)$  (right). Here also the total gas fraction is significantly correlated with both sSFR and  $12 + \log(O/H)$  (see the robust regression lines), although there are significant outliers in our sample for the metallicity trend, which is mainly dominated by the XMP galaxies. However, as in Fig. 12, there may be a co-dependence of the two axes because of the presence of  $M_{\text{star}}$  (left panel) and because of the stellar mass-metallicity correlation (right).

The H<sub>I</sub> depletion times,  $\tau_{dep}(H_I)$ , for our combined sample are shown in Fig. 14. There is more than an order-ofmagnitude scatter, but no systematic trend relative to sSFR. This result is similar to that found for more massive galaxies, and the horizontal dashed line in Fig. 14 corresponds to the roughly constant  $\tau_{dep}(H_I)$  of 3.4 Gyr found for the GASS galaxies by Schiminovich et al. (2010). Figure 14 shows a possible weak trend with O/H, which starts to affect  $\tau_{dep}(H_I)$  at  $12 + \log(O/H) \leq 8$ . This could be a consequence of the large fraction of H<sub>I</sub>-dominated galaxies at these metallicities (e.g., Krumholz 2013).



**Fig. 11.**  $M_{\text{H2}}/M_{\text{gas}} = M_{\text{H2}}/(M_{\text{HI}} + M_{\text{H2}})$  vs. sSFR (*left panel*) and vs. 12 + log(O/H) (*right*) for our sample and including data from the literature (Schruba et al. 2012; Bothwell et al. 2014); the *left panel* gives also the COLDGASS data (Huang & Kauffmann 2014). As in previous figures, the open triangles for Schruba et al. (2012) correspond to their compilation of data from the literature and the filled ones to their metal-poor dwarf observations.



**Fig. 12.**  $M_{\text{gas}} = (M_{\text{HI}} + M_{\text{H2}})/M_{\text{baryonic}}$  vs.  $M_{\text{star}}$ , including data from the literature (COLDGASS: Huang & Kauffmann 2014; Schruba et al. 2012; Filho et al. 2013; Bothwell et al. 2014). Our robust best-fit regression for these samples is shown as a dashed line.

In general, we find that galaxies are more gas rich at low stellar masses, at high sSFRs, and at low metallicities, although the metallicity trends tend to have more scatter. Similar conclusions have been reached by Bothwell et al. (2014) and by Filho et al. (2013), although for a more limited range in parameter space. Our combined samples span five orders of magnitude in  $M_{\text{star}}$ , a factor of ~10<sup>3</sup> in sSFR, and a factor of ~100 in oxygen abundance. The trends for total gas fraction are similar to those we find for the molecular gas depletion time  $\tau_{\text{dep}}$ : low-mass, metalpoor, and high sSFR galaxies tend to have shorter  $\tau_{\text{dep}}$ .  $\tau_{\text{dep}}$  decreases by a factor of ~100 over a variation of ~1000 in sSFR and a factor of ~10 in O/H. However, H I depletion times are roughly constant, although with large scatter, across this entire range of sSFR and O/H.

# 7. Discussion

The short molecular gas depletion times,  $\tau_{dep} \leq 100$  Myr, we find at the highest sSFRs ( $\geq 0.1$  Gyr<sup>-1</sup>) are comparable to starbursts at  $z \ge 1$ ; they are smaller than those found for "main-sequence" galaxies at similar redshifts (Genzel et al. 2015). Thus, the low-metallicity galaxies studied here may be low-mass analogues of starbursts observed at cosmological distances. This is not unlikely since most of them are likely merging or interacting systems (see Sect. 2.1). A starburst could enhance star-formation efficiency, namely the quantity of H<sub>2</sub> converted into stars over a free-fall time (e.g., Krumholz et al. 2009); mergers or interactions would be expected to drive gas into the central regions, causing higher gas densities which could, at least partly, compensate for the low metallicity through increased self shielding.

Independently of the origin of the short molecular gas depletion times, the secondary dependence of  $\tau_{dep}$  on  $M_{star}$  and/or metallicity is not straightforward to interpret. Because of the way we have derived the molecular gas depletion times, the determination of a power-law dependence of  $\alpha_{CO}$  on metallicity is intimately related to  $au_{dep}$ . To better understand whether short  $\tau_{dep}$  and its dependence on  $M_{star}$  and O/H are artefacts of our derivation of  $\alpha_{\rm CO}$ , here we investigate an alternative formulation of  $\alpha_{\rm CO}$ , and its potential relation with  $\tau_{\rm dep}$ . What we have defined as  $\tau_{dep}$  is the inverse of what Krumholz et al. (2011) call specific gas SFRs, SSFR<sub>gas</sub>(CO, H<sub>2</sub>, total), the ratio of SFR surface densities to the various gas components of the ISM:  $\Sigma_{gas}$ ,  $\Sigma_{H2}$ , and  $\Sigma_{HI}$ . Given a total gas mass surface density,  $\Sigma_{gas}$ , Krumholz et al. give precise predictions for the fraction of molecular gas mass in the form of  $H_2$ ,  $f_{H_2}$ , as well as the fraction of molecular gas where CO dominates over other forms of carbon,  $f_{CO}$ .

In their models (see also Krumholz et al. 2009; McKee & Krumholz 2010),  $f_{H_2}$  is a function of the ratio of the interstellar radiation field (ISRF) at the surface of the gas cloud,  $G'_0$ , and the hydrogen number volume density,  $n_H$ , as well as the dust optical depth to UV photons,  $\tau_{dust}$ .  $f_{H_2}$  depends on metallicity through  $\tau_{dust}$  which is assumed to be linearly proportional to  $Z/Z_{\odot}$ . The transition in a cloud of atomic and ionized carbon to CO is driven mainly by dust extinction, thus cloud optical depth. The fraction of gas mass where the carbon is in CO form,  $f_{CO}$ , depends on  $f_{H_2}$ , modified by an exponential function which depends only weakly on  $G'_0$ , and  $n_H$ , but strongly on metallicity through  $A_V$  (i.e.,  $\tau_{dust}$ ), assumed to vary linearly with  $Z/Z_{\odot}$  (for more details see Wolfire et al. 2010). In these models, SFR surface density depends on  $f_{H_2}$ .



**Fig. 13.**  $M_{gas} = (M_{HI} + M_{H2})/M_{baryonic}$  vs. sSFR (*left panel*) and vs. 12 + log(O/H) (*right*) for our sample and including data from the literature (Schruba et al. 2012; Filho et al. 2013; Bothwell et al. 2014); the *left panel* gives also the COLDGASS detections (Saintonge et al. 2011b). As in previous figures, the open triangles for Schruba et al. (2012) correspond to their compilation of data from the literature and the filled ones to their metal-poor dwarf observations. The regression lines shown in both panels correspond to our fit of the galaxies from Schruba et al. (2012), Filho et al. (2013), ALLSMOG, and our sample as described in the text (with all the uncertainties alluded to in Sect. 2.5).



**Fig. 14.**  $\tau_{dep}(H_I)$  vs. sSFR (*left panel*) and vs. 12 + log(O/H) (*right*) for our sample and including data from the literature (Schruba et al. 2012; Bothwell et al. 2014). As in previous figures, the open triangles for Schruba et al. (2012) correspond to their compilation of data from the literature and the filled ones to their metal-poor dwarf observations. The constant  $\tau_{dep}(H_I)$ , shown by a horizontal dashed line, corresponds to 3.4 Gyr as found for the GASS sample (Schiminovich et al. 2010).

and  $\Sigma_{\text{gas}}$ , and is defined by a roughly constant timescale, 2.6 Gyr. Thus, by assuming a characteristic value for  $G'_0/n_{\text{H}}$ , Krumholz et al. (2011) derive the dependence of the gas SSFRs on gas mass surface density,  $\Sigma_{\text{gas}}$ , and metallicity,  $Z/Z_{\odot}$ .

Figure 15 shows their predictions together with the data we have analyzed here. Although the models are formulated in terms of surface densities, to better compare with global values Krumholz et al. (2011) include a resolution correction factor (c), which we have adopted according to their prescription. Following Krumholz et al. (2011), the data points correspond to molecular gas masses,  $M_{\rm H2}$ (CO), inferred from CO luminosities using a constant conversion factor,  $\alpha_{\rm CO} = 4.4 M_{\odot}$ (K km s<sup>-1</sup> pc<sup>2</sup>)<sup>-1</sup>; this is the same  $\alpha_{\rm CO}$  as used up to now (see Saintonge et al. 2011b) but here with the inclusion of helium. The CO-inferred molecular mass has been derived assuming a total gas surface density  $\Sigma_{\rm gas} = 20 M_{\odot} \, {\rm pc}^{-2}$ ; as shown by the blue curves in Fig. 15, the exact value of  $\Sigma_{gas}$  is not critical. As emphasized by Krumholz et al. (2011), this inferred mass  $M_{\rm H2}(\rm CO)$  should not be interpreted as the total mass of molecular gas, but rather the mass of the molecular gas where the predominant form of carbon is CO. Also shown in Fig. 15 are the Krumholz et al. (2011) model predictions<sup>7</sup> as a function of metallicity,  $Z/Z_{\odot}$ , for three values of total gas column density (for more details see Krumholz et al. 2011). The models of CO-inferred H<sub>2</sub> mass differ from the true H<sub>2</sub> mass by a multiplicative factor that varies exponentially with  $Z/Z_{\odot}$  (because of the form of  $f_{\rm CO}$  described above, see also Wolfire et al. 2010; Bolatto et al. 2013). As in Krumholz et al. (2011), the SSFR<sub>gas</sub>(H<sub>2</sub>, CO) curves are shown

<sup>&</sup>lt;sup>7</sup> Here we continue to use the 8.69 solar calibration of Asplund et al. (2009) rather than the value of  $12 + \log(O/H) = 8.79$  used by Krumholz et al. (2011).



**Fig. 15.** Ratio of SFR and CO-inferred H<sub>2</sub> mass vs. relative metallicity as described in the text, including our data and data from the literature (COLDGASS: Schruba et al. 2012; Bothwell et al. 2014). As in previous figures, the open triangles for Schruba et al. (2012) correspond to their compilation of data from the literature and the filled ones to their metal-poor dwarf observations. Following Krumholz et al. (2011), here  $M_{\rm H2}$  is inferred with a constant  $\alpha_{\rm CO} = 4.4 M_{\odot} (\text{K km s}^{-1} \text{ pc}^2)^{-1}$ , as described in the text. Also shown are the predictions by Krumholz et al. (2011), showing three values for total gas surface density,  $\Sigma_{\rm gas}$ :  $5 M_{\odot} \text{ pc}^{-2}$ ,  $20 M_{\odot} \text{ pc}^{-2}$ , and  $80 M_{\odot} \text{ pc}^{-2}$  indicated by dashed, dotted, and solid lines, respectively. Total gas corresponds to red, purple to H<sub>2</sub>, and blue to CO.

only for metallicities and total gas surface densities that are sufficiently high that the  $H_2$  and CO mass fractions are non-zero. Like the data compilation presented by Krumholz et al. (2011), our data are also qualitatively consistent with the trends predicted by the models.

Figure 15 shows that the data and the models agree extremely well for the prediction of CO luminosity; however at significantly sub-solar metallicities,  $Z/Z_{\odot} \leq 0.2$ , there are some discrepancies relative to the H<sub>2</sub> prediction, with some data exceeding by a factor of 3 or more the expected variation of the  $\alpha_{CO}$  factor. This is shown in Fig. 16 where we compare the power-law dependence of  $\alpha_{CO}$  derived previously with the exponential variation predicted by the models. Unlike in Fig. 15 where  $M_{H2}$ (CO) depended on a constant  $\alpha_{CO}$ , here we derive  $M_{H2}$  from a metallicity-dependent  $\alpha_{CO}$ ; the left panel shows the power-law dependency  $\alpha_{CO} \propto Z/Z_{\odot}^{-1.9}$  as shown in previous figures, and the right panel the exponential dependency found by Wolfire et al. (2010) and Krumholz et al. (2011),  $\alpha_{CO} \propto \exp(-Z/Z_{\odot})$ . As discussed above, such a dependence arises through the assumption that  $\alpha_{CO}$  depends exponentially on the visual extinction within the clouds,  $A_V$ , and that  $A_V$  is proportional to  $Z/Z_{\odot}$ .

Because both  $M_{\rm H2}$  and SFR predicted by the Krumholz et al. (2011) models depend on  $f_{\rm H_2}$ , the expectation is that SSFR(H<sub>2</sub>) should be constant, independently of metallicity. Figure 16 shows that the data do not quantitatively follow this trend, but rather show a residual increase of SSFR(H<sub>2</sub>) toward low metallicities. Neither formulation (power-law or exponential) of  $\alpha_{\rm CO}$ gives a horizontal trend of SSFR(H<sub>2</sub>) as a function of  $Z/Z_{\odot}$ . The exponential  $\alpha_{\rm CO}$  seems to better approximate the models around solar metallicity, but overestimates  $\alpha_{\rm CO}$  at  $Z/Z_{\odot} \sim 0.1$ . On the other hand, the power-law  $\alpha_{\rm CO}$  reproduces well the lowest metallicity SSFR(H<sub>2</sub>) (perhaps by construction, given that our  $\alpha_{\rm CO}$ was heavily influenced by the lowest metallicity galaxy in our sample). In any case, it seems that the short depletion times, corresponding to high SSFR(gas) are not a result of our approach, nor is our power-law formulation of  $\alpha_{CO}$  significantly inferior to the exponential one.

As shown in Fig. 16, galaxies with  $Z/Z_{\odot} \leq 0.2$  are not always consistent with the model predictions for SSFR(H<sub>2</sub>). Because of the large span of  $\tau_{dep}$  [inverse of SSFR(H<sub>2</sub>)] in our sample, it could be that the roughly constant timescale in the models is an oversimplification; indeed the Krumholz models are not designed for starbursts with potentially short depletion times and a high-density ISM. Moreover, there is strong evidence that clumping factors are different for molecular and atomic phases (Leroy et al. 2013b), implying that the use of a single correction factor to compare the predictions by Krumholz et al. (2011) with observations is almost certainly an oversimplification.

There could also be other problems with the applicability of the model predictions to these metal-poor starbursts, since  $\alpha_{CO}$ could have additional parameter dependencies beyond metallicity. The assumption of constant  $G'_0/n_{\rm H}$  may not be valid at low metallicities; the Krumholz et al. (2011) models assume  $G'_0 = 1$ , but the average UV flux  $G'_0$  at low metallicities is expected to be harder and, at a given total stellar mass, more intense because of the warmer stellar temperatures. There could also be some problem with the assumption that dust opacities and the capacity of the gas to self-shield from UV radiation vary linearly with metallicity. As mentioned in Sect. 2, even at a similar low metallicity, galaxies can have dramatically different dust-to-gas and dust-tostars mass ratios (Hunt et al. 2014). Schneider et al. (2015) suggest that this difference is due to the differences in the density of the cool gas in the ISM, because of the increased efficiency of grain formation in high-density environments. Thus, in addition to metallicity, ISM volume density could also be an important parameter for governing both the H1-to-H2 and the H2-to-CO transitions in galaxies. However, the incidence of high ISM densities at low metallicities is not yet well established. More observations of the molecular ISM are needed for metal-poor galaxies, especially at  $Z/Z_{\odot} \leq 0.1-0.2$  where the validity of CO as a tracer of H<sub>2</sub> begins to be called into question.

### 8. Summary and conclusions

We have detected <sup>12</sup>CO(1–0) with IRAM 30-m observations of a sample of 8 metal-poor dwarf galaxies, including a tentative detection at a metallicity of 12 + log(O/H) = 7.7, ~0.1 Z/Z<sub>☉</sub>. By calculating stellar masses and SFRs for our sample, and comparing our observations with a compilation of data from the literature, we have extended the trend for molecular  $\tau_{dep}$  with stellar specific SFR by Saintonge et al. (2011b) to lower metallicities, lower stellar masses, and higher sSFRs. In agreement with Saintonge et al. (2011b) and Huang & Kauffmann (2014), we conclude that  $\tau_{dep}$  is not constant in galaxies; in our combined sample  $\tau_{dep}$  varies by a factor of 200 or more (from  $\leq 50$  Myr to ~10 Gyr) over a range of 10<sup>3</sup> in stellar sSFR and  $M_{star}$ .

By exploiting the correlation of  $\tau_{dep}$  with sSFR, we were able to constrain  $\alpha_{CO}$  as a function of metallicity, and found a dependence of  $\alpha_{CO} \propto Z/Z_{\odot}^{-2}$ , similar to that found by work based on dust-continuum measurements compared with gas mass (Leroy et al. 2011; Bolatto et al. 2011). In conjunction with atomic gas measurements taken from the literature, we used our formulation for  $\alpha_{CO}$  to infer total gas masses, and total baryonic masses. The gas mass relative to the total baryonic mass depends on  $M_{star}$ , stellar sSFR, and O/H; we found significant correlations for all three parameters in the sense that gas-mass fractions are higher for galaxies of lower stellar mass, lower metallicity and higher



Fig. 16. Ratio of SFR and CO-inferred H<sub>2</sub> mass vs. relative metallicity as described in the text, including our data and data from the literature (COLDGASS: Huang & Kauffmann 2014; Schruba et al. 2012; Bothwell et al. 2014). As in previous figures, the open triangles for Schruba et al. (2012) correspond to their compilation of data from the literature and the filled ones to their metal-poor dwarf observations. In the left panel,  $M_{\rm H2}$ is inferred with our formulation of  $\alpha_{CO}$  as a power-law dependence on O/H as described in the text; in the *right panel*, we use the exponentiallydependent  $\alpha_{CO}$  as found by Wolfire et al. (2010), Krumholz et al. (2011). Also shown are the predictions by Krumholz et al. (2011), showing three values for total gas surface density,  $\Sigma_{gas}$ : 5  $M_{\odot}$  pc<sup>-2</sup>, 20  $M_{\odot}$  pc<sup>-2</sup>, and 80  $M_{\odot}$  pc<sup>-2</sup> indicated by dashed, dotted, and solid lines, respectively. Red color shows total gas, purple H<sub>2</sub>, and blue CO.

sSFR. However, despite large scatter, the HI SFR depletion time is roughly constant for our combined sample,  $\sim 3.4$  Gyr (see also Schiminovich et al. 2010).

We have examined our data in the context of the starformation models by Krumholz et al. (2011), and compared the power-law formulation of  $\alpha_{\rm CO}$  with the theoretically motivated exponential formulation. The predicted ratios of SFR and gas mass compare extremely well for the CO data, but the  $SSFR(H_2)$ inferred from either form of  $\alpha_{CO}$  are inconsistent for  $Z/Z_{\odot} \leq 0.2$ . A residual trend of  $SSFR(H_2)$  remains even when it should be invariant with metallicity according to the models. The cause of the disagreement is not clear but could arise from the starburst nature of our metal-poor sample; more observations of molecular tracers at low metallicity are needed to perform a more extended comparison with model predictions.

Acknowledgements. We are grateful to the IRAM staff, both in Granada and at Pico Veleta, for ably managing the logistics and the telescope/receiver operations. The referee's comments were timely and insightful, and improved the manuscript. We thank the International Space Science Institute (Bern) for hospitality during the conception of this paper, and Amelie Saintonge and Mei-Ling Huang who kindly gave us their COLDGASS data in digital form. L.K.H. acknowledges support from PRIN-INAF 2012/13. We made use of the NASA/IPAC Extragalactic Database (NED).

# References

- Albrecht, M., Chini, R., Krügel, E., Müller, S. A. H., & Lemke, R. 2004, A&A, 414, 141
- Andrews, B. H., & Martini, P. 2013, ApJ, 765, 140
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, ARA&A, 47, 481
- Baldwin, J. A., Spinrad, H., & Terlevich, R. 1982, MNRAS, 198, 535
- Barone, L. T., Heithausen, A., Hüttemeister, S., Fritz, T., & Klein, U. 2000, MNRAS, 317, 649
- Beck, S. C., Turner, J. L., Langland-Shula, L. E., et al. 2002, AJ, 124, 2516
- Bigiel, F., Leroy, A., Walter, F., et al. 2008, AJ, 136, 2846
- Bigiel, F., Leroy, A. K., Walter, F., et al. 2011, ApJ, 730, L13
- Bolatto, A. D., Jackson, J. M., & Ingalls, J. G. 1999, ApJ, 513, 275
- Bolatto, A. D., Leroy, A. K., Rosolowsky, E., Walter, F., & Blitz, L. 2008, ApJ, 686.948

Bolatto, A. D., Leroy, A. K., Jameson, K., et al. 2011, ApJ, 741, 12

- Bolatto, A. D., Wolfire, M., & Leroy, A. K. 2013, ARA&A, 51, 207
- Bordalo, V., Plana, H., & Telles, E. 2009, ApJ, 696, 1668
- Bothwell, M. S., Wagg, J., Cicone, C., et al. 2014, MNRAS, 445, 2599
- Brinchmann, J., Charlot, S., White, S. D. M., et al. 2004, MNRAS, 351, 1151 Brinks, E., & Taylor, C. L. 1997, IAU Symp., 170, 406
- Brown, M. J. I., Moustakas, J., Smith, J.-D. T., et al. 2014, ApJS, 212, 18
- Buyle, P., Michielsen, D., de Rijcke, S., Ott, J., & Dejonghe, H. 2006, MNRAS, 373, 793
- Cairós, L. M., Caon, N., García-Lorenzo, B., et al. 2007, ApJ, 669, 251
- Calzetti, D., Wu, S.-Y., Hong, S., et al. 2010, ApJ, 714, 1256
- Cormier, D., Madden, S. C., Lebouteiller, V., et al. 2014, A&A, 564, A121
- de Grijs, R., Smith, L. J., Bunker, A., et al. 2004, MNRAS, 352, 263
- Dopita, M. A., Pereira, M., Kewley, L. J., & Capaccioli, M. 2002, ApJS, 143, 47
- Ekta, B., Pustilnik, S. A., & Chengalur, J. N. 2009, MNRAS, 397, 963
- Elmegreen, B. G., Rubio, M., Hunter, D. A., et al. 2013, Nature, 495, 487
- Engelbracht, C. W., Gordon, K. D., Rieke, G. H., et al. 2005, ApJ, 628, L29
- Engelbracht, C. W., Rieke, G. H., Gordon, K. D., et al. 2008, ApJ, 678, 804
- Filho, M. E., Winkel, B., Sánchez Almeida, J., et al. 2013, A&A, 558, A18
- Focardi, P., & Kelm, B. 2002, A&A, 391, 35
- Fox, J. 1997, Applied regression analysis, linear models, and related models (Thou- sand Oaks, CA: Sage publications)
- Fumagalli, M., Krumholz, M. R., & Hunt, L. K. 2010, ApJ, 722, 919
- Galliano, F., Dwek, E., & Chanial, P. 2008, ApJ, 672, 214
- Gao, Y., & Solomon, P. M. 2004, ApJ, 606, 271
- García-Burillo, S., Usero, A., Alonso-Herrero, A., et al. 2012, A&A, 539, A8
- Genzel, R., Tacconi, L. J., Combes, F., et al. 2012, ApJ, 746, 69
- Genzel, R., Tacconi, L. J., Lutz, D., et al. 2015, ApJ, 800, 20
- Georgakakis, A., Tsamis, Y. G., James, B. L., & Aloisi, A. 2011, MNRAS, 413,
- Gil de Paz, A., Madore, B. F., & Pevunova, O. 2003, ApJS, 147, 29
- Glover, S. C. O., & Mac Low, M.-M. 2011, MNRAS, 412, 337
- Gondhalekar, P. M., Johansson, L. E. B., Brosch, N., Glass, I. S., & Brinks, E. 1998, A&A, 335, 152
- Graciá-Carpio, J., García-Burillo, S., Planesas, P., Fuente, A., & Usero, A. 2008, A&A, 479, 703
- Gratier, P., Braine, J., Rodriguez-Fernandez, N. J., et al. 2010, A&A, 522, A3
- Grossi, M., Hunt, L. K., Madden, S. C., et al. 2015, A&A, 574, A126
- Guseva, N. G., Izotov, Y. I., & Thuan, T. X. 2000, ApJ, 531, 776
- Hirashita, H., & Hunt, L. K. 2004, A&A, 421, 555
- Houck, J. R., Charmandaris, V., Brandl, B. R., et al. 2004, ApJS, 154, 211
- Huang, M.-L., & Kauffmann, G. 2014, MNRAS, 443, 1329
- Hughes, A., Meidt, S. E., Colombo, D., et al. 2013, ApJ, 779, 46
- Hunt, L. K., Vanzi, L., & Thuan, T. X. 2001, A&A, 377, 66
- Hunt, L. K., Giovanardi, C., & Helou, G. 2002, A&A, 394, 873

- Hunt, L. K., Dyer, K. K., Thuan, T. X., & Ulvestad, J. S. 2004, ApJ, 606, 853
- Hunt, L. K., Dyer, K. K., & Thuan, T. X. 2005, A&A, 436, 837
- Hunt, L. K., Thuan, T. X., Sauvage, M., & Izotov, Y. I. 2006, ApJ, 653, 222 Hunt, L. K., Thuan, T. X., Izotov, Y. I., & Sauvage, M. 2010, ApJ, 712, 164
- Hunt, L., Magrini, L., Galli, D., et al. 2012, MNRAS, 427, 906
- Hunt, L. K., Testi, L., Casasola, V., et al. 2014, A&A, 561, A49
- Hunter, D. A., & Thronson, H. A., Jr. 1995, ApJ, 452, 238
- Hunter, D. A., Gallagher, J. S., & Rautenkranz, D. 1982, ApJS, 49, 53
- Hunter, D. A., van Woerden, H., & Gallagher, J. S., III 1994a, ApJS, 91, 79
- Hunter, D. A., O'Connell, R. W., & Gallagher, J. S., III 1994b, AJ, 108, 84
- Hunter, D. A., Rubin, V. C., Swaters, R. A., Sparke, L. S., & Levine, S. E. 2002, ApJ, 580, 194
- Israel, F. P. 1997, A&A, 328, 471
- Izotov, Y. I., & Thuan, T. X. 2004, ApJ, 602, 200
- Izotov, Y. I., Chaffee, F. H., Foltz, C. B., et al. 2001, ApJ, 560, 222
- Izotov, Y. I., Thuan, T. X., & Guseva, N. G. 2005, ApJ, 632, 210
- Izotov, Y. I., Thuan, T. X., & Stasińska, G. 2007, ApJ, 662, 15
- James, B. L., Tsamis, Y. G., & Barlow, M. J. 2010, MNRAS, 401, 759
- James, B. L., Tsamis, Y. G., Barlow, M. J., Walsh, J. R., & Westmoquette, M. S. 2013, MNRAS, 428, 86
- Johnson, K. E., Indebetouw, R., Watson, C., & Kobulnicky, H. A. 2004, AJ, 128, 610
- Johnson, K. E., Hunt, L. K., & Reines, A. E. 2009, AJ, 137, 3788
- Joy, M., & Lester, D. F. 1988, ApJ, 331, 145
- Karachentsev, I., Musella, I., & Grimaldi, A. 1996, A&A, 310, 722
- Kauffmann, G., Heckman, T. M., White, S. D. M., et al. 2003, MNRAS, 341, 33

- Kennicutt, R. C., Jr. 1998, ARA&A, 36, 189 Kennicutt, R. C., Jr., Armus, L., Bendo, G., et al. 2003, PASP, 115, 928 Kennicutt, R. C., Jr., Hao, C.-N., Calzetti, D., et al. 2009, ApJ, 703, 1672
- Kewley, L. J., & Ellison, S. L. 2008, ApJ, 681, 1183
- Kepley, A. A., Reines, A. E., Johnson, K. E., & Walker, L. M. 2014, AJ, 147, 43
- Kroupa, P. 2001, MNRAS, 322, 231
- Krumholz, M. R. 2013, MNRAS, 436, 2747
- Krumholz, M. R., McKee, C. F., & Tumlinson, J. 2009, ApJ, 699, 850
- Krumholz, M. R., Leroy, A. K., & McKee, C. F. 2011, ApJ, 731, 25
- Kuno, N., Sato, N., Nakanishi, H., et al. 2007, PASJ, 59, 117
- Lebouteiller, V., Cormier, D., Madden, S. C., et al. 2012, A&A, 548, A91
- Lee, H., Skillman, E. D., Cannon, J. M., et al. 2006, ApJ, 647, 970
- Lee, J. C., Gil de Paz, A., Tremonti, C., et al. 2009, ApJ, 706, 599
- Lelli, F., Verheijen, M., Fraternali, F., & Sancisi, R. 2012, A&A, 537, A72
- Lelli, F., Verheijen, M., & Fraternali, F. 2014, MNRAS, 445, 1694
- Leroy, A., Bolatto, A. D., Simon, J. D., & Blitz, L. 2005, ApJ, 625, 763
- Leroy, A., Cannon, J., Walter, F., Bolatto, A., & Weiss, A. 2007, ApJ, 663, 990
- Leroy, A. K., Walter, F., Brinks, E., et al. 2008, AJ, 136, 2782
- Leroy, A. K., Walter, F., Bigiel, F., et al. 2009a, AJ, 137, 4670
- Leroy, A. K., Bolatto, A., Bot, C., et al. 2009b, ApJ, 702, 352
- Leroy, A. K., Bolatto, A., Gordon, K., et al. 2011, ApJ, 737, 12
- Leroy, A. K., Walter, F., Sandstrom, K., et al. 2013a, AJ, 146, 19
- Leroy, A. K., Lee, C., Schruba, A., et al. 2013b, ApJ, 769, L12
- Li, G. K. 1985, Robust regression. In Exploring Data Tables, Trends, and Shapes, eds. D. C. Hoaglin, F. Mosteller, & J. W. Tukey (Wiley)
- Lipovetsky, V. A., Chaffee, F. H., Izotov, Y. I., et al. 1999, ApJ, 519, 177
- Madden, S. C., Galliano, F., Jones, A. P., & Sauvage, M. 2006, A&A, 446, 877
- Madden, S. C., Rémy-Ruyer, A., Galametz, M., et al. 2013, PASP, 125, 600
- Maloney, P., & Black, J. H. 1988, ApJ, 325, 389
- McGaugh, S. S., & Schombert, J. M. 2014, AJ, 148, 77
- McKee, C. F., & Krumholz, M. R. 2010, ApJ, 709, 308
- Micheva, G., Östlin, G., Bergvall, N., et al. 2013, MNRAS, 431, 102
- Minchin, R. F., Momjian, E., Auld, R., et al. 2010, AJ, 140, 1093 Moffett, A. J., Kannappan, S. J., Baker, A. J., & Laine, S. 2012, ApJ, 745, 34
- Moll, S. L., Mengel, S., de Grijs, R., Smith, L. J., & Crowther, P. A. 2007, MNRAS, 382, 1877
- Narayanan, D., Krumholz, M. R., Ostriker, E. C., & Hernquist, L. 2012, MNRAS, 421, 3127
- Norris, M. A., Meidt, S., Van de Ven, G., et al. 2014, ApJ, 797, 55

- Osterbrock, D. E., & Ferland, G. J. 2006, Astrophysics of gaseous nebulae and active galactic nuclei, 2nd, eds. D. E. Osterbrock, & G. J. Ferland (Sausalito, CA: University Science Books)
- Paradis, D., Dobashi, K., Shimoikura, T., et al. 2012, A&A, 543, A103
- Paturel, G., Theureau, G., Bottinelli, L., et al. 2003, A&A, 412, 57
- Pettini, M., & Pagel, B. E. J. 2004, MNRAS, 348, L59
- Pineda, J. L., Velusamy, T., Langer, W. D., et al. 2010, A&A, 521, L19
- Pineda, J. L., Langer, W. D., & Goldsmith, P. F. 2014, A&A, 570, A121
- Planck Collaboration XIX. 2011, A&A, 536, A19
- Plante, S., & Sauvage, M. 2002, AJ, 124, 1995
- Pustilnik, S. A., Brinks, E., Thuan, T. X., Lipovetsky, V. A., & Izotov, Y. I. 2001, AJ, 121, 1413
- Regan, M. W., Thornley, M. D., Helfer, T. T., et al. 2001, ApJ, 561, 218
- Reines, A. E., Johnson, K. E., & Hunt, L. K. 2008, AJ, 136, 1415
- Röllig, M., Ossenkopf, V., Jeyakumar, S., Stutzki, J., & Sternberg, A. 2006, A&A, 451, 917
- Rosolowsky, E., Engargiola, G., Plambeck, R., & Blitz, L. 2003, ApJ, 599, 258 Rubio, M., Lequeux, J., & Boulanger, F. 1993, A&A, 271, 9
- Sage, L. J., Salzer, J. J., Loose, H.-H., & Henkel, C. 1992, A&A, 265, 19
- Saintonge, A., Kauffmann, G., Kramer, C., et al. 2011a, MNRAS, 415, 32
- Saintonge, A., Kauffmann, G., Wang, J., et al. 2011b, MNRAS, 415, 61
- Salim, S., Rich, R. M., Charlot, S., et al. 2007, ApJS, 173, 267
- Salim, S., Lee, J. C., Ly, C., et al. 2014, ApJ, 797, 126
- Sandstrom, K. M., Leroy, A. K., Walter, F., et al. 2013, ApJ, 777, 5
- Sargent, W. L. W., & Searle, L. 1970, ApJ, 162, L155
- Schaerer, D., Contini, T., & Pindao, M. 1999, A&AS, 136, 35
- Schiminovich, D., Catinella, B., Kauffmann, G., et al. 2010, MNRAS, 408, 919 Schneider, R., Hunt, L. K., & Valiante, R. 2015, MNRAS, submitted
- Schruba, A., Leroy, A. K., Walter, F., et al. 2012, AJ, 143, 138
- Skibba, R. A., Engelbracht, C. W., Dale, D., et al. 2011, ApJ, 738, 89
- Smith, B. J., & Hancock, M. 2009, AJ, 138, 130
- Solomon, P. M., & Vanden Bout, P. A. 2005, ARA&A, 43, 677
- Steel, S. J., Smith, N., Metcalfe, L., Rabbette, M., & McBreen, B. 1996, A&A, 311, 721
- Swaters, R. A., van Albada, T. S., van der Hulst, J. M., & Sancisi, R. 2002, A&A, 390, 829
- Taylor, C. L., Brinks, E., Grashuis, R. M., & Skillman, E. D. 1995, ApJS, 99, 427
- Taylor, C. L., Kobulnicky, H. A., & Skillman, E. D. 1998, AJ, 116, 2746
- Temi, P., Brighenti, F., & Mathews, W. G. 2009, ApJ, 707, 890
- Thuan, T. X., & Izotov, Y. I. 2005, ApJS, 161, 240
- Thuan, T. X., Izotov, Y. I., & Lipovetsky, V. A. 1996, ApJ, 463, 120
- Thuan, T. X., Lipovetsky, V. A., Martin, J.-M., & Pustilnik, S. A. 1999, A&AS, 139

Vader, J. P., Frogel, J. A., Terndrup, D. M., & Heisler, C. A. 1993, AJ, 106, 1743

Vanzi, L., Rieke, G. H., Martin, C. L., & Shields, J. C. 1996, ApJ, 466, 150

Vanzi, L., Cresci, G., Sauvage, M., & Thompson, R. 2011, A&A, 534, A70

Westmoquette, M. S., Gallagher, J. S., & de Poitiers, L. 2010, MNRAS, 403,

Zhang, W., Kong, X., Li, C., Zhou, H.-Y., & Cheng, F.-Z. 2007, ApJ, 655, 851

A114, page 17 of 17

Vanzi, L., Hunt, L. K., Thuan, T. X., & Izotov, Y. I. 2000, A&A, 363, 493

Vanzi, L., Cresci, G., Telles, E., & Melnick, J. 2008, A&A, 486, 393

Walter, F., Brinks, E., de Blok, W. J. G., et al. 2008, AJ, 136, 2563

Wen, X.-Q., Wu, H., Zhu, Y.-N., et al. 2013, MNRAS, 433, 2946

Zibetti, S., Charlot, S., & Rix, H.-W. 2009, MNRAS, 400, 1181

Wolfire, M. G., Hollenbach, D., & McKee, C. F. 2010, ApJ, 716, 1191 Wu, Y., Charmandaris, V., Hao, L., Brandl, B. R., et al. 2006, ApJ, 639, 157

Thuan, T. X., Hunt, L. K., & Izotov, Y. I. 2008, ApJ, 689, 897

van Zee, L., Skillman, E. D., & Salzer, J. J. 1998, AJ, 116, 1186

Vanzi, L., Hunt, L. K., & Thuan, T. X. 2002, A&A, 390, 481

Vigroux, L., Stasińska, G., & Comte, G. 1987, A&A, 172, 15

van Zee, L. 2000, AJ, 119, 2757

van Zee, L., & Haynes, M. P. 2006, ApJ, 636, 214

van Zee, L. 2001, AJ, 121, 2003

Vanzi, L. 2003, A&A, 408, 523

Wilson, C. D. 1995, ApJ, 448, L97

1719