



<b>Publication Year</b>	2015
<b>Acceptance in OA @INAF</b>	2020-05-08T10:30:03Z
<b>Title</b>	New Near-infrared Period-Luminosity-Metallicity Relations for RR Lyrae Stars and the Outlook for Gaia
<b>Authors</b>	Muraveva, Tatiana; Palmer, M.; CLEMENTINI, Gisella; Luri, X.; Cioni, M. -R. L.; et al.
<b>DOI</b>	10.1088/0004-637X/807/2/127
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/24632">http://hdl.handle.net/20.500.12386/24632</a>
<b>Journal</b>	THE ASTROPHYSICAL JOURNAL
<b>Number</b>	807

## NEW NEAR-INFRARED PERIOD–LUMINOSITY–METALLICITY RELATIONS FOR RR LYRAE STARS AND THE OUTLOOK FOR GAIA\*

T. MURAVEVA<sup>1</sup>, M. PALMER<sup>2</sup>, G. CLEMENTINI<sup>1</sup>, X. LURI<sup>2</sup>, M.-R. L. CIONI<sup>3,4,5</sup>, M. I. MORETTI<sup>1,6</sup>, M. MARCONI<sup>7</sup>, V. RIPEPI<sup>7</sup>, AND  
S. RUBELE<sup>8</sup>

<sup>1</sup> INAF-Osservatorio Astronomico di Bologna, via Ranzani 1, I-40127, Bologna, Italy; [tatiana.muraveva@oabo.inaf.it](mailto:tatiana.muraveva@oabo.inaf.it)

<sup>2</sup> Dept. d’Astronomia i Meteorologia, Institut de Ciències del Cosmos, Universitat de Barcelona (IEEC-UB), Martí Franquès 1, E-08028 Barcelona, Spain

<sup>3</sup> Universität Potsdam, Institut für Physik und Astronomie, Karl-Liebknecht-Str. 24/25, D-14476 Potsdam, Germany

<sup>4</sup> Leibniz-Institut für Astrophysik Potsdam, An der Sternwarte 16, D-14482 Potsdam, Germany

<sup>5</sup> University of Hertfordshire, Physics Astronomy and Mathematics, College Lane, Hatfield AL10 9AB, UK

<sup>6</sup> Scuola Normale Superiore, Piazza dei Cavalieri 7, I-56126 Pisa, Italy

<sup>7</sup> INAF-Osservatorio Astronomico di Capodimonte, Salita Moiarriello 16, I-80131, Napoli, Italy

<sup>8</sup> INAF-Osservatorio Astronomico di Padova, Vicolo dell’Osservatorio 5, I-35122, Padova, Italy

*Received 2015 April 7; accepted 2015 May 21; published 2015 July 7*

### ABSTRACT

We present results of the analysis of 70 RR Lyrae stars located in the bar of the Large Magellanic Cloud (LMC). Combining the spectroscopically determined metallicity of these stars from the literature with precise periods from the OGLE III catalog and multi-epoch  $K_s$  photometry from the VISTA survey of the Magellanic Clouds system, we derive a new near-infrared period–luminosity–metallicity ( $PL_{K_s}Z$ ) relation for RR Lyrae variables. In order to fit the relation we use a fitting method developed specifically for this study. The zero-point of the relation is estimated two different ways: by assuming the value of the distance to the LMC and by using *Hubble Space Telescope* parallaxes of five RR Lyrae stars in the Milky Way (MW). The difference in distance moduli derived by applying these two approaches is  $\sim 0.2$  mag. To investigate this point further we derive the  $PL_{K_s}Z$  relation based on 23 MW RR Lyrae stars that had been analyzed in Baade–Wesselink studies. We compared the derived  $PL_{K_s}Z$  relations for RR Lyrae stars in the MW and LMC. Slopes and zero-points are different, but still consistent within the errors. The shallow slope of the metallicity term is confirmed by both LMC and MW variables. The astrometric space mission *Gaia* is expected to provide a huge contribution to the determination of the RR Lyrae  $PL_{K_s}Z$  relation; however, calculating an absolute magnitude from the trigonometric parallax of each star and fitting a  $PL_{K_s}Z$  relation directly to period and absolute magnitude leads to biased results. We present a tool to achieve an unbiased solution by modeling the data and inferring the slope and zero-point of the relation via statistical methods.

*Key words:* astrometry – distance scale – Magellanic Clouds – methods: data analysis – stars: statistics – stars: variables: RR Lyrae

### 1. INTRODUCTION

RR Lyrae stars are radially pulsating variables connected with low-mass helium-burning stars on the horizontal branch (HB) of the color–magnitude diagram. These objects are Population II stars, which are abundant in globular clusters and in the halos of galaxies. RR Lyrae stars are a perfect tool for studying the age, formation, and structure of their parent stellar system. Moreover, they are widely used for the determination of distances in the Milky Way (MW) and to Local Group galaxies.

RR Lyrae stars are primary distance indicators because of the existence of a narrow luminosity–metallicity ( $M_V - [\text{Fe}/\text{H}]$ ) relation in the visual band and period–luminosity–metallicity ( $PLZ$ ) relations in the infrared passbands. The near-infrared  $PL_{K_s}Z$  relation of RR Lyrae stars was originally discovered by Longmore et al. (1986) and later was the subject of study by many different authors (e.g., Bono et al. 2003, Catelan et al. 2004, Del Principe 2006, Sollima et al. 2006, Sollima et al. 2008, Borissova et al. 2009, Coppola et al. 2011, Ripepi et al. 2012a). The near-infrared  $PL_{K_s}Z$  relation has many advantages in comparison with the visual  $M_V - [\text{Fe}/\text{H}]$  relation. First of all, the luminosity of RR Lyrae stars in the

$K_s$  passband is less dependent on metallicity and interstellar extinction ( $A_{K_s} \sim 0.1A_V$ ). Furthermore, light curves of RR Lyrae stars in the  $K_s$  band have smaller amplitudes and are more symmetrical than optical light curves, making the determination of the mean  $K_s$  magnitudes easier and more precise.

In order to calibrate the  $PL_{K_s}Z$  relation a large sample of RR Lyrae stars is required, spanning a wide range of metallicities, for which accurate mean  $K_s$  and  $[\text{Fe}/\text{H}]$  measurements are available. We have selected 70 RR Lyrae variables in the Large Magellanic Cloud (LMC) with spectroscopically determined metallicities in the range of  $-2.06 < [\text{Fe}/\text{H}] < -0.63$  dex (Gratton et al. 2004). All of them have counterparts in the OGLE III catalog (Soszyński et al. 2009), therefore very precise periods are available. In order to increase the accuracy of the determination of mean  $K_s$  magnitudes, multi-epoch photometry is needed. For this reason we are using data from the near-infrared VISTA survey of the Magellanic Clouds System (VMC, Cioni et al. 2011), which is performing  $K_s$ -band observations of the whole Magellanic System in 12 (or more) epochs, while in many previous studies only single-epoch photometry from the 2MASS (Cutri et al. 2003) was used. To fit the  $PL_{K_s}Z$  relation we apply a fitting approach developed for the current study. This method takes into account errors in two dimensions, the intrinsic dispersion of the data and the possibility of inaccuracy in the formal error estimates.

\* Based on observations made with VISTA at ESO under programme ID 179, B-2003.

One main issue in the determination of distances with the RR Lyrae  $PL_{K_s}Z$  relation is the calibration of the zero-point. Trigonometric parallaxes remain the only direct method of determining distances to astronomical sources, free of any assumptions (such as, for instance, the distance to the LMC, etc.) and hence calibrating the  $PL_{K_s}Z$  zero-point. However, reasonably well estimated parallaxes exist, so far, only for five RR Lyrae variables in the MW observed by Benedict et al. (2011) with the *Hubble Space Telescope* Fine Guidance Sensor (*HST*/FGS). In this study we use both a global estimate of the LMC distance and the *HST* parallaxes in order to calibrate the zero-point of our  $PL_{K_s}Z$  relation based on LMC RR Lyrae stars. Furthermore, to check whether the RR Lyrae  $PL_{K_s}Z$  relation is universal and could thus be applied to measure distances in the MW and to other galaxies, we analyze a sample of 23 MW RR Lyrae stars, for which absolute magnitudes in the  $K$  and  $V$  passbands are available from the Baade–Wesselink (B–W) studies (e.g., Fernley et al. 1998b, and references therein). Based on these absolute magnitudes and applying our fitting approach we fit the RR Lyrae  $PL_{K_s}Z$  relation. Then we compare the  $PL_{K_s}Z$  relations derived for RR Lyrae stars in the MW and in the LMC.

*Gaia*, the ESA cornerstone mission launched in 2013 December, is expected to provide a great contribution to the determination of the RR Lyrae  $PL_{K_s}Z$  relation and to the definition of its zero-point in particular. The satellite is designed to produce the most precise three-dimensional map of the MW to date (Perryman et al. 2001) by measuring parallaxes of over one billion stars during its five-year mission, among which are thousands of RR Lyrae variables. In the current study we present a method that avoids the problems of the nonlinear transformation of trigonometric parallaxes (and negative parallaxes) to absolute magnitudes, and apply this method to fit the  $PL_{K_s}Z$  relation of the 23 MW RR Lyrae stars, based on simulated *Gaia* parallaxes.

In Section 2 we provide information about the 70 RR Lyrae stars in the LMC that form the basis of the present study. In Section 3 we present our method and results of fitting the RR Lyrae  $PL_{K_s}Z$  relation in the LMC and in the MW. In Section 4 we present the method to fit the  $PL_{K_s}Z$  relation with simulated *Gaia* parallaxes and apply this method to the 23 MW RR Lyrae stars analyzed in Section 3. Section 5 provides a summary of the results. In the appendices we present a detailed description of the fitting method that was developed for this study (Appendix A) and a compilation of metal abundances for the MW RR Lyrae stars (Appendix B).

## 2. DATA

Optical photometry for the LMC RR Lyrae stars discussed in this paper was obtained by Clementini et al. (2003) and Di Fabrizio et al. (2005) using the Danish 1.54 meter telescope in La Silla, Chile. Two different sky positions, hereafter called fields A and B, were observed. Both are located close to the bar of the LMC (Clementini et al. 2003, Di Fabrizio et al. 2005). As a result, accurate  $B$ ,  $V$  and  $I$  light curves tied to the Johnson-Cousins standard system and pulsation characteristics (period, epoch of maximum light, amplitudes and mean magnitudes) for 125 RR Lyrae stars were obtained (Di Fabrizio et al. 2005). Low-resolution spectra for 98 of these variables were collected by Gratton et al. (2004) using the FOcal Reducer/low dispersion Spectrograph instrument mounted at the ESO


VLT. They were used to derive metal abundances for individual stars by comparing the strength of the Ca II K line with that of the H lines (Preston 1959). For the calibration of the method, four clusters with metallicity in the range  $[-2.06; -1.26]$  dex were used. According to Gratton et al. (2004), the obtained metallicities are tied to a scale, which is, on average, 0.06 dex more metal-rich than the Zinn & West (1984) metallicity scale.

We cross-matched the sample of 98 RR Lyrae variables with known metallicities against the catalog of RR Lyrae stars observed by the OGLE III survey (Soszyński et al. 2009). The OGLE III catalog contains information about the position, photometric and pulsation properties of 24906 RR Lyrae stars in the LMC. We found that, respectively, 94, 2, and 2 objects are cross-identified with sources in the OGLE III catalog within a pairing radius of  $1''$ ,  $3''$ , and  $7''$ . The two stars with a counterpart at more than  $5''$  are OGLE-LMC-RRLYR-10345 and OGLE-LMC-RRLYR-10509; for these two objects we checked both the OGLE III finding charts and Gratton et al.'s (2004) Figure 5 (field B1) in order to understand if they are affected by any problem. Star OGLE-LMC-RRLYR-10345 is an isolated slightly elongated star without any clear blending problem, while star OGLE-LMC-RRLYR-10509 is very close to another source possibly making it more difficult to locate accurately the star center. Considering that Gratton et al. (2004) and OGLE III periods for these two stars agree within 0.5%, we kept these stars in our sample.

We compared the periods of the 98 RR Lyrae stars provided by Di Fabrizio et al. (2005) and those in the OGLE III catalog (Soszyński et al. 2009). For 96 objects the periods agree within  $\sim 2\%$ , while for two objects the periods differ significantly. For star A6332 the difference is of  $\sim 25\%$  and for star A5148 it is of  $\sim 37\%$  (star identifications are from Di Fabrizio et al. 2005). Moreover, star A5148 has been classified as a first-overtone RR Lyrae star (RRc) in the OGLE III catalog, and as a fundamental-mode RR Lyrae (RRab) by Di Fabrizio et al. (2005). Since accurately estimated periods and classifications play a key role in the current study, we discarded these two objects from the following analysis.

Seven objects (B2811, B4008, B3625, B2517, A2623, A2119, A10360) from the sample are classified as RRc by Di Fabrizio et al. (2005) and as second-overtone RR Lyrae stars (RRe) in the OGLE III catalog. We removed them from our analysis because of the uncertain classification. Furthermore, since one of the main purposes of the current research is to study the  $PL_{K_s}Z$  relation of RR Lyrae stars of ab- and c-types we discarded seven objects, which were classified as double-mode RR Lyrae stars (RRd) by Di Fabrizio et al. (2005): A7137, A8654, A3155, A4420, B7467, B6470, and B3347. This left us with a final sample of 61 RRab and 21 RRc stars, which all have a counterpart in the OGLE III catalog. The period search for the RR Lyrae stars in the OGLE III catalog was performed using an algorithm based on the Fourier analysis of the light curves (Soszyński et al. 2009). The uncertainties in the OGLE III periods for the 82 RR Lyrae stars in our sample are declared to be less than  $5 \times 10^{-6}$  days. Therefore we used the periods provided by the OGLE III catalog in order to fit the  $PL_{K_s}Z$  relation for our sample, and did not consider errors in the periods since they are negligible in comparison to the other uncertainties.

In order to derive mean  $K_s$  magnitudes for the RR Lyrae stars in our sample we used data from the VMC survey (Cioni



**Figure 1.** Examples of  $K_s$ -band light curves for RR Lyrae stars in our sample. Identification numbers are from Di Fabrizio et al. (2005), periods are from the OGLE III catalog (Soszyński et al. 2009) and are given in days.

et al. 2011). Started in 2009, the VMC survey covers a total area of  $116 \text{ deg}^2$  in the LMC with 68 contiguous tiles. The survey is obtaining  $YJK_s$  photometry. The  $K_s$ -band photometry is taken in time-series mode over 12 (or more) separate epochs and each single epoch reaches a limiting  $K_s$  magnitude  $\sim 19.3$  mag with a signal-to-noise ratio (S/N)  $\sim 5$  (see Figure 1 of Moretti et al. 2014). On the bright side, VMC is limited by saturation at  $K_s \sim 11.4$  mag. The majority of RR Lyrae stars in our sample are located within the VMC tile LMC 5\_5. Observations of the tile LMC 5\_5 were performed in 15 epochs taken in the period from 2010 October 30, to 2012 January 11. For two epochs of observation the ellipticity was too high, so these data were not considered in the analysis. Among the remaining 13 epochs there are 11 deep and 2 shallow epochs. Since shallow observations were obtained in good seeing conditions their S/Ns were enough to detect the RR Lyrae stars. In the following analysis we used all 13 available epochs to fit the light curves of the RR Lyrae stars. PSF photometry of the time-series data for this tile was performed on the homogenized epoch-tile images (Rubele et al. 2012) using the IRAF Daophot packages (Stetson et al. 1990). On each epoch-tile image the PSF model was

created using 2500 stars uniformly distributed; finally the Daophot ALLSTAR routine was used to perform the PSF photometry on all epoch images and time-series catalogs were correlated within a tolerance of one arcsec.

We have cross-matched our sample of 82 RR Lyrae stars against the PSF photometry catalog of the VMC tile LMC 5\_5. VMC counterparts for 71 objects were found within a pairing radius of  $1''$ . Among them, 70 sources have 13 epochs in the  $K_s$ -band, while for one object (B4749) we have observations only in 6 epochs. Six data points are not enough for a reliable fit of the light curve and, consequently, for the robust determination of the mean  $K_s$  magnitude, hence, we discarded this source from the following analysis and proceeded with the 70 RR Lyrae stars, for which 13 epochs in the  $K_s$ -band exist. We derived the mean  $K_s$  magnitudes of these 70 RR Lyrae stars by Fourier fitting the light curves with the GRaphical Analyzer of Time Series package (GRATIS, custom software developed at the Observatory of Bologna by P. Montegriffo; see, e.g., Clementini et al. 2000). To fit the light curves we discarded obvious outliers. Nevertheless, after the  $\sigma$ -clipping procedure, each source still had 11 or more data points. Examples of the  $K_s$  light curves are shown in Figure 1.

**Table 1**  
Properties of the 70 RR Lyrae Stars in the Bar of the LMC Analyzed in This Paper

Star	OGLE ID	R.A. (J2000)	Decl. (J2000)	Type	[Fe/H] (dex)	$\sigma_{[\text{Fe}/\text{H}]}$ (dex)	$P$ (days)	$\langle K_{s,0} \rangle$ (mag)	$\sigma_{(K_{s,0})}$ (mag)
A28665	OGLE-LMC-RRLYR-12944	5:22:06.55	-70:27:55.6	RRc	-0.63	0.24	0.3008299	18.450	0.046
A7864	OGLE-LMC-RRLYR-13857	5:23:39.25	-70:31:38.1	RRc	-1.36	0.22	0.3129458	18.550	0.055
B4946	OGLE-LMC-RRLYR-10621	5:18:11.08	-70:59:35.6	RRc	-1.11	0.25	0.3130142	18.394	0.054
A2636	OGLE-LMC-RRLYR-13548	5:23:09.09	-70:39:08.1	RRc	-1.61	0.29	0.3154437	18.562	0.045
A8837	OGLE-LMC-RRLYR-13326	5:22:45.70	-70:30:14.3	RRc	-1.52	0.22	0.3165579	18.660	0.089
A8622	OGLE-LMC-RRLYR-13164	5:22:28.93	-70:30:35.9	RRc	-1.44	0.28	0.3212334	18.426	0.032
A7231	OGLE-LMC-RRLYR-13680	5:23:22.42	-70:32:35.4	RRc	-1.46	0.26	0.3228047	18.236	0.051
A2234	OGLE-LMC-RRLYR-13479	5:23:01.47	-70:39:44.4	RRc	-1.53	0.18	0.3228060	18.292	0.044
A4388	OGLE-LMC-RRLYR-12614	5:21:31.67	-70:36:46.3	RRc	-1.33	0.27	0.3417737	18.411	0.048
A10113	OGLE-LMC-RRLYR-14046	5:24:00.38	-70:28:06.1	RRc	-1.52	0.25	0.3506618	18.197	0.036
B6255	OGLE-LMC-RRLYR-10111	5:17:17.88	-70:57:26.4	RRc	-1.52	0.16	0.3535596	18.305	0.038
B4179	OGLE-LMC-RRLYR-10142	5:17:19.95	-71:01:02.1	RRc	-1.53	0.27	0.3545232	18.150	0.034
A8812	OGLE-LMC-RRLYR-13150	5:22:26.44	-70:30:19.1	RRc	-1.23	0.24	0.3549660	18.281	0.036
B6715	OGLE-LMC-RRLYR-12593	5:21:29.33	-70:29:23.4	RRc	-1.39	0.18	0.3569006	18.308	0.032
A2024	OGLE-LMC-RRLYR-13572	5:23:11.02	-70:40:03.3	RRc	-1.62	0.26	0.3590534	18.246	0.043
B6164	OGLE-LMC-RRLYR-10612	5:18:10.17	-70:57:30.7	RRc	-1.88	0.22	0.3744821	18.039	0.045
A27697	OGLE-LMC-RRLYR-13012	5:22:14.03	-70:28:35.0	RRc	-1.33	0.25	0.3825700	18.030	0.023
A19450	OGLE-LMC-RRLYR-13841	5:23:37.95	-70:34:06.7	RRab	-0.76	0.13	0.3979182	18.481	0.071
B7064	OGLE-LMC-RRLYR-10708	5:18:18.63	-70:55:58.7	RRc	-2.03	0.20	0.4004744	18.029	0.043
B6957	OGLE-LMC-RRLYR-10702	5:18:18.08	-70:56:08.7	RRc	-1.48	0.18	0.4047399	18.027	0.043
B23502	OGLE-LMC-RRLYR-10509	5:18:00.25	-70:54:31.0	RRab	-1.55	0.14	0.4724681	18.243	0.076
A3061	OGLE-LMC-RRLYR-13704	5:23:25.18	-70:38:28.9	RRab	-1.26	0.12	0.4744410	18.415	0.041
B10811	OGLE-LMC-RRLYR-10684	5:18:16.01	-71:04:27.0	RRab	-1.42	0.20	0.4760753	18.197	0.036
B3400	OGLE-LMC-RRLYR-10072	5:17:14.51	-71:02:26.6	RRab	-1.45	0.24	0.4852148	18.346	0.092
A7325	OGLE-LMC-RRLYR-13855	5:23:39.13	-70:32:24.8	RRab	-1.18	0.26	0.4864544	18.223	0.046
B3033	OGLE-LMC-RRLYR-10659	5:18:14.04	-71:03:00.5	RRab	-1.26	0.21	0.4986975	18.130	0.066
B2055	OGLE-LMC-RRLYR-10108	5:17:17.44	-71:04:50.2	RRab	-1.70	0.23	0.5207746	18.254	0.074
A26525	OGLE-LMC-RRLYR-12811	5:21:52.50	-70:29:28.7	RRab	-1.41	0.22	0.5225029	18.168	0.053
A7211	OGLE-LMC-RRLYR-13092	5:22:21.17	-70:32:43.9	RRab	-1.33	0.19	0.5226857	18.193	0.041
A2767	OGLE-LMC-RRLYR-13634	5:23:17.75	-70:38:55.9	RRab	-1.37	0.08	0.5325871	18.054	0.036
B24089	OGLE-LMC-RRLYR-10345	5:17:43.51	-70:54:02.7	RRab	-1.48	0.16	0.5580613	18.094	0.069
A8788	OGLE-LMC-RRLYR-13678	5:23:22.41	-70:30:14.6	RRab	-1.61	0.21	0.5591710	18.197	0.036
A6398	OGLE-LMC-RRLYR-13294	5:22:40.76	-70:33:50.2	RRab	-1.40	0.30	0.5619466	17.957	0.027
A7247	OGLE-LMC-RRLYR-13708	5:23:25.58	-70:32:33.4	RRab	-1.38	0.21	0.5621512	18.045	0.049
A25301	OGLE-LMC-RRLYR-12638	5:21:34.00	-70:30:24.5	RRab	-1.58	0.27	0.5631146	18.268	0.051
A15387	OGLE-LMC-RRLYR-12603	5:21:30.43	-70:37:11.3	RRab	-1.81	0.12	0.5635914	18.049	0.052
B22917	OGLE-LMC-RRLYR-10713	5:18:19.10	-70:54:56.1	RRab	-1.29	0.16	0.5646803	18.179	0.054
A9245	OGLE-LMC-RRLYR-13536	5:23:07.67	-70:29:36.5	RRab	-1.27	0.18	0.5678763	18.013	0.035
A12896	OGLE-LMC-RRLYR-13330	5:22:46.15	-70:38:54.9	RRab	-1.53	0.10	0.5719281	18.143	0.035
A7609	OGLE-LMC-RRLYR-13941	5:23:48.39	-70:32:00.3	RRab	-1.63	0.11	0.5724984	18.023	0.046
B7442	OGLE-LMC-RRLYR-10082	5:17:15.73	-70:55:26.8	RRab	-1.58	0.11	0.5740274	18.096	0.048
A25362	OGLE-LMC-RRLYR-13848	5:23:38.53	-70:30:08.5	RRab	-1.39	0.15	0.5787944	18.033	0.052
B1907	OGLE-LMC-RRLYR-10638	5:18:12.36	-71:04:59.5	RRab	-1.70	0.26	0.5818283	17.915	0.036
A4974	OGLE-LMC-RRLYR-13372	5:22:51.26	-70:35:47.7	RRab	-1.36	0.10	0.5820430	17.992	0.054
B6798	OGLE-LMC-RRLYR-10044	5:17:11.37	-70:56:32.6	RRab	-1.40	0.23	0.5822610	17.910	0.114
B14449	OGLE-LMC-RRLYR-09999	5:17:05.37	-71:01:40.9	RRab	-1.70	0.13	0.5822854	18.118	0.071
A9494	OGLE-LMC-RRLYR-13354	5:22:49.26	-70:29:13.5	RRab	-1.69	0.28	0.5844615	17.874	0.035
A18314	OGLE-LMC-RRLYR-13353	5:22:49.13	-70:34:59.2	RRab	-1.42	0.18	0.5875708	18.093	0.030
A10487	OGLE-LMC-RRLYR-13126	5:22:24.61	-70:27:40.6	RRab	-1.49	0.11	0.5909585	18.030	0.016
A10214	OGLE-LMC-RRLYR-12609	5:21:31.14	-70:28:12.0	RRab	-1.48	0.12	0.5918196	17.904	0.065
A28066	OGLE-LMC-RRLYR-13765	5:23:30.10	-70:28:11.0	RRab	-1.44	0.17	0.5959296	18.007	0.060
A26821	OGLE-LMC-RRLYR-12831	5:21:53.95	-70:29:17.5	RRab	-1.37	0.13	0.5969220	18.130	0.047
B2249	OGLE-LMC-RRLYR-10061	5:17:13.06	-71:04:27.1	RRab	-1.56	0.15	0.6030630	17.999	0.050
A16249	OGLE-LMC-RRLYR-12960	5:22:08.27	-70:36:31.0	RRab	-1.87	0.12	0.6067385	18.060	0.045
A4933	OGLE-LMC-RRLYR-13175	5:22:30.05	-70:35:53.7	RRab	-1.48	0.12	0.6134920	17.768	0.027
A7734	OGLE-LMC-RRLYR-12956	5:22:07.86	-70:31:59.8	RRab	-1.40	0.15	0.6149615	17.888	0.036
A2525	OGLE-LMC-RRLYR-13788	5:23:32.45	-70:39:15.3	RRab	-2.06	0.14	0.6161452	17.964	0.051
A9154	OGLE-LMC-RRLYR-13494	5:23:02.93	-70:29:44.6	RRab	-1.66	0.14	0.6182903	17.972	0.029
B1408	OGLE-LMC-RRLYR-10067	5:17:13.84	-71:06:06.9	RRab	-1.70	0.11	0.6297088	18.021	0.013
A5589	OGLE-LMC-RRLYR-12968	5:22:09.60	-70:35:02.5	RRab	-1.60	0.13	0.6375745	17.948	0.035
A7468	OGLE-LMC-RRLYR-13176	5:22:30.06	-70:32:20.6	RRab	-1.55	0.11	0.6386908	18.043	0.041
A25510	OGLE-LMC-RRLYR-13002	5:22:13.43	-70:30:11.4	RRab	-1.72	0.11	0.6495506	17.713	0.038
A8720	OGLE-LMC-RRLYR-13956	5:23:50.19	-70:30:16.7	RRab	-1.88	0.34	0.6508174	17.847	0.037

**Table 1**  
(Continued)

Star	OGLE ID	R.A. (J2000)	Decl. (J2000)	Type	[Fe/H] (dex)	$\sigma_{[\text{Fe}/\text{H}]}$ (dex)	$P$ (days)	$\langle K_{s,0} \rangle$ (mag)	$\sigma_{(K_{s,0})}$ (mag)
B7063	OGLE-LMC-RRLYR-10973	5:18:44.05	-70:55:55.8	RRab	-1.49	0.14	0.6548698	17.867	0.023
B7620	OGLE-LMC-RRLYR-10541	5:18:03.58	-70:55:03.1	RRab	-2.05	0.12	0.6561602	17.689	0.034
A7477	OGLE-LMC-RRLYR-14068	5:24:02.97	-70:32:08.6	RRab	-1.67	0.28	0.6564084	17.955	0.017
A28293	OGLE-LMC-RRLYR-12758	5:21:46.13	-70:28:13.3	RRab	-1.74	0.10	0.6602890	17.979	0.050
A6426	OGLE-LMC-RRLYR-13196	5:22:32.51	-70:33:48.7	RRab	-1.59	0.09	0.6622400	17.868	0.038
A3948	OGLE-LMC-RRLYR-13285	5:22:40.40	-70:37:17.0	RRab	-1.46	0.12	0.6623845	17.944	0.036
A8094	OGLE-LMC-RRLYR-13306	5:22:43.06	-70:31:23.7	RRab	-1.83	0.12	0.7420663	17.890	0.033

**Note.** Columns report: (1) star identification from Di Fabrizio et al. (2005); (2) identification from the OGLE III catalog (Soszyński et al. 2009); (3) R.A. (OGLE); (4) decl. (OGLE); (5) type; (6) metallicity from Gratton et al. (2004); (7) metallicity error (Gratton et al. 2004); (8) period (OGLE); (9) dereddened mean  $K_s$  magnitude from the VMC data, determined from the analysis of the light curve with GRATIS; (10) Error of the mean  $K_s$  magnitude.

After deriving  $K_s$  mean magnitudes we performed the dereddening procedure. Clementini et al. (2003) estimated reddening values of  $E(B - V) = 0.116 \pm 0.017$  and  $0.086 \pm 0.017$  mag in LMC field *A* and *B*, respectively, using the method from Sturch (1966) and the colors of the edges of the instability strip defined by the RR Lyrae variables. Applying the coefficients from Cardelli et al. (1989) of  $A_K/A_V = 0.114$  and assuming a ratio of total to selective absorption of  $R_V = 3.1$ , we estimated the extinction in the  $K_s$ -band as

$$A_{K_s} = 0.35 \times E(B - V). \quad (1)$$

Table 1 summarizes the properties of the sample of 70 RR Lyrae stars that have a counterpart in the VMC catalog. The first and second columns give the identification of the stars in Di Fabrizio et al. (2005) and in the OGLE III catalog, respectively. The table also shows the coordinates and the classification of the stars from the OGLE III catalog, metallicities with errors from Gratton et al. (2004), and dereddened mean  $K_s$  magnitudes, determined with the GRATIS package, along with their errors.

### 3. $PL_{K_s}Z$ RELATION OF RR LYRAE STARS

#### 3.1. Method

Using the dereddened mean  $K_s$  magnitudes of the 70 RR Lyrae stars derived as described in Section 2, spectroscopically determined metallicities from Gratton et al. (2004), and accurately estimated periods from the OGLE III catalog (with RRc stars “fundamentalized” by adding 0.127 to the logarithm of the period) we can now fit the  $PL_{K_s}Z$  relation. The fit is performed using a fitting approach developed specifically for this work.

Fitting a line to data is a common exercise in science. Most common approaches use Minimum-Least-Squares methods; however, these are often based on assumptions that do not always hold for real observational data. The most basic methods assume that data are drawn from a thin line with errors, which are Gaussian, perfectly known, and exist in one axis only. These assumptions do not hold in the present case, as we have an unknown but potentially significant intrinsic dispersion, non-negligible errors in two dimensions ( $K_s$  and  $[\text{Fe}/\text{H}]$ ), and the possibility of inaccuracy in the formal error estimates (e.g., in the determination of the precision metallicity estimates).

We therefore follow the prescription of Hogg et al. (2010), who develop a method for fitting a line to data that avoids the

problems highlighted above by statistical modeling of the data. They present a method for use in two dimensions, which has been extended to three dimensions in this paper.

The method assumes that the data is drawn from a plane of the form


$$Ls(P, [\text{Fe}/\text{H}]) = A \log P + B [\text{Fe}/\text{H}] + C, \quad (2)$$

where  $A$  is the slope in the  $\log P$  axis,  $B$  is the slope in the metallicity  $[\text{Fe}/\text{H}]$  axis, and  $C$  is the intercept. We assume a uniform Gaussian intrinsic dispersion around the luminosity axis, plus the scatter caused by the Gaussian observational errors. The exact mathematical definition is given in Appendix A. The method utilizes adaptive Markov Chain Monte Carlo (MCMC) methods (Foreman-Mackey et al. 2013) to evaluate the posterior probability density function (PDF) of each parameter, given an input data set, and returns the maximum a posteriori probability estimate of each parameter, the formal error estimate, and the full posterior PDF. The formal error estimate is obtained from the 16% and 84% quartiles of the posterior PDF of the parameters, which give the  $1\sigma$  formal error estimate assuming that the posterior PDF is approximately normal. The free fit parameters are: the slope in  $\log P$ , the slope in metallicity, the zero-point, and the intrinsic dispersion perpendicular to the magnitude axis.

By applying this method we found the following relation between period, metallicity and mean apparent  $K_s$  magnitude:

$$K_{s,0} = (-2.73 \pm 0.25) \log P + (0.03 \pm 0.07) [\text{Fe}/\text{H}] + (17.43 \pm 0.01). \quad (3)$$

The intrinsic dispersion of the relation is found to be 0.01 mag. The rms deviation of the data around the relation, neglecting the intrinsic dispersion, is 0.1 mag. Since the reddening in the  $K_s$  band is negligible we suggest that the effects of the LMC depth cause the intrinsic dispersion of the relation. The left panel of Figure 2 presents the  $PL_{K_s}Z$  relation (Equation (3)) of the 70 LMC RR Lyrae stars in the period–luminosity–metallicity space, whereas the right panels show the projection of the  $PL_{K_s}Z$  on the  $\log(P) - K_s$  (top-right panel) and  $K_s - [\text{Fe}/\text{H}]$  (bottom right panel) planes. The gray lines in the figure are lines of equal metallicity (top-right) or equal period (bottom right). The method finds the relation (values of  $A$ ,  $B$ , and  $C$  for the relation  $K_s = A \log P + B [\text{Fe}/\text{H}] + C$ ) in the three dimensions ( $\log P$ ,  $K_s$ , and  $[\text{Fe}/\text{H}]$ ). Each of the gray lines in the top-right plot are therefore



**Figure 2.** Left panel:  $PL_{K_s}Z$  relation of the 70 LMC RR Lyrae stars (Equation (3)) analyzed in the paper, in the period–luminosity–metallicity space. Right panels: projections of the  $PL_{K_s}Z$  relation (Equation (3)) on the  $\log(P) - K_s$  (top) and  $K_s - [\text{Fe}/\text{H}]$  (bottom) planes. Gray lines represent lines of equal metallicity (top) and equal period (bottom). See the text for the details. Uncertainties in the  $K_s$  magnitude and  $[\text{Fe}/\text{H}]$  are omitted to simplify the figure, but they are provided in Table 1.

$K_s = A \log P + B [\text{Fe}/\text{H}] + C$  for the full range of periods, at the metallicity of each star (one line per star). Thus, by following the line up and down it is seen how  $K_s$  changes with period at some specific metallicity. The lines do not always cross the points on the diagram because the line is the result of the fit, and the points are affected by errors and intrinsic dispersion so they may be above or below the fit. In the bottom right plot the lines are  $K_s = A \log P + B [\text{Fe}/\text{H}] + C$  for the full range of metallicity with  $\log P$  taken from each star.

It is worth noting that we find a very small dependence of the  $K_s$  magnitude on metallicity. However, the metal abundance range spanned by the adopted sample does not reach the highest values (up to solar and supersolar) observed in the MW bulge and disk RR Lyrae populations. In order to study the effect of the adopted range of metallicities on the slope of the  $PL_{K_s}Z$  relation we derived this relation also for MW RR Lyrae stars. We discuss the results in Section 3.3.

### 3.2. Zero-point of the $PL_{K_s}Z$ Relation in the LMC

To use the derived  $PL_{K_s}Z$  relation for determining distances it is necessary to calibrate its zero-point. This can be done in a number of different ways. In this paper we follow two different approaches: the first one is based on adopting a value for the distance of the LMC; in the second one we use the absolute magnitudes of Galactic RR Lyrae stars for which trigonometric parallaxes have been measured with the *HST*/FGS. Both approaches have their advantages and disadvantages, we discuss them in the following sections.

#### 3.2.1. Zero-point Based on the LMC Distance

The LMC is widely considered to be the first rung of the cosmic distance ladder as it contains a large number of different distance indicators, such as Cepheids, RR Lyrae variables, eclipsing binaries (EBs), red giant branch stars, etc., allowing the galaxy distance to be determined using several independent techniques. Figure 8 of Benedict et al. (2002) shows an impressive summary of LMC distance moduli published during the ten years from 1992 to 2001. Values from 18.1 to 18.8 mag

were reported in the literature, with those smaller than 18.5 mag supporting the so-called “short” scale, and those larger than 18.5 mag, the “long” one. In more recent years the dramatic progress in the calibration of the different distance indicators has led the dispersion in LMC distance moduli to shrink significantly. Extreme values such as those listed in Benedict et al. (2002) are not very often seen in the recent literature (Clementini 2008). Still a general consensus on the LMC distance has not been fully reached yet. Moreover, there have been significant concerns about a possible “publication bias” affecting the distance to the LMC (Schaefer 2008; Rubele et al. 2012; Walker 2012). In particular, Schaefer (2008) claimed that from 2002 to 2007 June 31 independent papers reported new measurements of the distance of the LMC, and the new values clustered tightly around the value  $(m - M)_0 = 18.5 \pm 0.1$  mag, adopted by the *HST* Key Project on the extragalactic distance scale (Freedman et al. 2001). Schaefer (2008) considered the effects of the “publication bias” to be the most likely cause of the clustering of LMC distance measurements.

A number of studies on the compilation of distances to the LMC as derived from different distance indicators can be found in the literature of the last 15 years (e.g., Gibson 2000; Benedict et al. 2002; Clementini et al. 2003; Schaefer 2008; de Grijs et al. 2014). Clementini et al. (2003) analyzed the distance to the LMC measured using Population I and Population II standard candles and showed that all distance determinations converge within  $1\sigma$  error on a distance modulus  $(m - M)_0 = 18.515 \pm 0.085$  mag. The most recent compilation of LMC distance moduli is that of de Grijs et al. (2014) who compiled 233 separate distance determinations, published from 1990 March until 2013 December, and concluded that the canonical distance modulus of  $(m - M)_0 = 18.49 \pm 0.09$  mag may be used for all practical purposes. The compilation of de Grijs et al. (2014) includes the distance modulus of  $(m - M)_0 = 18.46 \pm 0.03$  estimated by Ripepi et al. (2012b) using LMC classical Cepheids observed by the VMC survey, and the recent determination of direct distances to eight long-period EBs in the LMC by Pietrzyński et al.

(2013), which is claimed to be accurate to within  $\sim 2\%$ :  $D_{\text{LMC}} = 49.97 \pm 0.19$  (stat)  $\pm 1.11$  (syst) kpc, corresponding to the distance modulus  $(m - M)_0 = 18.493 \pm 0.008$  (stat)  $\pm 0.047$  (syst) mag. Furthermore, the model fitting of the light curves of different classes of pulsating stars in the LMC, also based on different samples and hydrodynamical codes, provides values consistent with 18.5 mag (see Bono et al. 2002; Keller & Wood 2002, 2006; Marconi & Clementini 2005; McNamara et al. 2007).

The RR Lyrae stars in our sample are located in a relatively small area close to the center of the LMC bar. Neglecting depth/projection effects they can be considered as being all at the same distance from us and close to late-type EBs, which are all located relatively close to the barycenter of the LMC as derived by Pietrzyński et al. (2013). Therefore, in the following we adopt for the distance modulus of the LMC the value published by Pietrzyński et al. (2013). We subtracted this value from the dereddened mean  $K_s$  apparent magnitudes of our 70 RR Lyrae stars and derived absolute magnitudes in the  $K_s$  band ( $M_{K_s}$ ). Then by applying the technique described in Section 3.1 we derived the relation between  $K_s$ -band absolute magnitudes, periods, and metallicities, with the zero-point entirely based on the distance to the LMC by Pietrzyński et al. (2013):

$$M_{K_s} = (-2.73 \pm 0.25) \log P + (0.03 \pm 0.07) [\text{Fe}/\text{H}] - (1.06 \pm 0.01). \quad (4)$$

### 3.2.2. Zero-point Based on Trigonometric Parallaxes of Galactic RR Lyrae Stars

In order to obtain an estimate of the  $\text{PL}_{K_s}Z$  relation zero-point which is independent of the distance to the LMC and, in turn, be able to measure the distance to this galaxy from the RR Lyrae  $\text{PL}_{K_s}Z$  relation, it is necessary to know the absolute magnitude of the RR Lyrae stars with good accuracy. Trigonometric parallaxes remain the only direct method to measure distances and hence derive absolute magnitudes. Benedict et al. (2011) derived absolute trigonometric parallaxes for five Galactic RR Lyrae stars (RZ Cep, XZ Cyg, SU Dra, RR Lyr, and UV Oct) with the *HST*/FGS. With these parallaxes the authors estimated absolute magnitudes in the  $K_s$  and  $V$  passbands, corrected for interstellar extinction and Lutz-Kelker-Hanson bias (Lutz & Kelker 1973, Hanson 1979). Absolute magnitudes in the  $K_s$ -band, periods and metallicities from Benedict et al. (2011), and the slopes of the relation derived in Equation (3) were used in order to determine a zero-point from each of these five MW RR Lyrae stars. The metallicities in Benedict et al. (2011) are in the Zinn & West metallicity scale and were converted to the metallicity scale in Gratton et al. (2004) by adding 0.06 dex (see Section 2). The logarithm of the period of the RRc star RZ Cep was “fundamentalized” by adding 0.127. Then we calculated the weighted mean of the five zero-points, this corresponds to  $-1.27 \pm 0.08$  mag. The relation between absolute magnitudes, periods and metallicities with the zero-point based on the five MW RR Lyrae stars from Benedict et al. (2011) is:

$$M_{K_s} = (-2.73 \pm 0.25) \log P + (0.03 \pm 0.07) [\text{Fe}/\text{H}] - (1.27 \pm 0.08). \quad (5)$$

A recent analysis (A. Monson 2015, private communication) shows that there is likely a typo in Benedict et al.’s (2011) parallax for the RR Lyrae star RZ Cep. Hence, we excluded

this star from the sample and derived the zero-point based on parallaxes of the remaining four RR Lyrae stars (XZ Cyg, UV Oct, SU Dra, and RR Lyr):

$$M_{K_s} = (-2.73 \pm 0.25) \log P + (0.03 \pm 0.07) [\text{Fe}/\text{H}] - (1.25 \pm 0.06). \quad (6)$$

The situation improves, but there is still a difference of  $\sim 0.2$  mag between the two zero-point obtained based on the distance to the LMC (Equation (4)) and the one based on the *HST* parallaxes of four RR Lyrae stars (Equation (6)). In fact, if we apply our  $\text{PL}_{K_s}Z$  relation with zero-point calibrated on Benedict et al. (2011) parallaxes (Equation (6)) to determine the absolute magnitudes of the 70 RR Lyrae stars in our sample, we obtain a distance modulus for the LMC of  $(m - M)_0 = 18.68 \pm 0.10$  mag. This distance modulus is about 0.2 mag longer than the widely adopted value of  $(m - M)_0 = 18.5$  mag.

There are a number of possible explanations for the discrepancy between zero-points. First of all, we should remember that Pietrzyński et al. (2013) results have been called into question by Schaefer (2013) who, in addition to concerns regarding possible bandwagon effects, also pointed out that Pietrzyński et al.’s (2013) distance to the LMC differs significantly from the average distance inferred from four hot, early-type EBs,  $D = 47.1 \pm 1.4$  kpc ( $(m - M)_0 = 18.365 \pm 0.065$  mag), published by Guinan et al. (1998), Fitzpatrick et al. (2002, 2003), and Ribas et al. (2002). Furthermore, in using the late-type EBs to calibrate the RR Lyrae  $\text{PL}_{K_s}Z$  relation we have implicitly assumed that RR Lyrae stars and EBs are at the same distance from us. However, when pushing for distance comparisons at a few percent level the effects of sample size, spatial distribution, depth, and geometric projection become important and properly accounting for the internal structure of the LMC may become necessary. The RR Lyrae stars in our sample could be distributed along the whole depth of the LMC. Moreover, RR Lyrae stars and EBs from Pietrzyński et al. (2013) could reside in different sub-structures of the LMC, which could be the reason for the systematic error in the determination of the zero-point (see, e.g., Moretti et al. 2014 for different features of the LMC structure traced by classical Cepheids, RR Lyrae stars and hot EBs).

On the other hand, when calibrating the zero-point by applying parallaxes of the four MW RR Lyrae stars by Benedict et al. (2011) we implicitly assumed that the  $\text{PL}_{K_s}Z$  relation is the same in the MW and in the LMC, which may not be true (see Section 3.3). We may also wonder whether there might be unknown systematic errors affecting Benedict et al.’s parallaxes. These come from *HST* fields, which provide relative and not absolute trigonometric parallaxes. Absolute parallaxes of the reference stars in each field are estimated via a complex procedure of fitting the spectral type and luminosity class of each star. A general formal error of 0.5 mas is applied to the absolute parallax of the reference stars, equal for all stars in all fields, and without justification. This could result in miscalculated estimates of the precision of the final absolute parallax measurements of the four RR Lyrae stars. The Lutz-Kelker bias is corrected a posteriori. In this respect it is worth noting that, according to van Leeuwen (2007), the *Hipparcos* parallax of RR Lyrae itself, the only RR Lyrae variable for which the satellite measured the parallax with high precision ( $\pm 0.64$  mas), is about 0.31 mas smaller than Benedict



**Table 2**  
Properties of 23 Bright RR Lyrae Stars in the Milky Way

Star	R.A. (deg)	Decl. (deg)	$P$ (day)	[Fe/H] (dex)	$\sigma_{[\text{Fe}/\text{H}]}$ (dex)	$E(B - V)$ (mag)	$V$ (mag)	$K$ (mag)	$K_s$ (mag)	$M_V$ (mag)	$M_K$ (mag)	$\sigma_{M_V, M_K}$ (mag)
UU Cet	1.02135	-16.99764	0.606081	-1.38	0.08	0.015	12.08	10.85	10.837	0.62	-0.55	0.15
SW And	5.92954	29.40101	0.442279	-0.06	0.08	0.045	9.71	8.54	8.505	0.94	-0.07	0.15
RR Cet	23.03405	1.34173	0.553025	-1.38	0.08	0.015	9.73	8.56	8.520	0.68	-0.42	0.15
X Ari	47.12869	10.44590	0.651139	-2.50	0.08	0.18	9.57	7.95	7.941	0.57	-0.575	0.15
AR Per	64.32165	47.40018	0.425549	-0.34	0.16	0.31	10.51	8.66	8.642	0.87	-0.06	0.25
RX Eri	72.43455	-15.74118	0.587246	-1.63	0.16	0.03	9.69	8.42	8.429	0.66	-0.51	0.25
RR Gem	110.38971	30.88318	0.397316	-0.32	0.16	0.075	11.38	10.26	10.275	0.89	-0.02	0.25
TT Lyn	135.78245	44.58559	0.597438	-1.64	0.16	0.015	9.86	8.65	8.611	0.65	-0.55	0.15
T Sex	148.36833	2.05732	0.324698	-1.20	0.16	0.015	10.04	9.18	9.200	0.66	-0.07	0.25
RR Leo	151.93108	23.99176	0.452387	-1.37	0.16	0.015	10.73	9.70	9.730	0.76	-0.16	0.15
WY Ant	154.02061	-29.72845	0.574312	-1.32	0.16	0.06	10.87	9.64	9.674	0.55	-0.55	0.15
W Crt	171.62351	-17.91435	0.412013	-0.89	0.16	0.03	11.54	10.56	10.539	0.96	+0.08	0.15
TU UMa	172.45205	30.06733	0.557659	-1.38	0.16	0.015	9.82	8.67	8.660	0.70	-0.41	0.15
UU Vir	182.14613	-0.45676	0.475606	-0.64	0.16	0.015	10.56	9.51	9.414	0.80	-0.195	0.15
SW Dra	184.44429	69.51062	0.569670	-0.91	0.16	0.015	10.48	9.33	9.319	0.68	-0.38	0.15
RV Oct	206.63230	-84.40177	0.571130	-1.92	0.16	0.09	10.98	9.51	9.526	0.68	-0.40	0.15
TV Boo	214.15242	42.35992	0.312559	-2.31	0.16	0.015	10.97	10.22	10.248	0.58	-0.20	0.25
RS Boo	218.38839	31.75462	0.377337	-0.37	0.16	0.015	10.37	9.45	9.507	0.85	+0.00	0.25
VY Ser	232.75803	1.68382	0.714094	-1.71	0.08	0.03	10.13	8.78	8.826	0.61	-0.665	0.25
V445 Oph	246.17171	-6.54165	0.397023	+0.17	0.08	0.195	11.05	9.24	9.262	1.09	+0.30	0.25
TW Her	268.63000	30.41048	0.399601	-0.58	0.16	0.06	11.28	10.22	10.239	0.80	-0.07	0.15
AV Peg	328.01164	22.57483	0.390380	-0.03	0.16	0.06	10.50	9.36	9.346	1.10	+0.14	0.15
RV Phe	352.13106	-47.45362	0.596413	-1.69	0.16	0.015	11.94	10.72	10.768	0.86	-0.29	0.25

**Note.** The columns report: (1) name of the star; (2) R.A. (J2000) from SIMBAD database; (3) decl. (J2000) from SIMBAD database; (4) period from Fernley et al. (1998a); (5) metallicity from Clementini et al. (1995); (6) error in metallicity from Clementini et al. (1995); (7) reddening from Fernley et al. (1998a); (8)  $V$  magnitude from Fernley et al. (1998a); (9)  $K$  magnitude in the Johnson system from Fernley et al. (1998a); (10)  $K_s$  magnitude in the 2MASS system from Feast et al. (2008); (11) absolute magnitude in the  $V$  passband from Fernley (1994); (12) absolute magnitude in the  $K$  passband obtained from B–W analyses and corrected to  $p = 1.38$  (see the text for details); (13) errors in the absolute  $V$ ,  $K$  magnitudes.


et al.’s (2011) parallax for the star, although consistent with it within the errors, hence, the corresponding distance modulus is about 0.17 mag longer. In any case, a great contribution to the determination of the zero-point of the RR Lyrae  $PL_{K_s}Z$  relation is expected from the ESA astrometric satellite *Gaia*. We discuss this topic in Section 4.

### 3.3. $PL_{K_s}Z$ Relation of RR Lyrae Stars in the MW

In spite of many studies in the literature, it still remains unsettled whether the RR Lyrae  $PL_{K_s}Z$  is a universal relation. To investigate this matter we have derived the  $PL_{K_s}Z$  relation for RR Lyrae stars in the MW and compared it with the relation we have obtained in Section 3.2 for the LMC variables. To this end we selected 23 MW RR Lyrae stars which have absolute magnitudes known from B–W studies based on near-infrared data (Jones et al. 1988, 1992; Fernley et al. 1990; Liu & Janes 1990; Cacciari et al. 1992; Skillen et al. 1993; Fernley 1994, and references therein) and metallicities determined from abundance analysis of high-resolution spectra. Information about these 23 RR Lyrae stars is presented in Table 2. Star’s coordinates in the table are from the SIMBAD database; periods, apparent  $V$  and  $K$  magnitudes and reddening  $E(B - V)$  are from Fernley et al. (1998a). The sample contains two first-overtone mode RR Lyrae stars (namely, T Sex and TV Boo). As done for the LMC RRc stars, their periods were fundamentalized by adding +0.127 to the logarithm of the period. Absolute  $M_V$  and  $M_K$  magnitudes in Columns 10 and 12 were taken from the compilations of B–W results in Table 11 of Cacciari et al. (1992) and from Table 16

of Skillen et al. (1993) for the variable stars: WY Ant, W Crt, and RV Oct. According to Cacciari et al. (1992) the  $K$  magnitudes of the stars analyzed with the B–W method are in the Johnson photometric system. Following the discussion in Cacciari et al. (1992) and Skillen et al. (1993) we retained only 23 of the original lists of 29 field RR Lyrae stars analyzed with the B–W method, as stars which are likely to be evolved (DX Del, SU Dra, SS Leo, BB Pup, and W Tuc) were discarded. We also discarded DH Peg as there is suspicion that the star is a dwarf Cepheid (see Feast et al. 2008 and discussion therein). Furthermore, following Fernley (1994), original  $M_V$  and  $M_K$  values were revised (i) assuming for the  $p$  factor used to convert the observed pulsation velocity to true pulsation velocity in B–W analyses the value of  $p = 1.38$ , and (ii) multiple determinations of individual stars were averaged.

Metal abundances with related errors are needed to apply our fitting approach. Several different spectroscopic studies have targeted the stars in Table 2. In Appendix B we provide a summary of their results. The largest and most homogeneous samples are those by Clementini et al. (1995) and Lambert et al. (1996). These authors measured [Fe/H] abundances from high-resolution spectra for several of the stars in Table 2 and provided recalibrations of the  $\Delta S$  index (Preston 1959), from which metal abundances can be derived for the stars which lack abundance analysis. For the sake of homogeneity and ease of use in this paper we adopt metallicities and metallicities errors for the MW RR Lyrae stars as they are listed, ready for use, in Table 21 of Clementini et al. (1995). These [Fe/H] values are the average of the Fe I and Fe II abundances, adopting  $\log \epsilon(\text{Fe I}) = 7.56$  and  $\log \epsilon$



**Figure 3.** Left panel:  $PM_KZ$  relation of the 23 MW RR Lyrae stars (Equation (7)) in the period–luminosity–metallicity space. Right panel: projections of the  $PM_KZ$  relation (Equation (7)) on the  $\log(P) - M_K$  (top panel) and  $M_K - [\text{Fe}/\text{H}]$  (bottom panel) planes. Gray lines represent lines of equal metallicities (top panel) and periods (bottom panel). See the text for the details. Uncertainties in the  $M_K$  magnitude and  $[\text{Fe}/\text{H}]$  are omitted to simplify the figure, but they are provided in Table 2.

( $\text{Fe II}$ ) = 7.50 for the Sun. They are reported along with the related errors in columns 5 and 6 of Table 2. Metallicities in Lambert et al. (1996) were obtained from the  $\text{Fe II}$  abundance and adopting  $\log \epsilon(\text{Fe I}) = 7.51$  for the Sun. In Appendix B we present the PLZ relation obtained using Lambert et al.’s (1996) metallicities and our approach.

By applying our fitting approach (Section 3.1) to the 23 MW RR Lyrae stars we derived the following  $PL_KZ$  relation:

$$M_K = (-2.53 \pm 0.36)\log P + (0.07 \pm 0.04)[\text{Fe}/\text{H}] - (0.95 \pm 0.14). \quad (7)$$

The intrinsic dispersion of the relation is found to be 0.007 mag. The rms deviation of the data around the relation, neglecting the intrinsic dispersion, is 0.086 mag. It should be noted that the metallicities listed in Table 2 may differ slightly from the metal abundances used in the B–W analysis of these stars. However, this is not of great concern because the B–W based on near-infrared data is mildly affected by small changes in metallicity and reddening. We also point out that the rather large error of the  $\log P$  term in Equation (7) is largely driven by the large errors (0.15–0.25 mag) in the  $K$ -band absolute magnitudes from the B–W analyses. This is confirmed by the exercise with *Gaia* simulated parallaxes we present in Section 4.

The slope in  $[\text{Fe}/\text{H}]$  in Equation (7) is higher than the slope obtained for the LMC RR Lyrae stars (Equations (4), (6)), although they are still consistent within the respective errors. Equation (7) was derived over a wide range of metallicities  $[-2.5; 0.17]$  dex, but the slope of the metallicity term remains rather small. Thus, the relatively small metallicity range spanned by the LMC variables could be not responsible for the negligible dependence on metallicity of the RR Lyrae  $PL_KZ$  relation in the LMC.

The distribution of the 23 MW RR Lyrae stars in the period–luminosity–metallicity space and the projections of the  $PL_KZ$  relation (Equation (7)) on the  $\log(P) - M_K$  and  $M_K - [\text{Fe}/\text{H}]$  planes are shown in Figure 3. Gray lines are the same as in Figure 2 and are described in Section 3.1.

Some concern may arise since the  $K$  magnitudes of the 23 MW RR Lyrae stars used to derive Equation (7) are in the Johnson photometric system (see Cacciari et al. 1992), whereas for the LMC RR Lyrae variables we have  $K_s$  photometry in the VISTA system.<sup>9</sup> To address this issue we have reported in column 10 of Table 2 the average  $K_s$  magnitudes in the 2MASS system of the 23 MW RR Lyrae stars as provided by Feast et al. (2008). The difference with the Johnson  $K$  average magnitudes listed in column 9 is very small (of the order of about 0.03 mag, on average) and definitely much smaller than individual errors in the B–W  $K$ -band absolute magnitudes of the MW variables (0.15–0.25 mag), or errors in the  $K_s$  average apparent magnitudes of the LMC RR Lyrae stars (see column 10 of Table 1). Hence, we are confident that the difference in photometric system does not significantly affect our comparison.

### 3.4. Comparison with the Literature

The near-infrared  $PL_KZ$  relation of the RR Lyrae stars has been studied by several authors both from a theoretical and an observational point of view. Longmore et al.’s (1986) pioneering work was followed by Liu & Janes (1990), Skillen et al. (1993), and Jones et al. (1996). A comprehensive analysis of the IR properties of RR Lyrae stars was performed by Nemeč et al. (1994).

Some of the RR Lyrae  $PL_KZ$  relations available in the literature are presented in Table 3. Bono et al. (2003) derived the semi-theoretical relation presented in the first row of Table 3. This theoretical relation has been derived from an extended set of RR Lyrae nonlinear hydrodynamical models spanning a wide range of chemical compositions ( $Z$  from 0.0001 to 0.02, which approximately corresponds to  $[\text{Fe}/\text{H}]$  from  $-2.45$  to  $-0.15$  dex). Catelan et al. (2004) presented a theoretical calibration of the RR Lyrae  $PL_KZ$  relation based on

<sup>9</sup> The VISTA system is tied to the 2MASS photometry (Skrutskie et al. 2006), with the difference in  $K_s$  magnitude only mildly depending on the  $J - K_s$  color, and being of the order of 3–4 mmag for the typical  $J - K_s$  color of RR Lyrae stars.

**Table 3**  
PL<sub>K<sub>s</sub></sub>Z Relations from the Literature

Relation	<i>a</i>	<i>b</i>	ZP <sup>a</sup> from the Original Relation	ZP <sup>b</sup> from Benedict et al. (2011)
Theoretical or semi-theoretical relations				
Bono et al. (2003)	-2.101	0.231 ± 0.012	-0.770 ± 0.044	-0.58 ± 0.04
Catelan et al. (2004)	-2.353	0.175	-0.869	...
Empirical relations				
Dall’Ora et al. (2004)	-2.16 ± 0.09	...	...	-0.56 ± 0.02
Del Principe (2006) <sup>c,d</sup>	-2.71 ± 0.12	0.12 ± 0.04	...	-0.57 ± 0.02
Sollima et al. (2006) <sup>e</sup>	-2.38 ± 0.04	0.08 ± 0.11	-1.05 ± 0.13	-0.57 ± 0.03 <sup>f</sup>
Sollima et al. (2008) <sup>e</sup>	-2.38 ± 0.04	0.08 ± 0.11	-1.07 ± 0.11	-0.56 ± 0.02
Borissova et al. (2009) <sup>c,g</sup>	-2.11 ± 0.17	0.05 ± 0.07	-1.05	-0.56 ± 0.03
This paper (LMC) <sup>c,g</sup>	-2.73 ± 0.25	0.03 ± 0.07	-1.06 ± 0.01	-0.55 ± 0.06 <sup>i</sup>
This paper (MW) <sup>h</sup>	-2.53 ± 0.36	0.07 ± 0.04	-0.95 ± 0.14	-0.56 ± 0.06 <sup>i</sup>

#### Notes.

<sup>a</sup> Zero-point of the original relation from the literature in the form:  $a \log P + b[\text{Fe}/\text{H}] + \text{ZP}$ .

<sup>b</sup> Zero-point of the relation in the form:  $a(\log P + 0.28) + b([\text{Fe}/\text{H}] + 1.58) + \text{ZP}$ , as recalibrated by Benedict et al. (2011).

<sup>c</sup> Near-infrared photometry in the  $K_s$  band.

<sup>d</sup> Metallicity is on the Zinn & West (1984) metallicity scale.

<sup>e</sup> Metallicity is on the Carretta & Gratton (1997) metallicity scale.

<sup>f</sup> Zero-point was calibrated by Benedict et al. (2011) neglecting the metallicity term.

<sup>g</sup> Metallicity on the scale adopted by Gratton et al. (2004), which is, on average, 0.06 dex higher than Zinn & West (1984) scale.

<sup>h</sup> Metallicities from Clementini et al. (1995), they are on the scale of the high dispersion spectra (i.e., the Carretta et al. 2009).

<sup>i</sup> Zero-point was calibrated by us considering only four RR Lyrae stars (XZ Cyg, UV Oct, SU Dra, and RR Lyr) from Benedict et al. (2011) and excluding RZ Cep, since there are concerns about the parallax of this star.

synthetic HB models computed for several different metallicities, fully taking into account evolutionary effects besides the effect of chemical composition. They derived the relation:

$$M_K = -2.353 \log P + 0.175 \log Z - 0.597. \quad (8)$$

By using Equations (9) and (10) in Catelan et al. (2004) and assuming  $[\alpha/\text{Fe}] \sim 0.3$  (e.g., Carney 1996) we transformed Equation (8) into the form, presented in the second row of Table 3.

Dall’Ora et al. (2004) derived an empirical relation between apparent  $K$  magnitude and period for 21 RRab and 9 RRc stars in the LMC globular cluster Reticulum. Del Principe (2006) obtained the relation between apparent  $K_s$  magnitude, metallicity, and period from the analysis of RR Lyrae stars in the Galactic globular cluster  $\omega$  Cen.

Sollima et al. (2006) derived a PL<sub>K</sub>Z relation by analyzing 538 RR Lyrae stars in 15 Galactic clusters and in the LMC globular cluster Reticulum. This relation spans the metallicity range  $-2.15 < [\text{Fe}/\text{H}] < -0.9$  dex. Mean  $K$  magnitudes were estimated by combining 2MASS (Cutri et al. 2003) photometry and literature data. The zero-point was calibrated on RR Lyrae itself, whose distance modulus was derived using the *HST* trigonometric parallax measured for this star by Benedict et al. (2002). Sollima et al. (2008) presented *JKH* time-series photometry of RR Lyrae and derived a new zero-point of Sollima et al.’s (2006) PL<sub>K</sub>Z relation.

Borissova et al. (2009) presented near-infrared  $K_s$  photometry and spectroscopically measured metallicity for a sample of 50 field RR Lyrae stars in inner regions of the LMC. These authors had five measurements in the  $K_s$  passband for most of the stars in their sample and used templates from Jones et al. (1996) to fit the light curves and derive the mean  $K_s$  magnitudes. To improve statistics they added to their sample LMC RR Lyrae stars from Szweczyk et al.’s (2008) data set,

and derived the PL<sub>K<sub>s</sub></sub>Z relation based on the total sample of 107 LMC variables. The zero-point was calculated using Sollima et al.’s (2008) mean  $K$  magnitude, the reddening, and the trigonometric parallax of RR Lyrae.

Benedict et al. (2011) recalibrated all the literature relations listed in Table 3, but Catelan et al. (2004)’s one, by fitting to equations in the form:  $a(\log P + 0.28) + b([\text{Fe}/\text{H}] + 1.58) + \text{ZP}$ , the Lutz–Kelker–Hanson-corrected absolute magnitudes of the five MW RR Lyrae stars for which *HST* parallaxes are available. Since there are concerns about Benedict et al.’s (2011) RR Lyrae star (namely RZ Cep) we transformed the PL<sub>K</sub>Z relations of the LMC and MW RR Lyrae stars derived in this paper (Equations (4), (6)) to the form adopted by Benedict et al. (2011) and determined their zero-points on the basis of Benedict et al.’s (2011) *HST* parallaxes but excluding RZ Cep. The zero-points based on Benedict et al. (2011) parallaxes are presented in column 5 of Table 3, whereas the zero-point of our LMC PL<sub>K<sub>s</sub></sub>Z relation calculated by assuming the distance to the LMC (Section 3.2) and the zero-point of our MW PL<sub>K</sub>Z relation based on the B–W studies are (Section 3.3) are presented in Column 4.

The slope in the period of the RR Lyrae PL<sub>K</sub>Z relations (Column 2 of Table 3) differs significantly in the various studies. The slope we derived for the LMC RR Lyrae stars is in excellent agreement with that derived by Del Principe (2006), whereas the slope of the MW RR Lyrae PL<sub>K</sub>Z is in good agreement with that derived by Sollima et al. (2006, 2008).

The dependence on metallicity of the PL<sub>K</sub>Z relations (Column 3 of Table 3) also varies among different studies and generally is larger in the theoretical and semi-theoretical relations. The comparison of the metallicity dependence in the different empirical relations is complicated by the inhomogeneity of the metallicity scales adopted in these studies. Metallicities in Del Principe (2006) are in the Zinn & West

(1984) scale, whereas in Sollima et al. (2006, 2008) they are in the Carretta & Gratton (1997) scale. In the current study for the LMC RR Lyrae stars we used the metallicity scale defined by Gratton et al. (2004) which is also the scale adopted by Borissova et al. (2009). As discussed in Gratton et al. (2004) this scale is systematically 0.06 dex higher than the Zinn & West scale. This difference is small and systematic, hence it should not affect the results of this comparison. Finally, for the MW RR Lyrae stars we used the metallicities measured by Clementini et al. (1995). Because the spectroscopic [Fe/H] values in Clementini et al. (1995) are derived from high dispersion spectra analyzed using standard reduction procedures, the derived metallicities are on the scale of the high dispersion spectra (i.e., the Carretta et al. 2009 scale) and could be transformed to the Zinn & West scale using the relations provided in Carretta et al. (2009).

The slope in metallicity of the  $PL_{KZ}$  relation based on the LMC RR Lyrae stars is the smallest among the various studies listed in Table 3 and it is close to Borissova et al.'s (2009) slope. This is consistent with the two studies both involving LMC variables and using exactly the same metallicity scale. The slope in metallicity we found for the MW RR Lyrae stars is larger than that of the LMC RR Lyrae stars and, in spite of the difference in metallicity scales, it is very close to the slope obtained by Sollima et al. (2006) for RR Lyrae stars in globular clusters. However, taken at face value, the metallicity slopes of the empirical relations in Table 3 appear to be all rather small and in agreement to each other within the relative uncertainties, thus generally suggesting a mild dependence with on the RR Lyrae  $PL_{KZ}$ , independently of the specific environment.

#### 4. GAIA OBSERVATION OF RR LYRAE STARS IN THE MILKY WAY

The *Gaia* astrometric satellite will revolutionize many fields of astronomy (Perryman et al. 2001). Of particular importance will be its catalog of trigonometric parallaxes for more than one billion stars, with astrometric precision down to the  $\mu$  as level. Due to *Gaia*'s constant observation of the sky over the five year nominal mission, *Gaia* will repeatedly observe all stars brighter than its limiting magnitude, with an average of 70 observations per star. This will also make it possible for *Gaia* to discover and characterize many types of variables, including RR Lyrae stars and Cepheids.

*Gaia* is observing in the broad visual band *G* (Jordi et al. 2010) for its astrometric measurements, and is therefore not ideal for characterizing the RR Lyrae PLZ relation, which exists only in the infrared passbands. However, since *Gaia* will provide accurate parallaxes for an expected tens of thousands of MW RR Lyrae stars, it could serve as a perfect tool for the determination of the zero-point of the  $PL_{KZ}$  relation through a combination with external data sets. As it was discussed in Section 3.2.2, the current largest limiting factor in zero-point calibration of  $PL_{KZ}$  and  $M_V - [Fe/H]$  relations is the lack of a reliable and statistically significant sample of parallax measurements. The current state of the art is the sample of five RR Lyrae parallaxes from Benedict et al. (2011) using the *HST*. *Gaia* will improve this situation by several orders of magnitude in both precision and numbers of objects. Moreover, the distance to the LMC will be determined through the combination of *Gaia* parallaxes for a large sample of LMC bright stars, hence, a zero-point of the  $PL_{KZ}$  relation based on the distance to the LMC will be obtained with a high precision.

#### 4.1. Method

Using parallax data for the calibration of a PL relation is complicated by the presence of statistical biases (e.g., Lutz & Kelker 1973) and sample selection effects (e.g., Malmquist 1936). Nonlinear transformations on the parallax cause a highly asymmetric uncertainty on the absolute magnitude when calculated using parallax and apparent magnitude information via the relation:  $m - M = -5 - 5 \log(\varpi)$ , where  $\varpi$  is the parallax. Additionally, stars with a negative parallax measurement cannot be used to calculate an absolute magnitude, though they do contain information. For these reasons, calculating an absolute magnitude for each star and fitting a PL relation directly to period and absolute magnitude leads to a biased result.

An unbiased solution can be achieved through modeling the data and inferring the slope and zero-point of the relation via statistical methods. For a catalog of  $N$  stars we can define  $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N)$  where the vector ( $\mathbf{x}_i = m, l, b, P, \varpi, A$ ) describes the observed data on each object.  $P$  is the period,  $m$  the apparent magnitude,  $l$  and  $b$  the position,  $\varpi$  the parallax, and  $A$  the extinction. We can additionally define that the vector ( $\mathbf{x}_0 = m_0, l_0, b_0, P_0, r_0, A_0$ ) gives the "true" underlying object properties.

We assume that the stars follow a PL relation of the form  $M_0 = \rho \log P_0 + \delta$ , although this can be changed to include other terms, such as metallicity, as needed. We can therefore model the true absolute magnitudes of the population as being Normally distributed around the PL relation, with the dispersion describing the intrinsic scatter on the relation:

$$\varphi_M(\mathbf{x}_0 | \rho, \delta, \sigma_{PL}) = \frac{1}{\sigma_{PL} \sqrt{2\pi}} e^{-0.5 \left( \frac{M_0 - (\rho \log P_0 + \delta)}{\sigma_{PL}} \right)^2} \quad (9)$$

where  $\sigma_{PL}$  is the intrinsic dispersion of the PL relation. The parameters  $\rho$  and  $\delta$  are the slope and zero-point of the PL relation, which are to be found.

The true absolute magnitude is calculated through

$$M_0 = m_0 + 5 \log(\varpi_0) + 5 - A_0. \quad (10)$$

The observations are normally distributed around the true values with a standard deviation given by the formal error on the measurement

$$\mathcal{E}(\mathbf{x} | \mathbf{x}_0) = \frac{1}{\epsilon_\varpi \epsilon_m \epsilon_A (2\pi)^{3/2}} e^{-0.5 \left( \frac{\varpi - \varpi_0}{\epsilon_\varpi} \right)^2} e^{-0.5 \left( \frac{m - m_0}{\epsilon_m} \right)^2} \times e^{-0.5 \left( \frac{A - A_0}{\epsilon_A} \right)^2} \delta(l_0, b_0, P_0). \quad (11)$$

Assuming negligible errors on the position and period, the observations are described by a delta function. The terms  $\epsilon_\varpi$ ,  $\epsilon_m$ ,  $\epsilon_A$  are the formal errors on the parallax, magnitude, and extinction.

With the above models defined, the joint PDF for the observations is

$$\begin{aligned} \mathcal{P}(\mathbf{x}_i | \rho, \delta, \sigma_{PL}) \\ = \mathcal{S}(\mathbf{x}) \int_{\forall \mathbf{x}_0} \varphi_M(\mathbf{x}_0 | \rho, \delta, \sigma_{PL}) \mathcal{E}(\mathbf{x} | \mathbf{x}_0) d\mathbf{x}_0; \end{aligned} \quad (12)$$

the "true" parameters  $\mathbf{x}_0$  are never known and so these values are marginalized through integration. The term  $\mathcal{S}(\mathbf{x})$  is the selection function, which takes the probability of observing a

star into account, given the properties of the star and the instrument’s observational capabilities. To take the fact that *Gaia* is a magnitude limited sample into account, a step function is used with

$$S(x) = \begin{cases} 1, & \text{if } G < 20, \\ 0, & \text{otherwise.} \end{cases} \quad (13)$$

The Maximum Likelihood Estimation of the parameters is found by maximizing Equation (12) by varying the parameters  $(\alpha, \rho, \sigma_{PL})$ . This formulation avoids nonlinear transformations on error-affected data and includes a selection function which avoids the Malmquist bias.

#### 4.2. Simulated *Gaia* Data

In order to check the application of the method defined in Section 4.1 we have used the sample of 23 RR Lyrae stars in the MW discussed in Section 3.3 (see also Table 2). In order to investigate the performance of the *Gaia* satellite and the contents of the end-of-mission catalog, *Gaia*’s Data Processing and Analysis Consortium (DPAC) has a group working on the simulation of several aspects of the *Gaia* mission. One major product of this work is the *Gaia* Object Generator (GOG; Luri et al. 2014), designed to simulate both individual *Gaia* observations and the full contents of the end-of-mission catalog. GOG includes a full mathematical description of the nominal performance of the *Gaia* satellite, and is therefore capable of determining the expected precision in astrometric, photometric and spectroscopic observations. In general, the precision depends on the apparent magnitude of the star, its color, and its sky position, which affects the number and type of observations made (due to the *Gaia* scanning law).

To obtain a distance for each RR Lyrae star from the sample, we use

$$M_K = -2.53 \log P - 0.95, \quad (14)$$

as determined in Equation (7) to obtain an absolute magnitude (neglecting the metallicity term for simplicity). We then determine a distance by combining this absolute magnitude with the apparent magnitude and extinction as defined above. Color information as  $(V - I)$  is obtained from the *Hipparcos* catalog (Perryman & ESA 1997) where available. The apparent magnitude, position, color, and period data form the basis of a synthetic catalog of RR Lyrae stars, along with the distance obtained from the  $PM_KZ$  relation, and is used as the input catalog of “true” parameters for GOG.

GOG then creates simulated *Gaia* observations for our sample. We take the  $PM_KZ$  relation (Equation (14)) as true, as a study of the possible precision in  $PM_KZ$  calibration after the *Gaia* data will become available.

Using the fitting method described in Section 4.1 to the data including the simulated parallax observations and simulated errors applied to parallax and apparent magnitude, we find a  $PM_K$  relation of

$$M_K = (-2.531 \pm 0.038) \log P + (-0.95 \pm 0.01) \quad (15)$$

Comparison of these results to the input relation shows very good agreement. It proves that the fitting procedure given in Section 4.1 is accurate and unbiased. When *Gaia* parallaxes become available for the much larger sample of RR Lyrae stars, we will apply the described method to fit the  $PL_KZ$  relation of RR Lyrae variables in the MW. Moreover, precise distance to

the LMC obtained from the combination of *Gaia* parallaxes for a large sample of the bright LMC stars, will allow us to calibrate the zero-point of the  $PL_{K_s}Z$  relation based on the LMC RR Lyrae stars. This provides a hint of what will be possible to achieve with *Gaia* parallaxes.

## 5. SUMMARY

We studied a sample of 70 RR Lyrae stars in the LMC for which multi-epoch  $K_s$  photometry from the VMC survey, precise periods from the OGLE III catalog, and spectroscopically determined metallicities (Gratton et al. 2004) are available. There are 13 epoch data in the  $K_s$  band for all stars in the sample, which allowed us to determine mean  $K_s$  magnitudes with a great accuracy.

Specifically for this work we developed a fitting approach. This method has several advantages compared to the minimum-least-squares fitting, such as taking into account potentially significant intrinsic dispersion of the data, non-negligible errors in two dimensions, and the possibility of inaccuracies in the formal error estimates. We used this method to derive the  $PL_{K_s}Z$  relation of the 70 RR Lyrae stars in the LMC. Potentially the method could be used to fit any other sample of data.

The zero-point of the derived  $PL_{K_s}Z$  relation was estimated in two different ways: (i) by assuming the distance to the LMC determined by Pietrzyński et al. (2013); and (ii) by applying *HST* parallaxes of four MW RR Lyrae stars by Benedict et al. (2011). The zero-point derived using the MW RR Lyrae stars is 0.2 mag larger and consequently gives a longer distance to the LMC:  $(m - M) = 18.68 \pm 0.10$  mag. In future studies we suggest using the relation based on the precise distance to the LMC:

$$M_{K_s} = (-2.73 \pm 0.25) \log P + (0.03 \pm 0.07) [\text{Fe}/\text{H}] - (1.06 \pm 0.01). \quad (16)$$

We found a negligible dependence of the  $M_{K_s}$  on metallicity, which could be caused by the relatively small range in metallicity covered by the LMC RR Lyrae stars. Thus, we applied the fitting approach to 23 RR Lyrae stars in the MW, for which absolute  $M_K$  and  $M_V$  magnitudes are known from B–W studies. We derived the  $PL_{K_s}Z$  relation for MW RR Lyrae stars in the form

$$M_K = (-2.53 \pm 0.36) \log P + (0.07 \pm 0.04) [\text{Fe}/\text{H}] - (0.95 \pm 0.14). \quad (17)$$

Even though the metallicities of the MW RR Lyrae stars span a wide range  $[-2.5; 0.17]$  dex, the dependence on metallicity is relatively small and consistent, within the errors, with the slope in metallicity found for the LMC RR Lyrae variables. We concluded that the small range of metallicities does not cause the negligible dependence of the  $M_K$  on metallicity for the LMC RR Lyrae stars.

To solve the problem of the  $PL_{K_s}Z$  zero-point, a large sample of RR Lyrae stars with precisely determined parallaxes is necessary. A great contribution to this field is expected by the *Gaia* satellite. By using GOG we simulated *Gaia* parallaxes of 23 MW RR Lyrae stars with observational errors. We present a method for the calibration of the PL relation which avoids several of the problems that arise when using parallax data. The method was tested by deriving the  $PL_{K_s}$  relation based on the simulated *Gaia* parallaxes. When combined with metallicity

and photometry from other sources and a statistical tool such as the one developed in the present study, the extraordinarily large sample of *Gaia* parallaxes for RR Lyrae stars will allow us to estimate these relations with unprecedented precision.

This work was supported by the *Gaia* Research for European Astronomy Training (GREAT-ITN) Marie Curie network, funded through the European Union Seventh Framework Programme [FP7/2007-2013] under grant agreement No. 264895. This work was also supported by the MINECO (Spanish Ministry of Economy)—FEDER through grant ESP2013-48318-C2-1R. We thank the Cambridge Astronomy Survey Unit (CASU) and the Wide Field Astronomy Unit (WFAU) in Edinburgh for providing calibrated data products under the support of the Science and Technology Facility Council (STFC) in the UK. We are grateful to P. Montegriffo for the development and maintenance of the GRATIS software. We thank the members of the OGLE team for making their catalogs public.

## APPENDIX A BAYESIAN FITTING APPROACH

This method is based on the prescription of Hogg et al. (2010), extended into three dimensions and implemented in Python using an MCMC sampler to obtain parameter estimates along with their complete posterior PDF. Initially, we model the data as being drawn from a thin plane defined by

$$f(x, y) = A x + B y + C, \quad (18)$$

where  $A$  is the slope in the  $x$  axis,  $B$  is the slope in the  $y$  axis, and  $C$  is the intercept. In this initial model we assume that we have data in three axes,  $x$ ,  $y$  and  $z$ , with errors only in the  $z$  axis.

In this model, given an independent position  $(x_i, y_i)$ , an uncertainty  $\sigma_{z_i}$ , slopes  $A$  and  $B$ , and an intercept  $C$ , the frequency distribution  $p(z_i | x_i, y_i, \sigma_{z_i}, A, B, C)$  for  $z_i$  is

$$p(z_i | x_i, y_i, \sigma_{z_i}, A, B, C) = \frac{1}{\sqrt{2\pi} \sigma_{z_i}} \times \exp\left(-\frac{[z_i - A x_i - B y_i - C]^2}{2 \sigma_{z_i}^2}\right). \quad (19)$$

Therefore the likelihood is defined as

$$\mathcal{L} = \prod_{i=1}^N p(z_i | x_i, y_i, \sigma_{z_i}, A, B, C). \quad (20)$$

Taking the logarithm,

$$\ln \mathcal{L} = K - \sum_{i=1}^N \frac{[z_i - A x_i - B y_i - C]^2}{2 \sigma_{z_i}^2}, \quad (21)$$

which is effectively the least-squares solution.  $K$  is a normalization coefficient. Returning to Bayes rule it is possible to define

$$p(A, B, C | \{z_i\}_{i=1}^N, I) = \frac{p(\{z_i\}_{i=1}^N | A, B, C, I) p(A, B, C | I)}{p(\{z_i\}_{i=1}^N | I)}, \quad (22)$$

where  $\{z_i\}_{i=1}^N$  is all the data  $z_i$ ,  $I$  is all of the information of  $x$  and  $y$ ,  $\{x_i, y_i\}_{i=1}^N$ , along with the formal errors  $\{\sigma_{x_i}, \sigma_{y_i}, \sigma_{z_i}\}_{i=1}^N$ , plus any other prior information which may be available. In our case we use uninformative (uniform) priors, making our inference method analogous to Maximum Likelihood Estimation.

### A.1 Multiple Errors, no Dispersion

Because this case contains errors in more than one axis, they can be put together into a covariance tensor  $\mathbf{S}_i$

$$\mathbf{S}_i \equiv \begin{bmatrix} \sigma_{x_i}^2 & \sigma_{xy_i} & \sigma_{xz_i} \\ \sigma_{xy_i} & \sigma_{y_i}^2 & \sigma_{yz_i} \\ \sigma_{xz_i} & \sigma_{yz_i} & \sigma_{z_i}^2 \end{bmatrix}. \quad (23)$$

With errors in several dimensions, our observed data point  $(x_i, y_i, z_i)$  could have been drawn from any true point along the plane  $(x, y, z)$ . Making the probability of the data, given the model and the true position

$$p(x_i, y_i, z_i | \mathbf{S}_i, x, y, z) = \frac{1}{2\pi \sqrt{\det(\mathbf{S}_i)}} \times \exp\left(-\frac{1}{2} [\mathbf{Z}_i - \mathbf{Z}]^\top \mathbf{S}_i^{-1} [\mathbf{Z}_i - \mathbf{Z}]\right), \quad (24)$$

where we have implicitly made column vectors

$$\mathbf{Z} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}; \quad \mathbf{Z}_i = \begin{bmatrix} x_i \\ y_i \\ z_i \end{bmatrix}. \quad (25)$$

In only two dimensions (e.g.,  $y$  and  $z$ ), the slope (e.g.,  $B$ ) can be described by a unit vector  $\hat{\mathbf{v}}$  orthogonal to the line or linear relation (at any  $x$ )

$$\hat{\mathbf{v}} = \frac{1}{\sqrt{1+B^2}} \begin{bmatrix} -B \\ 1 \end{bmatrix} = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}, \quad (26)$$

where the angle  $\theta = \arctan B$  is made between the line and the  $y$  axis. The orthogonal displacement  $\Delta_i$  of each data point  $(y_i, z_i)$  from the line is given by

$$\Delta_i = \hat{\mathbf{v}}^\top \begin{bmatrix} y_i \\ z_i \end{bmatrix} - C \cos \theta. \quad (27)$$

Instead of extending fully into three dimensions, we will assume a negligible error in  $x$  (which will be the period, so that it has justifiably higher precision). The value of  $x$  can therefore be input directly into  $\Delta_i$  without worrying about the interplay between the other parameters:

$$\Delta_i = \hat{\mathbf{v}}^\top \begin{bmatrix} y_i \\ z_i \end{bmatrix} - (C \cos \theta + A x_i). \quad (28)$$

Assuming negligible errors in  $x$  also redefines the covariance matrix of the errors as

$$\mathbf{S}_i \equiv \begin{bmatrix} \sigma_{y_i}^2 & \sigma_{yz_i} \\ \sigma_{yz_i} & \sigma_{z_i}^2 \end{bmatrix}. \quad (29)$$

Similarly, each data point's covariance matrix  $\mathbf{S}_i$  projects down to an orthogonal variance  $\Sigma_i^2$  given by

$$\Sigma_i^2 = \hat{\mathbf{v}}^\top \mathbf{S}_i \hat{\mathbf{v}} \quad (30)$$

and then the log likelihood for  $(A, B, C)$  or  $(A, \theta, C \cos \theta)$  can be written as

$$\ln \mathcal{L} = K - \sum_{i=1}^N \frac{\Delta_i^2}{2 \Sigma_i^2}, \quad (31)$$

where  $K$  is some constant. This likelihood can be maximized to find  $A, B,$  and  $C$ .

### A.2 Dispersion

The final step is to introduce an intrinsic variance in the line,  $V$ , orthogonal to the line.

According to Hogg et al. (2010), each data point can be treated as being drawn from a projected distribution function that is a convolution of the projected uncertainty Gaussian, of variance  $\Sigma_i^2$  defined above, with the Gaussian intrinsic scatter of variance  $V$ . Therefore the likelihood becomes

$$\ln \mathcal{L} = K - \sum_{i=1}^N \frac{1}{2} \ln(\Sigma_i^2 + V) - \sum_{i=1}^N \frac{\Delta_i^2}{2[\Sigma_i^2 + V]}, \quad (32)$$

where again  $K$  is a constant and everything else is defined as above. We then solve for  $A, B, C,$  and  $V$  by maximizing the log likelihood. The optimization is performed using the adaptive MCMC sampler EMCEE developed by Foreman-Mackey et al. (2013). Any optimization algorithm (e.g., Nelder-Mead, Powell, etc.) will find the maximum of the log likelihood. MCMC was chosen due to the evaluation of the full posterior PDF of the parameters, which is useful for the determination of formal errors.

## APPENDIX B METALLICITIES FOR THE MW RR LYRAE STARS ANALYZED WITH THE B–W METHOD

In Table 4 we provide a summary of spectroscopic metal abundances ( $[\text{Fe}/\text{H}]$ ) derived for the field RR Lyrae stars analyzed with the B–W technique, by studies mainly based on high-resolution spectroscopic material. Exceptions are the values with reference 1, 2, and 4 that come from compilations which include metallicities measured from low resolution spectroscopic data and photometric indices (see the discussion in Section 4.2 of Cacciari et al. 1992).

By applying our fitting approach we derived the  $\text{PM}_K Z$  relation for 23 MW RR Lyrae stars described in Section 3.3, adopting as an alternative the metallicity values from Lambert et al. (1996):

$$M_K = (-2.66 \pm 0.36) \log P + (0.05 \pm 0.04) [\text{Fe}/\text{H}] - (1.00 \pm 0.15). \quad (33)$$

The intrinsic dispersion of the relation is found to be 0.007 mag. The rms deviation of the data around the relation, neglecting the intrinsic dispersion, is 0.090 mag. The slope in  $\log P$  of the relation based on Clementini et al.'s (1995) metallicities (Equation (7)) differs from the slope obtained using Lambert et al.'s (1996) metallicities (Equation (33)) however, the values are consistent within the errors.

**Table 4**  
Literature Metallicities of the MW RR Lyrae Stars Analyzed with the B–W Method

Star	$[\text{Fe}/\text{H}]$ (dex)	$\sigma_{[\text{Fe}/\text{H}]}$ (dex)	References	
UU Cet	-1.38	0.08	(5), average of Fe I and Fe II	
	-1.33	0.08	(3), average of Fe I and Fe II	
	-0.95	0.16	(7*), re-calibration of $\Delta S$ and $\Delta S$ value adopted by (7)	
	-1.45	0.16	(7*), re-calibration of $\Delta S$ in (7) and $\Delta S$ value adopted by (5)	
	-1.20	0.20	(4)	
	-1.28		(1, 2)	
	SW And	-0.06	0.08	(5), average of Fe I and Fe II
		-0.41	0.10	from Fe I in (7)
		-0.24	0.12	from Fe II in (7)
		-0.34	0.085	average of Fe I and Fe II in (7)
	( $s = 0.12$ )			
	-0.27	0.15	(11), from Fe I and solar abundance 7.51	
	-0.24	0.15	(11), from Fe II and solar abundance 7.51	
	-0.255	0.15	(11), average of Fe I and Fe II	
	+0.20	0.08	(3), average of Fe I and Fe II	
	-0.16		(9)	
	-0.15	0.15	(4)	
	-0.24		(1, 2)	
RR Cet	-1.38	0.08	(5), average of Fe I and Fe II	
	-1.62	0.15	(11), from Fe I	
	-1.49	0.15	(11), from Fe II	
	-1.18	0.09	(3), average of Fe I and Fe II	
	-1.61		(9)	
	-1.36	0.16	(7*), re-calibration of $\Delta S$ and $\Delta S$ value adopted by (7)	
	-1.25	0.10	(4)	
X Ari	-1.45		(1, 2)	
	-2.50	0.08	(5), average of Fe I and Fe II	
	-2.19	0.17	(6), from Fe I	
	-2.54	0.09	from Fe I in (7)	
	-2.75	0.08	from Fe II in (7)	
	-2.66	0.105	average of Fe I and Fe II in (7)	
		( $s = 0.15$ )		
		-2.74	0.09	(3), average of Fe I and Fe II
	-2.61		(8), from Fe I	
	-2.62		(8), from Fe II	
	-2.68		(9)	
	-2.20	0.10	(4)	
	-2.43		(1, 2)	
AR Per	-0.34	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value adopted in (5)	
	-0.23	0.07	from Fe I in (7)	
	-0.41	0.08	from Fe II in (7)	
	-0.31	0.09	average of Fe I and Fe II in (7)	
	-0.24	0.15	(11) from Fe I	
	-0.29	0.15	(11) from Fe II	
	-0.32		(9)	
	-0.30	0.20	(4)	
-0.30		(1, 2)		
RX Eri	-1.63	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value adopted in (5)	
	-1.98	0.16	(7*), re-calibration of $\Delta S$ and $\Delta S$ value adopted in (7)	
	-1.40	0.20	(4)	
	-1.33		(1, 2)	

**Table 4**  
(Continued)

Star	[Fe/H] (dex)	$\sigma_{[\text{Fe}/\text{H}]}$ (dex)	References
RR Gem	-0.32	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value adopted in (5)
	-0.44	0.16	(7*), re-calibration of $\Delta S$ and $\Delta S$ value adopted in (7)
	-0.30	0.25	(4)
	-0.29		(1, 2)
TT Lyn	-1.64	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value adopted in (5)
	-1.63	0.08	from Fe I in (7)
	-1.33	0.06	from Fe II in (7)
	-1.44	0.150 ( $s = 0.21$ )	average of Fe I and Fe II in (7)
	-1.64	0.15	(11), from Fe I
	-1.53	0.15	(11), from Fe II
	-1.41		(9)
	-1.35	0.20	(4)
	-1.56		(1, 2)
	-1.20	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ in (5)
T Sex	-1.75	0.12	from Fe I in (7)
	-1.50	0.09	from Fe II in (7)
	-1.59	0.125 ( $s = 0.18$ )	average of Fe I and Fe II in (7)
	-1.20	0.15	(4)
	-1.34		(1, 2)
	-1.37	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value in (5)
RR Leo	-1.54	0.11	from Fe I in (7)
	-1.17	0.10	from Fe II in (7)
	-1.34	0.185 ( $s = 0.26$ )	average of Fe I and Fe II in (7)
	-1.39		(9)
	-1.15	0.20	(4)
	-1.60		(1, 2)
WY Ant	-1.32	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value in (5)
	-1.96	0.10	(10), from Fe I
	-1.96	0.10	(10), from Fe II
	-1.96	0.10	average of Fe I and Fe II in (10) + errors from us
	-1.36	0.16	(7*), re-calibration of $\Delta S$ and $\Delta S$ value in (7)
	-1.25	0.20	(4*)
W Crt	-1.48		(1, 2)
	-0.89	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value in (5)
	-0.91	0.16	(7*), re-calibration of $\Delta S$ and $\Delta S$ value in (7)
	-0.70	0.20	(4*)
TU UMa	-0.54		(1, 2)
	-1.38	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value in (5)
	-1.64	0.08	from Fe I in (7)
	-1.45	0.08	from Fe II in (7)
	-1.55	0.095 ( $s = 0.13$ )	average of Fe I and Fe II in (7)
	-1.31	0.05 or 0.14	(6), from Fe I, error of 0.05 likely a typo
	-1.72	0.15	(11), from Fe I
-1.57	0.15	(11), from Fe II	

**Table 4**  
(Continued)

Star	[Fe/H] (dex)	$\sigma_{[\text{Fe}/\text{H}]}$ (dex)	References
UU Vir	-1.46		(9)
	-1.25	0.20	(4)
	-1.51		(1, 2)
	-0.64	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value in (5)
	-0.85	0.12	from Fe I in (7)
	-0.79	0.07	from Fe II in (7)
	-0.81	0.03 ( $s = 0.04$ )	average of Fe I and Fe II in (7)
	-0.90		(9)
	-0.50	0.15	(4)
	-0.87		(1, 2)
SW Dra	-0.91	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value in (5)
	-1.37	0.15	(11), from Fe I
	-1.28	0.15	(11), from Fe II
	-0.81	0.16	(7*), re-calibration of $\Delta S$ and $\Delta S$ value in (7)
	-1.14	0.20	(4)
	-1.12		(1, 2)
RV Oct	-1.92	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value in (5)
	-1.54	0.10	from Fe I in (10) errors from us
	-1.54	0.11	from Fe II in (10) errors from us
	-1.54	0.11	average of Fe I and Fe II in (10) errors from us
	-1.98	0.16	(7*), re-calibration of $\Delta S$ in (7) and $\Delta S$ value in (5)
	-1.75	0.20	(4*)
TV Boo	-1.71		(1, 2)
	-2.31	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value in (5)
	-2.36	0.16	(7*), re-calibration of $\Delta S$ in (7) and $\Delta S$ value in (5)
	-2.30	0.15	(4)
RS Boo	-2.44		(1, 2)
	-0.37	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value in (5)
	-0.55	0.11	from Fe I in (7)
	-0.39	0.08	from Fe II in (7)
	-0.45	0.08 ( $s = 0.11$ )	average of Fe I and Fe II in (7)
	-0.48	0.15	(11), from Fe I
VY Ser	-0.33	0.15	(11), from Fe II
	-0.40	0.25	(4)
	-0.36		(1, 2)
	-1.71	0.08	(5), average of Fe I and Fe II
	-2.09	0.08	from Fe I in (7)
	-1.76	0.06	from Fe II in (7)
V445 Oph	-1.88	0.165 ( $s = 0.23$ )	average of Fe I and Fe II in (7)
	-1.71	0.07	(3), average of Fe I and Fe II
	-2.00	0.15	(11), from Fe I
	-1.90	0.15	(11), from Fe II
	-1.80	0.15	(4)
	-1.79		(1, 2)
	+0.17	0.08	(5), average of Fe I and Fe II
+0.13	0.10	(3), average of Fe I and Fe II	
	-0.26	0.16	(7*), re-calibration of $\Delta S$ in (7) and $\Delta S$ value in (7)
	+0.24		(9)
	-0.30	0.25	(4)



**Table 4**  
(Continued)

Star	[Fe/H] (dex)	$\sigma_{[\text{Fe}/\text{H}]}$ (dex)	References
	-0.19		(1, 2)
TW Her	-0.58	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value in (5)
	-0.56	0.16	(7*), re-calibration of $\Delta S$ and $\Delta S$ value in (7)
	-0.50	0.15	(4)
	-0.69		(1, 2)
AV Peg	-0.03	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value in (5)
	-0.35	0.09	from Fe I in (7)
	-0.04	0.06	from Fe II in (7)
	-0.14	0.155	average of Fe I and Fe II in (7)
	0.00	0.20	(4)
	-0.08		(1, 2)
RV Phe	-1.69	0.16	(5*), re-calibration of $\Delta S$ and $\Delta S$ value in (5)
	-1.75	0.16	(7*), re-calibration of $\Delta S$ and $\Delta S$ value in (7)
	-1.35	0.25	(4)
	-1.69		(1, 2)

**Note.** (1) [Fe/H] from Table 1 in Fernley et al. (1998a); (2) [Fe/H] values from Feast et al. (2008); (3) [Fe/H] values from Nemec et al. (2013) using the values from the VWA analysis as they are listed in column (9) of Table 7 in that paper. They are the average of the Fe I and Fe II abundances. (4) [Fe/H] values from Table 11 of Cacciari et al. (1992); (4\*) [Fe/H] values from Table 16 of Skillen et al. (1993); (5) [Fe/H] values from abundance analysis of high-resolution spectra performed by Clementini et al. (1995). Values are the average of the Fe I and Fe II measurements adopting for the solar abundance:  $\log \epsilon(\text{Fe I}) = 7.56$  and  $\log \epsilon(\text{Fe II}) = 7.50$ , respectively (see Table 12 of Clementini et al. 1995); (5\*) [Fe/H] values obtained from Clementini et al. (1995) re-calibration of the  $\Delta S$  index. (6) [Fe/H] values from Table 7 of Pancino et al. (2015). Abundances are from Fe I averaging values from different spectra of the same star as described at the end of the paper Section 4.4. The error for the abundance of TU Uma is likely a typo; according to the paper Table 6 likely should be 0.14 dex; (7) Metal abundances from Lambert et al. (1996). [Fe/H] values were derived from the photometric determinations in the paper Table 3 adopting for the Sun  $\log \epsilon(\text{Fe}) = 7.51$  (according to what stated in the footnotes of the paper Table 5); (7\*) [Fe/H] values obtained from Lambert et al.'s (1996) re-calibration of the  $\Delta S$  index (Equation (3) in that paper), which was derived by these authors using the Fe II abundances and  $\Delta S$  from Blanco (1992). Lambert et al. (1996) does not provide errors for the metallicities from  $\Delta S$ , hence we adopted an error of 0.16 dex, as done by Clementini et al. (1995) for their metallicities from  $\Delta S$ ; (8) [Fe/H] values from Table 10 of Haschke et al. (2012); (9) [Fe/H] values abundance analysis performed by Wallerstein & Huang (2010), no errors are provided; (10) [Fe/H] values from For et al. (2011) obtained as the average weighted by errors of the values in the paper Table 5. Errors are the sum in quadrature of the individual errors in Table 5 divided by the square root of  $N$  (with  $N$  number of measurements, i.e.: 11 for WY Ant and 17 for RV Oct). The average value from Fe I for WY Ant published in Table 11 of For et al. (2011). is  $-1.95$  dex while we find  $-1.96$  dex with our procedure. We also list the abundances from Fe I and Fe II separately, with errors calculated as per the above procedure: (11) [Fe/H] values from Fernley & Barnes (1996). They are from Fe II with values taken from the paper Table 4(b). Those from Fe I are taken from the paper Table 4(a). Errors are estimated by the authors to be of  $\pm 0.15$  dex in [Fe/H].

## REFERENCES

Benedict, G. F., McArthur, B. E., Feast, M. W., et al. 2011, *AJ*, 142, 187  
 Benedict, G. F., McArthur, B. E., Fredrick, L. W., et al. 2002, *AJ*, 123, 473

Blanco, V. M. 1992, *AJ*, 104, 734  
 Bono, G., Caputo, F., Castellani, V., et al. 2003, *MNRAS*, 344, 1097  
 Bono, G., Castellani, V., & Marconi, M. 2002, *ApJL*, 565, L83  
 Borissova, J., Rejkuba, M., Minniti, D., Catelan, M., & Ivanov, V. D. 2009, *A&A*, 502, 505  
 Cacciari, C., Clementini, G., & Fernley, J. A. 1992, *ApJ*, 396, 219  
 Cardelli, J. A., Clayton, C., & Mathis, J. S. 1989, *AJ*, 345, 245  
 Carney, B. W. 1996, *PASP*, 108, 900  
 Carretta, E., Bragaglia, A., Gratton, R., D'Orazi, V., & Lucatello, S. 2009, *A&A*, 508, 695  
 Carretta, E., & Gratton, R. G. 1997, *A&AS*, 121, 95  
 Catelan, M., Pritzl, B. J., & Smith, H. A. E. 2004, *ApJS*, 154, 633  
 Cioni, M.-R. L., Clementini, G., Girardi, L., et al. 2011, *A&A*, 527, A116  
 Clementini, G. 2008, in IAU Symposium 256, The Magellanic System: Stars, Gas, and Galaxies, ed. J. T. van Loon & J. M. Oliveira (Cambridge: Cambridge Univ. Press), 373  
 Clementini, G., Di Tomaso, S., Di Fabrizio, L., et al. 2000, *AJ*, 120, 2054  
 Clementini, G., Gratton, R., Merighi, R., et al. 1995, *AJ*, 110, 2319  
 Clementini, G., Gratton, R. G., Bragaglia, A., et al. 2003, *AJ*, 125, 1309  
 Coppola, G., Dall'Orta, M., Ripepi, V., et al. 2011, *MNRAS*, 416, 1056  
 Cutri, R. M., Krutskie, M. F., van Dyk, S., et al. 2003, 2MASS All Sky Catalog of Point Sources (Pasadena, CA: NASA/IPAC)  
 Dall'Orta, M., Storm, J., Bono, G., et al. 2004, *ApJ*, 610, 269  
 De Grijs, R., Wicker, J. E., & Bono, G. 2014, *AJ*, 147, 122  
 Del Principe, M. 2006, *AJ*, 652, 362  
 Di Fabrizio, L., Clementini, G., Maio, M., et al. 2005, *A&A*, 430, 603  
 Feast, M. W., Clifton, D. L., Kinman, T. D., van Leeuwen, F., & Whitelock, P. A. 2008, *MNRAS*, 386, 2115  
 Fernley, J. A. 1994, *A&A*, 284, L16  
 Fernley, J. A., & Barnes, T. G. 1996, *A&A*, 312, 957  
 Fernley, J. A., Barnes, T. G., Skillen, I., et al. 1998a, *A&A*, 330, 515  
 Fernley, J. A., Carney, B. W., Skillen, I., Cacciari, C., & Janes, K. 1998b, *MNRAS*, 293, L61  
 Fernley, J. A., Skillen, I., Jameson, R. F., et al. 1990, *MNRAS*, 247, 287  
 Fitzpatrick, E. L., Ribas, I., Guinan, E. F., Maloney, F. P., & Claret, A. 2003, *ApJ*, 587, 685  
 Fitzpatrick, E. L., Ribas, I., Guinan, E. F., et al. 2002, *ApJ*, 564, 260  
 For, B.-Q., Sneden, C., & Preston, G. W. 2011, *ApJS*, 197, 29  
 Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306  
 Freedman, W. J., Madore, B. F., Gibson, B. K., et al. 2001, *ApJ*, 553, 47  
 Gibson, B. K. 2000, *MmSAI*, 71, 693  
 Gratton, R. G., Bragaglia, A., Clementini, G., et al. 2004, *A&A*, 421, 937  
 Guinan, E. F., Fitzpatrick, E. L., Dewarf, L. E., et al. 1998, *ApJL*, 509, L21  
 Hanson, R. B. 1979, *MNRAS*, 186, 875  
 Haschke, R., Grebel, E. K., Frebel, A., et al. 2012, *AJ*, 144, 88  
 Hogg, D. W., Bovy, J., & Lang, D. 2010, arXiv:1008.4686  
 Jones, R. V., Carney, B. W., & Fulbright, J. P. 1996, *PASP*, 108, 877  
 Jones, R. V., Carney, B. W., & Latham, D. W. 1988, *ApJ*, 332, 206  
 Jones, R. V., Carney, B. W., Storm, J., & Latham, D. W. 1992, *ApJ*, 386, 646  
 Jordi, C., Gebran, M., Carrasco, J. M., et al. 2010, *A&A*, 523, A48  
 Keller, S. C., & Wood, P. R. 2002, *ApJ*, 578, 144  
 Keller, S. C., & Wood, P. R. 2006, *ApJ*, 642, 834  
 Lambert, D. L., Heath, J. E., Lemke, M., & Drake, J. 1996, *ApJS*, 103, 183  
 Liu, T., & Janes, K. A. 1990, *ApJ*, 354, 273  
 Longmore, A. J., Fernley, J. A., & Jameson, R. F. 1986, *MNRAS*, 220, 279  
 Luri, X., Palmer, M., Arenou, F., et al. 2014, *A&A*, 566, A119  
 Lutz, T. E., & Kelker, D. H. 1973, *PASP*, 85, 573  
 Malmquist, K. 1936, *Stockholms Ob. Medd.*, 12, 26  
 Marconi, M., & Clementini, G. 2005, *AJ*, 129, 2257  
 McNamara, D. H., Clementini, G., & Marconi, M. 2007, *AJ*, 133, 2752  
 Moretti, M. I., Clementini, G., Muraveva, T., et al. 2014, *MNRAS*, 437, 2702  
 Nemec, J. M., Cohen, J. G., Ripepi, V., et al. 2013, *ApJ*, 773, 181  
 Nemec, J. M., Nemec, A. F. L., & Lutz, T. E. 1994, *AJ*, 108, 222  
 Pancino, E., Britavskiy, N., Romano, D., et al. 2015, *MNRAS*, 447, 2404  
 Perryman, M. A. C., de Boer, K. S., Gilmore, G., et al. 2001, *A&A*, 369, 339  
 Perryman, M. A. C., & ESA 1997, The Hipparcos and Tycho Catalogues (ESA SP-1200; Noordwijk: ESA)  
 Pietrzyński, G., Graczyk, D., Gieren, W., et al. 2013, *Natur*, 495, 76  
 Preston, G. W. 1959, *ApJ*, 130, 507  
 Ribas, I., Fitzpatrick, E. L., Maloney, F. P., Guinan, E. F., & Udalski, A. 2002, *ApJ*, 574, 771  
 Ripepi, V., Moretti, M. I., Clementini, G., et al. 2012a, *Ap&SS*, 341, 51  
 Ripepi, V., Moretti, M. I., Marconi, M., et al. 2012b, *MNRAS*, 424, 1807

- Rubele, S., Kerber, L., Girardi, L., et al. 2012, *A&A*, **537**, A106
- Schaefer, B. E. 2008, *AJ*, **135**, 112
- Schaefer, B. E. 2013, *Natur*, **495**, 51
- Skillen, I., Fernley, J. A., Stobie, R. S., & Jameson, R. F. 1993, *MNRAS*, **265**, 301
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, **131**, 1163
- Sollima, A., Cacciari, C., & Valenti, E. 2006, *MNRAS*, **372**, 1675
- Sollima, A., Cacciari, C., Arkharov, A. A. H., et al. 2008, *MNRAS*, **384**, 1583
- Soszyński, I., Udalski, A., Szymański, M. K., et al. 2009, *AcA*, **59**, 1
- Staetson, P. B., Davis, L. E., & Crabtree, D. R. 1990, in ASP Conf. Ser. 8, CCDs, in Astronomy ed. G. H. Jacoby (San Francisco, CA: ASP), 289
- Sturch, C. 1966, *ApJ*, **143**, 774
- Szewczyk, O., Pietrzyński, G., Gieren, W., et al. 2008, *AJ*, **136**, 272
- Walker, A. R. 2012, *Ap&SS*, **341**, 43
- Wallerstein, G., & Huang, W. 2010, *MmSAI*, **81**, 952
- van Leeuwen, F. 2007, *Hipparcos*, the New Reductions of the Raw Data (Berlin: Springer)
- Zinn, R., & West, M. J. 1984, *ApJS*, **55**, 45