



<b>Publication Year</b>	2015
<b>Acceptance in OA @INAF</b>	2020-05-11T18:04:15Z
<b>Title</b>	Real-time imaging of density ducts between the plasmasphere and ionosphere
<b>Authors</b>	Loi, Shyeh Tjing; Murphy, Tara; Cairns, Iver H.; Menk, Frederick W.; Waters, Colin L.; et al.
<b>DOI</b>	10.1002/2015GL063699
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/24710">http://hdl.handle.net/20.500.12386/24710</a>
<b>Journal</b>	GEOPHYSICAL RESEARCH LETTERS
<b>Number</b>	42

## RESEARCH LETTER

10.1002/2015GL063699

## Key Points:

- Vast arrays of regularly spaced, field-aligned ducts are common at midlatitudes
- Widefield, low-frequency radio telescopes are powerful probes of the ionosphere
- New imaging technique allows regional-scale monitoring of plasma dynamics in 3-D

## Supporting Information:

- Movie S2
- Movie S1
- Text S1 and Movies S1 and S2 captions

## Correspondence to:

S. T. Loi,  
sloi5113@uni.sydney.edu.au

## Citation:

Loi, S. T., et al. (2015), Real-time imaging of density ducts between the plasmasphere and ionosphere, *Geophys. Res. Lett.*, *42*, 3707–3714, doi:10.1002/2015GL063699.

Received 3 MAR 2015

Accepted 21 APR 2015

Accepted article online 11 MAY 2015

Published online 25 MAY 2015

## Real-time imaging of density ducts between the plasmasphere and ionosphere


Shyeh Tjing Loi<sup>1,2</sup>, Tara Murphy<sup>1,2</sup>, Iver H. Cairns<sup>1</sup>, Frederick W. Menk<sup>3</sup>, Colin L. Waters<sup>3</sup>, Philip J. Erickson<sup>4</sup>, Cathryn M. Trott<sup>2,5</sup>, Natasha Hurley-Walker<sup>5</sup>, John Morgan<sup>5</sup>, Emil Lenc<sup>1,2</sup>, André R. Offringa<sup>2,6,7</sup>, Martin E. Bell<sup>2,8</sup>, Ronald D. Ekers<sup>8</sup>, B. M. Gaensler<sup>1,2,9</sup>, Colin J. Lonsdale<sup>4</sup>, Lu Feng<sup>10</sup>, Paul J. Hancock<sup>2,5</sup>, David L. Kaplan<sup>11</sup>, G. Bernardi<sup>12,13,14</sup>, J. D. Bowman<sup>15</sup>, F. Briggs<sup>2,6</sup>, R. J. Cappallo<sup>4</sup>, A. A. Deshpande<sup>16</sup>, L. J. Greenhill<sup>13</sup>, B. J. Hazelton<sup>17</sup>, M. Johnston-Hollitt<sup>18</sup>, S. R. McWhirter<sup>4</sup>, D. A. Mitchell<sup>2,8</sup>, M. F. Morales<sup>17</sup>, E. Morgan<sup>10</sup>, D. Oberoi<sup>19</sup>, S. M. Ord<sup>2,5</sup>, T. Prabu<sup>16</sup>, N. Udaya Shankar<sup>16</sup>, K. S. Srivani<sup>16</sup>, R. Subrahmanyam<sup>2,16</sup>, S. J. Tingay<sup>2,5</sup>, R. B. Wayth<sup>2,5</sup>, R. L. Webster<sup>2,20</sup>, A. Williams<sup>5</sup>, and C. L. Williams<sup>10</sup>

<sup>1</sup>Sydney Institute for Astronomy, School of Physics, University of Sydney, Sydney, New South Wales, Australia, <sup>2</sup>ARC Centre of Excellence for All-sky Astrophysics, Redfern, New South Wales, Australia, <sup>3</sup>School of Mathematical and Physical Sciences, University of Newcastle, Callaghan, New South Wales, Australia, <sup>4</sup>MIT Haystack Observatory, Westford, Massachusetts, USA, <sup>5</sup>International Centre for Radio Astronomy Research, Curtin University, Bentley, Western Australia, Australia, <sup>6</sup>Research School of Astronomy and Astrophysics, Australian National University, Canberra, Australian Capital Territory, Australia, <sup>7</sup>Netherlands Institute for Radio Astronomy, Dwingeloo, Netherlands, <sup>8</sup>CSIRO Astronomy and Space Science, Epping, New South Wales, Australia, <sup>9</sup>Dunlap Institute for Astronomy and Astrophysics, University of Toronto, Toronto, Ontario, Canada, <sup>10</sup>Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA, <sup>11</sup>Department of Physics, University of Wisconsin-Milwaukee, Milwaukee, Wisconsin, USA, <sup>12</sup>Square Kilometre Array South Africa, Cape Town, South Africa, <sup>13</sup>Harvard-Smithsonian Center for Astrophysics, Cambridge, Massachusetts, USA, <sup>14</sup>Department of Physics and Electronics, Rhodes University, Grahamstown, South Africa, <sup>15</sup>School of Earth and Space Exploration, Arizona State University, Tempe, Arizona, USA, <sup>16</sup>Raman Research Institute, Bangalore, India, <sup>17</sup>Department of Physics, University of Washington, Seattle, Washington, USA, <sup>18</sup>School of Chemical and Physical Sciences, Victoria University of Wellington, Wellington, New Zealand, <sup>19</sup>National Centre for Radio Astrophysics, Tata Institute for Fundamental Research, Pune, India, <sup>20</sup>School of Physics, University of Melbourne, Parkville, Victoria, Australia

**Abstract** Ionization of the Earth's atmosphere by sunlight forms a complex, multilayered plasma environment within the Earth's magnetosphere, the innermost layers being the ionosphere and plasmasphere. The plasmasphere is believed to be embedded with cylindrical density structures (ducts) aligned along the Earth's magnetic field, but direct evidence for these remains scarce. Here we report the first direct wide-angle observation of an extensive array of field-aligned ducts bridging the upper ionosphere and inner plasmasphere, using a novel ground-based imaging technique. We establish their heights and motions by feature tracking and parallax analysis. The structures are strikingly organized, appearing as regularly spaced, alternating tubes of overdensities and underdensities strongly aligned with the Earth's magnetic field. These findings represent the first direct visual evidence for the existence of such structures.

## 1. Introduction

The plasmasphere is a toroidal region within the Earth's magnetosphere that is filled with plasma from the ionosphere, which is a photoionized layer near the surface of the Earth [Goldstein, 2006]. The existence of field-aligned ducts in the plasmasphere is widely accepted as the explanation for ground-based detections of whistler-mode waves, which may be guided along such ducts [Ohta et al., 1996; Bakharev et al., 2010]. Wave-particle interactions that accelerate and precipitate particles into the atmosphere occur efficiently within these ducts, so their existence aids the removal of energetic particles from the magnetosphere and thereby influences global magnetospheric-ionospheric coupling and energetics [Sonwalkar, 2006]. Duct properties have been studied mainly through indirect, ground-based observations of whistlers from both natural sources (lightning) and man-made VLF transmitters [Singh et al., 1998; Ganguli et al., 2000; Clilverd et al., 2003]. Since the guiding structures cannot be detected independently of the whistlers, observations that use lightning-generated whistlers are necessarily intermittent, and those that use man-made whistlers




are spatially restricted. Direct detections of plasmaspheric ducts have been achieved by satellite in situ measurements [Sonwalkar *et al.*, 1994], but these and also indirect satellite-based measurements such as sounding [Darrouzet *et al.*, 2009; Yearby *et al.*, 2011; Karpachev *et al.*, 2014] are heavily restricted by spacecraft orbital patterns. Duct morphologies have thus been difficult to establish.

The Murchison Widefield Array (MWA) [Lonsdale *et al.*, 2009; Tingay *et al.*, 2013] is a low-frequency (80–300 MHz) radio telescope located in Western Australia at 26°42'12''S, 116°40'15''E (geographic), and a geomagnetic latitude of 38.6°S (McIlwain  $L$  parameter [McIlwain, 1961] of 1.6). Although chiefly intended for studying the cosmos, it is also an exquisitely sensitive probe of the near-Earth plasma. Novel design concepts include its wide field of view (FoV) and high-cadence imaging capability. A multitude of unresolved celestial radio sources (mostly radio galaxies and quasars) back-illuminate the plasma, allowing one to probe fluctuations at high spatial completeness by measuring the angular distribution of source refractive shifts. Since interferometers measure angular positions using baseline phase differences, they are insensitive to the constant offset component of the electron column density (“total electron content” (TEC)), which adds an equal phase to all antennas and cancels out. Rather, the refraction-induced angular shift  $\Delta\theta$  of a radio source depends on the transverse gradient  $\nabla_{\perp}$  TEC in the TEC toward the source and is given by [Thompson *et al.*, 2001]

$$\Delta\theta = -\frac{1}{8\pi^2} \frac{e^2}{\epsilon_0 m_e v^2} \nabla_{\perp} \text{TEC}. \quad (1)$$

**Figure 1.** The vector fields of celestial source offsets for the (a) first, (b) middle, and (c) last snapshots, overplotted with the geomagnetic field lines (black solid lines) from a single  $L$  shell. Field line positions were computed based on the Australian Geomagnetic Reference Field model [Geoscience Australia, 2010] at an altitude of 600 km. Assumed to be locally straight and parallel, their apparent convergence arises from a perspective distortion due to their steep ( $-60^\circ$ ) inclination. The center of each arrow marks the time-averaged position (used as a reference point) of a celestial radio source, and the arrow represents its displacement vector. Arrow lengths have been scaled to 50 times the actual displacement distance. Blue and red denote arrows with positive and negative  $x$  components (westward and eastward), respectively. North is up and east is to the left as per the astronomical convention, and  $x = y = 0^\circ$  marks the location of the zenith. Units along the  $x$  and  $y$  axes correspond to angular distance across the MWA FoV ( $1^\circ \leftrightarrow 10$  km at 600 km altitude).



**Figure 2.** The vector divergence plot for the middle snapshot (i.e., the divergence of the vector field shown in Figure 1b), with the duct structure clearly visible. Positive divergence regions (relative overdensities) are represented by red-yellow hues, while negative divergence regions (relative underdensities) are represented by blue-cyan hues. White represents either values close to zero (within the FoV) or non-numeric (outside the FoV). Geomagnetic field lines have been overlotted as black solid lines, as for Figure 1.

date sources to a number of quality restrictions and then cross matched the remainder with the National Radio Astronomy Observatory Very Large Array (VLA) Sky Survey (NVSS) catalog [Condon *et al.*, 1998], a published database of celestial radio sources, retaining only those candidates with counterparts in NVSS. Cross matching was performed by identifying the nearest NVSS source to the candidate source within the search radius (3 arc min) and associating it to the latter.

We took the reference position of a source, uniquely identified by its NVSS catalog name, to be the average of its celestial coordinates (measured by AEGEAN) over all snapshots. Then for each snapshot, we computed the angular offset vector of each source from its reference position. The distributions of these offsets (equivalently the TEC gradient field) are shown in Figure 1 for several snapshots (see Movie S1 in the supporting information for an animation of the full data set). Strikingly organized bands of arrows are visible, with a characteristic band separation of  $\sim 2^\circ$ . These reflect a spatially oscillating gradient field, implying alternating bands of underdensities and overdensities. The bands are tightly aligned with the Earth's magnetic field. Taking the divergence of the vector field reveals the duct structure more prominently. This is shown in Figure 2 for a representative snapshot (see Movie S2 for an animation of the full data set).


The measured offsets display a  $\lambda^2$  proportionality (Figure 3a) consistent with equation (1), where  $\lambda$  is the observing wavelength. Source motions exceed measurement errors and are often larger than the width of the telescope point spread function (119 arc sec at 183 MHz). Quasi-sinusoidal oscillations of radio sources are apparent; the time series for one source, with errors, is shown in Figure 3b. Note that celestial sources drift across the sky much faster than do the structures, which are almost stationary above the MWA. Their characteristic TEC gradient (a measure of the amplitude), obtained from the mean angular offset using equation (1), increases over time and roughly doubles in 1.5 h (Figure 3c). This is comparable to the growth timescales predicted and reported for whistler ducts in the literature [Singh *et al.*, 1998; McCormick, 2002].

We estimated the altitude, motion, and orientation of the structures by using the MWA as a stereo camera. The MWA consists of 128 receiving elements ("tiles") spread over an area about 3 km across (Figure 4a). By dividing the array east-west and imaging using the two groups of tiles separately, we measured a parallax shift (Figure 4b) that implies a characteristic altitude of  $570 \pm 40$  km ( $L = 1.8$ ), averaged over the interval and the FoV. This is within the  $L$  range previously inferred for similar structures [Jacobson and Erickson, 1993; Hoogeveen and Jacobson, 1997].

Here  $e$  and  $m_e$  are the electron charge and mass,  $\epsilon_0$  is the vacuum permittivity, and  $\nu$  is the radio observing frequency. The negative sign indicates that the direction of refraction is toward decreasing TEC.

## 2. Observations and Results

We analyzed MWA data comprising 46 snapshots, each integrated for 2 min at a frequency of 183 MHz over a 30.72 MHz bandwidth, of a  $30^\circ \times 50^\circ$  (N-S  $\times$  E-W) patch of sky near zenith. These were recorded over a 1.5 h interval on 15 October 2013 between 1346 and 1517 UTC (premidnight local time), in which the telescope was tracking celestial coordinates RA =  $0^\circ$ , Dec =  $-27^\circ$  (J2000). We measured the refractive shifts of the  $\sim 1000$  celestial sources visible in each  $\sim 30^\circ$  wide snapshot by fitting to the positions of the same sources seen in each snapshot and computing offsets from the time-averaged positions. In more detail, the source-finding software AEGEAN [Hancock *et al.*, 2012] was first used to locate intensity peaks (candidate radio sources) in the images. We subjected the candi-




**Figure 3.** (a) Wavelength ( $\lambda$ ) dependence of the size of the offset for five bright radio sources, whose flux densities are shown in the figure legend ( $1 \text{ Jy} \equiv 10^{-26} \text{ W m}^{-2} \text{ Hz}^{-1}$ ). Each set of points represents measurements taken in a single snapshot of time. Straight lines represent best fits to measurements at four different frequencies within the 30 MHz observing bandwidth, constrained to pass through the origin. (b) East-west component of the displacement of one of the sources in Figure 3a as a function of time. Different colors represent measurements at four independent frequencies. Dashed lines indicate the width of the telescope point spread function. Error bars in Figures 3a and 3b represent position fitting errors. (c) Time dependence of the average magnitude of the TEC gradient over all sources in a snapshot, for images formed using the full 30 MHz band.

A parallax analysis is only meaningful for structures that are localized in altitude. However, the MWA measures the spatial derivative of a column-integrated quantity and so is sensitive to irregularities over a wide range of altitudes. An argument against the structures being extended in altitude, and thus for the applicability of the parallax analysis, comes from the observation that the bands are still prominent up to zenith angles of  $\zeta \sim 20^\circ$ . If they were extended in altitude (like vertical sheets), then oblique lines of sight would pierce multiple sheets, causing the features to blend into one another and wash out. Clear bands in the  $\nabla_{\perp} \text{TEC}$  vector field exist out to  $\zeta \sim 20^\circ$ , implying that the irregularities cannot be extended in altitude by more than  $\cot \zeta \approx 3$  times the interband separation, i.e., that their horizontal and vertical cross-sectional widths are comparable. This further supports the cylindrical duct interpretation, as opposed to vertical sheet-like structures. The contribution to the TEC from the smooth plasma above and below the irregularity layer cancels away when the horizontal spatial derivative is taken, and so the situation is observationally equivalent to a thin layer of plasma embedded with irregularities that is bounded above and below by vacuum. This supports the validity of the parallax technique and explains why we obtained a well-defined altitude upon measurement.

Computing the parallax separately over the first and second halves of the interval establishes a downward drift from  $720 \pm 90 \text{ km}$  to  $470 \pm 40 \text{ km}$  with time. Splitting the data spatially into northern and southern halves of the FoV reveals larger altitudes to the north. This is consistent with the steep magnetic inclination at the MWA site ( $-60^\circ$ ) and implies that the structures stretch between  $\sim 400$  and  $1000 \text{ km}$  within the FoV, likely extending above this. Thus, they bridge the topside ionosphere and inner plasmasphere, and so the transition from a neutral- to plasma-dominated medium. These prop-

erties are consistent with the altitudes that whistler ducts are expected to occupy at night [Sonwalkar, 2006]. The physical spacings of the tubes are then  $10\text{--}50 \text{ km}$ , and associated TEC fluctuations between about  $0.1$  and



**Figure 4.** (a) Layout of the 128 MWA tiles, with hollow square and circle markers highlighting the two groups of tiles used for measuring the parallax shift. Central core tiles (crosses) were excluded to maximize the effective baseline (~890 m), leaving 37 tiles for the east half and 35 for the west. (b) Histograms of the parallax shifts measured for the east and west halves, each from 92 independent measurements (46 snapshots  $\times$  two instrumental polarizations). The relative displacement of the two histograms represents a parallax of  $5.3 \pm 0.4$  arc min.

for the Earth’s magnetic field, radial motions of ducts and their  $L$  shell values can be deduced from whistler spectrograms, but zonal drifts cannot be similarly established [Saxton and Smith, 1989; Carpenter and Smith, 2001].

Claims of regularly spaced, regional-scale, field-aligned density structures with properties similar to those seen here have been made based on observations by the Very Large Array (VLA) radio interferometer [Jacobson and Erickson, 1993; Helmboldt et al., 2012a, 2012b] and the Los Alamos plasmaspheric drift radio interferometer [Jacobson et al., 1996; Hoogeveen and Jacobson, 1997]. Although these studies could measure zonal drifts, identification of the structures relied on fitting plane waves to sparsely sampled measurements, a model whose suitability could not be independently verified. The assumption that the irregularities were field aligned, together with a model for the Earth’s magnetic field, is required to infer their altitudes. Many of these assumptions are unnecessary here. The sampling completeness of the MWA surpasses that of the VLA and Los Alamos instruments by a factor of  $\sim 100$ , making it the first radio telescope capable of imaging the plasma. This has allowed us to visualize for the first time the regional-scale, field-aligned nature of these density structures and to demonstrate a clear and remarkable spatial periodicity in their layout, without any prior assumptions about their morphology. Unlike these and other ground-based studies, our parallax technique to establish the altitudes requires no knowledge of the Earth’s magnetic field. By removing the reliance on wave transmission for detection of the ducts, they may be monitored passively and continuously for long durations, including periods when structural distortions or imperfections may prevent them from sustaining wave propagation.

0.7 total electron content units (TECU) ( $1 \text{ TECU} = 10^{16} \text{ el m}^{-2}$ ), again in agreement with inferred properties of whistler ducts [Sonwalkar, 2006].

### 3. Discussion

#### 3.1. Previous Capabilities

While the hypothesized existence of field-aligned density ducts accounts for a large body of experimental data [Ohta et al., 1996; Singh et al., 1998; Carpenter and Smith, 2001], it has not been previously possible to directly verify that large, cylindrical density structures greatly extended along the Earth’s magnetic field lines exist or to track their motions precisely and continuously over regional scales. Satellite in situ observations [Sonwalkar et al. 1994] can only measure densities at single points and cannot instantaneously probe the regional-scale density distribution. Topside sounding data and whistler spectrograms have been interpreted in terms of signal guidance by field-aligned ducts [Darrouzet et al., 2009; Yearby et al., 2011], but such methods rely on assumptions about wave propagation [Singh et al., 1998], and detections are only possible under suitable conditions. They also suffer from spatial restrictions imposed by satellite orbital paths and the placements of ground-based receivers. With a model

### 3.2. Formation Mechanism

Plasmaspheric density irregularities appear more often during geomagnetically disturbed periods, but the underlying formation mechanisms are not well established [Hoogeveen and Jacobson, 1997; Darrouzet et al., 2009]. Geomagnetic conditions during the interval were mildly unsettled with a  $K_p$  index of 2, the observations taking place in the recovery phase of a moderate storm (maximum ring current of  $Dst = -45$  nT near 04 UTC, 24 h maximum  $K_p$  of 4). Theoretical studies indicate that ripples and undulations in the plasmaspheric electron density may be produced by an interchange instability that preferentially occurs just after periods of strong magnetospheric convection [Sazykin et al., 2004; Buzulukova et al., 2008]. Such ripples have been observed on a global scale in satellite observations [Goldstein, 2005]. Although below the spatial scale of the grids used in those simulation studies, smaller duct-like structures such as those seen in our data may form if this mechanism continues to operate on smaller spatial scales. We note that field-aligned structures of lesser prominence and multiplicity often appear in MWA data under quieter geomagnetic conditions. They are visible on around half of all nights inspected so far, most of which (with the exception of the current data set) were not obtained during periods of storm time activity.

Organized azimuthal structures in the outer plasmasphere have been attributed to  $\mathbf{E} \times \mathbf{B}$  convection under the influence of ultralow frequency (ULF) standing waves [Adrian et al., 2004]. Although the spatial scales of ULF waves greatly exceed those observed here, small-scale density structures can grow by flux tube interchange under the imposition of a much larger-scale electric field [Rodger et al., 1998]. The steady downward transport established above is consistent with  $\mathbf{E} \times \mathbf{B}$  convection for a westward electric field of  $\sim 1$  mV m<sup>-1</sup>, a reasonable value for periods of substorm activity [Carpenter and Smith, 2001]. Quasiperiodic, field-aligned structures called plasma bubbles form through a related process, where traveling ionospheric disturbances (TIDs) trigger the Rayleigh-Taylor instability [Buhari et al., 2014]. However, plasma bubble formation is only viable near the equator where near-horizontal geomagnetic field lines can support the plasma against gravity and not at midlatitudes. The fluctuations are unlikely to be TIDs themselves, which have a dispersion relation [Shiokawa et al., 2013] grossly inconsistent with the spatial and temporal properties measured here. Furthermore, TID-like fluctuations routinely appear in MWA data, and these exhibit markedly different properties from those seen here.

Spatially varying electric fields in the plasmasphere may cause localized compressions, squeezing plasma down selected flux tubes [Park and Banks, 1974]. Secondary ionization from particle precipitation may also produce flux tube selective densifications. Although natural precipitation events are largely confined to the auroral zones, the powerful VLF navigation beacon at North West Cape ( $\sim 400$  km north of the MWA) is known to cause electron precipitation concentrated toward the south [Parrot et al., 2007]. A Fourier analysis of our data reveals the existence of structures drifting in opposite directions. Components with larger spatial frequencies are observed to drift eastward, while those with smaller spatial frequencies drift westward. We measure drift speeds within 2% of corotation, consistent with these being in the inner plasmasphere where corotation is relatively strict [Galvan et al., 2010]. The regular spacing of the structures could have arisen from a Kelvin-Helmholtz instability driven by shear in the plasma and thereafter amplified through an interchange instability.

### 4. Conclusion and Outlook

Despite the seeming ubiquity of field-aligned ducts in magnetospheric systems (e.g., Io-Jupiter [Imai et al., 1992] and the Sun [Duncan, 1979]), self-organization processes in plasmas have been difficult to isolate observationally. Our results demonstrate that widefield radio telescopes such as the MWA are powerful quantitative tools for studying their formation, dynamics, and morphology. Radio telescopes differ fundamentally from many approaches for probing the ionosphere and plasmasphere in that they measure density gradients rather than absolute density. Their insensitivity to the constant offset component of the TEC makes them particularly suitable for studying plasma density disturbances/irregularities and permits them to probe regions above the peak electron density with ease, since they are not significantly shielded by the denser underlying plasma.

The ground-based feature tracking and altitude triangulation capabilities we have demonstrated here offer valuable opportunities for the real-time, regional-scale monitoring of inner magnetospheric structures and dynamics on a near-continuous basis, unconstrained by the limitations of spacecraft orbits or the propagation of whistlers. Unlike ground-based whistler observations, which only allow for radial motions to be measured, the MWA can track both radial and horizontal motions, thereby allowing bulk plasma drifts to be

characterized in three dimensions. Its potential to perform 3-D reconstruction of density structures can provide empirical constraints on the plasma distribution both along and across magnetic flux tubes. This may be useful, for example, for specifying appropriate boundary conditions in simulation studies of global plasma flows [Tu *et al.*, 2006].

#### Acknowledgments

The data supporting this paper are available upon request submitted via email to the corresponding author at sloi5113@uni.sydney.edu.au. This scientific work makes use of the Murchison Radio-astronomy Observatory, operated by CSIRO. We acknowledge the Wajarri Yamatji people as the traditional owners of the Observatory site. Support for the MWA comes from the U.S. National Science Foundation (grants AST-0457585, PHY-0835713, CAREER-0847753, AST-0908884, and AST-1412421), the Australian Research Council (LIEF grants LE0775621 and LE0882938), the U.S. Air Force Office of Scientific Research (grant FA9550-0510247), and the Centre for All-sky Astrophysics (an Australian Research Council Centre of Excellence funded by grant CE110001020). Support is also provided by the Smithsonian Astrophysical Observatory, the MIT School of Science, the Raman Research Institute, the Australian National University, and the Victoria University of Wellington (via grant MED-E1799 from the New Zealand Ministry of Economic Development and an IBM Shared University Research grant). The Australian Federal government provides additional support via the Commonwealth Scientific and Industrial Research Organisation (CSIRO), National Collaborative Research Infrastructure Strategy, Education Investment Fund, and the Australia India Strategic Research Fund, and Astronomy Australia Limited, under contract to Curtin University. We acknowledge the iVEC Petabyte Data Store, the Initiative in Innovative Computing and the CUDA Center for Excellence sponsored by NVIDIA at Harvard University, and the International Centre for Radio Astronomy Research (ICRAR), a Joint Venture of Curtin University and The University of Western Australia, funded by the Western Australian State government. We thank D. B. Melrose for discussions regarding physical interpretations.

The Editor thanks Vijay Harid and an anonymous reviewer for their assistance in evaluating this paper.

#### References

- Adrian, M. L., D. L. Gallagher, and L. A. Avakov (2004), IMAGE EUV observation of radially bifurcated plasmaspheric features: First observations of a possible standing ULF waveform in the inner magnetosphere, *J. Geophys. Res.*, *109*, A01203, doi:10.1029/2003JA009974.
- Bakharev, P. V., T. M. Zaboronkova, A. V. Kudrin, and C. Krafft (2010), Whistler waves guided by density depletion ducts in a magnetoplasma, *Plasma Phys. Rep.*, *36*(11), 919–930, doi:10.1134/S1063780X10110012.
- Buhari, S. M., M. Abdullah, A. M. Hasbi, Y. Otsuka, T. Yokoyama, M. Nishioka, and T. Tsugawa (2014), Continuous generation and two-dimensional structure of equatorial plasma bubbles observed by high-density GPS receivers in Southeast Asia, *J. Geophys. Res. Space Physics*, *119*, 10,569–10,580, doi:10.1002/2014JA020433.
- Buzulukova, N., M.-C. Fok, T. E. Moore, and D. M. Ober (2008), Generation of plasmaspheric undulations, *Geophys. Res. Lett.*, *35*, L13105, doi:10.1029/2008GL034164.
- Carpenter, D. L., and A. J. Smith (2001), The study of bulk plasma motions and associated electric fields in the plasmasphere by means of whistler-mode signals, *J. Atmos. Sol. Terr. Phys.*, *63*, 1117–1132, doi:10.1016/S1364-6826(00)00217-0.
- Cliilverd, M. A., et al. (2003), In situ and ground-based intercalibration measurements of plasma density at  $L = 2.5$ , *J. Geophys. Res.*, *108*(A10), 1365, doi:10.1029/2003JA009866.
- Condon, J. J., W. D. Cotton, E. W. Greisen, Q. F. Yin, R. A. Perley, G. B. Taylor, and J. J. Broderick (1998), The NRAO VLA sky survey, *Astron. J.*, *115*, 1693–1716.
- Darrouzet, F., et al. (2009), Plasmaspheric density structures and dynamics: Properties observed by the CLUSTER and IMAGE missions, *Space Sci. Rev.*, *145*(1–2), 55–106, doi:10.1007/s11214-008-9438-9.
- Duncan, R. A. (1979), Wave ducting of solar metre-wave radio emission as an explanation of fundamental/harmonic source coincidence and other anomalies, *Sol. Phys.*, *63*, 389–398, doi:10.1007/BF00174543.
- Galvan, D. A., M. B. Moldwin, B. R. Sandel, and G. Crowley (2010), On the causes of plasmaspheric rotation variability: IMAGE EUV observations, *J. Geophys. Res.*, *115*, A01214, doi:10.1029/2009JA014321.
- Ganguli, G., M. A. Reynolds, and M. W. Liemohn (2000), The plasmasphere and advances in plasmaspheric research, *J. Atmos. Sol. Terr. Phys.*, *62*, 1647–1657, doi:10.1016/S1364-6826(00)00117-6.
- Geoscience Australia (2010), Australian Geomagnetic Field Reference Values, Geoscience Australia. [Available at <http://www.ga.gov.au/metadata-gateway/metadata/record/71108/>].
- Goldstein, J. (2005), Coupled response of the inner magnetosphere and ionosphere on 17 April 2002, *J. Geophys. Res.*, *110*, A03205, doi:10.1029/2004JA010712.
- Goldstein, J. (2006), Plasmasphere response: Tutorial and review of recent imaging results, *Space Sci. Rev.*, *124* (1–4), 203–216, doi:10.1007/s11214-006-9105-y.
- Hancock, P. J., T. Murphy, B. M. Gaensler, A. Hopkins, and J. R. Curran (2012), Compact continuum source finding for next generation radio surveys, *Mon. Not. R. Astron. Soc.*, *422*(2), 1812–1824, doi:10.1111/j.1365-2966.2012.20768.x.
- Helmboldt, J. F., T. J. W. Lazio, H. T. Intema, and K. F. Dymond (2012a), A new technique for spectral analysis of ionospheric TEC fluctuations observed with the Very Large Array VHF system: From QP echoes to MSTIDs, *Radio Sci.*, *47*, RS0L02, doi:10.1029/2011RS004787.
- Helmboldt, J. F., W. M. Lane, and W. D. Cotton (2012b), Climatology of midlatitude ionospheric disturbances from the Very Large Array Low-frequency Sky Survey, *Radio Sci.*, *47*, RS5008, doi:10.1029/2012RS005025.
- Hoogeveen, G. W., and A. R. Jacobson (1997), Improved analysis of plasmasphere motion using the VLA radio interferometer, *Ann. Geophys.*, *15*, 236–245, doi:10.1007/s00585-997-0236-6.
- Imai, K., L. Wang, and T. D. Cart (1992), A model for the production of Jupiter's decametric modulation lanes, *Geophys. Res. Lett.*, *19*(9), 953–956.
- Jacobson, A. R., and W. C. Erickson (1993), Observations of electron density irregularities in the plasmasphere using the VLA radio interferometer, *Ann. Geophys.*, *11*(10), 869–888.
- Jacobson, A. R., G. Hoogeveen, R. C. Carlos, and M. C. Kelley (1996), Observations of inner plasmasphere irregularities with a satellite-beacon radio-interferometer array, *J. Geophys. Res.*, *101*(A9), 19,665–19,682, doi:10.1029/96JA01253.
- Karpachev, A. T., G. A. Zhabankov, V. P. Kuleshova, and V. A. Telegin (2014), Guided radio-wave propagation in the equatorial ionosphere according to the topside sounding onboard Interkosmos-19, *J. Atmos. Sol. Terr. Phys.*, *121*, 89–97, doi:10.1016/j.jastp.2014.10.008.
- Lonsdale, C. J., et al. (2009), The Murchison widefield array: Design overview, *Proc. IEEE*, *97*(8), 1497–1506, doi:10.1109/JPROC.2009.2017564.
- McCormick, R. J. (2002), Reconsidering the effectiveness of quasi-static thunderstorm electric fields for whistler duct formation, *J. Geophys. Res.*, *107*(A11), 1396, doi:10.1029/2001JA009219.
- McIlwain, C. E. (1961), Coordinates for mapping the distribution of magnetically trapped particles, *J. Geophys. Res.*, *66*(11), 3681–3691, doi:10.1029/JZ066i011p03681.
- Ohta, K., T. Kitagawa, N. Shima, M. Hayakawa, and R. L. Dowden (1996), Characteristics of mid-latitude whistler ducts as deduced from ground-based measurements, *Geophys. Res. Lett.*, *23*(23), 3301–3304, doi:10.1029/96GL03253.
- Park, C. G., and P. M. Banks (1974), Influence of thermal plasma flow on the mid-latitude nighttime F2 layer: Effects of electric fields and neutral winds inside the plasmasphere, *J. Geophys. Res.*, *79*(31), 4661–4668, doi:10.1029/JA079i031p04661.
- Parrot, M., J. A. Sauvaud, J. J. Berthelier, and J. P. Lebreton (2007), First in-situ observations of strong ionospheric perturbations generated by a powerful VLF ground-based transmitter, *Geophys. Res. Lett.*, *34*, L11111, doi:10.1029/2007GL029368.
- Rodger, J., N. R. Thomson, and L. Dowden (1998), Are whistler ducts created by thunderstorm electrostatic fields?, *J. Geophys. Res.*, *103*(A2), 2163–2169, doi:10.1029/97JA02927.
- Saxton, J. M., and A. J. Smith (1989), Quiet time plasmaspheric electric fields and plasmasphere-ionosphere coupling fluxes at  $L=2.5$ , *Planet. Space Sci.*, *37*(3), 283–293, doi:10.1016/0032-0633(89)90025-1.
- Sazykin, S., R. A. Wolf, R. W. Spiro, T. I. Gombosi, D. L. D. Zeeuw, and M. F. Thomsen (2004), Interchange instability in the inner magnetosphere associated with geosynchronous particle flux decreases, *Geophys. Res. Lett.*, *31*, L05803, doi:10.1029/2003GL019191.
- Shiokawa, K., M. Mori, Y. Otsuka, S. Oyama, S. Nozawa, S. Suzuki, and M. Connors (2013), Observation of nighttime medium-scale travelling ionospheric disturbances by two 630-nm airglow imagers near the auroral zone, *J. Atmos. Sol. Terr. Phys.*, *103*, 184–194, doi:10.1016/j.jastp.2013.03.024.

- Singh, R. P., A. K. Singh, and D. K. Singh (1998), Plasmaspheric parameters as determined from whistler spectrograms: A review, *J. Atmos. Sol. Terr. Phys.*, *60*(5), 495–508, doi:10.1016/S1364-6826(98)00001-7.
- Sonwalkar, V. S. (2006), The influence of plasma density irregularities on whistler-mode wave propagation, in *Geospace Electromagnetic Waves and Radiation*, vol. 191, pp. 141–190, Springer, Berlin.
- Sonwalkar, V. S., U. S. Inan, T. F. Bell, R. A. Helliwell, V. M. Chmyrev, Y. P. Sobolev, O. Y. Ovcharenko, and V. Selegej (1994), Simultaneous observations of VLF ground transmitter signals on the DE 1 and COSMOS 1809 satellites: Detection of a magnetospheric caustic and a duct, *J. Geophys. Res.*, *99*(A9), 17,511–17,522, doi:10.1029/94JA00866.
- Thompson, A. R., J. M. Moran, and G. W. Swenson Jr. (2001), *Interferometry and Synthesis in Radio Astronomy*, 2nd ed., John Wiley, Weinheim, Germany, doi:10.1002/9783527617845.
- Tingay, S. J., et al. (2013), The Murchison Widefield Array: The square kilometre array precursor at low radio frequencies, *Publ. Astron. Soc. Aust.*, *30*, e007, doi:10.1017/pasa.2012.007.
- Tu, J., P. Song, B. W. Reinisch, J. L. Green, and X. Huang (2006), Empirical specification of field-aligned plasma density profiles for plasmasphere refilling, *J. Geophys. Res.*, *111*, A06216, doi:10.1029/2005JA011582.
- Yearby, K. H., M. A. Balikhin, Y. V. Khotyaintsev, S. N. Walker, V. V. Krasnoselskikh, H. S. C. K. Alleyne, and O. Agapitov (2011), Ducted propagation of chorus waves: Cluster observations, *Ann. Geophys.*, *29*(9), 1629–1634, doi:10.5194/angeo-29-1629-2011.