



Publication Year	2015
Acceptance in OA @INAF	2020-05-27T11:56:56Z
Title	X-Ray High-resolution Spectroscopy Reveals Feedback in a Seyfert Galaxy from an Ultra-fast Wind with Complex Ionization and Velocity Structure
Authors	Longinotti, A. L.; Krongold, Y.; Guainazzi, M.; GIROLETTI, MARCELLO; PANESSA, Francesca; et al.
DOI	10.1088/2041-8205/813/2/L39
Handle	http://hdl.handle.net/20.500.12386/25224
Journal	THE ASTROPHYSICAL JOURNAL LETTERS
Number	813

X-RAY HIGH-RESOLUTION SPECTROSCOPY REVEALS FEEDBACK IN A SEYFERT GALAXY FROM AN ULTRA-FAST WIND WITH COMPLEX IONIZATION AND VELOCITY STRUCTURE

A. L. LONGINOTTI^{1,2,3}, Y. KRONGOLD², M. GUAINAZZI^{3,4}, M. GIROLETTI⁵, F. PANESSA⁶,
E. COSTANTINI⁷, M. SANTOS-LLEO³, AND P. RODRIGUEZ-PASCUAL³

¹ Catedrática CONACYT—Instituto Nacional de Astrofísica, Óptica y Electrónica, Luis E. Erro 1, Tonantzintla, Puebla, C.P. 72840, México

² Instituto de Astronomía, Universidad Nacional Autónoma de México, Apartado Postal 70264, 04510 México D.F., México

³ ESAC, P.O. Box, 78 E-28691 Villanueva de la Cañada, Madrid, Spain

⁴ Institute of Space and Astronautical Science, 3-1-1 Yoshinodai, Chuo-ku, Sagami-hara, Kanagawa, Japan

⁵ INAF Osservatorio di Radioastronomia, via Gobetti 101, I-40129 Bologna, Italy

⁶ INAF—Istituto di Astrofisica e Planetologia Spaziali di Roma (IAPS), Via del Fosso del Cavaliere 100, I-00133 Roma, Italy

⁷ SRON Netherlands Institute for Space Research, Sorbonnelaan 2, 3584 CA Utrecht, The Netherlands

Received 2015 October 3; accepted 2015 October 28; published 2015 November 6

ABSTRACT

Winds outflowing from active galactic nuclei (AGNs) may carry significant amounts of mass and energy out to their host galaxies. In this paper we report the detection of a sub-relativistic outflow observed in the narrow line Seyfert 1 galaxy IRAS 17020+4544 as a series of absorption lines corresponding to at least five absorption components with an unprecedented wide range of associated column densities and ionization levels and velocities in the range of 23,000–33,000 km s⁻¹, detected at X-ray high spectral resolution ($E/\Delta E \sim 1000$) with the ESA’s observatory *XMM-Newton*. The charge states of the material constituting the wind clearly indicate a range of low to moderate ionization states in the outflowing gas and column densities that are significantly lower than observed in highly ionized ultra-fast outflows. We estimate that at least one of the outflow components may carry sufficient energy to substantially suppress star formation and heat the gas in the host galaxy. IRAS 17020+4544 therefore provides an interesting example of feedback by a moderately luminous AGN that is hosted in a spiral galaxy, a case barely envisaged in most evolution models, which often predict that feedback processes take place in massive elliptical galaxies hosting luminous quasars in a post-merger phase.

Key words: accretion, accretion disks – galaxies: Seyfert – line: identification

1. INTRODUCTION

The effect of active galactic nucleus (AGN) winds on the larger scale environment, which is commonly referred to as “AGN feedback,” has far-reaching effects on the evolution of the host galaxy properties (Di Matteo et al. 2005; Hopkins & Elvis 2010). For this reason, quasar feedback has been extensively postulated as a fundamental mechanism for regulating the relation of supermassive black holes with their host galaxies. The recent reports of sub-relativistic X-ray winds in two very bright quasars (Nardini et al. 2015; Tombesi et al. 2015) corroborate the role of highly ionized accretion disk winds as a source of feedback. Indeed, the X-ray spectra of some AGNs carry the signature of gas outflowing at sub-relativistic speeds ($v \geq 0.1c$), so highly ionized that the only dominant ions are helium-like and hydrogen-like ionic species. These systems are known as “ultra-fast outflows” (UFOs) and they are observed in the Fe K band at $E \geq 7$ keV. Several papers reported on UFOs hosted in individual AGNs (Pounds et al. 2003; Chartas et al. 2009; Pounds & Vaughan 2011; Lanzuisi et al. 2012; Pounds 2014), and statistical studies show that UFOs are detected in 30%–40% of nearby AGNs (Tombesi et al. 2010, 2012; Gofford et al. 2013). The approximate ranges of mass outflow rates (0.01 – $1 M_{\odot} \text{ yr}^{-1}$) and kinetic energies (10^{42} – $10^{45} \text{ erg s}^{-1}$) may be in good agreement with theoretical predictions for black hole winds (King 2010). However, the very existence of such winds has been debated within the community (Kaspi & Behar 2006; Gallo & Fabian 2013; Laha et al. 2014; Zoghbi et al. 2015) because the intrinsic limitations in the CCD data resolution ($E/\Delta E \sim 30$ – 40) and the often uncertain identification of the

atomic transition do not allow an unambiguous—i.e., model-independent—confirmation that these features represent a bona fide UFO.

IRAS 17020+4544 is a narrow line Seyfert 1 spiral galaxy at redshift $z=0.0604$, with a black hole mass of $M_{\text{BH}} \sim 5.9 \times 10^6 M_{\odot}$ (Wang & Lu 2001), an X-ray luminosity of $\sim 1.5 \times 10^{44} \text{ erg s}^{-1}$, and a flux of $1.6 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$ in 0.3–10 keV. The bolometric luminosity of $\sim 5.2 \times 10^{44} \text{ erg s}^{-1}$, estimated assuming a conservative bolometric correction $k=10$ (Marconi et al. 2004), implies that the source is accreting at a high accretion rate ($L_{\text{BOL}}/L_{\text{EDD}} \sim 0.7$).

2. DATA REDUCTION AND SPECTRAL ANALYSIS

XMM-Newton observed IRAS 17020+4544 for 160 ks on 2014 January 23 and 25 (OBSIDs 0721220101 0721220301), and twice in 2004 for 40 ks (0206860101 0206860201). The X-ray data include the spectra from the Reflection Grating Spectrometer (RGS; den Herder et al. 2001), and the CCD spectra from the European Photon Imaging Camera (EPIC pn; Strüder et al. 2001). No background flares due to high-energy particles are present in the observations. RGS data were processed by the standard System Analysis Software (SAS 13.5.0) tool *rgsproc*. Due to the minor flux variability and the lack of spectral changes within the two *XMM* OBSIDs of each year ($\Delta\Gamma=3\%$, $\Delta\text{Flux}=9\%$), RGS spectra were combined in one single spectrum by the SAS tool *rgscombine* in order to maximize the signal-to-noise ratio. EPIC pn event files were obtained by applying standard filtering criteria through the SAS task *epproc*. Source and background spectra were extracted from circular regions of $40''$. Background regions were selected




Figure 1. RGS spectra fitted with the model including a power-law continuum absorbed by four “slow” warm absorber components. The inset shows the spectral portion mostly affected by the ultra-fast outflow. The spectrum was binned by five instrumental channels for display purposes.

in source-free portions of the CCD. Response matrices were provided by the SAS tools *arfgen* and *rmfgen*. The software used for the spectral analysis was XSPEC v.12.8.2h.

2.1. The Continuum and the Warm Absorbers

The RGS spectral analysis was carried out on the unbinned spectrum and the Cash statistics (Cash 1979) was applied in the spectral fitting process. Fit parameters are quoted with 1σ error bars throughout the paper. The X-ray continuum in 7–35 Å was parameterized by a power law modified by Galactic absorption (Kalberla et al. 2005) of column density $N_{\text{H}} = 2 \times 10^{20} \text{ cm}^{-2}$. The residuals of the spectrum against the best-fit continuum show a forest of narrow absorption features in the soft X-ray band imprinted by ionized gas along the line of sight to the AGN. We first modeled the warm absorber that was already known in this source (Komossa & Bade 1998). To this purpose we employed the self-consistent photoionization code PHASE (Krongold et al. 2003). Each PHASE component describes one layer of ionized gas characterized by a particular set of parameters: ionization state U^8 , column density, covering factor, turbulent velocity, and outflow velocity. To compute the absorption spectrum we assumed the ionization balance produced by the incoming ionizing radiation emitted by the source; therefore we constructed the spectral energy distribution (SED) for IRAS 17020+4544 using data from the NASA Extragalactic Database and from *XMM-Newton*. For fitting the warm absorber four such absorbing components were required by the data (A. L. Longinotti et al. 2015, in preparation). The velocity component along the line of sight of the four absorbers is found to be of the order of $10^{2-3} \text{ km s}^{-1}$. The photon index of the power law is $\Gamma = 2.8 \pm 0.04$, as in typical narrow line Seyfert 1 sources. The model including the power-law

continuum and the four ionized absorbers yields a fit statistics of $C_{\text{stat}} = 3258$ for 2752 degrees of freedom in the 7–35 Å band (Figure 1).

Once the warm absorber was well parameterized we noticed the presence of additional absorption features, particularly pronounced in the 18–23 Å band (see inset of Figure 1). Initially, we tested for the presence of individual absorption lines by adding narrow Gaussian lines with negative intensity to the continuum+warm absorbers model. The line widths are fixed to 0.1 eV and their positions are free. Table 1 reports the list of the absorption lines detected above a 2σ threshold, along with their positions and the identification of the atomic transition. The improvement in C -statistics is measured for the addition of 2 degrees of freedom (dof) and in the spectral band 7–35 Å. The results reported in Table 1 show that additional absorption is required by the data, thus, supported by this evidence, we proceeded to model it in a self-consistent manner.

2.2. Modeling the Ultra-fast Outflow

The analysis was then restricted to the spectral region where residuals are most prominent (18–23 Å), which is the least contaminated by the warm absorber (see Figure 1), which imprints only two unblended absorption lines from O VII and O VIII in this spectral range. The statistics for the model with a local power-law continuum and the four warm absorbers fitted in this band is $C_{\text{stat}}/\text{dof} = 512/478$. We start by adding to this model a fast outflow component modeled by PHASE. Four additional such components were included in the model. After validating the statistical improvement of each of them, five UFO components are required by the fit (see Table 2). In contrast to adding Gaussian lines at random positions that are not descriptive of the global ionization state of the gas, the use of PHASE guarantees that the absorption is modeled self-consistently. The lines are narrow and unresolved by RGS, so the turbulent velocity was fixed to 50 km s^{-1} . All UFO

⁸ $U = \frac{Q}{4\pi R^2 n_e c}$, where Q is the luminosity of ionizing photons, n_e and R are the electron density and the distance of the wind from the X-ray source, respectively.

Table 1List of Absorption Lines Individually Detected above a 2σ Threshold

Obs λ (Å)	Intensity (10^{-5} ph cm $^{-2}$ s $^{-1}$)	ΔC_{stat}	Sign. σ	Line ID
21.18 $^{+0.02}_{-0.02}$	-2.7 $^{+0.60}_{-0.60}$	34	4.5	O VII
22.03 $^{+0.04}_{-0.02}$	-3.01 $^{+0.89}_{-0.73}$	32	3.4	O IV
18.60 $^{+0.03}_{-0.03}$	-1.54 $^{+0.55}_{-0.55}$	19	2.8	O VIII
21.67 $^{+0.03}_{-0.03}$	-2.17 $^{+0.80}_{-0.80}$	21	2.7	O V
21.83 $^{+0.03}_{-0.01}$	-2.28 $^{+0.92}_{-0.92}$	18	2.5	O III
21.55 $^{+0.03}_{-0.02}$	-1.76 $^{+0.85}_{-0.91}$	8	2.0	O IV
20.93 $^{+0.02}_{-0.03}$	-1.61 $^{+0.83}_{-0.82}$	10	2.0	O VI

Note. Identifications were made by matching this phenomenological fit to the physical model in Table 2.

components have outflow velocities that are consistent with the lines being displaced by $v \sim 0.1c$. The final fit statistics for the best-fit model that includes the power-law continuum, the four warm absorber components, plus 5 UFO components, is $C_{\text{stat}}/\text{dof} = 409/463$. Figure 2 shows that our analysis is fully capable of separating the two sets of lines from the slow and fast outflowing gas without ambiguity. When the best-fit model is extended to the full RGS band down to 35 Å, the detection of the UFOs is confirmed, therefore we are confident that these features cannot be mistaken with the absorption lines from the warm absorber at lower wavelengths. We also tested the robustness of the UFO detection by estimating its significance using a local continuum in the 18–23 Å band that does not include the warm absorber. All five components are detected with the same significance. Figure 3 shows the RGS spectrum fitted by the best-fit model.

Although an alternative scenario is unlikely since the patterns of the line series observed in our data are very well fitted by our self-consistent model, we have considered the presence of local absorption ($z = 0$) as a possible alternative to the ultra-fast wind. Fitting a PHASE component with a velocity outflow fixed within a range that was consistent with the absorption being local did not provide a satisfactory fit to the spectrum because the positions of the absorption lines, if interpreted at $z = 0$, are not consistent with known ionic transitions. In particular, most of the local low-ionization absorption (Gatuzz et al. 2013) at $z = 0$ lies in the range 22–24 Å, i.e., above the region where the bulk of the UFOs lines are detected. Likewise, the absorption lines cannot be attributed to intervening material between us and the galaxy, as the measured outflow velocity is significantly larger than the receding velocity of IRAS 17020+4544 (18,000 km s $^{-1}$).

2.2.1. XMM-Newton Data of 2004

We tested the presence of the fast outflow in the *XMM-Newton* archival data of IRAS 17020+4544 obtained in 2004. We kept the baseline model with the same continuum and the warm absorber as the overall spectral shape and flux level in 2004 are very similar to those inferred in 2014. This yielded a fit statistics of $C_{\text{stat}}/\text{dof} = 514/485$ in 18–23 Å. When the ultra-fast outflow was added to the model, only component A (i.e., the most significant in the 2014 data) is detected in this data set, with an improvement of $\Delta C_{\text{stat}} = 21$. The parameters of this outflow component are consistent with the ones reported for 2014. The other four components are not detected but we cannot statistically exclude their presence in these data. This

test shows that at least one Component (A) of the outflow seems persistent over a 10-year timescale.

2.3. Monte Carlo Methods

To provide a rigorous estimate of the statistical significance of the outflow detected in 2014, Monte Carlo simulations were performed. Following the approach that was taken for modeling the UFO in the RGS spectra, the restricted band 18–23 Å was considered. As a baseline model for the synthetic data sets we assumed the (continuum+ 4 warm absorbers) model for testing the significance of Component A. For the other UFO components we always included all the previous UFO components in the model as done in the real spectral fitting, i.e., we included continuum+ 4 warm absorbers+UFO A for testing the significance of UFO B, continuum+ 4 warm absorbers+UFO A+UFO B for testing the significance of UFO C, etc. For each UFO we produced 1000 simulated spectra with the XSPEC package. Each realization was folded through the same response matrix employed in the spectral fitting and random noise was added. Therefore each spectrum corresponds to a background-subtracted RGS data set with the same photon statistics as the real data set. In each simulated spectrum the improvement in C -statistics after adding a UFO component to the baseline model was measured and recorded. We checked that the results were distributed according to a Gaussian distribution and this conclusion led to the significance quoted in Table 2. The significance of Component C was estimated by tying the wind velocity between Components B and C to be the same within the errors. Given the two very similar outflow velocities, the information on the velocity of Component B is used as a prior for assessing the significance of Component C.

2.4. EPIC pn Spectral Analysis

We searched for a potential high-ionization counterpart of the RGS UFO in the 2014 EPIC pn spectrum. The data fitted in the 2–10 keV band with a steep power-law continuum ($\Gamma = 2.25 \pm 0.08$) show a prominent Fe K α line in emission that was fitted phenomenologically with a Gaussian profile. The resulting peak energy is consistent with emission from ionized material ($E = 6.95 \pm 0.10$ keV), the line width is $\sigma = 0.26^{+0.21}_{-0.10}$ keV, and the Equivalent Width is 150^{+85}_{-55} eV. We then added a negative narrow Gaussian line at the position of the Fe xxvi atomic transition (6.97 keV rest-frame), blueshifted by the corresponding velocity measured in the RGS UFO (i.e., $E \sim 7.2$ keV). No absorption line is detected at this position. The 90% upper limit on the line EW is 14 eV and is significantly lower than the average intensity of Fe K UFOs (Tombsi et al. 2010). We then explored the potential presence of absorption lines at higher energies by adding two narrow Gaussians with free position to the power-law continuum. Two features are marginally detected at $E = 8.75 \pm 0.13$ keV and $E = 9.39 \pm 0.10$ keV, with respective intensities of 49 ± 43 eV and $EW = 64 \pm 50$ eV. Although the signal-to-noise in this region of the EPIC pn spectrum is low due to the sharp decrease of the *XMM-Newton* telescopes' effective area above 7 keV and to the steep intrinsic spectrum of IRAS 17020+4544, the positions of the two absorption lines are consistent with being Fe xxv and Fe xxvi blueshifted by a velocity of $v \sim 0.34c$.

Table 2
Parameters of the Five UFO Components Detected in the RGS Spectrum

UFO Component Index	$\log U$ (erg cm s^{-1})	$\text{Log } N_{\text{H}}$ (cm^{-2})	v_{out} (km s^{-1})	Statistics ΔC_{stat}	Significance
Comp (A)	$-0.39^{+0.30}_{-0.15}$	$21.47^{+0.18}_{-0.21}$	23640^{+150}_{-60}	45	9.0σ
Comp (B)	$-1.99^{+0.33}_{-0.26}$	$20.42^{+0.21}_{-0.58}$	27200^{+240}_{-240}	26	5.3σ
Comp (C)	$2.58^{+0.17}_{-0.85}$	$23.99^{+0}_{-1.86}$	27200^{+300}_{-270}	10	3.6σ
Comp (D)	$0.33^{+1.79}_{-0.40}$	$21.42^{+0.84}_{-1.28}$	25300^{+210}_{-180}	12	2.6σ
Comp (E)	$-2.92^{+0.51}_{-0.14}$	$19.67^{+0.34}_{-0.36}$	33900^{+360}_{-270}	10	2.0σ

Note. The statistical improvement (fifth column) refers to the addition of each PHASE component to the model comprising the continuum, the warm absorbers, and the previous UFO components. The significance is estimated through Monte Carlo methods.




Figure 2. Features from the warm absorber (included in the model) and from the UFO (not included) do not overlap with each other. Instrumental features (in blue) do not hamper the detection of absorption intrinsic to the source. Labels are drawn from Table 2's physical model.




Figure 3. Portion of the RGS spectrum most affected by the fast wind absorption and modeled by the best fit reported in Table 2.

3. RESULTS

3.1. A Stratified Multi-component Ultra-fast Outflow

The simultaneous presence of three UFO components, required to have such high significance (9.0σ , 5.3σ , and 3.6σ) is, to our knowledge, unprecedented. More importantly,

the newly discovered wind presents a much more complex nature than envisaged so far for UFOs. Components B and C are separated by four decades in ionizations and more than three decades in column density, yet they are outflowing at the same speed.

Most theoretical models of black hole winds predict that winds at a sub-relativistic speed are bound to reach an ionization level so high that they only produce absorption in the Fe K band, e.g., the shock outflow model (King 2010). However, the unquestionable presence of a sub-relativistic wind at low-ionization in IRAS 17020+4544 breaks the linear correlation between ionization and velocity of the wind that is predicted by this model and that inspired the unification scenario for UFO and warm absorbers postulated by Tombesi et al. (2013). We note that this scenario has already been questioned (Gupta et al. 2013, 2015).

The newly discovered multi-component outflow could be interpreted as gas condensed in clumps with distinct properties (v , U , N_{H}) embedded in a high-velocity, high-ionization wind as the one described in the shock outflow model. However, it would be difficult to explain how these clumps may survive for $t \geq 10$ years, as inferred by the comparison of the *XMM-Newton* spectra in 2014 and 2004. More likely, the observed absorption lines are revealing the cooling region envisaged in models of black hole outflows right after the first shock between the wind and the surrounding material (King 2010). In this model, the accretion disk wind launched with velocity v_{out} suffers an isothermal shock with the circumnuclear gas, which produces the effect of slowing the wind to a velocity $v \sim \frac{v_{\text{out}}}{4}$. The properties of the two tentative highly ionized absorption lines reported in Section 2.4 are consistent with the predictions of this model. Although the structure of the outflowing system in IRAS 17020+4544 requires a more elaborate explanation, the observed velocities of the wind seem to be in excellent agreement with this scenario: $v_{\text{out}} \sim 0.34c = 102,000 \text{ km s}^{-1}$ (EPIC pn measurement) and $v \sim 0.08\text{--}0.11c = 24,000\text{--}33,000 \text{ km s}^{-1}$ (RGS measurement). Deeper X-ray observations will certainly help to shed light on the presence and properties of the highly ionized features.

An additional point of interest of the outflow in IRAS 17020+4544 comes from observations in the radio band. Recent data obtained by our group (M. Giroletti et al. 2015, in preparation), along with a re-analysis of archival data (Gu & Chen 2010), confirm the presence of outflowing material on a scale of about 10 pc, with a speed lower than $\sim 0.2c$, potentially in good agreement with the X-ray velocities. The presence of the radio jet/outflow suggests that the wind may be magnetically driven or produced by the interaction of the radio jet with the surrounding material. On the other hand, the high accretion rate

of the source ($L_{\text{BOL}}/L_{\text{EDD}} \sim 0.7$) is in agreement with models of radiatively driven outflows (King 2010).

3.1.1. Comparison with Other Results

Detection of ultra-fast outflows in high-resolution spectra of local AGNs with moderate ionization was already reported. Pounds (2014) reports a secure detection of only one component in the RGS data of PG1211+143 since the second outflow may well be associated with local absorption due to the coincidence of its velocity with the systemic velocity of the AGN. The claim of a dual relativistic outflow in Ark 564 (Gupta et al. 2013) is based on measurement of individual lines with velocities that are inconsistent with each other, therefore it is difficult to associate them with the same outflowing system. Finally, only one absorber was detected above the threshold of 99.99% in the *Chandra* grating spectra of Mrk 590 (Gupta et al. 2015). While all of these results have certainly contributed to foster the search for UFOs, none of them have unveiled an ultra-fast outflow constituted by several ionization and velocity components like the one discovered in IRAS 17020+4544. Our result therefore represents the first unambiguous discovery of a fast ($v \sim 0.1 c$) multi-component AGN outflow in X-ray high-resolution spectroscopic data, ultimately settling the open controversy on their existence.

3.2. Energetics of the Ultra-fast Outflow and Consequences for Feedback

The energetics of the outflow were estimated under the common assumption that the outflow velocity is larger or equal to the escape velocity at the launch radius $r = \frac{2GM_{\text{BH}}}{v_{\text{out}}^2}$. Taking into account the lowest and highest velocities measured in the wind (corresponding to Components A and E, respectively) the launching radius is limited to be lower than $(1.4\text{--}2.8) \times 10^{14}$ cm. Following Tombesi et al. (2015) and Crenshaw & Kramer (2012) the expression $\dot{M}_{\text{out}} = 4\pi\mu r N_{\text{H}} v_{\text{out}} m_p C_f$ can be used to estimate the mass outflow rate in spherical geometry, with m_p and μ corresponding to the mass of the proton and to the mean atomic mass per particle ($\mu = 1.4$), C_f as the covering fraction of the outflow, and N_{H} and v_{out} as the column density and the velocity of the flow measured in our data. By expressing these two quantities, respectively, in units of 10^{21} cm^{-2} and $10,000 \text{ km s}^{-1}$, this formula can be conveniently parameterized as $\dot{M}_{\text{out}} = 7.3 \times 10^{-4} \frac{M_{\odot}}{\text{yr}} \frac{N_{\text{H}21}}{v_{\text{out}(10,000)}} C_f \text{ gr s}^{-1}$, which yields a relevant mass outflow rate only for Component C, i.e., $\dot{M}_{\text{out}}(\text{C}) \sim 0.26 C_f M_{\odot} \text{ yr}^{-1}$. The energy outflow rate can be estimated by starting with $\dot{E} = \frac{1}{2} \dot{M}_{\text{out}} v_{\text{out}}^2$, where the previous expression for \dot{M}_{out} can be substituted in order to express the energy outflow rate in terms of the column density and the velocity of the wind: $\dot{E} = 2.31 \times 10^{40} N_{\text{H}21} v_{\text{out}}(10,000) C_f \text{ erg s}^{-1}$. Considering the bolometric luminosity, we can express $\frac{\dot{E}}{L_{\text{BOL}}} = \frac{\dot{E}}{10^{42} \text{ erg s}^{-1}}$, which for Component C yields $\frac{\dot{E}(\text{C})}{L_{\text{BOL}}} = 11\% C_f$.

The study of the properties of this outflow may hold important clues for elucidating the contribution of accretion disk winds to AGN feedback. Even if the outflow covers a

small part of the solid angle, the energy rate is sufficient to power AGN feedback according to the required minimum value (Hopkins & Elvis 2010) of $\sim 0.5\%$. Most notably, the AGN luminosity in IRAS 17020+4544 is ~ 100 times lower than in the sources where the most compelling evidence for black hole feedback produced by X-ray highly ionized fast winds was found (Nardini et al. 2015; Tombesi et al. 2015). Deep optical images (Ohta et al. 2007) show that this AGN is hosted by a spiral barred galaxy with no signs of interactions. Hence, IRAS 17020+4544 is the first case of an undisturbed field spiral galaxy where feedback from accretion disk winds can regulate star formation and drive the fundamental relation between the stellar velocity dispersion in the galaxy bulge and the mass of the central supermassive black holes (Ferrarese & Merritt 2000). These findings will certainly offer new insights into the physical mechanisms at play during the phase when galaxy evolution is regulated by black hole feedback.

This paper is based on observations obtained with *XMM-Newton*, an ESA science mission with instruments and contributions directly funded by ESA Member States and NASA. We thank the anonymous referee. We acknowledge support from the ESAC Faculty, *INTEGRAL* ASI/INAF n. 2013-025.R.O, and CONACYT n.233263. Y.K. acknowledges support from grant DGAPA PAIIPIIT IN104215 and CONACYT grant 168519. This paper is dedicated to Prof. Maurizio Longinotti, Haematologist.

REFERENCES

- Cash, W. 1979, *ApJ*, **228**, 939
 Chartas, G., Saez, C., Brandt, W. N., Giustini, M., & Garmire, G. P. 2009, *ApJ*, **706**, 644
 Crenshaw, D. M., & Kraemer, S. B. 2012, *ApJ*, **753**, 75
 den Herder, J. W., Brinkman, A. C., Kahn, S. M., et al. 2001, *A&A*, **365**, L7
 Di Matteo, T., Springel, V., & Hernquist, L. 2005, *Natur*, **433**, 604
 Ferrarese, L., & Merritt, D. 2000, *ApJL*, **539**, L9
 Gallo, L. C., & Fabian, A. C. 2013, *MNRAS*, **434**, L66
 Gatuzz, E., García, J., Mendoza, C., et al. 2013, *ApJ*, **768**, 60
 Gofford, J., Reeves, J. N., Tombesi, F., et al. 2013, *MNRAS*, **430**, 60
 Gu, M., & Chen, Y. 2010, *AJ*, **139**, 2612
 Gupta, A., Mathur, S., & Krongold, Y. 2015, *ApJ*, **798**, 4
 Gupta, A., Mathur, S., Krongold, Y., & Nicastro, F. 2013, *ApJ*, **772**, 66
 Hopkins, P. F., & Elvis, M. 2010, *MNRAS*, **401**, 7
 Kalberla, P. M. W., Burton, W. B., Hartmann, D., et al. 2005, *A&A*, **440**, 775
 Kaspi, S., & Behar, E. 2006, *ApJ*, **636**, 674
 King, A. R. 2010, *MNRAS*, **402**, 1516
 Komossa, S., & Bade, N. 1998, *A&A*, **331**, L49
 Krongold, Y., Nicastro, F., Brickhouse, N. S., et al. 2003, *ApJ*, **597**, 832
 Laha, S., Guainazzi, M., Dewangan, G. C., Chakravorty, S., & Kembhavi, A. K. 2014, *MNRAS*, **441**, 2613
 Lanzuisi, G., Giustini, M., Cappi, M., et al. 2012, *A&A*, **544**, A2
 Marconi, A., Risaliti, G., Gilli, R., et al. 2004, *MNRAS*, **351**, 169
 Nardini, E., Reeves, J. N., Gofford, J., et al. 2015, *Sci*, **347**, 860
 Ohta, K., Aoki, K., Kawaguchi, T., & Kiuchi, G. 2007, *ApJS*, **169**, 1
 Pounds, K. A. 2014, *MNRAS*, **437**, 3221
 Pounds, K. A., Reeves, J. N., King, A. R., et al. 2003, *MNRAS*, **345**, 705
 Pounds, K. A., & Vaughan, S. 2011, *MNRAS*, **413**, 1251
 Strüder, L., Briel, U., Dennerl, K., et al. 2001, *A&A*, **365**, L18
 Tombesi, F., Cappi, M., Reeves, J. N., & Braito, V. 2012, *MNRAS*, **422**, L1
 Tombesi, F., Cappi, M., Reeves, J. N., et al. 2010, *A&A*, **521**, A57
 Tombesi, F., Cappi, M., Reeves, J. N., et al. 2013, *MNRAS*, **430**, 1102
 Tombesi, F., Meléndez, M., Veilleux, S., et al. 2015, *Natur*, **519**, 436
 Wang, T., & Lu, Y. 2001, *A&A*, **377**, 52
 Zoghbi, A., Miller, J. M., Walton, D. J., et al. 2015, *ApJL*, **799**, L24