



Publication Year	2017
Acceptance in OA @INAF	2020-07-24T08:39:40Z
Title	Evidence for a radiatively driven disc-wind in PDS 456?
Authors	Matzeu, G. A.; Reeves, J. N.; BRAITO, Valentina; NARDINI, EMANUELE; McLaughlin, D. E.; et al.
DOI	10.1093/mnras/slx129
Handle	http://hdl.handle.net/20.500.12386/26606
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
Number	472

Evidence for a radiatively driven disc-wind in PDS 456?

G. A. Matzeu,^{1,2*} J. N. Reeves,^{2,3} V. Braitto,¹ E. Nardini,⁴ D. E. McLaughlin,²
A. P. Lobban,² F. Tombesi^{5,6,7} and M. T. Costa²

¹INAF – Osservatorio Astronomico di Brera, Via Bianchi 46, I-23807 Merate (LC), Italy

²Astrophysics Group, School of Physical and Geographical Sciences, Keele University, Keele, Staffordshire ST5 5BG, UK

³Center for Space Science and Technology, University of Maryland Baltimore County, 1000 Hilltop Circle, Baltimore, MD 21250, USA

⁴INAF – Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, I-50125 Firenze, Italy

⁵X-ray Astrophysics Laboratory, NASA/Goddard Space Flight Center, Greenbelt, MD 20771, USA

⁶Department of Astronomy and CRESST, University of Maryland, College Park, MD 20742, USA

⁷Department of Physics, University of Rome ‘Tor Vergata’, Via della Ricerca Scientifica 1, I-00133 Rome, Italy

Accepted 2017 August 11. Received 2017 August 11; in original form 2017 July 4

ABSTRACT

We present a newly discovered correlation between the wind outflow velocity and the X-ray luminosity in the luminous ($L_{\text{bol}} \sim 10^{47} \text{ erg s}^{-1}$) nearby ($z = 0.184$) quasar PDS 456. All the contemporary *XMM–Newton*, *NuSTAR* and *Suzaku* observations from 2001–2014 were revisited and we find that the centroid energy of the blueshifted Fe K absorption profile increases with luminosity. This translates into a correlation between the wind outflow velocity and the hard X-ray luminosity (between 7 and 30 keV) where we find that $v_w/c \propto L_{7-30}^\gamma$ where $\gamma = 0.22 \pm 0.04$. We also show that this is consistent with a wind that is predominately radiatively driven, possibly resulting from the high Eddington ratio of PDS 456.

Key words: black hole physics – galaxies: active – galaxies: nuclei – quasars: individual: (PDS 456) – X-rays: galaxies.

1 INTRODUCTION

Blueshifted Fe K shell absorption lines, observed at > 7 keV in the rest-frame X-ray spectra of active galactic nucleus (AGN), were first discovered in luminous quasars (e.g. Chartas, Brandt & Gallagher 2003; Pounds et al. 2003; Reeves, O’Brien & Ward 2003). The high velocities inferred, i.e. $v_w \gtrsim 0.1c$, imply an association with an accretion disc-wind. The importance of these winds is supported by their frequent detection, as they are observed in the X-ray spectra of approximately 40 per cent of AGN (Tombesi et al. 2010; Gofford et al. 2013), suggesting that their geometry is characterized by a wide opening angle as recently confirmed in the luminous quasar PDS 456 by Nardini et al. (2015, hereafter N15). These outflows are characterized by considerably high column densities ($N_{\text{H}} \sim 10^{23} \text{ cm}^{-2}$) and a mean velocity (v_w) $\sim 0.1c$ (Tombesi et al. 2010) up to mildly relativistic values of ~ 0.2 – $0.4c$ in the most extreme cases (e.g. Chartas et al. 2002; Reeves et al. 2009; Tombesi et al. 2015). The high velocities and high columns can result in a large amount of mechanical power, possibly exceeding the 0.5–5 per cent of the bolometric luminosity (L hereafter), as required for significant AGN feedback (King 2003; King & Pounds 2003; Di Matteo, Springel & Hernquist 2005; Hop-

kins & Elvis 2010). In principle, the physical properties of these fast winds can provide more insight into the mechanism through which they are driven/accelerated. The possible scenarios are, *radiatively driven winds* (e.g. Proga, Stone & Kallman 2000; Proga & Kallman 2004) and/or *magneto-centrifugally driven winds* (MHD wind hereafter, e.g. Ohsuga et al. 2009; Kazanas et al. 2012). Most outflow studies so far mainly concentrated on radiatively driven winds (e.g. Sim et al. 2010; Reeves et al. 2014; Hagino et al. 2015; Nomura & Ohsuga 2017). However, recently Fukumura et al. (2017) successfully applied an MHD wind model to the micro-quasar GRO J1655–40 suggesting that a similar mechanism applies across the black hole mass and luminosity scales.

The quasar PDS 456, located at redshift $z = 0.184$ (Torres et al. 1997) is the most luminous ($L \sim 10^{47} \text{ erg s}^{-1}$; Simpson et al. 1999; Reeves et al. 2000) radio-quiet quasar in the local Universe ($z < 0.3$). Since the first *XMM–Newton* (40 ks) observation carried out in 2001 (Reeves et al. 2003), PDS 456 has showed the presence of a persistent deep absorption trough in the Fe K band above 7 keV. This can be associated with highly ionized Fe K-shell absorption with a corresponding outflow velocity of $v_w \sim 0.25$ – $0.3c$ (Reeves et al. 2009, 2014; Gofford et al. 2014; N15; Matzeu et al. 2016, hereafter M16), which originates from an accretion disc-wind. In this Letter, we present a new correlation between the disc-wind velocity derived from the Fe K profile and the intrinsic X-ray luminosity, which we interpret in the framework

* E-mail: gabriele.matzeu@brera.inaf.it

Table 1. Summary of the *Suzaku*, *XMM–Newton* and *NuSTAR* observations of PDS 456 performed from 2001 to 2014. The net exposure for *NuSTAR* is for a single FPMA/B detector while *XMM–Newton* and *Suzaku* net exposures are based on the EPIC-pn and individual XIS CCD detectors, respectively. References: (1) Reeves et al. (2003), (2) Behar et al. (2010), (3) Reeves et al. (2009), (4) Reeves et al. (2014), (5) Gofford et al. (2014), (6) M16, (7) N15.

Obs.	Satellite	Start date (UT)	Exposure (ks)	Ref.
2001	<i>XMM–Newton</i>	2001-02-26 09:44	44.0	(1)
2007	<i>XMM–Newton</i>	2007-09-12 01:07	179.0	(2)
2007	<i>Suzaku</i>	2007-02-24, 17:58	190.6	(3)
2011	<i>Suzaku</i>	2011-03-16, 15:00	125.6	(4)
2013a	<i>Suzaku</i>	2013-02-21, 21:22	182.3	(5, 6)
2013b	<i>Suzaku</i>	2013-03-03, 19:43	164.8	
2013c	<i>Suzaku</i>	2013-03-08, 12:00	108.3	
A	<i>XMM–Newton</i>	2013-08-27, 04:41	95.7	(7)
	<i>NuSTAR</i>	2013-08-27, 03:41	43.8	
B	<i>XMM–Newton</i>	2013-09-06, 03:24	95.7	
	<i>NuSTAR</i>	2013-09-06, 02:56	43.0	
C	<i>XMM–Newton</i>	2013-09-15, 18:47	102.0	
	<i>NuSTAR</i>	2013-09-15, 17:56	44.0	
D	<i>XMM–Newton</i>	2013-09-20, 02:47	93.0	
	<i>NuSTAR</i>	2013-09-20, 03:06	58.6	
E	<i>XMM–Newton</i>	2014-02-26, 08:03	100.3	
	<i>NuSTAR</i>	2014-02-26, 08:16	109.5	

of a radiatively driven wind. Values of $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $\Omega_{\Lambda_0} = 0.73$ are assumed throughout this work and errors are quoted at the 1σ confidence level or 90 per cent where otherwise stated.

2 WIND ANALYSIS

We utilize all the available *XMM–Newton*, *Suzaku* and *NuSTAR* data sets, concentrating on the X-ray band above 2 keV fitted with a simple power-law continuum modified by Galactic absorption ($N_{\text{H}} = 2.0 \times 10^{21} \text{ cm}^{-2}$) and a single layer of neutral partial covering absorption to account for any spectral curvature with velocity set to be equal to the fast wind’s outflow velocity (see M16). A broad Fe K emission line at 7 keV was also included, where the width was fixed to that measured by the absorption line when the latter is modelled with a simple Gaussian. The X-ray spectra were adopted from previous work (see Table 1 for details). In Fig. 1, the residuals of the Fe K absorption profiles as a ratio to a power-law continuum are plotted for 4 of the 12 observations with different X-ray luminosities (in decreasing order). It is evident that the centroid energy increases across the different epochs with increasing luminosity.

As an initial test, we investigate how the energy of the Fe K profile responds to the ionizing luminosity, measured in either the 7–30 keV band (L_{7-30} hereafter) or the 2–10 keV band (L_{2-10}) between all the sequences. The 7–30 keV band was chosen as it is above the Fe K absorption edge threshold and is where the primary continuum emerges. This luminosity is absorption corrected and is either inferred directly from the *NuSTAR* data or from extrapolating the best-fitting model above 10 keV, for those sequences where no *NuSTAR* observation is available.¹ Monte Carlo simulations were performed in order to estimate the uncertainties on the absorption-

¹ The extrapolation can be considered as a good assumption, given that all the *NuSTAR* observations of PDS 456 indicate a continuum shape of a simple power law ($\Gamma = 2.2\text{--}2.6$) without any strong hard excess.




Figure 1. Residuals to the power-law continuum fitted between 2 and 10 keV band from four selected observations of PDS 456 decreasing in X-ray luminosity. This shows how the centroid energy of the Fe K absorption profile generally decreases in energy with decreasing luminosity from top to bottom. The vertical dotted lines and the grey shadowed areas indicate the position of the centroid and the 90 per cent uncertainty (see Table 2).

corrected intrinsic luminosity. For each observation, 1000 simulations were performed with the `xspec` command `fakeit`, assuming the best-fitting models obtained from the spectral analysis of the actual data. Each simulated spectrum was fitted again to take into account the uncertainties on the null hypothesis, allowing the main spectral parameters vary. These are the photon index and normalization of the continuum, the covering fraction and column density of the partial covering absorber, as well as the column and outflow velocity of the ionized absorber. With the new model, a second simulation was run and fitted, from which we derived the distribution of the luminosities, which were well approximated by a Gaussian distribution. The subsequent 1σ uncertainty was used as the input to the correlation analysis. We also extracted the simulated distributions of the main spectral parameters to check that they were representative of the uncertainties of the actual best-fitting models. Thus, the derived errors on the luminosities fully account for the uncertainties on each of the free spectral parameters.

We first parametrize the blueshifted absorption lines with a Gaussian profile, whose main parameters are tabulated in Table 2. In Fig. 2, we show the centroid energy versus L_{7-30} which becomes more blueshifted at higher luminosities suggesting a positive correlation with the intrinsic luminosity. Thus, from the centroid energy, we can infer the corresponding outflow velocity (v_w/c) of the material, assuming it is mainly associated with $\text{FeXXVI Ly}\alpha$ rather than $\text{Fe XXV He}\alpha$ (see Table 2). The top panel of Fig. 3 shows the outflow velocity derived from the Gaussian fit against L_{7-30} plotted in log–log space. Subsequently, the data points were fitted with a standard regression line of the form $\log(v_w/c) = m \log(L_{7-30}) + \log(C)$, performed through the bivariate correlated errors and intrinsic scatter algorithm (BCES; Akritas & Bershady 1996) which is one of the most common methods which takes into account errors in both x and y values. The regression line produced a gradient of $m = 0.20 \pm 0.05$

Table 2. Summary of the observations and the corresponding properties of the Fe K absorption features. These are obtained by fitting with a Gaussian profile (columns 1–3) and then subsequently with *xSTAR* (columns 4–6). From the former, we measured with the respective 90 percent errors on (1) the rest-frame energy, (2) the inferred outflow velocity and (3) the equivalent width of each line. By modelling with *xSTAR*, we measured the outflow velocity (4), while from the fits, the column density ranges from $\log(N_{\text{H}}/\text{cm}^{-2}) = 23.16^{+0.28}_{-0.23}$ – $24.06^{+0.18}_{-0.21}$ and ionization ranges from $\log(\xi/\text{erg cm s}^{-1}) = 5.1^{+0.11}_{-0.21}$ – $6.33^{+0.08}_{-0.19}$. In columns (5) and (6), we report the intrinsic luminosities and their corresponding 1σ errors in the 2–10 keV and 7–30 keV bands, respectively.

Satellite	Obs.	E_{rest} (1) (keV)	Gaussian v_w/c (2)	EW (3) (eV)	<i>xSTAR</i> v_w/c (4)	L_{2-10} keV (5) ($\times 10^{44}$ erg s $^{-1}$)	L_{7-30} keV (6) ($\times 10^{44}$ erg s $^{-1}$)
<i>XMM-Newton</i> and <i>NuSTAR</i>	A	$9.25^{+0.18}_{-0.11}$	$0.286^{+0.017}_{-0.010}$	-88 ± 47	$0.299^{+0.012}_{-0.012}$	7.94 ± 0.26	5.22 ± 0.11
	B	$9.06^{+0.19}_{-0.20}$	$0.268^{+0.019}_{-0.020}$	-295 ± 63	$0.278^{+0.016}_{-0.020}$	3.57 ± 0.31	1.68 ± 0.06
	C	$9.12^{+0.11}_{-0.09}$	$0.275^{+0.011}_{-0.009}$	-359 ± 55	$0.286^{+0.006}_{-0.007}$	4.51 ± 0.26	2.60 ± 0.08
	D	$9.16^{+0.12}_{-0.12}$	$0.277^{+0.012}_{-0.012}$	-256 ± 60	$0.292^{+0.008}_{-0.007}$	5.10 ± 0.31	2.93 ± 0.08
	E	$8.91^{+0.11}_{-0.11}$	$0.268^{+0.019}_{-0.020}$	-405 ± 60	$0.269^{+0.008}_{-0.009}$	3.68 ± 0.35	1.81 ± 0.06
<i>Suzaku</i>	2007	$9.26^{+0.15}_{-0.13}$	$0.287^{+0.015}_{-0.013}$	-235 ± 74	$0.296^{+0.012}_{-0.012}$	4.35 ± 0.34	2.32 ± 0.12
	2011	$9.07^{+0.18}_{-0.20}$	$0.270^{+0.018}_{-0.020}$	-460 ± 151	$0.266^{+0.011}_{-0.011}$	4.14 ± 0.53	2.21 ± 0.19
	2013a	$8.85^{+0.33}_{-0.29}$	$0.248^{+0.035}_{-0.030}$	> -214	$0.248^{+0.030}_{-0.020}$	3.53 ± 0.35	1.88 ± 0.14
	2013b	$8.86^{+0.09}_{-0.09}$	$0.249^{+0.010}_{-0.010}$	-390 ± 81	$0.255^{+0.010}_{-0.010}$	2.90 ± 0.36	1.55 ± 0.16
	2013c	$8.70^{+0.07}_{-0.08}$	$0.234^{+0.008}_{-0.009}$	-568 ± 82	$0.240^{+0.010}_{-0.010}$	2.77 ± 0.38	1.46 ± 0.17
<i>XMM-Newton</i>	2001	$9.84^{+0.19}_{-0.16}$	$0.335^{+0.018}_{-0.019}$	-337 ± 94	$0.340^{+0.018}_{-0.015}$	10.5 ± 0.07	4.97 ± 0.28
	2007	$9.51^{+0.14}_{-0.14}$	$0.309^{+0.013}_{-0.013}$	> -71	$0.299^{+0.026}_{-0.034}$	5.55 ± 0.29	3.67 ± 0.14




Figure 2. Plot showing how the centroid energy of the Fe K features correlates with L_{7-30} in PDS 456. The distribution of the data favours a positive linear correlation between the two quantities. *Suzaku* (green circles), *XMM-Newton* and *NuSTAR* (blue stars) and *XMM-Newton* (red squares).

where the shaded magenta area indicates the slope dispersion (at 1σ confidence level). A comparable regression fit is found with the intrinsic 2–10 keV luminosity (L_{2-10}), with $m = 0.22 \pm 0.04$ (see Table 3).

The Fe K absorption features were then modelled with a more physically motivated self-consistent *xSTAR* grid, previously adopted in N15 and M16, generated using the UV to hard X-ray spectral energy distribution (SED) of PDS 456. The *xSTAR* model allows the column and/or the ionization to adjust between observations, as well as the net outflow velocity. The corresponding v_w/c values of all 12 observations derived from the *xSTAR* model are listed in Table 2 and plotted against L_{7-30} on the bottom panel in Fig. 3. The linear regression fit gives a gradient of $m = 0.22 \pm 0.04$ consistent with the Gaussian analysis. Similarly, the correlation with L_{2-10} also gives a consistent result of $m = 0.24 \pm 0.03$. From both




Figure 3. Top: log–log plot showing the correlation between the outflow velocity, derived from fitting the Fe K features with a Gaussian profile and the intrinsic 7–30 keV continuum luminosity (L_{7-30}). The red line shows the linear regression fitted to the data with 1σ uncertainty shaded. Bottom: the outflow velocity is now measured by fitting the Fe K profile with an *xSTAR* grid where the linear regression produced a gradient of 0.22 ± 0.04 . These positive correlations suggest that when the source is more luminous the disc-wind is faster.

Table 3. Results of the linear regressions between the measured outflow velocities (measured from both Gaussian profile and *xstar* grids) and the intrinsic luminosities. m and Δm are the gradient and the respective 1σ errors, respectively, whereas χ^2_ν is the reduced χ^2/dof . P_{null} is the null hypothesis probability that there is no linear correlation.

x	y	m	Δm	χ^2_ν	P_{null}
From Gaussian					
$\log(L_{2-10})$	$\log(v_w/c)$	0.22	0.04	0.84	6.70×10^{-7}
$\log(L_{7-30})$		0.20	0.05	1.08	
From <i>xstar</i>					
$\log(L_{2-10})$	$\log(v_w/c)$	0.24	0.03	0.82	8.29×10^{-8}
$\log(L_{7-30})$		0.22	0.04	1.01	

the correlations shown in Fig. 3, it is clear that the outflow velocity of the highly ionized material is indeed responding to the intrinsic luminosity, where the most luminous epochs of PDS 456 are characterized by a faster outflow (e.g. $v_w/c = 0.296 \pm 0.012$ in the 2007 *Suzaku* observation and $v_w/c = 0.340^{+0.018}_{-0.015}$ in the 2001 *XMM-Newton* observation). On the other hand, the low-luminosity 2013 *Suzaku* observations (i.e. 2013b or 2013c, see Tables 1 and 2) present overall the slowest outflows with $v_w/c = 0.255 \pm 0.010$ and $v_w/c = 0.240 \pm 0.010$, respectively. Thus, an important question is: how the changes in luminosity have a direct impact on the outflow velocity in the disc-wind observed in PDS 456?

3 DISCUSSION

3.1 A radiatively driven wind?

The positive correlation between the outflow velocity and the ionizing luminosity of PDS 456, may imply that the wind is radiatively driven. The high ionization derived from the wind seen at Fe K in PDS 456, with ionization parameters typically of $\log(\xi/\text{erg cm s}^{-1}) \sim 5-6$ (e.g. N15), might result in a small line opacity, due to only Fe XXV–XXVI. While at first sight, this suggests that the dominant interaction mechanism within the photon field is through Thomson scattering (e.g. King & Pounds 2003), the opacity may be further boosted by the presence of lower ionization X-ray absorption within the wind. This lower ionization gas towards PDS 456 has been observed in the form of the clumpy X-ray partial covering absorber with size scales of $\sim 10s$ of R_g from its variability (M16), as well as through the presence of broad absorption profiles revealed in the soft X-ray band (Reeves et al. 2016). Furthermore, blueshifted UV emission and absorption lines from the wind are also present in PDS 456 (O’Brien et al. 2005), similar to those in broad absorption line (BAL) quasars. Such gas would boost the radiation force acting on the wind, via line driving (see Hagino et al. 2016), resulting in an increase in the force multiplier factor.

To quantify further, we start from the definition of the net force (i.e. radiative minus gravitational force) acting on an electron–ion pair within the wind:

$$F_{\text{net}} = \mu m_p v(r_w) \frac{dv}{dr_w} = \frac{L \sigma}{4\pi r_w^2 c} - \frac{GM_{\text{BH}} \mu m_p}{r_w^2}, \quad (1)$$

where $\mu = n_{\text{H}}/n_{\text{e}} \sim 1.2$ is a constant factor accounting for cosmic elemental abundances, m_p is the proton mass, r_w is the radial distance from the central X-ray source and $\sigma = \sigma_{\text{PE}} + \sigma_{\text{T}}$ is

the total cross-section (photoelectric plus Thomson). Rearranging equation (1) gives the equation of motion within the wind:

$$v(r_w) dv = \left(\frac{L \sigma}{4\pi r_w^2 \mu m_p c} - \frac{GM_{\text{BH}}}{r_w^2} \right) dr_w. \quad (2)$$

Now, we define $r_w = R_{\text{in}}$ as the launch radius and $v = v_i$ as the initial velocity of the disc-wind. At larger radii, we assume that $r_w \gg R_{\text{in}}$ and hence $r_w \rightarrow \infty$ and $v \rightarrow v_w$ (i.e. the terminal speed at large radii). Integrating equation (2) between these limits and rearranging gives

$$v_w^2 - v_i^2 = \frac{2GM_{\text{BH}}}{R_{\text{in}}} \left(\frac{L \sigma}{4\pi GM_{\text{BH}} \mu m_p c} - 1 \right). \quad (3)$$

On the right-hand side, the first term is the escape velocity of the particle and the second term is essentially the local L/L_{Edd} within the wind, where the condition $\sigma > \sigma_{\text{T}}$ results from the possible contribution of line driving within the wind, giving a force multiplier factor > 1 . Thus, if we simplify the above equation such that $v_w^2 \gg v_i^2$ and that locally $L > L_{\text{Edd}}$ (e.g. as a result of the force multiplier), we get a dependence between the outflow velocity, the luminosity and R_{in} :

$$\frac{v_w}{c} \sim \left(\frac{\sigma}{2\pi \mu m_p c^3} \right)^{1/2} k_{7-30}^{1/2} L_{7-30}^{1/2} R_{\text{in}}^{-1/2} \propto k_{7-30}^{1/2} L_{7-30}^{1/2} R_{\text{in}}^{-1/2}. \quad (4)$$

Here, $L = k_{7-30} L_{7-30}$, where k_{7-30} is the bolometric correction factor ($\gtrsim 100$ for PDS 456).

Our relationship expressed in equation (4) between the outflow velocity and the ionizing luminosity is ‘steeper’ than what is observed in Fig. 3. There are two possibilities that may explain this; first the wind launch radius, R_{in} , may not be constant with luminosity, or k_{7-30} is variable. In the first scenario as the luminosity increases, the innermost parts of the wind can become fully ionized. Subsequently at higher luminosities, the innermost fastest streamlines of the wind become effectively unobservable. Given that the ionization parameter varies as $\xi \propto L r_w^{-2}$, the effective radius at which the wind is observable increases with luminosity, leading to the ‘flattening’ of the slope as observed.

In the latter case, the terminal velocity should be proportional to the whole bolometric luminosity rather than just what is directly measured in the hard X-ray band. Furthermore, all of the *XMM-Newton* observations of PDS 456 clearly show that the UV flux is considerably less variable than in the X-ray band (see fig. 2 in Matzeu et al. 2017). As a result, k_{7-30} might be variable as a function of luminosity which again may ‘flatten’ the expected correlation. While it is not currently possible to quantitatively test the relation between v_w/c and L (as only the *XMM-Newton* observations have simultaneous optical/UV photometry), the resulting correlation between v_w/c and L is at least qualitatively consistent with a steeper slope of ~ 0.5 . Ongoing optical/UV and X-ray monitoring of PDS 456 with *Swift* will allow us to measure the bolometric correction as a function of luminosity.

One final question is whether it is plausible to radiatively accelerate the wind up to $0.3c$. From equation (3) and taking $L \sim L_{\text{Edd}}$ for PDS 456, for a reasonable launch radius of $30 R_g$ (given the rapid wind variability, M16), only a modest force multiplier factor of $\tau = \sigma/\sigma_{\text{T}} = 2.5$ is required. Furthermore, if PDS 456 is mildly super-Eddington (e.g. $2L_{\text{Edd}}$), then τ approaches unity. Thus, the observed velocities can easily be reproduced within this framework. Note that a comparable positive correlation between the outflow velocity and luminosity has been recently found by Gofford et al. (2015) and Fiore et al. (2017), the latter from collating a sample of known AGN outflows.

3.2 MHD launching scenario

Alternatively, could MHD winds reproduce the above correlation? While to date there is no expected correlation in the MHD wind scenarios (e.g. Ohsuga et al. 2009; Kazanas et al. 2012), the observed velocity can still depend on the continuum photon index, the SED shape and in particular the UV to X-ray slope, α_{ox} .

Indeed Fukumura et al. (2010) predicted that sources with steeper α_{ox} values (i.e. X-ray quiet compared to the UV) should be characterized by faster disc-winds. This is the result of the shape of the ionizing SED. When the X-ray luminosity increases compared to the UV, the innermost parts of the wind become fully ionized. As a result, only the slower streamlines of the wind are observed, launched from further out. In this scenario, we may expect the outflow velocity to decrease with increasing X-ray luminosity, the opposite of what is observed in PDS 456.

None the less, the strong positive correlation in Fig. 3 does not preclude that MHD processes play an important role in providing the initial lift of the wind material off the disc (i.e. v_i). In this regard, we have some evidence that confirms this scenario, where in the low-luminosity observation of PDS 456 in 2013 with *Suzaku*, a strong X-ray flare increased the luminosity by a factor of ~ 4 in just ~ 50 ks which was likely to be magnetically driven (M16). The disc-wind became stronger after the flare was observed, supporting a scenario whereby MHD processes contributed at least to the initial ejection of the material off the disc, only for it to be then accelerated by the intense radiation pressure. Other AGNs that accrete near the Eddington limit, e.g. in luminous quasars and narrow line Seyfert 1 galaxies such as APM 08279+5255 (Chartas et al. 2003), PG1211+143 (Pounds et al. 2003), IRAS F11119+3257 (Tombesi et al. 2015), 1H 0707–495 (Hagino et al. 2016), IRAS 13224–3809 (Parker et al. 2017) are also likely to be promising candidates for radiatively driven winds. However, the contribution of MHD winds are likely to be more important at the lower L/L_{Edd} ratio end of the population, where radiation driving would be difficult without large force multiplier factors.

4 CONCLUSIONS

We analysed all 12 X-ray observations of the fast disc-wind in PDS 456 from 2001 to 2014, observed through its Fe K absorption profile. For the first time, we find a positive correlation between the wind velocity and the intrinsic hard X-ray luminosity. This provides evidence that the accretion disc-wind in PDS 456 is most likely to be radiatively driven, with higher luminosities helping to accelerate the wind to higher terminal velocities.

ACKNOWLEDGEMENTS

We thank the anonymous referee for their helpful report. GM and VB acknowledge support from the Italian Space Agency (ASI INAF NuSTAR I/037/12/0). JR, DML and AL acknowledge the support of STFC. EN is supported by EU's Marie Skłodowska-Curie grant no. 664931. FT acknowledges support by the Programma per Giovani Ricercatori - 2014 'Rita Levi Montalcini'.

REFERENCES

- Akritas M. G., Bershadsky M. A., 1996, *ApJ*, 470, 706
 Behar E., Kaspi S., Reeves J., Turner T. J., Mushotzky R., O'Brien P. T., 2010, *ApJ*, 712, 26
 Chartas G., Brandt W. N., Gallagher S. C., Garmire G. P., 2002, *ApJ*, 579, 169
 Chartas G., Brandt W. N., Gallagher S. C., 2003, *ApJ*, 595, 85
 Di Matteo T., Springel V., Hernquist L., 2005, *Nature*, 433, 604
 Fiore F. et al., 2017, *A&A*, 601, A143
 Fukumura K., Kazanas D., Contopoulos I., Behar E., 2010, *ApJ*, 715, 636
 Fukumura K., Kazanas D., Shrader C., Behar E., Tombesi F., Contopoulos I., 2017, *Nat. Astron.*, 1, 0062
 Gofford J., Reeves J. N., Tombesi F., Braitto V., Turner T. J., Miller L., Cappi M., 2013, *MNRAS*, 430, 60
 Gofford J. et al., 2014, *ApJ*, 784, 77
 Gofford J., Reeves J. N., McLaughlin D. E., Braitto V., Turner T. J., Tombesi F., Cappi M., 2015, *MNRAS*, 451, 4169
 Hagino K., Odaka H., Done C., Gandhi P., Watanabe S., Sako M., Takahashi T., 2015, *MNRAS*, 446, 663
 Hagino K., Odaka H., Done C., Tomaru R., Watanabe S., Takahashi T., 2016, *MNRAS*, 461, 3954
 Hopkins P. F., Elvis M., 2010, *MNRAS*, 401, 7
 Kazanas D., Fukumura K., Behar E., Contopoulos I., Shrader C., 2012, *Astron. Rev.*, 7, 92
 King A., 2003, *ApJ*, 596, L27
 King A. R., Pounds K. A., 2003, *MNRAS*, 345, 657
 Matzeu G. A., Reeves J. N., Nardini E., Braitto V., Costa M. T., Tombesi F., Gofford J., 2016, *MNRAS*, 458, 1311 (M16)
 Matzeu G. A., Reeves J. N., Nardini E., Braitto V., Turner T. J., Costa M. T., 2017, *MNRAS*, 465, 2804
 Nardini E. et al., 2015, *Science*, 347, 860 (N15)
 Nomura M., Ohsuga K., 2017, *MNRAS*, 465, 2873
 O'Brien P. T., Reeves J. N., Simpson C., Ward M. J., 2005, *MNRAS*, 360, L25
 Ohsuga K., Mineshige S., Mori M., Kato Y., 2009, *PASJ*, 61, L7
 Parker M. L. et al., 2017, *Nature*, 543, 83
 Pounds K. A., Reeves J. N., King A. R., Page K. L., O'Brien P. T., Turner M. J. L., 2003, *MNRAS*, 345, 705
 Proga D., Kallman T. R., 2004, *ApJ*, 616, 688
 Proga D., Stone J. M., Kallman T. R., 2000, *ApJ*, 543, 686
 Reeves J. N., O'Brien P. T., Vaughan S., Law-Green D., Ward M., Simpson C., Pounds K. A., Edelson R., 2000, *MNRAS*, 312, L17
 Reeves J. N., O'Brien P. T., Ward M. J., 2003, *ApJ*, 593, L65
 Reeves J. N. et al., 2009, *ApJ*, 701, 493
 Reeves J. N. et al., 2014, *ApJ*, 780, 45
 Reeves J. N., Braitto V., Nardini E., Behar E., O'Brien P. T., Tombesi F., Turner T. J., Costa M. T., 2016, *ApJ*, 824, 20
 Sim S. A., Miller L., Long K. S., Turner T. J., Reeves J. N., 2010, *MNRAS*, 404, 1369
 Simpson C., Ward M., O'Brien P., Reeves J., 1999, *MNRAS*, 303, L23
 Tombesi F., Cappi M., Reeves J. N., Palumbo G. G. C., Yaqoob T., Braitto V., Dadina M., 2010, *A&A*, 521, A57
 Tombesi F., Meléndez M., Veilleux S., Reeves J. N., González-Alfonso E., Reynolds C. S., 2015, *Nature*, 519, 436
 Torres C. A. O., Quast G. R., Coziol R., Jablonski F., de la Reza R., Lépine J. R. D., Gregório-Hetem J., 1997, *ApJ*, 488, L19

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.