



Publication Year	2017
Acceptance in OA@INAF	2020-08-27T14:36:54Z
Title	An orientation-based unification of young jetted active galactic nuclei: the case of 3C 286
Authors	Berton, Marco; FOSCHINI, LUIGI; CACCIANIGA, Alessandro; Ciroi, Stefano; Congiu, Enrico; et al.
DOI	10.3389/fspas.2017.00008
Handle	http://hdl.handle.net/20.500.12386/26899
Journal	FRONTIERS IN ASTRONOMY AND SPACE SCIENCES
Number	4



An Orientation-Based Unification of Young Jetted AGN: The Case of 3C 286

Marco Berton^{1,2*}, Luigi Foschini², Alessandro Caccianiga², Stefano Ciroi¹, Enrico Congiu^{1,2}, Valentina Cracco¹, Michele Frezzato¹, Giovanni La Mura¹ and Piero Rafanelli¹

¹ Dipartimento di Fisica e Astronomia "G. Galilei," Università di Padova, Padova, Italy, ² Brera Astronomical Observatory (INAF), Merate, Italy

OPEN ACCESS

Edited by:

Paola Marziani,
Astronomical Observatory of Padua,
National Institute for Astrophysics
(INAF), Italy

Reviewed by:

Alenka Negrete,
National Autonomous University of
Mexico, Mexico
Victor Manuel Patiño Álvarez,
National Institute of Astrophysics,
Optics and Electronics, Mexico

*Correspondence:

Marco Berton
marco.berton@unipd.it

Specialty section:

This article was submitted to
Cosmology,
a section of the journal
Frontiers in Astronomy and Space
Sciences

Received: 22 May 2017

Accepted: 05 July 2017

Published: 25 July 2017

Citation:

Berton M, Foschini L, Caccianiga A,
Ciroi S, Congiu E, Cracco V,
Frezzato M, La Mura G and Rafanelli P
(2017) An Orientation-Based
Unification of Young Jetted AGN: The
Case of 3C 286.
Front. Astron. Space Sci. 4:8.
doi: 10.3389/fspas.2017.00008

In recent years, the old paradigm according to which only high-mass black holes can launch powerful relativistic jets in active galactic nuclei (AGN) has begun to crumble. The discovery of γ -rays coming from narrow-line Seyfert 1 galaxies (NLS1s), usually considered young and growing AGN harboring a central black hole with mass typically lower than $10^8 M_{\odot}$, indicated that also these low-mass AGN can produce powerful relativistic jets. The search for parent population of γ -ray emitting NLS1s revealed their connection with compact steep-spectrum sources (CSS). In this proceeding we present a review of the current knowledge of these sources, we present the new important case of 3C 286, classified here for the first time as NLS1, and we finally provide a tentative orientation based unification of NLS1s and CSS sources.

Keywords: Active Galactic Nuclei (AGN), narrow-line Seyfert 1 galaxies, compact steep-spectrum sources, relativistic jets, unification models

1. OVERVIEW

Relativistic jets are usually thought to be a product of accretion onto a central supermassive black hole (SMBH) (e.g., Blandford and Znajek, 1977; Blandford and Payne, 1982). In active galactic nuclei (AGN), the interplay between the accretion disk and the SMBH is likely to produce powerful and collimated bipolar outflows. According to the unified model of jetted-AGN (Urry and Padovani, 1995), the two classes of blazars, BL Lacertae objects (BL Lacs) and flat-spectrum radio quasars (FSRQs), are double-lobed radio galaxies, with Fanaroff-Riley (FR) morphological types I and II, respectively, in which the line of sight falls inside the relativistic jet aperture cone. In recent years, this scenario has been slightly revised in a more physical way. BL Lacs and FSRQs indeed seem to reflect two different accretion modes. While BL Lacs have an inefficient accretion onto their central SMBH and a low density environment, FSRQs have instead a strong disk accreting efficiently and a photon- and matter-rich environment (Best and Heckman, 2012). This dichotomy led to the distinction between high-excitation radio galaxies and low-excitation radio galaxies (HERGs and LERGs, respectively) classified according, for example, to the [O III]/H α flux ratio (Laing et al., 1994), and corresponding, respectively, to FSRQ and BL Lac objects when observed at small angles (although exceptions are known, Giommi et al., 2012, see also the review by Foschini, 2017).

Another important aspect of jetted-AGN is that for many years they were thought to be produced only by the most massive black holes harbored in giant passive elliptical galaxies (Laor, 2000; Böttcher and Dermer, 2002; Marscher, 2009). However, the discovery of gamma-ray emission coming from the AGN class known as narrow-line Seyfert 1 galaxies (NLS1s) proved this paradigm

to be wrong (Abdo et al., 2009a,b; Foschini et al., 2015). Low mass AGN, indeed, are also able to launch relativistic jets, whose power is lower because of the well-known scaling of the black hole mass (Heinz and Sunyaev, 2003; Foschini, 2011). This result was confirmed by the identification of low-mass sources among flat-spectrum radio quasars (FSRQs, Ghisellini, 2016), and by the discovery of relativistic jets associated with spiral galaxies (e.g., Keel et al., 2006; Mao et al., 2015).

NLS1s are the most prominent class of low-mass AGN. Classified according to the $\text{FWHM}(H\beta) < 2,000 \text{ km s}^{-1}$ and the ratio between $[\text{O III}]/H\beta < 3$, they are unobscured AGN with relatively low black hole mass ($\lesssim 10^8 M_{\odot}$) and high Eddington ratio, which accounts for the narrowness of the permitted lines and other observational properties (for a review, see Komossa, 2008). A well-known interpretation for these peculiar features is that NLS1s are young AGN still growing and evolving (Mathur, 2000), or sources rejuvenated by a recent merger (Mathur et al., 2012). While a large fraction of NLS1s is radio-quiet, some of them are radio-loud and, as previously mentioned, harbor powerful relativistic jets (Yuan et al., 2008). These radio-loud NLS1s (RLNLS1s) are characterized by a strong and very fast variability consistent with the low-mass scenario, and particularly evident in those with a flat radio spectrum (Foschini et al., 2015).

Another class of radio-loud AGN is known as compact steep-spectrum sources (CSS, see the review by O’Dea, 1998). The radio morphology of these objects is typically characterized by fully developed radio lobes and by a small linear size, with jets confined within the host galaxy. Many authors interpret these properties as signs of young age (Fanti et al., 1995): different estimates of this parameter indeed revealed that they could be radio sources younger than 10^5 years (Owsianik and Conway, 1998; Murgia et al., 1999). CSS sources might eventually evolve to giant double-lobed radio galaxies, or simply switch off because of disk instabilities, going through multiple activity phases during their lifetime (e.g., Czerny et al., 2009; Orienti, 2015). As regular radio galaxies, CSS sources can also be divided in HERGs and LERGs according to their optical spectrum (Kunert-Bajraszewska and Labiano, 2010; Kunert-Bajraszewska et al., 2010).

2. CSS SOURCES AND NLS1S

A possible link between NLS1s and CSS was first suggested by Oshlack et al. (2001), who remarked how the RLNLS1 PKS 2004-447 radio morphology was consistent with that of a CSS source. In the following years, many authors reached the same conclusion (Gallo et al., 2006; Komossa et al., 2006; Yuan et al., 2008; Caccianiga et al., 2014; Schulz et al., 2015). In particular Gu et al. (2015) carried out a VLBA survey on 14 NLS1s at 5 GHz, finding that essentially all RLNLS1s with a steep radio spectrum have a compact morphology, just like CSS. Caccianiga et al. (2017), focusing on the steep-spectrum RLNLS1 SDSS J143244.91+301435.3, found a spectral turnover at low frequency and a small linear size, confirming that this object can be classified as CSS as well.

Beside the morphological similarities, there are other hints that point toward a unification between CSS sources and RLNLS1s. This was investigated under three different points of view in Berton et al. (2016a), using two complete samples of CSS/HERGs and flat-spectrum RLNLS1s (F-NLS1s) limited to $z = 0.6$. A first indication of similarity between these two samples was provided by the V/V_{max} test, which revealed that both of them are consistent with no evolution up to their redshift limit. Moreover, the 1.4 GHz luminosity function (LF) of both samples was derived. Following Urry and Padovani (1995), the effect of relativistic beaming was applied to the observed LF of CSS/HERG to predict the expected shape of the LF of the sources observed along the jet direction. The resulting LF was in good agreement with the observed LF of F-NLS1 thus indicating that at least some CSS/HERGs might be RLNLS1s observed at large angles.

Finally, a comparison between the black hole mass distributions of both samples unveiled again a very good agreement. We repeated here the analysis on the black hole mass distributions on a larger sample of 60 NLS1s. In **Figure 1** we show the distribution of the CSS/HERGs sample by Berton et al. (2016a), compared to all of the masses derived for flat- and steep-spectrum RLNLS1s in Foschini et al. (2015) (42 objects) and Berton et al. (2015) (17 sources), and to that of 3C 286 (see Section 3). It is immediately evident from the plot that 3C 286 is located in the same region as NLS1s. To test the similarity of the distributions, we applied the Kolmogorov-Smirnov test (K-S). The null hypothesis is that the two distributions are originated by the same population. As a rejection threshold, we used a p -value lower than 0.10. Considering CSS/HERGs, the null hypothesis cannot be rejected when comparing them to both flat- and steep-spectrum RLNLS1s, with p -values 0.82 and 0.39, respectively. Testing RLNLS1s as a whole, the p -value is 0.68, again too high to reject the null hypothesis. The mass distribution of CSS/HERGs is therefore statistically consistent




FIGURE 1 | Black hole mass vs. Eddington ratio distribution. Flat-spectrum RLNLS1s are indicated by green triangles, the red circles indicate CSS/HERG sources, the black squares indicate steep-spectrum RLNLS1s, and the orange star represents 3C 286.

with that of RLNLS1s, in agreement with the previously shown results.

3. 3C 286

We present here another interesting case of CSS/HERG, that of 3C 286 ($z = 0.85$). This object is a well-known CSS classified for the first time by Peacock and Wall (1982). It is often used as a calibrator at centimeter wavelengths for both total flux and linear polarization. It shows both the turnover in the radio spectrum around 300 MHz, a spectral index between 1.4 and 50 GHz of $\alpha = -0.61$. It also exhibits the compact morphology typical of CSS: while it remains unresolved at kpc scales, a core-jet structure is visible on pc scale. The jet inclination has been estimated of 48° with respect to the line of sight (e.g., see An et al., 2017, and references therein).

The first optical spectrum of this radio source was obtained by Schmidt (1962), who did not recognize the strongly redshifted Mg II $\lambda 2798$ line. Since its redshift determination (Burbidge and Burbidge, 1969), a significant improvement has been made in optical with the SDSS-BOSS survey (Dawson et al., 2013), which obtained a new spectrum extended up to 10000\AA . The $H\beta$ line, now visible, has a FWHM of $(1811 \pm 169) \text{ km s}^{-1}$, a ratio $[\text{O III}]/H\beta$ of 1.03 ± 0.05 , and Fe II multiplets are clearly visible in the spectrum (see Figure 2). The corresponding uncertainties were calculated with a Monte Carlo method described by Berton et al. (2016b). These parameters clearly indicate that 3C 286 can be classified as NLS1.

We derived its black hole mass following the method described by Foschini et al. (2015). After correcting for redshift, we subtracted the continuum emission by fitting it with a power




FIGURE 2 | Section of the optical spectrum of 3C 286, obtained from the SDSS-BOSS survey. **Top panel:** black solid line is the region between 4,000 and 5,500 Å restframe continuum subtracted, the red dashed line represents the Fe II template, while the blue solid line is the spectrum after the iron subtraction. **Bottom panel:** black solid line indicates the $H\beta$ region, the three Gaussians used to reproduce the line profile are indicated by the green, blue and purple solid line, and the red dashed line represents their sum.

law, plus the Fe II multiplets using an online software (Kovačević et al., 2010; Shapovalova et al., 2012)¹ as shown in top panel of Figure 2. We then fitted $H\beta$ profile using three Gaussians, one to reproduce the narrow component and two to reproduce the broad component (see bottom panel of Figure 1). The narrow component flux was fixed to be 1/10 of $[\text{O III}] \lambda 5007$ (Véron-Cetty et al., 2001) and its FWHM to be the same as $[\text{O II}] \lambda 3727$ ($\sim 600 \text{ km s}^{-1}$). After subtracting the $H\beta$ narrow component, we derived the second-order moment of the broad component σ as a proxy for velocity (Peterson et al., 2004), we obtained the broad-line region (BLR) radius by means of $H\beta$ luminosity. Under the hypothesis of virialized system, the black hole mass is $1.5 \times 10^8 M_\odot$ and the Eddington ratio, defined as the ratio between the bolometric luminosity and the Eddington luminosity of the black hole, is 0.37. Both of these values are consistent with those derived for other radio-loud NLS1s (Foschini et al., 2015, see Figure 1).

Interestingly, this object is a γ -ray emitter, identified in the 3rd Fermi catalog with the source 3FGL J1330.5+3023 (Ackermann et al., 2015)². 3C 286 is therefore the third γ -ray detected steep-spectrum RLNLS1 after RX J2314.9+2243 (Komossa et al., 2015) and B3 1441+476 (Liao et al., 2015). This is a very interesting result, since the number of misaligned NLS1s detected in γ -rays appears to be significantly higher than in other AGN classes. Indeed, only ~ 10 radiogalaxies are known to be γ -ray emitters (Rieger, 2016), and their corresponding beamed population of FSRQs and BL Lacs is made of 1144 sources. Conversely, among NLS1s, currently three misaligned γ -ray sources are known, out of nine oriented objects. The physical interpretation of this phenomenon is not clear yet, but interaction between the jet and the interstellar medium inside the host galaxy could in principle produce the observed γ -ray emission (Migliori et al., 2014).

The classification of this source has another important implication on NLS1s nature. Some authors (e.g., Decarli et al., 2008) suggested that NLS1s are not objects with a low-mass black hole and a high accretion rate. If their BLR had a disk-like shape, when observed pole-on the permitted lines would appear as narrow because of the lack of Doppler broadening. NLS1 would be just low-inclination sources rather AGN with a low-mass central SMBH. The large observing angle (48°) estimated by An et al. (2017) for 3C 286 seems to contradict this hypothesis, at least for this particular object and other RLNLS1s sharing the same property (e.g., Mrk 783, Congiu et al., 2017).

4. YOUNG JETTED-AGN UNIFICATION

As previously mentioned, CSS sources are usually considered as young radiogalaxies still growing (Fanti et al., 1995). The previous results seem to point out that these objects can often be jetted-NLS1s observed at large angles. Of course, this does not mean that all CSS sources are misaligned NLS1s, because some objects probably have a larger black hole mass. The opposite instead might be true, since many of the jetted-NLS1 observed at large angles studied so far appear as CSS/HERGs. An important conclusion can be inferred from this result. If CSS/HERGs are

¹http://servo.aob.rs/FeII_AGN/,

²<http://www.asdc.asi.it/fermi3fgl/>.

young sources, and RLNLS1s are CSS sources, they should be young objects as well. This result, just like the high inclination of 3C 286, is in contrast with the vision of NLS1s as a pole-on view of a type 1 AGN.

According to the unified model of radio-loud AGN (Urry and Padovani, 1995), the parent population of FSRQs are FR II radio galaxies, that is when an FR II is observed along its relativistic jet, it appears as a FSRQ. In a similar way, FR I radio galaxies constitute the parent population of BL Lacs. This interpretation might not be completely appropriate, since there are FR I sources that can be associated with FSRQs, and FR II which appear as BL Lacs. Recently it was suggested that a more accurate unification is between FSRQs and all the FR radio galaxies optically classified as HERG (FR_{HERG}), while BL Lacs should be associated to FR with a LERG-type optical spectrum (FR_{LERG}, Giommi et al., 2012). This revised unification is more physics-based, since it accounts for the different accretion mechanisms (strong disk for FSRQs, weak disk for BL Lacs, Best and Heckman, 2012). CSS/HERGs, in this picture, are the young stage of FR_{HERG} objects. Therefore, if flat-spectrum RLNLS1s are beamed CSS/HERGs, and FSRQs are beamed FR_{HERG}, F-NLS1s could be the young stage of FSRQs (Foschini et al., 2015; Berton et al., 2016a, see **Figure 3**).

This is in agreement with the finding that flat-spectrum RLNLS1s are the low-mass tail of FSRQs (Foschini et al., 2015; Paliya et al., 2016). Indeed, the jet power scales with the black hole mass (Heinz and Sunyaev, 2003; Foschini, 2011) and, as expected, the jet power of RLNLS1s is lower than that observed in FSRQs. Once rescaled for the central mass, though, the normalized power is essentially the same, indicating that the launching mechanism is the same (Foschini et al., 2015). Finally, as shown in Berton et al. (2016a), the radio luminosity function of flat-spectrum RLNLS1s suggests that they might be the low-luminosity tail of FSRQs, indicating again an evolutionary link between these two classes.

All these results can be summarized in the scheme of **Figure 3**. A young jetted-AGN with a strong accretion disk and a photon-rich environment in the central engine surroundings, when observed along its relativistic jets, can appear as a flat-spectrum RLNLS1, characterized by a low black hole mass and a low jet luminosity and power. When the inclination angle increases, the object appears first as a steep-spectrum RLNLS1, with an extended radio emission and a relatively small linear size, that can be classified as a CSS/HERG. Finally, when the line of sight intercepts the molecular torus surrounding the nucleus, the object appears as a type 2 AGN in optical and as a CSS in radio. For older jetted-AGN, instead, the usual unified model holds, with FR_{HERG} (type 1 or 2 quasars in optical, according to the inclination) as parent population of FSRQs.

Such a scheme is probably true only in a statistical sense. Objects like 3C 286 are still fairly rare, and might represent an exception. Moreover, it is unlikely that all type 1 (unobscured) young AGN fit the NLS1 classification. Several other non-NLS1 unobscured AGN with a low-mass black hole exist and were found among γ -ray emitting sources (e.g., Shaw et al., 2012; Ghisellini, 2016). These objects do not appear as NLS1s even when observed at larger angles and in fact, as expected, not all




FIGURE 3 | Unification scheme of jetted-AGN with high accretion rate with respect to the Eddington limit and a high-density photon-rich environment. On the left side, young and smaller sources (NLS1s and CSS sources), compared to older and larger objects (FSRQs and FR_{HERG}).

type 1 CSS/HERGs can be classified as NLS1s. However, despite these notable exceptions, the scheme of **Figure 3** seems to be consistent with several observations.

This orientation-based unification could be confirmed by means of RLNLS1s and CSS/HERGs host galaxy observations. NLS1s usually have a late-type host galaxy (Crenshaw et al., 2003), and some RLNLS1s indeed seem to share the same property (Antón et al., 2008; Caccianiga et al., 2015; Kotilainen et al., 2016; Olguín-Iglesias et al., 2017). CSS sources in general are believed to be instead hosted by early-type hosts (e.g., Best et al., 2005). However, some studies pointed out that their host galaxy can be a spiral as well (Morganti et al., 2011), and that HERGs and LERGs can live in different morphological types, with HERGs showing more late-type hosts (Best and Heckman, 2012). Further observations are needed to better investigate this crucial aspect of young AGN.

5. SUMMARY

In this proceeding we presented a review of the current knowledge about young jetted-AGN, showing as well new results strictly connected to this topic and a tentative orientation-based unification model of these sources. We provided a new classification for the well-known CSS source 3C 286 which, thanks to new observations in the near-infrared, can be now

classified as a NLS1 galaxy. This result confirms that RLNLS1s can often appear, in radio, as CSS sources. Moreover, we confirmed that the black hole mass distribution of CSS/HERGs is very similar to that of RLNLS1s. Along with other results already published in the literature, these findings seem to strengthen the scheme according to which flat-spectrum RLNLS1s are CSS/HERGs observed along their relativistic jet, that can appear as a steep-spectrum RLNLS1 when observed at intermediate angles, and as a type 2 radio galaxy when obscured. RLNLS1s and CSS/HERGs then might represent the young and growing stage of FSRQs and FR_{HERG} radio galaxies, respectively, and constitute a part of the young AGN population. New observations on larger samples, particularly aimed at determining the host galaxy morphology, are needed to provide a final confirmation to this unification model.

FUNDING

This work has been partially supported by PRIN INAF 2014 *Jet and astro-particle physics of gamma-ray blazars* (PI F. Tavecchio).

AUTHOR CONTRIBUTIONS

MB coordinated the project, analyzed the spectroscopic data, contributed to the development of the model, and wrote the manuscript. LF initiated the project, contributed to the development of the model, and provided a critical review of the manuscript. AC contributed to the development of the model, and provided a critical review of the manuscript. SC, EC, VC,

MF, GL, and PR helped collecting the data and provided a critical review of the manuscript.

ACKNOWLEDGMENTS

We are grateful to Mrs. Lucia Zarantonello for drawing the sketch of the young unification scheme. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. Funding for the Sloan Digital Sky Survey has been provided by the Alfred P. Sloan Foundation, and the U.S. Department of Energy Office of Science. The SDSS web site is <http://www.sdss.org>. SDSS-III is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS-III Collaboration including the University of Arizona, the Brazilian Participation Group, Brookhaven National Laboratory, Carnegie Mellon University, University of Florida, the French Participation Group, the German Participation Group, Harvard University, the Instituto de Astrofísica de Canarias, the Michigan State/Notre Dame/JINA Participation Group, Johns Hopkins University, Lawrence Berkeley National Laboratory, Max Planck Institute for Astrophysics, Max Planck Institute for Extraterrestrial Physics, New Mexico State University, University of Portsmouth, Princeton University, the Spanish Participation Group, University of Tokyo, University of Utah, Vanderbilt University, University of Virginia, University of Washington, and Yale University.

REFERENCES

- Abdo, A. A., Ackermann, M., Ajello, M., Axelsson, M., Baldini, L., Ballet, J., et al. (2009a). Fermi/Large Area Telescope Discovery of Gamma-Ray Emission from a Relativistic Jet in the Narrow-Line Quasar PMN J0948+0022. *Astrophys. J.* 699, 976–984. doi: 10.1088/0004-637X/699/2/976
- Abdo, A. A., Ackermann, M., Ajello, M., Axelsson, M., Baldini, L., Ballet, J., et al. (2009b). Multiwavelength Monitoring of the Enigmatic Narrow-Line Seyfert 1 PMN J0948+0022 in 2009 March–July. *Astrophys. J.* 707, 727–737. doi: 10.1088/0004-637X/707/1/727
- Ackermann, M., Ajello, M., Atwood, W. B., Baldini, L., Ballet, J., Barbiellini, G., et al. (2015). The Third Catalog of Active Galactic Nuclei Detected by the Fermi Large Area Telescope. *Astrophys. J.* 810:14. doi: 10.1088/0004-637X/810/1/14
- An, T., Lao, B.-Q., Zhao, W., Mohan, P., Cheng, X.-P., Cui, Y.-Z., et al. (2017). Parsec-scale jet properties of the gamma-ray quasar 3C 286. *Mon. Not. R. Astron. Soc.* 466, 952–959. doi: 10.1093/mnras/stw2887
- Antón, S., Browne, I. W. A., and Marchã, M. J. (2008). The colour of the narrow line Sy1-blazar 0324+3410. *Astron. Astrophys.* 490, 583–587. doi: 10.1051/0004-6361/20078926
- Berton, M., Caccianiga, A., Foschini, L., Peterson, B. M., Mathur, S., Terreran, G., et al. (2016a). Compact steep-spectrum sources as the parent population of flat-spectrum radio-loud narrow-line Seyfert 1 galaxies. *Astron. Astrophys.* 591:A98. doi: 10.1051/0004-6361/201628171
- Berton, M., Foschini, L., Ciroi, S., Cracco, V., La Mura, G., Di Mille, F., et al. (2016b). [O III] line properties in two samples of radio-emitting narrow-line Seyfert 1 galaxies. *Astron. Astrophys.* 591:A88. doi: 10.1051/0004-6361/201527056
- Berton, M., Foschini, L., Ciroi, S., Cracco, V., La Mura, G., Lister, M. L., et al. (2015). Parent population of flat-spectrum radio-loud narrow-Line Seyfert 1 galaxies. *Astron. Astrophys.* 578:A28. doi: 10.1051/0004-6361/201525691
- Best, P. N., and Heckman, T. M. (2012). On the fundamental dichotomy in the local radio-AGN population: accretion, evolution and host galaxy properties. *Mon. Not. R. Astron. Soc.* 421, 1569–1582. doi: 10.1111/j.1365-2966.2012.20414.x
- Best, P. N., Kauffmann, G., Heckman, T. M., Brinchmann, J., Charlot, S., Ivezić, Ž., et al. (2005). The host galaxies of radio-loud active galactic nuclei: mass dependences, gas cooling and active galactic nuclei feedback. *Mon. Not. R. Astron. Soc.* 362, 25–40. doi: 10.1111/j.1365-2966.2005.09192.x
- Blandford, R. D., and Payne, D. G. (1982). Hydromagnetic flows from accretion discs and the production of radio jets. *Mon. Not. Roy. Astron. Soc.* 199, 883–903. doi: 10.1093/mnras/199.4.883
- Blandford, R. D., and Znajek, R. L. (1977). Electromagnetic extraction of energy from Kerr black holes. *Mon. Not. Roy. Astron. Soc.* 179, 433–456.
- Böttcher, M., and Dermer, C. D. (2002). An evolutionary scenario for blazar unification. *Astrophys. J.* 564, 86–91. doi: 10.1093/mnras/179.3.433
- Burbidge, G. R., and Burbidge, E. M. (1969). Red-shifts of Quasi-stellar Objects and Related Extragalactic Systems. *Nature* 222, 735–741. doi: 10.1038/222735a0
- Caccianiga, A., Antón, S., Ballo, L., Dallacasa, D., Della Ceca, R., Fanali, R., et al. (2014). SDSS J143244.91+301435.3: a link between radio-loud narrow-line Seyfert 1 galaxies and compact steep-spectrum radio sources? *Mon. Not. R. Astron. Soc.* 441, 172–186. doi: 10.1093/mnras/stu508
- Caccianiga, A., Antón, S., Ballo, L., Foschini, L., Maccacaro, T., Della Ceca, R., et al. (2015). WISE colours and star formation in the host galaxies of radio-loud narrow-line Seyfert 1. *Mon. Not. R. Astron. Soc.* 451, 1795–1805. doi: 10.1093/mnras/stv939
- Caccianiga, A., Dallacasa, D., Antón, S., Ballo, L., Berton, M., Mack, K.-H., et al. (2017). SDSSJ143244.91+301435.3 at VLBI: a compact radio galaxy in a narrow-line Seyfert 1. *Mon. Not. R. Astron. Soc.* 464, 1474–1480. doi: 10.1093/mnras/stw2471

- Congiu, E., Berton, M., Giroletti, M., Antonucci, R., Caccianiga, A., Kharb, P., et al. (2017). Kiloparsec-scale emission in the narrow-line Seyfert 1 galaxy Mrk 783. *Astron. Astrophys.* 603:A32. doi: 10.1051/0004-6361/201730616
- Crenshaw, D. M., Kraemer, S. B., and Gabel, J. R. (2003). The host galaxies of narrow-line Seyfert 1 galaxies: evidence for bar-driven fueling. *Astron. J.* 126, 1690–1698. doi: 10.1086/377625
- Czerny, B., Siemiginowska, A., Janiuk, A., Nikiel-Wroczyński, B., and Stawarz, L. (2009). Accretion disk model of short-timescale intermittent activity in young radio sources. *Astrophys. J.* 698, 840–851. doi: 10.1088/0004-637X/698/1/840
- Dawson, K. S., Schlegel, D. J., Ahn, C. P., Anderson, S. F., Aubourg, É., Bailey, S., et al. (2013). The Baryon oscillation spectroscopic survey of SDSS-III. *Astron. J.* 145:10. doi: 10.1088/0004-6256/145/1/10
- Decarli, R., Dotti, M., Fontana, M., and Haardt, F. (2008). Are the black hole masses in narrow-line Seyfert 1 galaxies actually small? *Mon. Not R. Astron. Soc.* 386, L15–L19. doi: 10.1111/j.1745-3933.2008.00451.x
- Fanti, C., Fanti, R., Dallacasa, D., Schilizzi, R. T., Spencer, R. E., and Stanghellini, C. (1995). Are compact steep-spectrum sources young? *Astron. Astrophys.* 302:317.
- Foschini, L. (2011). Accretion and jet power in active galactic nuclei. *Res. Astron. Astrophys.* 11, 1266–1278. doi: 10.1088/1674-4527/11/11/003
- Foschini, L. (2017). What we talk about when we talk about blazars? *Front. Astron. Space Sci.* 4:6. doi: 10.3389/fspas.2017.00006
- Foschini, L., Berton, M., Caccianiga, A., Ciroi, S., Cracco, V., Peterson, B. M., et al. (2015). Properties of flat-spectrum radio-loud narrow-line Seyfert 1 galaxies. *Astron. Astrophys.* 575:A13. doi: 10.1051/0004-6361/201424972
- Gallo, L. C., Edwards, P. G., Ferrero, E., Kataoka, J., Lewis, D. R., Ellingsen, S. P., et al. (2006). The spectral energy distribution of PKS 2004-447: a compact steep-spectrum source and possible radio-loud narrow-line Seyfert 1 galaxy. *Mon. Not R. Astron. Soc.* 370, 245–254. doi: 10.1111/j.1365-2966.2006.10482.x
- Ghisellini, G. (2016). The Blazar Sequence 2.0. *Galaxies* 4:36. doi: 10.3390/galaxies400036
- Giommi, P., Padovani, P., Polenta, G., Turriziani, S., D’Elia, V., and Piranomonte, S. (2012). A simplified view of blazars: clearing the fog around long-standing selection effects. *Mon. Not R. Astron. Soc.* 420, 2899–2911. doi: 10.1111/j.1365-2966.2011.20044.x
- Gu, M., Chen, Y., Komossa, S., Yuan, W., Shen, Z., Wajima, K., et al. (2015). The radio properties of radio-loud narrow-line Seyfert 1 galaxies on parsec scales. *Astrophys. J.* 221:3. doi: 10.1088/0067-0049/221/1/3
- Heinz, S., and Sunyaev, R. A. (2003). The non-linear dependence of flux on black hole mass and accretion rate in core-dominated jets. *Mon. Not R. Astron. Soc.* 343, L59–L64. doi: 10.1046/j.1365-8711.2003.06918.x
- Keel, W. C., White, R. E. III, Owen, F. N., and Ledlow, M. J. (2006). The spiral host galaxy of the double radio source 0313-192. *Astron. J.* 132, 2233–2242. doi: 10.1086/508340
- Komossa, S. (2008). “Narrow-line Seyfert 1 Galaxies,” in *Revista Mexicana de Astronomía y Astrofísica Conference Series*, Vol. 32 of *Revista Mexicana de Astronomía y Astrofísica Conference Series* (Huatulco, MX), 86–92.
- Komossa, S., Voges, W., Xu, D., Mathur, S., Adorf, H.-M., Lemson, G., et al. (2006). Radio-loud Narrow-Line Type 1 Quasars. *Astron. J.* 132, 531–545. doi: 10.1086/505043
- Komossa, S., Xu, D., Fuhrmann, L., Grupe, D., Yao, S., Fan, Z., et al. (2015). What powers the radio-loud narrow-line Seyfert 1 galaxy RX J2314.9+2243?: a view onto its central engine from radio to X-rays. *Astron. Astrophys.* 574:A121. doi: 10.1051/0004-6361/201424814
- Kotilainen, J. K., León-Tavares, J., Olguín-Iglesias, A., Baes, M., Anórve, C., Chavushyan, V., et al. (2016). Discovery of a pseudobulge galaxy launching powerful relativistic jets. *Astrophys. J.* 832:157. doi: 10.3847/0004-637X/832/2/157
- Kovačević, J., Popović, L. Č., and Dimitrijević, M. S. (2010). Analysis of Optical Fe II Emission in a Sample of Active Galactic Nucleus Spectra. *Astrophys. J.* 189, 15–36. doi: 10.1088/0067-0049/189/1/15
- Kunert-Bajraszewska, M., Gawroński, M. P., Labiano, A., and Siemiginowska, A. (2010). A survey of low-luminosity compact sources and its implication for the evolution of radio-loud active galactic nuclei - I. Radio data. *Mon. Not R. Astron. Soc.* 408, 2261–2278. doi: 10.1111/j.1365-2966.2010.17271.x
- Kunert-Bajraszewska, M., and Labiano, A. (2010). A survey of low-luminosity compact sources and its implication for the evolution of radio-loud active galactic nuclei - II. Optical analysis. *Mon. Not R. Astron. Soc.* 408, 2279–2289. doi: 10.1111/j.1365-2966.2010.17300.x
- Laing, R. A., Jenkins, C. R., Wall, J. V., and Unger, S. W. (1994). “Spectrophotometry of a complete sample of 3CR radio sources: implications for unified models,” in *The Physics of Active Galaxies*, Vol. 54 of *Astronomical Society of the Pacific Conference Series* (Canberra), 201.
- Laor, A. (2000). On Black Hole Masses and Radio Loudness in Active Galactic Nuclei. *Astrophys. J.* 543, L111–L114. doi: 10.1086/317280
- Liao, N.-H., Liang, Y.-F., Weng, S.-S., Gu, M.-F., and Fan, Y.-Z. (2015). Discovery of γ -ray emission from a steep radio spectrum NLS1 B3 1441+476. *ArXiv e-prints*. arXiv:1510.05584
- Mao, M. Y., Owen, F., Duffin, R., Keel, B., Lacy, M., Momjian, E., et al. (2015). J1649+2635: a grand-design spiral with a large double-lobed radio source. *Mon. Not R. Astron. Soc.*, 446, 4176–4185. doi: 10.1093/mnras/stu2302
- Marscher, A. P. (2009). Jets in active galactic nuclei. *ArXiv e-prints*. arXiv:0909.2576
- Mathur, S. (2000). Narrow-line Seyfert 1 galaxies and the evolution of galaxies and active galaxies. *Mon. Not R. Astron. Soc.* 314, L17–L20. doi: 10.1046/j.1365-8711.2000.03530.x
- Mathur, S., Fields, D., Peterson, B. M., and Grupe, D. (2012). Supermassive Black Holes, Pseudobulges, and the Narrow-line Seyfert 1 Galaxies. *Astrophys. J.* 754:146. doi: 10.1088/0004-637X/754/2/146
- Migliori, G., Siemiginowska, A., Kelly, B. C., Stawarz, L., Celotti, A., and Begelman, M. C. (2014). Jet Emission in Young Radio Sources: A Fermi Large Area Telescope Gamma-Ray View. *Astrophys. J.* 780:165. doi: 10.1088/0004-637X/780/2/165
- Morganti, R., Holt, J., Tadhunter, C., Ramos Almeida, C., Dicken, D., Inskip, K., et al. (2011). PKS 1814-637: a powerful radio-loud AGN in a disk galaxy. *Astron. Astrophys.* 535:A97. doi: 10.1051/0004-6361/201117686
- Murgia, M., Fanti, C., Fanti, R., Gregorini, L., Klein, U., Mack, K.-H., et al. (1999). Synchrotron spectra and ages of compact steep spectrum radio sources. *Astron. Astrophys.* 345, 769–777.
- O’Dea, C. P. (1998). The Compact Steep-Spectrum and Gigahertz Peaked-Spectrum Radio Sources. *Astron. Soc. Pacific*, 110, 493–532. doi: 10.1086/316162
- Olguín-Iglesias, A., Kotilainen, J. K., León Tavares, J., Chavushyan, V., and Anórve, C. (2017). Evidence of bar-driven secular evolution in the gamma-ray narrow-line Seyfert 1 galaxy FBQS J164442.5+261913. *Mon. Not R. Astron. Soc.* 467, 3712–3722. doi: 10.1093/mnras/stx022
- Orienti, M. (2015). Radio properties of Compact Steep Spectrum and GHz-Peaked Spectrum radio sources. *ArXiv e-prints*. arXiv:1511.00436
- Oshlack, A. Y. K. N., Webster, R. L., and Whiting, M. T. (2001). A Very Radio Loud Narrow-Line Seyfert 1: PKS 2004-447. *Astrophys. J.* 558, 578–582. doi: 10.1086/322299
- Owsianik, I., and Conway, J. E. (1998). First detection of hotspot advance in a Compact Symmetric Object. Evidence for a class of very young extragalactic radio sources. *Astron. Astrophys.* 337, 69–79.
- Paliya, V. S., Rajput, B., Stalin, C. S., and Pandey, S. B. (2016). Broadband Observations of the Gamma-Ray Emitting Narrow Line Seyfert 1 Galaxy SBS 0846+513. *Astrophys. J.* 819:121. doi: 10.3847/0004-637X/819/2/121
- Peacock, J. A., and Wall, J. V. (1982). Bright extragalactic radio sources at 2.7 GHz. II - Observations with the Cambridge 5-km telescope. *Mon. Not R. Astron. Soc.* 198, 843–860. doi: 10.1093/mnras/198.3.843
- Peterson, B. M., Ferrarese, L., Gilbert, K. M., Kaspi, S., Malkan, M. A., Maoz, D., et al. (2004). Central Masses and Broad-Line Region Sizes of Active Galactic Nuclei. II. A Homogeneous Analysis of a Large Reverberation-Mapping Database. *Astrophys. J.* 613, 682–699. doi: 10.1086/423269
- Rieger, F. M. (2016). Gamma-Rays from Non-Blazar AGN. *ArXiv e-prints*. arXiv:1611.02986
- Schmidt, M. (1962). Spectrum of a Stellar Object Identified with the Radio Source 3c 286. *Astrophys. J.* 136:684. doi: 10.1086/147424
- Schulz, R., Kreikenbohm, A., Kadler, M., Ojha, R., Ros, E., Stevens, J., et al. (2015). The Gamma-Ray Emitting Radio-Loud Narrow-Line Seyfert 1 Galaxy PKS 2004-447 II. The Radio View. *ArXiv e-prints*. arXiv:1511.02631
- Shapovalova, A. I., Popović, L. Č., Burenkov, A. N., Chavushyan, V. H., Ilić, D., Kovačević, A., et al. (2012). Spectral Optical Monitoring of the Narrow-line Seyfert 1 Galaxy Ark 564. *Astrophys. J.* 202:10. doi: 10.1088/0067-0049/202/1/10

- Shaw, M. S., Romani, R. W., Cotter, G., Healey, S. E., Michelson, P. F., Readhead, A. C. S., et al. (2012). Spectroscopy of Broad-line Blazars from 1LAC. *Astrophys. J.* 748:49. doi: 10.1088/0004-637X/748/1/49
- Urry, C. M., and Padovani, P. (1995). Unified Schemes for Radio-Loud Active Galactic Nuclei. *Publ. Astron. Soc. Pacific* 107:803. doi: 10.1086/133630
- Véron-Cetty, M.-P., Véron, P., and Gonçalves, A. C. (2001). A spectrophotometric atlas of Narrow-Line Seyfert 1 galaxies. *Astron. Astrophys.* 372, 730–754. doi: 10.1051/0004-6361:20010489
- Yuan, W., Zhou, H. Y., Komossa, S., Dong, X. B., Wang, T. G., Lu, H. L., et al. (2008). A Population of Radio-Loud Narrow-Line Seyfert 1 Galaxies with Blazar-Like Properties? *Astrophys. J.* 685:801. doi: 10.1086/591046

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Berton, Foschini, Caccianiga, Ciroi, Congiu, Cracco, Frezzato, La Mura and Rafanelli. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.