



<b>Publication Year</b>	2017
<b>Acceptance in OA</b>	2020-09-14T06:49:21Z
<b>Title</b>	A study of the cold cores population in the Perseus star-forming regions
<b>Authors</b>	PEZZUTO, Stefano, Fiorellino, E., BENEDETTINI, Milena, SCHISANO, EUGENIO, ELIA, Davide Quintino, André, P., Könyves, V., Ladjelate, B., Di Francesco, J., Piccotti, L., Herschel Gould Belt Survey Consortium
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/27341">http://hdl.handle.net/20.500.12386/27341</a>
<b>Journal</b>	MEMORIE DELLA SOCIETA ASTRONOMICA ITALIANA
<b>Volume</b>	88



# A study of the cold cores population in the Perseus star-forming regions

S. Pezzuto<sup>1</sup>, E. Fiorellino<sup>1,2</sup>, M. Benedettini<sup>1</sup>, E. Schisano<sup>1</sup>, D. Elia<sup>1</sup>, P. André<sup>3</sup>,  
V. Könyves<sup>3</sup>, B. Ladjelate<sup>3</sup>, J. Di Francesco<sup>4,5</sup>,  
L. Piccotti<sup>6,1</sup>, and the Herschel Gould Belt Survey Consortium

<sup>1</sup> INAF – IAPS, via Fosso del Cavaliere 100, I-00133 Roma, Italy  
e-mail: pezzuto@iaps.inaf.it

<sup>2</sup> Dip. di Fisica, Uni. di Roma “Sapienza”, Piazzale Aldo Moro 5, I-00185 Roma, Italy

<sup>3</sup> Laboratoire AIM, CEA/DSM-CNRS-Université Paris Diderot, IRFU/Service d’Astrophysique, CEA Saclay, 91191 Gif-sur-Yvette, France

<sup>4</sup> NRC Herzberg Astronomy and Astrophysics, 5071 West Saanich Rd, Victoria, BC, V9E 2E7, Canada

<sup>5</sup> Department of Physics and Astronomy, University of Victoria, 3800 Finnerty Road, Victoria, BC, Canada V8P 5C2

<sup>6</sup> Observatorio Astronómico Ramón María Aller, P.O. Box 197, Santiago de Compostela, Spain

**Abstract.** As part of the Herschel Gould Belt survey, the Perseus star-forming cloud was observed with the Herschel PACS and SPIRE instruments. Source catalogs are preliminary, as well as the here presented core mass function.


## 1. Introduction

The complex of star-forming regions in Perseus is located at distances of  $\sim 250\text{--}300$  pc. It hosts a number of well-known sites of active star formation like NGC 1333, L1448, L1455, B1, IC348. Perseus was observed as part of the Herschel Gould Belt survey (HGBS, André et al. 2010) which aims at obtaining a complete census of prestellar cores and Class 0 sources in the closest star-forming regions. The survey was carried out with the Herschel (Pilbratt et al. 2010) instruments PACS (Poglitsch et al. 2010) and SPIRE (Griffin et al. 2010). These data have been already exploited in a few papers: Sadavoy et al. (2012) made a multiwavelength study of a few young sources in B1-


E; Pezzuto et al. (2012) reported on the analysis of the SED of B1-bS and B1-bN, two first hydrostatic cores candidates; Sadavoy et al. (2014) identified a first list of Class 0 protostars. In Figure 1 we show the column density map derived from our observations.

## 2. Sources extraction and photometry

Compact sources have been identified and their flux measured with *getsources* (Men’shchikov et al. 2012). Not-reliable sources have been removed and the resulting list has been cross-checked with external databases (NED, WISE, Spitzer, Simbad) to remove possible contaminants. A detailed description of the selec-



**Fig. 1.** The column density map of the Perseus complex; black areas are denser. The molecular hydrogen column density runs from  $2.5 \cdot 10^{20} \text{ cm}^{-2}$  up to  $1.4 \cdot 10^{23} \text{ cm}^{-2}$ .



**Fig. 2.** The CMF for Perseus: the Salpeter law  $dN \propto M^{-1.35}$ , arbitrarily scaled, is shown for comparison.

tion procedure can be found in Könyves et al. (2015, hereinafter Paper I). A tentative list of 838 sources has been generated; 94% are robust identifications since they are also found with another code (*CuTE*x, Molinari et al. 2011). In particular, 37% are spatially coincident, within  $6''$ , and 57% fall within the ellipti-

cal size of the sources computed by *getsources* at  $250 \mu\text{m}$ .

### 3. Conclusions

For all the 743 (89%) starless cores having reliable SED fits (see Paper I), in Figure 2 we show a preliminary CMF ( $dN/d\log M$ ); 62% (461 out of 743) are provisionally identified as gravitationally bound. The ongoing analysis is finalized to make the catalog more robust, by identifying and excluding spurious sources, especially at the low-mass end of the CMF.

### References

- André, P., et al. 2010, *A&A*, 518, L102
- Griffin, M. J., et al. 2010, *A&A*, 518, L3
- Könyves, V. et al. 2015 *A&A*, 584, A91
- Men'shchikov, A., et al. 2012, *A&A*, 542, A81
- Molinari, S., et al. 2011, *A&A*, 530, A133
- Pezzuto, S., et al. 2012, *A&A*, 547, A54
- Pilbratt, G. L., et al. 2010, *A&A*, 518, L1
- Poglitsch, A., et al. 2010, *A&A*, 518, L2
- Sadavoy, S. I., et al. 2012, 540, A10
- Sadavoy, S. I., et al. 2014, *ApJ*, 787, L18