



<b>Publication Year</b>	2018
<b>Acceptance in OA @INAF</b>	2020-10-13T13:52:20Z
<b>Title</b>	Methanol masers and magnetic field in IRAS18089-1732
<b>Authors</b>	Dall'Olio, Daria; Vlemmings, W. H. T.; Surcis, Gabriele; Beuther, H.; Lankhaar, B.; et al.
<b>DOI</b>	10.1017/S174392131701016X
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/27763">http://hdl.handle.net/20.500.12386/27763</a>
<b>Series</b>	PROCEEDINGS OF THE INTERNATIONAL ASTRONOMICAL UNION
<b>Number</b>	vol. 13, S336

# Methanol masers and magnetic field in IRAS18089-1732

Daria Dall’Olio<sup>1</sup>, W. H. T. Vlemmings<sup>1</sup>, G. Surcis<sup>2</sup>, H. Beuther<sup>3</sup>, B.  
Lankhaar<sup>1</sup>, M. V. Persson<sup>1</sup>, A. M. S. Richards<sup>4</sup> and E. Varenius<sup>4</sup>

<sup>1</sup>Department of Space, Earth and Environment, Chalmers University of Technology, Onsala  
Space Observatory, Observatorievägen 90, 43992 Onsala, Sweden  
email: [daria.dalolio@chalmers.se](mailto:daria.dalolio@chalmers.se)

<sup>2</sup>INAF–Osservatorio Astronomico di Cagliari, Via della Scienza 5, 09047 Selargius, Italy

<sup>3</sup>Max-Planck-Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany

<sup>4</sup>Jodrell Bank Centre for Astrophysics, Department of Physics and Astronomy, University of  
Manchester, M139PL Manchester, UK


**Abstract.** Theoretical simulations have shown that magnetic fields play an important role in massive star formation: they can suppress fragmentation in the star forming cloud, enhance accretion via disc and regulate outflows and jets. However, models require specific magnetic configurations and need more observational constraints to properly test the impact of magnetic fields. We investigate the magnetic field structure of the massive protostar IRAS18089-1732, analysing 6.7 GHz methanol maser MERLIN observations. IRAS18089-1732 is a well studied high mass protostar, showing a hot core chemistry, an accretion disc and a bipolar outflow. An ordered magnetic field oriented around its disc has been detected from previous observations of polarised dust. This gives us the chance to investigate how the magnetic field at the small scale probed by masers relates to the large scale field probed by the dust.

**Keywords.** stars: formation, stars: magnetic field, masers, polarization

---

## 1. Introduction

The importance of the magnetic field in high mass star formation (HMSF) is not yet fully clear and there are still many open questions concerning its role in the accretion processes and generation of jets and outflows. In the past few years, masers have been successfully used to study the magnetic field at few AU scales around massive protostars. Thanks to their narrow and strong spectral lines and through their polarized emissions, it is possible to reconstruct the morphology and the strength along the line of sight of the magnetic field, by measuring linear polarization angles and Zeeman splitting (Vlemmings et al. 2010; Surcis et al. 2015). This can be done on scale comparable to circumstellar discs ( $\sim 100$  au). In order to improve our models and to build a complete picture of the magnetic field evolution, we need more observational constraints; in particular we need to verify that the magnetic field at small scales probed by masers traces the field at larger scales, probed, for example, by the dust, and not small-scale fluctuations. Currently, few observations of both masers and dust polarisation exist towards the same regions (e.g. Surcis et al. 2014). IRAS18089-1732 is a well studied high mass star forming region, showing a hot core chemistry and a disk-outflow system. Previous observations of polarized dust made with SMA revealed an ordered magnetic field oriented around the disk at large scales ( $\sim 5000$  au, Beuther et al. 2010).



**Figure 1.** Masers identified in March (left) and April (right). The zoom is made on the region marked by the dashed grey boxes. Triangles and circles represent masers. The different sizes of the triangles and the circles represent the intensity. Line segments mark the direction of the polarisation angle for the maser features that show linear polarisation. The average direction of the resulting magnetic field  $\Phi_B$  obtained for two groups of masers is indicated in the bottom right corners of each panel.


## 2. Masers and dust probe the same large-scale magnetic field

From the analysis of a three-epoch MERLIN observation of the 6.7 GHz  $\text{CH}_3\text{OH}$  (methanol) masers, Dall’Olio et al. 2017 identified two groups separated in velocity and polarisation angles as showed in Fig. 1. Masers that are tracing the disc (triangles) show a velocity range, between  $30.036.4 \text{ km s}^{-1}$ , and an orientation of the magnetic field  $62^\circ \pm 3^\circ$ . The group of masers that are tracing a cloud of gas close to the outflow (circles) shows a velocity range between  $37.739.2 \text{ km s}^{-1}$ , and an orientation of the magnetic field  $14^\circ \pm 4^\circ$ . From the analysis of the group of masers generated in the same region of the disc, Dall’Olio et al. 2017 obtained the first polarised map of the masers for IRAS 18089-1732 and showed that the small-scale magnetic field probed by the masers is consistent with the large-scale magnetic field traced by the dust. This confirms that methanol masers trace the large scale field, and that the large scale field component, even at the AU scale of the masers, dominates over any small scale field fluctuations Fig. 2.

*We acknowledge funding from the ERC under the European Union’s Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreement n. 614264.*

## References

- Beuther, H., Vlemmings, W. H. T., Rao, R., & van der Tak, F. F. S. 2010, ApJ, 724, L113  
 Dall’Olio, D., Vlemmings, W. H. T., Surcis, G., et al. 2017, ArXiv [arXiv:1708.02961]  
 Surcis, G., Vlemmings, W. H. T., van Langevelde, H. J., et al. 2015, A&A, 578, A102  
 Surcis, G., Vlemmings, W. H. T., van Langevelde, H. J., Moscadelli, L., & Hutawarakorn Kramer, B. 2014, A&A, 563, A30  
 Vlemmings, W. H. T., Surcis, G., Torstenson, K. J. E., & van Langevelde, H. J. 2010, MNRAS, 404, 134



**Figure 2.** The figure shows that the magnetic field in the maser regions (triangles and long dashed segments) presents the same orientation as observed from the dust tracing the disc (contours and isolated short segments). The orientation is preserved in the two epochs March (left panel) and April (right panel).