

Publication Year	2018
Acceptance in OA@INAF	2020-11-19T11:26:02Z
Title	Identification of low luminosity high redshift galaxies using galaxy clusters as cosmic telescopes
Authors	Delgado-Correal, Camilo; Rosati, Piero; Bartosch Caminha, Gabriel; GRILLO, CLAUDIO; MERCURIO, AMATA; et al.
DOI	10.1142/9789813226609_0242
Handle	http://hdl.handle.net/20.500.12386/28443

Identification of low luminosity high redshift galaxies using galaxy clusters as cosmic telescopes

Camilo Delgado-Correal*, Piero Rosati and Gabriel Bartosch Caminha

Universita degli Studi di Ferrara, Via Saragat 1, I-44122 Ferrara, Italy *E-mail: delgado@fe.infn.it

Claudio Grillo

Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen Juliane Maries Vej 30, DK-2100 Copenhagen, Denmark E-mail: grillo@dark-cosmology.dk

Amata Mercurio

INAF - Osservatorio Astronomico di Capodimonte Via Moiariello 16, I-80131 Napoli, Italy E-mail: mercurio@na.astro.it

Italo Balestra

University Observatory Munich Scheinerstrasse 1, 81679 Munich, Germany E-mail: balestra@mpe.mpg.de

Mario Nonino

Via G. B. Tiepolo 11, I-34143, Trieste, Italy E-mail: nonino@oats.inaf.it

Eros Vanzella

INAF - Osservatorio Astronomico di Bologna Via Ranzani 1, 40127 Bologna, Italia E-mail: eros.vanzella@oabo.inaf.it

Current models of structure formation suggest that the first galaxies formed at z > 10when the universe was < 500 Myr old. The detection and characterization of galaxies at these early epochs is therefore critical to estimate the star formation rate density and their contribution to the reionization. The CLASH project (Cluster Lensing And Supernova survey with Hubble) combines an HST Treasury program to obtain panchromatic (ACS + WFC3) imaging of 25 carefully selected massive clusters, with other multiwavelength observations, including a large spectroscopic campaign with VLT/VIMOS. Gravitational lensing, which is particularly powerful in several CLASH clusters, boost the efficiency of finding low-luminosity (i.e. L < L*) galaxies, which are thought to play a critical role in reionizing the Universe beyond $z \sim 10$. In this letter, we will give some highlights of the construction of a sample of ~ 200 magnified lensed galaxies at 2 < z < 7, collected as part of the CLASH-VLT project, whose spectro-photometric data can be used to characterize the physical properties of the low-luminosity galaxy population at high-z.

Keywords: High-z galaxies; cluster galaxies; magnification map; gravitational lensing.

1. Building an array of cosmic telescopes

The epoch of reionization marks a major phase transition of the Universe, during which the intergalactic space became transparent to UV photons. Determining when this occurred, the physical processes involved and the sources of ionizing radiation represents one of the major goals in observational cosmology. The production of ionizing radiation is most probably driven by star formation and/or nuclear activity, but their global contribution to the ionizing background is still matter of debate.¹ Irrespective of the nature of ionizing radiation, the general consensus is that the faint sources dominate the ionizing background.^{2,3} This implicitly assumes that ionizing photons are not trapped in faint sources and can escape. It is therefore important to push studies into low luminosity regimes (L < 0.1L*, which corresponds to $m_{1500} > 27.1$ at z = 3).



Fig. 1. HST gallery of 25 clusters from the CLASH survey (boxes indicate the 13 included in the CLASH-VLT survey).

Searching for low luminosities high redshift galaxies, thus complementing blank field studies, is a primary goal of surveys carried out through lensing clusters. Several spectroscopically confirmed high-z galaxies have recently been identified in deep field searches, e.g. Refs. 8, 9, 10, 11, 12 and 13. Gravitational lensing offer a unique opportunity to push these studies to significantly fainter luminosities, before the the advent of JWST and giant ground-based telescopes in the next decade. Our collaboration has recently produced high precision lensing models of CLASH clusters, thanks to the redshift measurement of many multiply lensed systems from the CLASH-VLT spectroscopic campaign,⁴ which are critical for strong lensing mass reconstruction techniques. This effort has complemented photometric redshift information readily available from the multi-band HST survey (524 HST orbits of Hubble with more 1000 hours observations of 25 clusters in 16 filters from the CLASH program).⁵ CLASH-VLT has used the VIMOS wide-field spectrograph to obtain ~ 30,000 redshifts in 12 CLASH Clusters at z = 0.2 - 0.6 in the south, with the spectroscopic identification of 500 - 1000 members per cluster. For example, in the case of MACS J0416.1 – 2403, we identified ~ 800 clusters members⁶ and 30 multiple images with redshifts between 1.6 and 3.2.⁷ With these velocity data, we can measure the total mass density profile from the dynamical analysis with unprecedented accuracy (using the Jean equation) and compare it with the mass profiles derived from weak and strong lensing analyses, as well as X-ray hydrostatic masses from Chandra data.

2. Exploiting magnification to probe fainter fluxes



Fig. 2. Distribution of rest-frame UV absolute magnitudes and redshifts for samples of galaxies drawn from blank deep fields and lensed field surveys (CLASH-VLT).

Over the last decade, several groups have used galaxy clusters as cosmic lenses to investigate properties of distant sources, ¹⁵ most of them are galaxies, ^{14,16,17} in some rare cases quasars or more recently even transient objects, including a multiply imaged, highly magnified supernova, named *Refsdal*, lensed by MACS J1149.6 + 2223

cluster.¹⁸ The accuracy with which the reappearance of the time-delayed image of Refsdal has been predicted by some strong lensing models,¹⁹ involving positions, flux magnifications and time delays, underscores the high-level of precision that lensing models have reached when supported by abundant spectroscopic information.

As a result of our lensing models, accurate magnification maps (e.g. Refs. 20, 21) can be used to derived intrinsic (unlensed) properties of lensed galaxies as faint as $(M \sim -15)$, i.e. ~ 5 mag fainter than M^{*} at high redshifts²²(see Fig. 2). By fitting stellar population synthesis models to the spectral energy distribution of these galaxies and using their spectra, we can derive in such unexplored luminosity regime: stellar masses, star formation rates, gas/dust content, and physical parameters of their interstellar medium (e.g. Ref. 23). These new datasets will help us to make significant progress in our understanding of galaxy formation at early times.

Acknowledgments

C.D.C. is supported by the Erasmus Mundus Joint Doctorate Program by Grants Number 2014-0707 from the agency EACEA of the European Commission and G.B.C. is supported by the CAPES-ICRANET program through the grant BEX 13946/13-7.

References

- 1. F. Fontanot, et al. MNRAS 438, 2097 (2014).
- 2. P. Madau, F. Haardt ApJL 813, L8 (2014).
- 3. J. Wise, et al. MNRAS 442, 2560 (2014).
- 4. P. Rosati, et al. The Messenger 158, 48 (2014).
- 5. M. Postman, et al. ApJS 199, 25 (2012).
- 6. I. Balestra et al. arXiv:1511.02522, (2015).
- 7. C. Grillo, et al. ApJ 800, 38 (2015).
- 8. P. Oesch, et al. ApJL 804, L30 (2015).
- 9. A. Zitrin, et al. ApJL 810, L12 (2015).
- 10. S. Finkelstein, et al. Nature 502, 24 (2013).
- 11. M. Schenker, et al. ApJ 744, 179 (2012).
- 12. E. Vanzella, et al. ApJ 695, 1163 (2009).
- 13. O. Le Fèvre et al. A & A 559, A14 (2013).
- 14. D. Coe, et al. ApJ **762**, 32 (2013).
- 15. J-P. Kneib, P. Natarajan Astron Astrophys Rev 19, 47 (2011).
- 16. L. Bradley et al. ApJ **792**, 76 (2014).
- 17. A. Monna, et al. MNRAS 438, 1417 (2014).
- 18. T. Treu, et al. ApJ 817, 60 (2016).
- 19. C. Grillo, et al. arXiv:1511.04093, (2015).
- 20. G. Caminha, et al. arXiv:1512.04555, (2015).
- 21. A. Zitrin, et al. ApJ 801, 44 (2015).
- 22. R. Bouwens, et al. ApJ 803, 34 (2015).
- 23. E. Vanzella, et al. A & A 576, A116 (2015).