



Publication Year	2018
Acceptance in OA	2021-02-16T16:09:52Z
Title	LOFAR discovery of radio emission in MACS J0717.5+3745
Authors	Bonafede, A., Brügger, M., Rafferty, D., Zhuravleva, I., Riseley, C. J., van Weeren, R. J., Farnes, J. S., Vazza, F., Savini, F., Wilber, A., Botteon, A., Brunetti, G., CASSANO, Rossella, Ferrari, C., De Gasperin, F., Orrú, E., Pizzo, R. F., Röttgering, H. J. A., Shimwell, T. W.
Publisher's version (DOI)	10.1093/mnras/sty1121
Handle	http://hdl.handle.net/20.500.12386/30412
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
Volume	478

LOFAR discovery of radio emission in MACS J0717.5+3745

A. Bonafede,^{1,2,3*} M. Brüggen,² D. Rafferty,² I. Zhuravleva,^{4,5} C. J. Riseley,⁶
 R. J. van Weeren,⁷ J. S. Farnes,^{8,9} F. Vazza,^{1,2,3} F. Savini,² A. Wilber,² A. Botteon,^{1,3}
 G. Brunetti,¹ R. Cassano,¹ C. Ferrari,¹⁰ F. de Gasperin,^{2,7} E. Orrú,¹¹ R. F. Pizzo,¹¹
 H. J. A. Röttgering⁷ and T. W. Shimwell^{7,11}

¹INAF - Istituto di Radioastronomia, Bologna Via Gobetti 101, I-40129 Bologna, Italy

²Hamburger Sternwarte, Universität Hamburg, Gojenbergsweg 112, D-21029, Hamburg, Germany

³Dipartimento di Fisica e Astronomia, Università di Bologna, via P. Gobetti 93/2, I-40129, Bologna, Italy

⁴Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, 452 Lomita Mall, Stanford, California 94305-4085, USA

⁵Department of Physics, Stanford University, 382 Via Pueblo Mall, Stanford, California 94305-4060, USA

⁶CSIRO Astronomy & Space Science, 26 Dick Perry Avenue, Kensington, WA, 6151, Australia

⁷Leiden Observatory, Leiden University, PO Box 9513, NL-2300 RA Leiden, The Netherlands

⁸Oxford e-Research Centre (OeRC), Department of Engineering Science, University of Oxford, Oxford, OX1 3QG, UK

⁹Department of Astrophysics/IMAPP, Radboud University, PO Box 9010, NL-6500 GL Nijmegen, the Netherlands

¹⁰Observatoire de la Côte d'Azur, Laboratoire Lagrange UMR 7293. Bd. de l'Observatoire, CS 34229 F-06304 Nice, France

¹¹ASTRON, the Netherlands Institute for Radio Astronomy, Postbus 2, NL-7990 AA Dwingeloo, the Netherlands

Accepted 2018 April 27. Received 2018 April 23; in original form 2018 February 26

ABSTRACT

We present results from Low-Frequency Array (LOFAR) and Giant Meterwave Radio Telescope (GMRT) observations of the galaxy cluster MACSJ0717.5+3745. The cluster is undergoing a violent merger involving at least four sub-clusters, and it is known to host a radio halo. LOFAR observations reveal new sources of radio emission in the Intra-Cluster Medium: (i) a radio bridge that connects the cluster to a head-tail radio galaxy located along a filament of galaxies falling into the main cluster, (ii) a 1.9 Mpc radio arc that is located north-west of the main mass component, (iii) radio emission along the X-ray bar that traces the gas in the X-rays south-west of the cluster centre. We use deep GMRT observations at 608 MHz to constrain the spectral indices of these new radio sources, and of the emission that was already studied in the literature at higher frequency. We find that the spectrum of the radio halo and of the relic at LOFAR frequency follows the same power law as observed at higher frequencies. The radio bridge, the radio arc, and the radio bar all have steep spectra that can be used to constrain the particle acceleration mechanisms. We argue that the radio bridge could be caused by the re-acceleration of electrons by shock waves that are injected along the filament during the cluster mass assembly. Despite the sensitivity reached by our observations, the emission from the radio halo does not trace the emission of the gas revealed by X-ray observations. We argue that this could be due to the difference in the ratio of kinetic over thermal energy of the intra-cluster gas, suggested by X-ray observations.

Key words: galaxies: clusters: individual: MACSJ0717+3745 – radiation mechanism: non thermal – acceleration of particles – techniques: interferometri.

1 INTRODUCTION

The intra-cluster medium (ICM) of galaxy clusters is filled with a weakly magnetised plasma that can contain relativistic electrons emitting synchrotron radiation over Mpc scale. The emission is

classified as radio haloes and radio relics depending on its morphological properties: radio haloes are found at cluster centres and are mostly connected with major mergers (Buote 2001; Cuciti et al. 2015, but see also Bonafede et al. 2014 and Sommer et al. 2017 for some outliers), while radio relics usually have an arc-like morphology and are found at cluster peripheries (van Weeren et al. 2010).

* E-mail: abonafede@ira.inaf.it

Table 1. MACSJ0717.5+3745.

Name	MACSJ0717.5+3745	ref.
RA [J2000]	07h17m30.9s	
DEC[J2000]	37d45′30″	
z	0.546	E01
M_{500}	$(11.5 \pm 0.5) \times 10^{14} M_{\odot}$	PUC
M_{vir}	$(3.5 \pm 0.6 \times 10^{15} M_{\odot})$	U14
$L_X[0.1 - 2.4\text{keV}]$	$(2.74 \pm 0.03) \times 10^{45} \text{ erg/s}$	E07

E01: Ebeling, Edge & Henry (2001), PUC: Planck Collaboration et al. (2016), U14: Umetsu et al. (2014), E07: Ebeling et al. (2007).

Mergers between galaxy clusters can dissipate up to 10^{64} ergs of energy in the ICM, and the current theoretical models predict that a fraction of this energy may be channeled into the (re)acceleration of cosmic ray electrons (CRe) and magnetic field amplification. Turbulence injected by mergers could re-accelerate a population of low-energy CRe and produce radio haloes, while shock waves that propagate in the ICM during mergers could amplify the magnetic field and (re)accelerate CRe producing radio relics. We refer the reader to the reviews by Brügggen et al. (2011), Feretti et al. (2012) and Brunetti & Jones (2014) for more details.

MACSJ0717.5+3745 (hereinafter MACSJ0717) is a very complex system. It is located at redshift $z = 0.546$, and since its discovery, it has been the subject of several observational campaigns (Edge et al. 2003; Ma et al. 2008; Bonafede et al. 2009; van Weeren et al. 2009; Medezinski et al. 2013; Sayers et al. 2013; Umetsu et al. 2014; Limousin et al. 2016; van Weeren et al. 2016; Adam et al. 2017). The main properties of the cluster are listed in Table 1.

X-ray and optical observations show a complex merger involving at least four sub-clusters (Limousin et al. 2016). The ICM temperature shows strong gradients, with the eastern part significantly hotter than the western part. The hottest region in the south-east reaches ~ 20 keV, while recent *Chandra* data suggest the presence of a cold front in the N-NE region (van Weeren et al. 2017). The X-ray emission reveals a V-shaped structure, associated with the main mass component. To the north-west of this component, a bullet-like structure is associated with a second sub-cluster. On the south-east, a bar-shaped structure coincides with two more sub-clusters (see Figs 1 and 2).

In the South-East of the cluster, a 19 Mpc long filament of galaxies was found by Ebeling, Barrett & Donovan (2004) and confirmed by Jauzac et al. (2012). Recently, deep *Chandra* observations have detected the part of the filament that is close to the cluster, and found a galaxy group of $\sim 10^{13} M_{\odot}$ embedded in the filament (Ogrea et al. 2018, submitted). A head-tail radio galaxy (hereinafter HT radio galaxy) at $z = 0.5399$ is found along the filament, between the cluster and the X-ray detected group (Ebeling, Ma & Barrett 2014). In Fig. 1, the X-ray emission is shown and the different components of the system are labelled.

We refer the reader to Ebeling et al. (2014), Limousin et al. (2016), Medezinski et al. (2013), and van Weeren et al. (2017) for a detailed analysis of the cluster X-ray emission and dynamical state.

The cluster hosts a powerful radio halo (Bonafede et al. 2009; van Weeren et al. 2009), that is asymmetric and whose largest linear size is more than 1.4 Mpc. Within the halo, the cluster also hosts a bright-polarized filament or radio relic, aptly named the ‘chair-shaped’ filament by Pandey-Pommier et al. (2013). For simplicity, we will refer to this structure as a relic. The relic is polarized at the ~ 17 per cent level at 4.9 GHz, while the halo also shows polarization at the 2–7 per cent at 1.4 GHz (Bonafede et al. 2009). Such polarized

emission – which is common for relics – is unusual for haloes, and may be related to the peculiar dynamical state of the system.

The emission from the radio halo roughly follows the bar and V-shaped structures detected in the X-rays, while no emission was found in the western part of the cluster (Bonafede et al. 2009; van Weeren et al. 2009). Deep Jansky Very Large Array (VLA) observations have recently found several radio filaments on scales 100–300 kpc, departing from the halo towards the NE and the NW, and at least a few of these are located in the cluster outskirts (van Weeren et al. 2017).

Because of its high radio power, this is one of the few radio haloes that can be imaged by the existing interferometers at frequencies higher than 1.4 GHz. Hence, it is a primary target to study the spectral properties of the radio emission over a large frequency range. van Weeren et al. (2017) found an average spectral index¹ of ~ -1.3 to -1.4 by fitting a straight power law through flux measurements at 1.5, 3.0, and 5.5 GHz. This is in agreement with previous results by Bonafede et al. (2009), and Pandey-Pommier et al. (2013), obtained with shallower and lower frequencies observations, respectively.

In this paper, we present new low-frequency observations of the cluster obtained with the Low-Frequency Array (LOFAR; van Haarlem et al. 2013). Our aim is to constrain the spectral properties of the diffuse emission to gain insights on the (re)acceleration processes in this complex system, and to search for additional emission in the western part of the cluster where, despite the presence of hot gas and dynamical activity, no radio emission has been detected.

The rest of the paper is as follows: in Section 2, we describe the radio observations and the main steps of the data reduction. In Section 3, we analyse the results of the LOFAR and Giant Metrewave Radio Telescope (GMRT) observations, and discuss the spectral properties of the system. A combined radio and X-ray analysis is reported in Section 4, and we conclude in Section 5. Throughout the paper we use a Λ CDM cosmological model with $H_0 = 69.6 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.286$, and $\Omega_{\Lambda} = 0.714$. At the cluster redshift the angular to linear scale is 6.459 kpc/arcsec.

2 OBSERVATIONS AND DATA REDUCTION

2.1 LOFAR

The cluster was observed on 2013 March 19 for a total of 5 h, using the LOFAR High Band Antenna (HBA) stations in the HBA_DUAL_INNER mode. A total of 61 antennas were present (13 Remote stations, and 24 Core stations, each split into two). Observations covered the frequency range 111–182 MHz using 366 sub-bands. 3C286 was observed in the frequency range 115–176 MHz using 310 sub-bands and was used as calibrator. Both the cluster and the calibrator data were taken with a sampling time of 2 s. Each sub-band of 0.195 MHz bandwidth was recorded with 64 channels. Data were initially flagged by the observatory using AOFLAGGER (Offringa, van de Gronde & Roerdink 2012) and then averaged down to 4 channels per sub-band and 5 and 4 s sampling for the cluster and calibrator, respectively.

Data were calibrated using the facet calibration approach (van Weeren et al. 2016). We refer the reader to van Weeren et al. (2016) for a detailed description, and outline here only the main steps.

Calibrator data were further averaged to 10 s and two channels per sub-band. The stations CS013HBA and CS032HBA were flagged and the data were calibrated against a source model, following the

¹Throughout this paper we define the spectral index α as $S(\nu) \propto \nu^{\alpha}$.




Figure 1. Colours: X-ray emission from *Chandra* in the band 0.2–5 keV. Contours: radio emission from LOFAR at 147 MHz. The beam is $19' \times 18'$. The rms noise (σ) is $0.16 \text{ mJy beam}^{-1}$. Contours start at 4σ and are spaced by a factor 2. The contour at -4σ is dashed. The main components of the cluster emission in the X-rays and radio are labelled in yellow and white, respectively.




Figure 2. Subaru *b*, *v*, and *z* composite image overlaid on to X-ray emission from *Chandra* (cyan) in the band 0.2–5 keV, and radio emission from LOFAR at 147 MHz (orange). The image has illustrative purposes only. A black circle is superposed on a bright star to mask it. Quantitative values for the X-ray and radio emission are shown in Figs 1 and 3, respectively.

Scaife & Heald (2012) flux density scale. XX and YY gains were determined, together with the rotation angle to account for possible differential Faraday rotation, that was found to be negligible. Clock offsets for each station were derived from these solutions, fitting

for Clock delays and differential TEC (Total Electron Content). Amplitude gains, clock offsets, and instrumental XX-YY phase-offsets were applied to the target data to set the initial flux scale, and to correct for instrumental effects, respectively.

Then, the target data were calibrated in phase against a Global Sky Model, that is derived from several radio surveys [specifically, the VLA Low-frequency Sky Survey, VLSS (Lane et al. 2014), the Westerbork Northern Sky Survey, WENSS (Rengelink et al. 1997), and the Northern VLA Sky Survey, NVSS (Condon et al. 1998)].

The first set of images at intermediate resolution ($\sim 30''$) were created using `WSCLEAN` (Offringa et al. 2014; Offringa & Smirnov 2017), grouping the data in chunks of ~ 2 MHz each. Images were corrected for the station beam at the phase centre. The model components were subtracted from the UV data, and new images at lower resolution ($\sim 2'$) were created, including sources up to 20 degrees from the target centre. The model component list was updated with the clean components found in the low-resolution images. After this step, we are left with a list of model components for the field, and an almost empty UV data set, as required by the *Factor* pipeline,² that performs the facet calibration.

The LOFAR HBA field of view has been divided into 50 facets. Each facet is set to contain a source brighter than 0.1 Jy and smaller than 2 arcmin, that is used as facet calibrator. UV data from baselines shorter than 80λ have not been used during self-calibration and deconvolution. Our aim is to derive the direction-dependent gains in the direction of the target. To do this, we first need to minimize the artefacts from bright sources around the target. The contamination due to artefacts from nearby sources is modest, and indeed the cluster itself is the second brightest source in the field after B3 0704+384. Hence, we first derived direction-dependent gains in the direction of B3 0704+384. These gains have been applied to the facet and a new model has been derived and subtracted from the visibilities. Direction-dependent gains were then derived for the target facet using the cluster as calibrator. We also checked that deriving direction-dependent gains for the facets around the target before processing the target facet did not lead to a better calibration for the target, because these facets are affected by residual artefacts and calibration errors from the cluster facet. Direction-dependent gains are derived through several self-calibration cycles, using a multi-resolution algorithm. In this procedure, we initially image the data at 20 arcsec resolution and progressively increase the resolution to ~ 5 arcsec. Facets are processed as follows: data are phase-shifted towards the centre of the facet calibrator and further averaged in frequency to speed up the calibration process. The model components of the facet calibrator are added back to the visibilities and several cycles of Stokes I phase, and TEC self-calibration are performed on a 10 s time-scale. Finally, some rounds of complex gain self-calibration are performed.

After calibration, data were imaged using the Briggs weighting scheme (Briggs 1995) setting the parameter *robust* = -0.25 to suppress the sidelobes. Gaussian tapers were applied to decrease the weight of long baselines and better image the extended emission. Images were taken with `WSClean` using the multi-scale and multi-frequency deconvolution mode implemented in the code. Imaging parameters are listed in Table 2, and the images are shown in Fig. 3.

Flux densities have been checked against the TIFR GMRT Sky Survey (TGSS) (Intema et al. 2017), and we found that the LOFAR flux densities are consistent within the calibration errors. For consistency with previous works, we adopt a conservative 15 per cent flux density uncertainty (Shimwell et al. 2016; van Weeren et al. 2017; Wilber et al. 2018; Savini et al. 2018).

2.2 GMRT observations

A 12 h long observation at 608 MHz was performed with the GMRT on 2011 June 5.

Data have been recorded in a spectral mode, using 256 channels having a width of 130 kHz each, for a total bandwidth of 33 MHz. The integration time was set to 8 s. We processed the observation using the Source Peeling and Atmospheric Modelling (SPAM) tool (Intema et al. 2009) to take into account direction-dependent effects. The main steps are outlined below, and we refer the reader to Intema et al. (2009) for further details.

The sources 3C147 and 3C286 were observed for 25 min at the beginning and at the end of the observing block, respectively, and used to correct for the bandpass and to set the absolute flux scale, following the Scaife & Heald (2012) flux scale. Strong radio-frequency interference (RFI) were removed from the data using statistical outlier flagging tools, and much of the remaining low-level RFI was modelled and subtracted from the data using `OBIT` (Cotton 2008). After RFI removal, data were averaged down to 24 channels, to speed up the following steps and, at the same time to avoid significant bandwidth smearing during imaging. To correct the phase gains of the target field, we started from a global sky model (see Section 2.1). SPAM permits to correct for ionospheric effects, and remove direction-dependent gain errors, reaching thermal-noise limited images. Within SPAM, imaging is done with `AIPS` using the wide-field imaging technique to compensate for the non-complanarity of the array. The presence of strong sources in the field of view enables one to derive directional-dependent gains for each of them (similar to the peeling technique) and to use these gains to fit a phase-screen over the entire field of view. After ionospheric corrections, sources outside the inner 8 arcmin were subtracted to facilitate the imaging steps.

Data have been imaged with `CASA` (McMullin et al. 2007) using different weighting schemes and Gaussian UV-tapers to achieve different resolutions. The final images have been corrected for the GMRT primary beam response, and are shown in Fig. 4. Imaging parameters are listed in Table 2. We assume a 10 per cent error on the absolute flux scale.

3 RESULTS

Both LOFAR and GMRT observations detect new emission that was not detected by previous, shallower radio observations. Because of the different sensitivities of the two instruments towards large-scale emission, we first analyse the observations separately, and then perform a spectral index study.

3.1 Radio emission at 147 MHz

The main result of the LOFAR observations is the discovery of additional emission west of the halo (radio arc) around the X-ray bar, and south-east of the halo in the direction of the accreting sub-group along the intergalactic filament (bridge). The halo emission is more extended than previously found by VLA and GMRT observations (Bonafede et al. 2009; Pandey-Pommier et al. 2013; van Weeren et al. 2017) and it extends beyond the relic in the east and south-east directions (Fig. 1). In Fig. 1, the radio emission at 147 MHz is shown in contours, and the new features are labelled.

A foreground Fanaroff Riley type I (FRI) radio galaxy ($z = 0.1546$) has been identified to the south-east of the cluster. Its lobes are prominent in the VLA and GMRT observations (see Fig. 4), but are almost undetected in the LOFAR image because

²<https://github.com/lofar-astron/factor>.

Table 2. Images.

Image	Weighting scheme	UV-Taper	Restoring beam	rms noise mJy beam ⁻¹	Fig. of merit
LOFAR HBA HR	Briggs, robust = -1	–	5.8'' × 4.6''	0.13	3
LOFAR HBA LR	Briggs, robust = -0.25	10''	19'' × 18''	0.16	3
GMRT 608 HR	Briggs, robust = 0.25	–	6.3'' × 5.6''	0.03	4
GMRT 608 LR	Briggs, robust = -0.25	20''	18.3'' × 17.4''	0.10	4
For spectral index image – UV-range > 500 λ					
LOFAR HBA	Uniform	10	10'' × 10''	0.25	5, top panel
GMRT 608	Uniform	10	10'' × 10''	0.07	5, top panel
LOFAR HBA	Uniform	30	30'' × 30''	0.50	5, bottom panel
GMRT 608	Uniform	30	30'' × 30''	0.17	5, bottom panel




Figure 3. Left-hand panel: LOFAR image at 147 MHz at high resolution (5.8'' × 4.6''). The rms noise is 0.13 mJy beam⁻¹. The blue circles and ellipses indicate the sources embedded in the diffuse emission that were masked. Right-hand panel: LOFAR image at 147 MHz at low resolution (19'' × 18''). The noise is 0.16 mJy beam⁻¹. Blue polygons mark the regions used to separate the different components of the radio emission (bridge, bar, arc). In both panels contours start at 4σ and are spaced by a factor 2. Contour at -4σ are dashed. The restoring beams are shown in the bottom-left corner of the two panels.




Figure 4. Left-hand panel: GMRT image at 608 MHz at high resolution (6.3'' × 5.6''). The rms noise is 30 μJy beam⁻¹. Right-hand panel: GMRT image at 608 MHz at low resolution (18.3'' × 17.4''). The noise is 0.1 mJy beam⁻¹. The blue circles and ellipses indicate the sources embedded in the diffuse emission that have been masked. In both panels, contours start at 4σ and are spaced by a factor 2. Contour at -4σ are dashed. The restoring beams are shown in the bottom-left corner of the two panels.

of the combined effect of their spectral index ($\alpha_{1\text{GHz}}^{6.5\text{GHz}} \sim 0.6$, van Weeren et al. 2017) and low-surface brightness (~ 0.2 mJy beam $^{-1}$ at 5 GHz, van Weeren et al. 2017).

The radio arc has a total flux density of 49 ± 7 mJy. Assuming $\alpha = -1.3$ for the k -correction, this flux density corresponds to a power $P_{147\text{MHz}} = (7 \pm 1) \times 10^{25}$ W Hz $^{-1}$. The largest angular size of the emission is ~ 4.9 arcmin, corresponding to ~ 1.9 Mpc. The arc is located at a projected distance of ~ 1.2 arcmin (460 kpc) from the main mass component, between the W sub-cluster and the V-shaped emission visible in the X-rays. The radio emission from the radio arc does not follow the X-ray emission from the gas in the same region of the cluster. The structure could be seen in projection on to the cluster centre, and could be associated with a merger or accretion shock. In this case, it could be classified as a relic. However, given the complex structure of the whole radio emission and having no information about projection effects, any conclusion would be speculative.

South-east of the X-ray bar, hints of a new radio bridge are found, connecting the radio halo to the HT radio galaxy located along the optical filament ~ 3 arcmin from the X-ray centre.³ Bridges connecting radio haloes to HT radio galaxies have been found in few other clusters already (the Coma cluster Giovannini et al. 1993) suggesting that the fossil electrons from the tail are (re)accelerated by phenomena connected with the merger. However, this case is somewhat different. In fact, the lobes of the HT radio galaxy are pointing in the opposite direction with respect to the bridge. The properties of the radio bridge – in connection with the gas properties of the filament – are further analysed in Section 4.1.

van Weeren et al. (2017) have found that the radio emission south-east of the main mass concentration roughly follows the X-ray bar. LOFAR observations reveal further emission covering the entire bar and extending beyond it.

3.2 Radio emission at 608 MHz

The GMRT observation allows us to reach a sensitivity that is a factor ~ 2 deeper with respect to the data published so far at this frequency (van Weeren et al. 2009). In Fig. 4, the emission at 608 MHz is shown at two different resolutions, obtained with the imaging parameters listed in Table 2. The GMRT image at low resolution shows diffuse emission that was previously undetected at this frequency: the halo appears more extended in the NW direction, and towards S. No emission is detected corresponding to the radio arc and to the bridge visible in the LOFAR images. However, we note that the halo extension towards NW, which is also partially detected in the VLA image at 1.4 GHz (Bonafede et al. 2009), is not detected by LOFAR.

In Table 3, we list the flux densities and sizes of the radio components in the cluster both at LOFAR and GMRT frequencies. These are derived from the low-resolution images (see Table 2 for details), above the 4σ contour, and masking the discrete sources embedded in the diffuse emission. Since the boundaries of the relic and of the bar cannot be easily separated from the halo component, we also list

³As no evidence for radio emission from the bridge was found in the VLA or in the GMRT observations (Bonafede et al. 2009; van Weeren et al. 2017; Pandey-Pommier et al. 2013), and as the background noise of the images are not uniform, even a detection at $\sim 10\sigma$ could be partially affected by calibration artefacts. Hence, deeper observations with other instruments would be required to confirm with higher confidence the properties of the radio bridge and of the radio arc.

the properties of the total radio emission, which includes the halo, the relic, and the radio bar. In Figs 3 and 4, we show the regions used to compute the flux densities listed in Table 3.

3.3 Spectral properties of the radio emission

Using LOFAR and GMRT observations, we have produced spectral index maps of the cluster radio emission. LOFAR and GMRT observations have been imaged using the same UV-range, uniform weighting scheme, and a Gaussian taper as listed in Table 2. The minimum baseline has been chosen to have a dense sampling of the GMRT data, and it is particularly critical here, given the different frequency and baseline lengths of the two interferometers.

We have used two tapering functions with different FWHM of 10 and 30 arcsec in order to analyse the spectral index variations, and to constrain the spectral index of the diffuse emission, respectively. To compute the spectral index image, LOFAR and GMRT images have been convolved with a Gaussian beam to achieve the exact same resolution. Both images have been blanked at 2σ and a spectral index map has been computed. The spectral index images are shown in Fig. 5. The errors on the spectral index have been computed according to

$$\alpha_{err} = \frac{1}{\ln(\nu_1/\nu_2)} \sqrt{\left(\frac{\Delta S_1}{S_1}\right)^2 + \left(\frac{\Delta S_2}{S_2}\right)^2}, \quad (1)$$

where ΔS_i takes into account both the flux density errors ($\delta S_i \times S_i$) and the image noises (σ_i).

The high-resolution spectral index image shows that the cluster diffuse emission is steep, in agreement with previous works by Bonafede et al. (2009) and van Weeren et al. (2009, 2017). We detect a steep spectrum region between the relic and the foreground FRI radio galaxy with a mean value $\langle \alpha_{608\text{MHz}}^{147\text{MHz}} \rangle = -2.2 \pm 0.2$. This region could be contaminated by the past emission of a steep-spectrum source detected at higher frequency by van Weeren et al. (2017). The Narrow Angle Tail radio galaxy at the cluster centre shows a steepening of the spectral index along the tail from $\alpha_{608\text{MHz}}^{147\text{MHz}} \sim -0.7$ in the core down to $\alpha_{608\text{MHz}}^{147\text{MHz}} \sim -2.3$, consistent with the behaviour at higher frequencies reported by van Weeren et al. (2017). Further out along the relic the spectral index becomes flatter ($\alpha_{608\text{MHz}}^{147\text{MHz}} \sim -1.3$ to -1), possibly indicating that the aged electrons from the radio galaxy have been re-energised by a shock (van Weeren et al. 2017).

To compute an average spectral index of the radio halo and of the radio relic, we have blanked the sources embedded in the diffuse emission. The average spectral index of the halo is $\langle \alpha_{608\text{MHz}}^{147\text{MHz}} \rangle = -1.4 \pm 0.1$. We note that this value is the emission detected both at 608 and 147 MHz and it is not representative of the whole emission detected by LOFAR or GMRT. The average spectral index of the relic is $\langle \alpha_{608\text{MHz}}^{147\text{MHz}} \rangle = -1.1 \pm 0.1$. These values are consistent with those reported in the literature by Bonafede et al. (2009) and van Weeren et al. (2009, 2017) within the errors, and indicate that the spectrum does not change significantly at LOFAR frequencies.

The new features detected by LOFAR (i.e. the radio arc, the radio bridge, and the radio bar) are not visible in the GMRT image. This is likely due to a combination of their steep spectrum, weak surface brightness, and large angular extent that is filtered out by the GMRT. Imaging the data using the same UV-range and restoring beam allows us to investigate this. In Fig. 5, the spectral index at low resolution is shown, together with the LOFAR contours. From this image, we can conclude that: (i) the spectrum of the emission E of the relic is steep ($\langle \alpha_{608\text{MHz}}^{147\text{MHz}} \rangle = -1.7 \pm 0.2$). (ii) Most of the radio arc is not detected in the GMRT image, because of both

Table 3. MACSJ0717.5+3745: details on radio components.

Radio component	$S_{147\text{MHz}}$ Jy	$S_{608\text{MHz}}$ Jy	Size at 147 MHz* arcsec – kpc	Size at 608 MHz* arcsec – kpc	$\alpha_{608\text{MHz}}^{147\text{MHz}}$
Total	0.9 ± 0.1	0.21 ± 0.03			
Filament/Relic	0.50 ± 0.08	0.18 ± 0.03	125–800	125–807	-1.1 ± 0.1
Halo	0.37 ± 0.06	0.034 ± 0.005	160–1030	160–1030	-1.4 ± 0.1
Bridge	$(1.3 \pm 0.5) \times 10^{-2}$	–	65–400	–	< -1.4
Bar	0.020 ± 0.003	$(1.7 \pm 0.2) \times 10^{-3}$	70–450	10–65	< -1.6
Arc	0.049 ± 0.007	–	300–1900	–	< -1.3

Col1: Name of the radio source; Col 2 and 3 : Flux density from the LOFAR LR and GMRT LR image; Col 4 and Col 5: Maximum angular and linear size of the source in LOFAR and GMRT. *For the Halo, the parameter D_H , defined as $\sqrt{D_{max} \times D_{min}}$ is given, where D_{max} and D_{min} refer to the maximum and minimum scale, respectively. All quantities are projected. Col. 6: spectral index between 608 MHz and 147 MHz. Note that the spectral index refers to the regions as specified in the text, and not to the entire size of the radio component listed in Col 3 and 4.




Figure 5. Top: LOFAR–GMRT spectral index image (left) and associated errors (right) at the resolution of 10 arcsec. Bottom: LOFAR–GMRT spectral index image (left) and associated errors (right) at the resolution of 30 arcsec. Contours display the LOFAR stokes I image, starting at 4σ and increasing by a factor 2 each. The -4σ contours are plotted with dotted lines.

its low-surface brightness and large-scale size. Indeed, only the brightest patch of emission is seen by LOFAR once the baselines shorter than 500λ are excluded from imaging. In this region, we can put a limit $\langle \alpha_{608 \text{ MHz}}^{147 \text{ MHz}} \rangle < -1.3$. (iii) The radio bridge and the radio bar are not detected in the GMRT image because of their steep spectrum. We can put a limit on the spectral index in these regions $\langle \alpha_{608 \text{ MHz}}^{147 \text{ MHz}} \rangle < -1.6$ and $\langle \alpha_{608 \text{ MHz}}^{147 \text{ MHz}} \rangle < -1.4$ in the radio bar and radio bridge, respectively. All the limits to the spectra computed above consider the mean LOFAR surface brightness and 2σ noise of the GMRT image.

The emission north-west of the radio halo, detected by GMRT observations, is not visible in the LOFAR image. As the LOFAR image is dominated by the bright halo and radio arc, we cannot exclude that the emission is not visible because of deconvolution artefacts. Alternatively, this emission would need a spectral index $\alpha \geq -1.3$ to fall below the LOFAR sensitivity.

4 RADIO AND X-RAY EMISSION

Using deep *Chandra* observations of the cluster (Ogrea et al. 2018, submitted; van Weeren et al. 2017), we can derive constraints on the particle acceleration mechanisms that produce the radio emission in the cluster centre and outskirts. In this section, we perform a joint radio and X-ray analysis of the radio bridge and the radio halo.

4.1 The radio bridge

The detection of radio emission along the filament connecting the HT radio galaxy with the main cluster allows us to constrain the non-thermal properties at the outskirts of clusters. The radio emission is detected in a region between the main cluster and a subgroup that has a temperature of ~ 3 keV and an X-ray luminosity of $\sim 10^{43}$ ergs s^{-1} in the band 0.1–2.4 keV. The group is located at 2 Mpc south-east from the main cluster and it is likely at its first infall towards the cluster (Ogrea et al. 2018, submitted). The portion of the filament between the group and the cluster is overdense by a factor 100–150 with respect to the critical density of the Universe at the redshift of the cluster. This part of the filament has a temperature of $1.6_{-0.3}^{+0.5}$ keV and a density of $\sim 10^{-4} \text{ cm}^{-3}$ (Ogrea et al. 2018, submitted).

Being within $r_{100} - r_{150}$, the radio emission in the bridge is probing a region that is gravitationally bound to the main cluster, where the magnetic field has likely been compressed and amplified, erasing all signatures from a primordial seed (Dolag, Bartelmann & Lesch 2002; Miniati & Beresnyak 2015, and ref. therein). The detection of radio emission indicates that relativistic electrons are present in this region. The central galaxy of the group is radio loud, but being at its first infall on to the main cluster, it is unlikely that it contributes to the radio emission in the bridge.

During the accretion of matter on to the main cluster, shock waves are injected in the ICM, that can heat the gas, and accelerate particles through Fermi-I type mechanisms, like Diffusive Shock Acceleration (DSA; Drury 1983).

Using the constraints on the gas temperature and density derived from X-ray observations (Ogrea et al. 2018, submitted), we investigate here whether and under which conditions the radio emission in the bridge can be produced by shock (re)acceleration through DSA.

We start by investigating a simple ‘single zone’ model for the shock, in which we characterize the entire bridge region with a single value for the gas density, gas temperature, and magnetic field. We

numerically solve the time-dependent evolution of the energy distribution of relativistic electrons under the following conditions: (i) shocks accelerate electrons through DSA, and the resulting energy distribution of the particles is a power law in energy ($\frac{dN}{dE} \propto E^{-\delta}$), with δ that depends on the injection Mach number, \mathcal{M}_{inj} , according to $\delta = 2(\mathcal{M}^2 + 1)/(\mathcal{M}^2 - 1)$, e.g. Sarazin (1999). (ii) Particles undergo energetic losses due to synchrotron and Inverse Compton, as well as collisional losses. (iii) Electrons might be re-accelerated by a second shock, shortly before the epoch of our radio observation. In the linear acceleration regime, the particle post-shock spectrum after re-acceleration will be (Markevitch et al. 2005; Kang, Ryu & Jones 2012)

$$\frac{dN}{d\gamma} = (\delta + 2)\gamma^{-\delta} \int_{\gamma_{\text{min}}}^{\gamma} \frac{dN_a}{d\gamma} \gamma^{\delta-1} d\gamma, \quad (2)$$

where γ is the Lorentz gamma factor of electrons, γ_{min} is the minimum γ factor of the particle injected by the first shock after their ageing, and $\frac{dN_a}{d\gamma}$ is the spectrum of the aged electrons. The scenarios where particles are re-accelerated by a second shock are labelled with *+re* in Table 4 and Fig. 6.

The time-dependent diffusion-loss equation of cosmic ray electrons (Kardashev 1962; Sarazin 1999) is solved with the Chang & Cooper (1970) finite difference scheme, using 5×10^4 energy bins of $\Delta\gamma = 10$ in the $2 \leq \gamma \leq 5 \cdot 10^4$ energy range, and a fixed timestep of 10 Myr.

As the size of the filament in the X-rays is bigger than the size of the radio bridge, we cannot disentangle whether the average values of density and temperature in the filament are pre- or post-shock values. Hence, we have tested both scenarios (labelled with *pre* – and *post* –, respectively, in Table 4 and Fig. 6). We have applied jump conditions as a function of the assumed Mach number (\mathcal{M}_{inj}) to recover the pre-shock values of n_e and T . In the re-acceleration scenarios, the Mach number of the second shock (\mathcal{M}_{re}) is used to compute the pre-shock density and temperature values. The kinetic energy flux, Φ , through the shock surface is proportional to $\Phi \propto \rho_{\text{pre}} \mathcal{M}^3 c_{\text{s,pre}}^3$, where ρ_{pre} is the pre-shock gas mass density and $c_{\text{s,pre}}$ is the pre-shock sound speed (Vazza et al. 2015).


We have investigated different combinations of magnetic fields, Mach numbers, and times of injection and re-acceleration needed to reproduce the flux density of the radio bridge at 147 MHz and the radio spectral index ($\alpha_{608 \text{ MHz}}^{147 \text{ MHz}} \leq -1.4$). In Table 4, we list the main parameters of our model: the post-shock gas density and temperature values (n_{post} and T_{post}), the Mach number of the first shock (\mathcal{M}_{inj}) at the epoch t_{inj} , and the efficiency of the first shock acceleration $\xi_{e, \text{inj}}$. The resulting energy in relativistic electrons is $E_{e, \text{inj}}$.

For the re-acceleration models, we also list the Mach number of the re-accelerating shock (\mathcal{M}_{re}) active at the epoch t_{re} . We list in the table also the magnetic field of the radio bridge, B , the flux density at 147 MHz, $S_{147 \text{ MHz}}$, the radio spectral index $\alpha_{608 \text{ MHz}}^{147 \text{ MHz}}$, and the predicted flux density at 50 MHz, $S_{50 \text{ MHz}}$. The radio emission is obtained by numerically integrating the synchrotron emission from the final distribution of accelerated particles (Ginzburg & Syrovatskii 1965). Fig. 6 shows the expected flux densities for the above models as a function of the observing frequency.

Although all models are tailored to reproduce the observed flux density and spectral index limit, some of them can be ruled out: single injection scenarios (i.e. pre and post models), as well as *pre + re* scenario require either a large injection efficiency, which is troublesome for DSA (Vazza et al. 2015), and/or high values of temperature and density that are not compatible with the observational constraints.

Table 4. Model parameter for our simulation of the radio bridge (see Section 4.1 for more details).

run ID	n_{post} cm^{-3}	T_{post} K	\mathcal{M}_{inj}	$\log_{10}(E_{\text{e, inj}})$ erg	$\xi_{\text{e, inj}}$	t_{inj} Gyr	\mathcal{M}_{re}	t_{re} Gyr	B μG	$S_{147\text{MHz}}$ mJy	$\alpha_{608\text{MHz}}^{147\text{MHz}}$	$S_{50\text{MHz}}$ mJy
pre	$1.2 \cdot 10^{-3}$	$1.6 \cdot 10^8$	5.0	53.45	10^{-3}	-0.06	-	-	3.0	13.0	-1.43	49.4
pre+re	$9 \cdot 10^{-4}$	$5.3 \cdot 10^7$	15.0	53.90	10^{-4}	-0.9	2.5	-0.02	1.0	13.0	-1.51	60.0
post	$3 \cdot 10^{-4}$	$1.9 \cdot 10^7$	5.0	53.35	$3 \cdot 10^{-2}$	-0.06	-	-	3.0	13.0	-1.42	47.0
post+re	$3 \cdot 10^{-4}$	$1.9 \cdot 10^7$	15.0	52.54	$1.6 \cdot 10^{-3}$	-0.9	3.0	-0.04	3.0	13.0	-1.42	46.0
post+re2	$3 \cdot 10^{-4}$	$1.9 \cdot 10^7$	25.0	53.59	$1.6 \cdot 10^{-2}$	-1.0	2.2	-0.05	3.0	13.0	-1.40	48.7


Figure 6. Top panel: predicted radio spectra for our models of the radio bridge, as in Table 4. Bottom panel: Same as top panel, but with radio emission normalized to the model pre to highlight the differences among the models.

The *post + re* scenarios is the only one, among those investigated here, that could reproduce the radio bridge flux density with reasonable values of the model parameters. In this model, the second shock has a Mach number $\mathcal{M}_{\text{re}} = 3$, that would produce radio emission on a scale L_{rad} which is much smaller than the projected size of the bridge (see equation 15 in Kang et al. 2012). Hence, one should assume that the shock propagates with a very small angle with respect to the line of sight, perpendicular to the filament main axis. Such ‘transversal’ shocks are observed in cosmological simulations in filaments that connect interacting clusters (Vazza et al. 2015). In this case, the observed spectrum will be the superposition of different populations of electrons (re)accelerated at slightly different times as the shock propagates through the radio bridge (Sarazin 1999).

We have resimulated this scenario by considering the emission from the superposition of the different populations of electrons (scenario *post + re2* in Table 4). While most parameters are unchanged

(see the last row of Table 4), one needs to assume a larger energy of reaccelerated electrons to compensate for the cooling losses of the layers that have been accelerated first. The predicted spectrum in this case becomes flatter at higher frequencies (Fig. 6), leaving to future observations the possibility to better investigate this scenario.

4.2 Radio halo and X-ray emission

Theoretical models for the formation of radio haloes would expect that radio emission approximately follows the X-ray emission from the gas (Brunetti & Jones 2014, and ref. therein). This is observed in some clusters (Govoni et al. 2001) while it is not true in other cases (Abell 1132, Wilber et al. 2018). The radio halo in MACSJ0717 is probably the most striking case where radio emission is offset from the X-ray emission. In the turbulent re-acceleration scenario, this would require a different energy in turbulence and magnetic field in regions with and without radio emission, and/or the presence of a seed population of electrons only in the former region.

While the latter hypothesis is hard to verify, the amount of energy in turbulence in different cluster regions can be estimated through the amplitude of gas density fluctuations measured from X-ray observations. In stratified cluster atmospheres, the amplitude of gas density fluctuations, $\frac{\delta\rho_k}{\rho}$, and one-component velocity, V_{1k} , are proportional to each other at each wavenumber k within the inertial range of scales, namely:

$$\frac{\delta\rho_k}{\rho} = \eta \frac{V_{1k}}{c_s}, \quad (3)$$

where c_s is the sound speed of the gas and $\eta = 1.0 \pm 0.3$ is the proportionality coefficient calculated from cosmological simulations of galaxy clusters (Zhuravleva et al. 2014, see also Zhuravleva et al. 2015 for applications). A similar method has been recently used by Eckert et al. (2017). Using this approach, we compare the amplitude of the density fluctuations in the regions of the cluster with and without radio halo emission and derive information on the spectrum of the velocity field in the two regions. We have processed the *Chandra* data published by van Weeren et al. (2017), and analysed the cluster image in 0.5–3.5 keV band. This band is chosen because the X-ray surface brightness is almost independent on the gas temperature. In order to remove a first-order global density gradient, we have fitted the radial profile of the X-ray surface brightness with a spherically symmetric β model, and divided the image by this model. We have computed the power spectrum of the X-ray surface brightness fluctuations using a modified Δ –variance method (Churazov et al. 2012). Following Churazov et al. (2012), the X-ray surface brightness fluctuations have been analysed using a 2D power spectrum approach, and the resulting spectrum in 2D has been then converted to a 3D power spectrum of gas density fluctuations. The cluster X-ray emission is complex, and a spherically symmetric β -model is not an accurate description of the gas distribution. None the less, it permits to remove a first-order density




Figure 7. Red region: amplitude of the gas density fluctuations as a function of the wavenumber k derived for the whole cluster emission (white circle in the inset). Purple and light blue regions: amplitude of the gas density fluctuations computed for the east and west part, respectively.

gradient. The results that follow depend on the underlying model. In Section 4.3, we discuss how our assumption affects the results.

In Fig. 7, we show the amplitude of density fluctuations as a function of wavenumber k in the two regions of the clusters. On a scale of ~ 350 kpc, the average amplitude of density fluctuations is 0.77 ± 0.09 in the region of the halo and 0.55 ± 0.05 in the region without radio halo. This gives a ratio of ~ 1.4 between the two. If equation 3 holds, neglecting the differences in density, and assuming that c_s is the same in the two regions, we can conclude that the ratio of kinetic over thermal energy is twice as large in the region with radio emission than in the region without. However, there are indications that the temperature in the two regions is different by a factor ~ 1.4 (van Weeren et al. 2017), which would translate in a factor 3 of ratio of kinetic over thermal energy in the regions with and without radio emission. Interestingly, the ratio of the average radio power at 147 MHz in the regions with and without radio emission is more than a factor 40 (at 1σ). Neglecting effects due to different magnetic fields and/or populations of seed particles in the two regions, our results suggest that the power emitted by electrons in the radio halo at 147 MHz has a super-linear scaling with the gas kinetic energy.

4.3 Underlying model of the X-ray surface brightness.

The results obtained in the previous section depend on the assumptions we have made on the X-ray surface brightness distribution of the cluster. We have modelled the X-ray surface brightness using a spherically symmetric β -model, that is a good representation for virialized systems. As the cluster is in a very active merger state, non-negligible departures from a spherically symmetric β -model are expected. This can be seen from the X-ray image shown in Fig. 1, and also in Fig. 9, where we show the residuals of the X-ray surface brightness after division by the spherically symmetric β -model.

To check how the asymmetry of the gas distribution affects the density amplitude measurements, we repeated the analysis of




Figure 8. Same as Fig. 7 but considering a patched β -model for the cluster density distribution. Red region: amplitude of the gas density fluctuations as a function of the wavenumber k derived for the whole cluster emission. Purple and light blue regions: amplitude of the gas density fluctuations computed for the east and west part, respectively.

Section 4.2 considering a different β -model (so-called ‘patched’ β -model, Zhuravleva et al. 2015), that is elongated in the SE-NW direction. Our patched β -model is defined as in Zhuravleva et al. (2015), i.e. $I_{pm} = I_{\beta} S_{\sigma} [I_X / I_{\beta}]$, where I_{β} is the spherically symmetric β -model, I_X is the cluster X-ray surface brightness, $S_{\sigma}[\cdot]$ is the Gaussian smoothing with the smoothing window size σ . We choose $\sigma = 50$, and the resulting patched β -model is shown in Fig. 9 (top right panel). In the same figure, we also plot the residuals of the X-ray surface brightness distribution, obtained by dividing the X-ray image by the patched β -model (bottom right panel). In Fig. 8, we show the amplitude of the gas density fluctuations as a function of the wavenumber k , obtained assuming a patched β -model instead of a spherically symmetric β -model. The amplitude of the gas density fluctuations is suppressed in both the east and west regions, indicating a strong dependence on the underlying model.

On a scale of ~ 350 kpc, the average amplitude of density fluctuations is now 0.47 ± 0.02 in the region of the halo and 0.41 ± 0.04 in the region without radio halo. Hence, they are consistent within 1σ . If we only consider the mean value, we obtain a ratio of ~ 1.13 for the two regions, hence v_{kin} that is 30 per cent higher in the region of the radio halo.

This analysis indicates that our results depend on the model of gas density that we use. Either using a spherically symmetric and a patched β -model, the amplitude of the gas density fluctuations is higher in the east region than in the west region, although when we consider the patched β -model the difference is only marginal, and the amplitudes are consistent within 1σ .

Constraining the dependence of the radio power of the gas kinetic energy would give important constraints for theoretical models of halo formation (Brunetti & Lazarian 2016). The analysis we have performed here suggests a super-linear scaling of the radio power with respect to the gas kinetic energy, as well as a strong dependence of the results on the underlying model for the cluster gas density distribution.




Figure 9. Top panels: Normalized spherically symmetric β -model (left) and patched β -model (right) used to derive a first-order density gradient of the gas. Bottom panels: residuals of the X-ray emission after division by the spherically symmetric β -model (left) and patched β -model (right).

5 CONCLUSIONS

MACSJ0717 is undergoing a violent merger that involves at least four sub-clusters. The radio emission is complex and shows unique features that are visible from 147 MHz up to 5 GHz. We have presented new results from LOFAR and GMRT observations of the galaxy cluster MACSJ0717, and using X-ray observations, we have derived new constraints on the particle acceleration processes in the cluster centre and outskirts. Our results can be summarized as follows:

(i) LOFAR observations at 147 MHz reveal new emission from the ICM: (i) a radio arc located north-west of the cluster centre and extending for 1.9 Mpc in the north-south direction; (ii) a radio bridge connecting the main cluster to an HT radio galaxy located in the direction of a 19 Mpc-long filament of galaxies; (iii) a radio bar that traces the X-ray bar observed south of the main mass component.

(ii) Using GMRT observations at 608 MHz, we have constrained the spectra of the radio arc ($\alpha_{608\text{ MHz}}^{147\text{ MHz}} < -1.3$), the radio bridge ($\alpha_{608\text{ MHz}}^{147\text{ MHz}} < -1.4$), and the radio bar ($\alpha_{608\text{ MHz}}^{147\text{ MHz}} < -1.4$). The spectra of the radio halo and of the radio relic do not show a significant departure from the power law observed at higher frequencies and already studied in the literature.

(iii) We have investigated under which conditions the radio bridge can originate from electron re-acceleration by a weak Mach number shock. A ‘transversal’ shock moving perpendicular to the filament main axis can explain the properties of the radio emission, although some fine tuning of the parameters is required.

(iv) The radio halo at LOFAR frequencies is more extended than previously observed. None the less, the radio emission does not follow the X-ray emission of the gas in the western part of the cluster. Assuming that the spectrum of density fluctuations – as deduced from *Chandra* observations – traces the spectrum of the gas velocity, data suggest a different ratio of kinetic over thermal energy

in the regions with and without radio halo. This result depends on the model we assume for the gas density distribution, and lacks a robust statistical significance. Deeper observations, as well as a more accurate modelling for the cluster density distribution would be required to investigate this point in more detail.

ACKNOWLEDGEMENTS

Annalisa Bonafede acknowledges financial support from the ERC-Stg DRANOEL, no 714245. FV acknowledges financial support from the ERC-Stg MAGCOW, no.714196. RJvW and HJAR acknowledge support from the ERC Advanced Investigator programme NewClusters 321271 and the VIDI research programme with project number 639.042.729, which is financed by the Netherlands Organization for Scientific Research (NWO). We thank Prof. Daniele Dallacasa for useful discussions, and the referee for useful comments. This work had made use of the Lofar Solution Tool (LoSoTo), developed by F. de Gasperin. F.d.G. is supported by the VENI research programme with project number 1808, which is financed by the Netherlands Organization for Scientific Research (NWO). We thank the staff of the GMRT that made these observations possible. GMRT is run by the National Centre for Astrophysics of the Tata Institute of Fundamental Research.

This research has made use of the NASA/IPAC Extragalactic Database (NED), that is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Adam R. et al., 2017, preprint ([arXiv: 1712.01836](https://arxiv.org/abs/1712.01836))
 Bonafede A. et al., 2009, *A&A*, 503, 707
 Bonafede A. et al., 2014, *MNRAS*, 444, L44
 Briggs D. S., 1995, PhD thesis, ‘High fidelity deconvolution of moderately resolved sources, New Mexico Institute of Mining Technology, Socorro, New Mexico, USA
 Brunetti G., Jones T. W., 2014, *Int. J. Mod. Phys. D*, 23, 1430007
 Brunetti G., Lazarian A., 2016, *MNRAS*, 458, 2584
 Brüggén M., Bykov A., Ryu D., Röttgering H., 2011, *SSR*, p. 138
 Buote D. A., 2001, *ApJ*, 553, L15
 Chang J. S., Cooper G., 1970, *J. Comput. Phys.*, 6, 1
 Churazov E. et al., 2012, *MNRAS*, 421, 1123
 Condon J. J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B., Broderick J. J., 1998, *AJ*, 115, 1693
 Cotton W. D., 2008, *PASP*, 120, 439
 Cuciti V., Cassano R., Brunetti G., Dallacasa D., Kale R., Etori S., Venturi T., 2015, *A&A*, 580, A97
 Dolag K., Bartelmann M., Lesch H., 2002, *A&A*, 387, 383
 Drury L., 1983, *Space Sci. Rev.*, 36, 57
 Ebeling H., Barrett E., Donovan D., 2004, *ApJ*, 609, L49
 Ebeling H., Barrett E., Donovan D., Ma C.-J., Edge A. C., van Speybroeck L., 2007, *ApJ*, 661, L33
 Ebeling H., Edge A. C., Henry J. P., 2001, *ApJ*, 553, 668
 Ebeling H., Ma C.-J., Barrett E., 2014, *ApJS*, 211, 21
 Eckert D., Gaspari M., Vazza F., Gastaldello F., Tramacere A., Zimmer S., Etori S., Paltani S., 2017, *ApJ*, 843, L29
 Edge A. C., Ebeling H., Bremer M., Röttgering H., van Haarlem M. P., Rengelink R., Courtney N. J. D., 2003, *MNRAS*, 339, 913
 Feretti L., Giovannini G., Govoni F., Murgia M., 2012, *A&AR*, 20, 54
 Ginzburg V. L., Syrovatskii S. I., 1965, *ARA&A*, 3, 297
 Giovannini G., Feretti L., Venturi T., Kim K.-T., Kronberg P. P., 1993, *ApJ*, 406, 399
 Govoni F., Feretti L., Giovannini G., Böhringer H., Reiprich T. H., Murgia M., 2001, *A&A*, 376, 803

- Intema H. T., Jagannathan P., Mooley K. P., Frail D. A., 2017, *A&A*, 598, A78
- Intema H. T., van der Tol S., Cotton W. D., Cohen A. S., van Bemmell I. M., Röttgering H. J. A., 2009, *A&A*, 501, 1185
- Jauzac M. et al., 2012, *MNRAS*, 426, 3369
- Kang H., Ryu D., Jones T. W., 2012, *ApJ*, 756, 97
- Kardashev N. S., 1962, *SvA*, 6, 317
- Lane W. M., Cotton W. D., van Velzen S., Clarke T. E., Kassim N. E., Helmboldt J. F., Lazio T. J. W., Cohen A. S., 2014, *MNRAS*, 440, 327
- Limousin M. et al., 2016, *A&A*, 588, A99
- Ma C.-J., Ebeling H., Donovan D., Barrett E., 2008, *ApJ*, 684, 160
- Markevitch M., Govoni F., Brunetti G., Jerius D., 2005, *ApJ*, 627, 733
- McMullin J. P., Waters B., Schiebel D., Young W., Golap K., 2007, in Shaw R. A., Hill F., Bell D. J., eds, *ASP Conf. Ser.*, 376, San Francisco, p. 127
- Medezinski E. et al., 2013, *ApJ*, 777, 43
- Miniati F., Beresnyak A., 2015, *Nature*, 523, 59
- Offringa A. R., Smirnov O., 2017, *MNRAS*, 471, 301
- Offringa A. R., van de Gronde J. J., Roerdink J. B. T. M., 2012, *A&A*, 539, A95
- Offringa A. R. et al., 2014, *MNRAS*, 444, 606
- Ogrea , et al., 2018, submitted
- Pandey-Pommier M., Richard J., Combes F., Dwarakanath K. S., Guiderdoni B., Ferrari C., Sirothia S., Narasimha D., 2013, *A&A*, 557, A117
- Planck Collaboration et al., 2016, *A&A*, 594, A27
- Rengelink R. B., Tang Y., de Bruyn A. G., Miley G. K., Bremer M. N., Roettgering H. J. A., Bremer M. A. R., 1997, *A&AS*, 124, 259
- Sarazin C. L., 1999, *ApJ*, 520, 529
- Savini F. et al., 2018, *MNRAS*, 474, 5023
- Sayers J. et al., 2013, *ApJ*, 778, 52
- Scaife A. M. M., Heald G. H., 2012, *MNRAS*, 423, L30
- Shimwell T. W. et al., 2016, *MNRAS*, 459, 277
- Sommer M. W., Basu K., Intema H., Pacaud F., Bonafede A., Babul A., Bertoldi F., 2017, *MNRAS*, 466, 996
- Umetsu K. et al., 2014, *ApJ*, 795, 163
- van Haarlem M. P. et al., 2013, *A&A*, 556, A2
- van Weeren R. J., Röttgering H. J. A., Brügger M., Cohen A., 2009, *A&A*, 505, 991
- van Weeren R. J., Röttgering H. J. A., Brügger M., Hoeft M., 2010, *Science*, 330, 347
- van Weeren R. J. et al., 2016, *ApJ*, 817, 98
- van Weeren R. J. et al., 2016, *ApJS*, 223, 2
- van Weeren R. J. et al., 2017, *ApJ*, 835, 197
- Vazza F., Eckert D., Brügger M., Huber B., 2015, *MNRAS*, 451, 2198
- Vazza F., Ferrari C., Brügger M., Bonafede A., Gheller C., Wang P., 2015, *A&A*, 580, A119
- Wilber A. et al., 2018, *MNRAS*, 473, 3536
- Zhuravleva I. et al., 2014, *ApJ*, 788, L13
- Zhuravleva I. et al., 2015, *MNRAS*, 450, 4184

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.