



Publication Year	2019
Acceptance in OA @INAF	2021-02-26T17:08:06Z
Title	HARPS-N radial velocities confirm the low densities of the Kepler-9 planets
Authors	BORSATO, LUCA; Malavolta, Luca; Piotto, G.; Buchhave, L. A.; Mortier, A.; et al.
DOI	10.1093/mnras/stz181
Handle	http://hdl.handle.net/20.500.12386/30657
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
Number	484

HARPS-N radial velocities confirm the low densities of the Kepler-9 planets

L. Borsato¹,^{1,2}★ L. Malavolta^{1,2} G. Piotto^{1,2} L. A. Buchhave³ A. Mortier⁴,
 K. Rice^{5,6} A. C. Cameron⁷ A. Coffinet⁸ A. Sozzetti⁹ D. Charbonneau¹⁰
 R. Cosentino^{11,12} X. Dumusque⁸ P. Figueira¹³ D. W. Latham¹⁰ M. Lopez-Morales¹⁰
 M. Mayor⁸ G. Micela¹⁴ E. Molinari^{11,15} F. Pepe⁸ D. Phillips¹⁰ E. Poretti^{11,16}
 S. Udry⁸ and C. Watson¹⁷

¹Dipartimento di Fisica e Astronomia ‘Galileo Galilei’, Università di Padova, vicolo dell’Osservatorio 3, Padova I-35122, Italy

²INAF – Osservatorio Astronomico di Padova, vicolo dell’Osservatorio 5, Padova I-35122, Italy

³DTU Space, National Space Institute, Technical University of Denmark, Elektrovej 327, DK-2800 Lyngby, Denmark

⁴Astrophysics Group, Cavendish Laboratory, University of Cambridge, J.J. Thomson Avenue, Cambridge CB3 0HE, UK

⁵SUPA, Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK

⁶Centre for Exoplanet Science, University of Edinburgh, Peter Guthrie Tait Road, Edinburgh EH9 3FD, UK

⁷Centre for Exoplanet Science, SUPA, School of Physics and Astronomy, University of St Andrews, St Andrews KY16 9SS, UK

⁸Observatoire de Genève, Université de Genève, 51 ch. des Maillettes, CH-1290 Versoix, Switzerland

⁹INAF – Osservatorio Astrofisico di Torino, Via Osservatorio 20, I-10025 Pino Torinese, Italy

¹⁰Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

¹¹INAF – Fundación Galileo Galilei, Rambla José Ana Fernández Pérez 7, E-38712 Breña Baja, Spain

¹²INAF – Osservatorio Astrofisico di Catania, Via S. Sofia 78, I-95123 Catania, Italy

¹³European Southern Observatory (ESO), Alonso de Cordova 3107, Vitacura, Santiago, Chile

¹⁴INAF – Osservatorio Astronomico di Palermo, Piazza del Parlamento 1, I-90134 Palermo, Italy

¹⁵INAF – Osservatorio di Cagliari, via della Scienza 5, I-09047 Selargius, CA, Italy

¹⁶INAF – Osservatorio Astronomico di Brera, Via E. Bianchi 46, I-23807 Merate, Italy

¹⁷Astrophysics Research Centre, School of Mathematics and Physics, Queen’s University Belfast, University Rd, Belfast BT7 1NN, UK

Accepted 2019 January 13. Received 2018 December 20; in original form 2018 November 21

ABSTRACT

We investigated the discrepancy between planetary mass determinations using the transit timing variations (TTVs) and radial velocities (RVs), by analysing the multiplanet system Kepler-9. Despite being the first system characterized with TTVs, there are several discrepant solutions in the literature, with those reporting lower planetary densities being apparently in disagreement with high-precision RV observations. To resolve this, we gathered HARPS-N RVs at epochs that maximized the difference between the predicted RV curves from discrepant solutions in the literature. We also reanalysed the full *Kepler* data set and performed a dynamical fit, within a Bayesian framework, using the newly derived central and duration times of the transits. We compared these results with the RV data and found that our solution better describes the RV observations, despite the masses of the planets being nearly half that presented in the discovery paper. We therefore confirm that the TTV method can provide mass determinations that agree with those determined using high-precision RVs. The low densities of the planets place them in the scarcely populated region of the super-Neptunes/inflated sub-Saturns in the mass–radius diagram.

Key words: techniques: radial velocities – techniques: spectroscopic – stars: fundamental parameters – stars: individual: Kepler-9.

1 INTRODUCTION

One of the most important accomplishments of the *Kepler* mission (Borucki et al. 2011) is the demonstration that transit timing variation (TTV) is a powerful tool to estimate the masses of planets

* E-mail: luca.borsato@unipd.it

around stars (Agol et al. 2005; Holman & Murray 2005) that are too faint for a proper radial velocity (RV) follow up. Notable, early examples are the characterization of the two Saturn-like planets around Kepler-9 (Holman et al. 2010), a system of five low-mass, small-size planets around Kepler-11 (Lissauer et al. 2011), and the three-planet system around Kepler-18 (Cochran et al. 2011). However, with the increasing number of well-characterized, low-mass planets, a marked difference in the density distribution of planets with TTV- and RV-derived masses has started to appear. This suggests the presence of an intrinsic problem with one of the two techniques (Weiss & Marcy 2014). Subsequent studies on individual systems involving both TTV and RVs, such as WASP-47 (Becker et al. 2015; Weiss et al. 2017), K2-19 (Barros et al. 2015; Nespral et al. 2017), and Kepler-19 (Malavolta et al. 2017a), as well as ensemble studies on system with different characteristics (Jontof-Hutter et al. 2016) and statistical analysis on simulated observations (Steffen 2016; Mills & Mazeh 2017), showed that both techniques lead to similar results, and the discrepancy in planetary density is likely the result of an observational bias. In some cases, inconsistencies between TTV and RV masses still persist, as for KOI-94d where the dynamical mass (Masuda et al. 2013) is half the mass obtained by high-precision RVs (Weiss et al. 2013). For this reason, it is important to analyse as many systems as possible with both techniques.

In this paper, we focus on the planetary system around Kepler-9, a faint ($V = 13.9$) Sun-like star. From the analysis of the three quarters of *Kepler* data, Holman et al. (2010, hereafter H10) identified two transiting Saturn-size planets with periods and radii of $P_b = 19.24$ d, $R_b = 9.4 R_\oplus$ and $P_c = 38.91$ d, $R_c = 9.2 R_\oplus$, respectively, and another transiting body validated by Torres et al. (2011) as a super-Earth size planet with period and radius of $P_d = 1.59$ d, $R_d = 1.64 R_\oplus$. Using TTVs coupled with six RV measurements obtained with Keck-HIRES, they determined a mass of $80.0 \pm 4.1 M_\oplus$ for Kepler-9b and $54.3 \pm 4.1 M_\oplus$ for Kepler-9c. A subsequent work by Borsato et al. (2014, hereafter B14) nearly halved the mass determinations, $M_b = 43.5 \pm 0.6 M_\oplus$ and $M_c = 29.8 \pm 0.6 M_\oplus$.¹ The new analysis was performed using time of transits (T_0 s) extracted from 12 *Kepler* quarters, but the HIRES RVs were excluded because the combined fit was not particularly good. Both results were obtained using a two-planet model, since the TTV amplitude induced by Kepler-9d (Torres et al. 2011) is expected to be only tens of seconds, and is, hence, too low to be measured in the *Kepler* long-cadence data (Holman et al. 2010). Very recently, these results were confirmed by Freudenthal et al. (2018) within the project Kepler Object of Interest Network (KOINet; von Essen et al. 2018). Using a photodynamical model, they analysed the photometric data of all the 17 *Kepler* quarters and 13 new ground-based light curves. The Keck-HIRES data are not consistent with their solution, as in B14, and the discrepancy has been ascribed to stellar activity.

The two Saturn-like planets in the Kepler-9 system belong to the small group of planets whose masses can be obtained dynamically by modelling the TTVs and whose RV signals are detectable with current facilities. However, the two sets of solutions actually available in the literature for this system are either partially inconsistent with transit timings obtained after the publication of the discovery paper (H10) or with high-precision RVs (B14 and following analyses), and this inconsistency has never been dealt

with in any work.² For this reason, we decided to observe the target with HARPS-N, the high-precision velocimeter mounted at the Telescopio Nazionale Galileo (La Palma) to understand which of the two solutions is more consistent with an independent set of RVs. At the same time, we aim to improve the literature values by reanalysing all the 17 *Kepler* quarters and provide a more robust estimate on the error bars of the orbital parameters. In this work, we describe the observational strategy we pursued with HARPS-N within the Guaranteed Time Observations programme of the HARPS-N Collaboration, the comparison of RVs with the literature solutions, and the details on the determination of the updated orbital parameters.

2 OBSERVATIONAL STRATEGY

With a magnitude of $V = 13.9$, mass determination of Kepler-9b and c is a very challenging task even for HARPS-N. The planets have an expected RV semi-amplitude of $K_b \simeq 19 \text{ m s}^{-1}$ and $K_c \simeq 10 \text{ m s}^{-1}$ from the H10 solution, and $K_b \simeq 10 \text{ m s}^{-1}$, $K_c \simeq 6 \text{ m s}^{-1}$ from the B14 solution. This is comparable with the expected RV error of $\simeq 11 \text{ m s}^{-1}$ for a 30 min exposure. An independent RV determination of the mass of the planets at 5σ would be extremely time consuming, especially in the case of the low-mass scenario, and it would compete against other *Kepler* targets more fitting to the science goal of the HARPS-N Collaboration (e.g. precise mass determination of super-Earth and mini-Neptune planets), in the limited window visibility of the *Kepler* field during a night. For this target, we specifically designed an observational strategy that could allow us to distinguish between the two proposed solutions. We first propagated the solution of H10 and B14 to cover the observing season, using the dynamical integrator embedded in TRADES³ (Borsato et al. 2014). For consistency, we used the same stellar mass of the two papers. Within the nights allocated to HARPS-N Collaboration, we selected those epochs in which the difference between the H10 and B14 expected RVs was at its maximum (Fig. 1). To reduce the CCF noise associated with a single epoch without introducing systematic errors due to the variation of the barycentric RV correction within the exposure time, we gathered – whenever possible – two consecutive 30 min exposures. Following this strategy, we obtained a total of 16 epochs divided in 30 exposures of 1800 s (in two nights only one exposure was taken), with an average signal-to-noise ratio of 16 at 5500 Å and an average internal error of 11.6 m s^{-1} per exposure. RVs were corrected for Moon contamination following the recipe described in Malavolta et al. (2017a) and successfully applied by Osborn et al. (2017). Table B1 lists the final RV measurements.

3 STELLAR PARAMETERS

We followed the same approach described in Malavolta et al. (2018) to determine the mass, radius, and density of the star. We started by measuring the photospheric parameters of the target with three different techniques on a spectrum obtained by stacking all the HARPS-N exposures (signal-to-noise ratio $\simeq 90$). Using CCFpams⁴ (Malavolta et al. 2017b), we obtained

²Wang et al. (2018) presented 21 new KECK-Hires RVs spanning the transit of Kepler-9b, but they assumed a dynamical model similar to that of B14 to model the data.

³Available at <https://github.com/lucaborsato/trades>.

⁴Available at <https://github.com/LucaMalavolta/CCFpams>.

¹The authors acknowledged that bootstrap-derived error bars were likely underestimated.




Figure 1. Scheduling plot: predicted RV model for H10 (blue) and B14 (red), with 2.5 and 10 m s^{-1} uncertainties (light and dark shaded areas, respectively), used to schedule the HARPS-N observations (white–black circles).

$T_{\text{eff}} = 5836 \pm 51$ K, $\log g = 4.50 \pm 0.10$, and $[\text{Fe}/\text{H}] = 0.04 \pm 0.04$. The ARES + MOOG⁵ approach (e.g. Mortier et al. 2014) returned $T_{\text{eff}} = 5827 \pm 35$ K, $\log g = 4.46 \pm 0.04$, and $[\text{Fe}/\text{H}] = 0.12 \pm 0.03$. Finally, with the Stellar Parameters Classification tool (SPC; Buchhave et al. 2012, 2014), we obtained $T_{\text{eff}} = 5750 \pm 50$ K, $\log g = 4.45 \pm 0.10$, and $[\text{Fe}/\text{H}] = -0.02 \pm 0.08$. All the reported errors are internal only.

For each set of stellar parameters, we then determined the stellar mass and radius using isochrones (Morton 2015), with posterior sampling performed by MultiNest (Feroz & Hobson 2008; Feroz, Hobson & Bridges 2009; Feroz et al. 2013). We provided as input the parallax of the target from the *Gaia* DR2 catalogue ($p = 1.563 \pm 0.017$ mas, $d = 640 \pm 7$ pc; Gaia Collaboration 2016, 2018) with the correction suggested by Stassun & Torres (2018) of $-82 \pm 33 \mu\text{as}$, plus the photometry from the Two Micron All Sky Survey (2MASS; Cutri et al. 2003; Skrutskie et al. 2006). For stellar models, we used both MESA Isochrones and Stellar Tracks (MIST; Choi et al. 2016; Dotter 2016; Paxton et al. 2011) and the Dartmouth Stellar Evolution Database (Dotter et al. 2008). We performed the analysis on the photospheric parameters obtained with HARPS-N as well as the literature values obtained by H10, Huber et al. (2014), Petigura et al. (2017), and Wang et al. (2018). For all methods, we assumed $\sigma_{T_{\text{eff}}} = 75$ K, $\sigma_{\log g} = 0.10$, and $\sigma_{[\text{Fe}/\text{H}]} = 0.05$ as a good estimate of the systematic errors, when the provided errors were lower than these values.

From the median and standard deviation of all the posterior samplings, we obtained $M_{\star} = 1.022_{-0.029}^{+0.039} M_{\odot}$ and $R_{\star} = 0.96 \pm 0.02 R_{\odot}$. We derived the stellar density $\rho_{\star} = 1.16_{-0.09}^{+0.08} \rho_{\odot}$ directly from the posterior distributions of M_{\star} and R_{\star} . Following Lovis et al. (2011), from the individual HARPS-N

Table 1. Astrophysical parameters of the star.

Parameter	Value	Unit
2MASS alias	J19021775+3824032	–
α_{J2000}	19:02:17.76	hms
δ_{J2000}	+38:24:03.2	dms
R_{\star}	0.958 ± 0.020	R_{\odot}
M_{\star}	$1.022_{-0.039}^{+0.029}$	M_{\odot}
ρ_{\star}	$1.16_{-0.09}^{+0.08}$	ρ_{\odot}
$\log(L_{\star}/L_{\odot})$	$-0.038_{-0.027}^{+0.026}$	–
T_{eff}	5774 ± 60	K
$\log g$	$4.49_{-0.03}^{+0.02}$	–
$[\text{Fe}/\text{H}]$	0.05 ± 0.07	–
$p^{(a)}$	1.643 ± 0.037	mas
distance	614 ± 13	pc
A_V	$0.10_{-0.07}^{+0.10}$	mag
$B-V$	0.70 ± 0.13	mag
age	$2.0_{-1.3}^{+2.0}$	Gyr
$\log R'_{\text{HK}}$	-4.67 ± 0.09	–

Note. ^(a)*Gaia* parallax corrected for systematic offset as suggested by Stassun & Torres (2018).

exposures we measured a $\log R'_{\text{HK}}$ index of -4.67 ± 0.09 , consistent with the young age of the star ($2.0_{-1.3}^{+2.0}$ Gyr) derived from the fit of the isochrones (e.g. Pace 2013). The astrophysical parameters of the star are summarized in Table 1, where the temperature, gravity, and metallicity are those obtained from the posterior distributions to take into account the constraint from *Gaia* parallax. The stellar density determined in this work agrees very well with the value derived in Freudenthal et al. (2018) with the dynamical analysis of the light curve.

4 ANALYSIS OF THE KEPLER DATA

We downloaded the Kepler-9 data from the MAST⁶ covering the full 17 quarters of the mission in long (LC) and short (SC) cadence. We used the Presearch Data Conditioning (PDC) fluxes instead of Simple Aperture Photometry (SAP), because the handling of systematic trends and errors was out of the purpose of this work. We created a normalized full light curve by dividing the PDC flux of each quarter by its median value. We selected carefully the portion of the transits and the out-of-transit, in particular when the light curve shows transit events of Kepler-9b and Kepler-9c very close to each other.

For each light curve, we fitted the following parameters: $\log_{10} \rho_{\star}$, where ρ_{\star} is the stellar density in kg m^{-3} , \sqrt{k} where k is the ratio between the planetary and stellar radii, \sqrt{b} where b is the impact parameter,⁷ the central time of the transit (T_0), a quadratic limb-darkening (LD) law with the parameters q_1 and q_2 introduced in Kipping (2013), a linear trend (with a_0 as the intercept and a_1 the angular coefficient), and $\log_2 \sigma_j$, where σ_j is a jitter term to add in quadrature to the errors of the *Kepler* light curve. We used an asymmetric prior for the $\log_{10} \rho_{\star}$ based on the stellar mass and radius values determined in Section 3, while we used a uniform prior for the other parameters (see Table 2); we kept fixed the period (P), the eccentricity, and argument of pericentre (values from B14 solution).

⁵ARESv2: available at <http://www.astro.up.pt/~sousasag/ares/> (Sousa et al. 2015); MOOG: available at <http://www.as.utexas.edu/~chris/moog.html> (Snedden 1973).

⁶Mikulsky Archive for Space Telescope, data release 25 (DR25).

⁷We used the equation of b from Winn (2010) and Kipping (2010) that takes into account non-zero eccentricity (e) and the argument of pericentre (ω).

Table 2. Table of the priors, boundaries, and final solution of the fitted parameters for transit analysis and the derived parameters.

Parameter	Prior type	Boundaries (min, max)	Kepler-9b	Kepler-9c
$\log_{10}\rho_*$	G	[0., 6.]	$3.196^{+0.046}_{-0.060}$	$3.188^{+0.048}_{-0.067}$
\sqrt{k}	U	[0., 1.]	$0.2786^{+0.0033}_{-0.0030}$	$0.2750^{+0.0033}_{-0.0027}$
\sqrt{b}	U	$[0., \sqrt{2}]$	$0.767^{+0.042}_{-0.035}$	$0.861^{+0.024}_{-0.022}$
T_0	U	$[T_0^{\text{guess}} \pm 1.5 \times T_{14}]$	–	–
q_1	U	[0., 1.]	$0.43^{+0.17}_{-0.26}$	$0.48^{+0.20}_{-0.36}$
q_2	U	[0., 1.]	$0.18^{+0.20}_{-0.18}$	$0.18^{+0.16}_{-0.18}$
a_0	U	[–100., 100.]	–	–
a_1	U	[–100., 100.]	–	–
$\log_2\sigma_j$	U	$[\log_2(< \sigma_i > \times 10^{-4}), 0.]$	–	–
<i>Derived transit model</i>				
ρ_* (ρ_\odot)	–	–	$1.12^{+0.11}_{-0.16}$	$1.10^{+0.11}_{-0.17}$
a/R_*	–	–	$31.3^{+1.1}_{-1.5}$	$49.8^{+1.7}_{-2.6}$
a (au)	–	–	$0.143^{+0.007}_{-0.006}$	$0.227^{+0.012}_{-0.008}$
k	–	–	$0.0776^{+0.0019}_{-0.0015}$	$0.0756^{0.0018}_{-0.0014}$
R_p (R_\oplus)	–	–	$8.29^{+0.54}_{-0.43}$	$8.08^{+0.54}_{-0.41}$
b	–	–	$0.59^{0.06}_{-0.05}$	$0.74^{+0.04}_{-0.04}$
i ($^\circ$)	–	–	$88.9^{+0.1}_{-0.2}$	$89.1^{+0.1}_{-0.1}$
T_{14} (min)	–	–	$254.6^{+7.9}_{-3.5}$	$273.9^{+7.3}_{-7.3}$
u_1	–	–	$0.24^{+0.23}_{-0.24}$	$0.26^{+0.20}_{-0.26}$
u_2	–	–	$0.41^{0.11}_{-0.41}$	$0.43^{+0.14}_{-0.43}$
<i>Linear ephemeris</i>				
T_0 (BJD _{TDB}), this work	–	–	2454977.512 ± 0.065	2454968.84 ± 0.20
P_{ephem} (days), this work	–	–	19.2460 ± 0.0015	38.9492 ± 0.0093

Note. Prior type U means Uniform, while G means Gaussian (it could be an asymmetric Gaussian). The stellar density prior has been computed as asymmetric Gaussian from M_* and R_* . T_0^{guess} has been obtained from the selection of each transit light curve, while the T_{14} is the transit total duration (equation 30 in Kipping 2010). Following Kipping (2013), we checked that the values of q_1 and q_2 were mapped to physical values of the quadratic LD coefficients u_1 and u_2 . The linear trend coefficients (a_0 , a_1) are bounded to very high values just to prevent singular behaviour. The minimum value for the $\log_2\sigma_j$ has been computed taking into account the mean value of the photometric errors (σ_i) of the portion of the light curve. We fixed the period, eccentricity, and argument of pericentre of both planets at the values of the best fit of B14.

We ran a Differential Evolution algorithm (Storn & Price 1997, pyDE⁸) and then a Bayesian analysis of each selected light curve around each transit by using the affine-invariant ensemble sampler (Goodman & Weare 2010) for Markov Chain Monte Carlo (MCMC) implemented within the emcee package (Foreman-Mackey et al. 2013) and modelling each transit with batman (Kreidberg 2015). We took into account the long exposure time of the LC data oversampling⁹ the transit model; for consistency, we used the same oversampling also for the SC data.

We fit each transit portion for both planets b and c for the SC with 50 walkers for 25 000 generations with pyDE and then with 50 000 steps with emcee. Then we discarded as burn-in the first 25 000 steps after checking the convergence of the chains with the Gelman–Rubin (GR) statistics (Gelman & Rubin 1992, $R = 1.01$). For each transit, we obtained a final posterior after applying a thinning factor of 100, i.e. a pessimistic estimate of the autocorrelation time of the chains. All the posteriors from different transit fits were merged to obtain the final posterior of the parameters $\log_{10}\rho_*$, \sqrt{k} , \sqrt{b} , q_1 , and q_2 .

For each transit and merged posterior, we computed the high density interval (HDI¹⁰) at 68.27 per cent (equivalent to 1σ error) for the fitted parameters and other physical quantities of interest

derived from them. As parameter estimation of each transit we selected the parameter set that maximizes the likelihood (maximum likelihood estimation, MLE) within the HDI, while we computed the median of the merged posterior distributions. The values of \sqrt{k} , q_1 , and q_2 extracted from the SC merged posteriors have been used as priors for the LC analysis (same number of walkers, generations, steps, and burn-in). When we did not use the priors from the SC, we found that the mean transit duration for LC was 10σ longer than the SC for both planets.

See Figs 2 and 3 for the river plots of planets b and c, respectively, showing the data and the computed model of each transit in LC and SC and the TTV effect with respect to a linear ephemeris.

As final parameters from the light curve analysis, we decided to use the mean between the SC and LC median values of the merged posteriors; we associated as lower uncertainty the lower value between the SC and LC and the greater values for the upper uncertainty. The results are summarized in Table 2.

5 ORBITAL PARAMETERS FROM DYNAMICAL ANALYSIS

We extract from the LC and SC analysis the central time, T_0 , and the total duration, T_{14} (defined as the difference between fourth and first contact time and computed with equation 30 in Kipping 2010), of each transit; we kept the time and duration from SC when present, otherwise we used the analysis of the LC. We assigned as symmetric errors of T_0 s and T_{14} s the maximum between the lower and upper

⁸We used the python implementation pyDE available at <https://github.com/hpparvi/PyDE>.

⁹We used an oversampling with a subexposure time of 9 s.

¹⁰Also indicated as high-density region or high-probability region.




Figure 2. River plot for Kepler-9b: the LC (and SC when available) of each transit has been plotted with colour code depending on the normalized magnitude (transit with darker colour, best-fitting model in orange). The light curves are sorted vertically by each epoch (or transit number with respect to a reference time) as a function of the phase (in hours) with respect to a linear ephemeris (vertical line at zero) showing the TTV effect.




Figure 3. Same as Fig. 2, but for Kepler-9c.

values of the HDI. The measured central time and duration from each transit, in SC or LC mode, are available in Table A1.

For each planet, we used TRADES to fit the following parameters: the mass of the planet in units of stellar mass, M_p/M_* , the planetary period P_p , the eccentricity vector components $\sqrt{e_p} \cos(\omega_p)$ and $\sqrt{e_p} \sin(\omega_p)$, where e_p is the eccentricity and ω_p the argument of pericentre of the planet,¹¹ the inclination vector components $i \cos(\Omega_p)$ and $i \sin(\Omega_p)$, where i is the orbital inclination and Ω_p the longitude of ascending node, and the mean longitude of the planet λ_p , defined as the sum of the argument of pericentre, the

¹¹We stress the difference with ω_* as defined by Eastman, Gaudi & Agol (2013), that is $\omega_p = \omega_* + 180^\circ$.

longitude of the ascending node, and the mean anomaly \mathcal{M}_p . All the dynamical parameters are computed at the epoch of reference $\text{BJD}_{\text{TDB}} = 2455088.212$.

We used uniform priors with broad but physically motivated boundaries: planetary periods are constrained within 2 d of the value of the linear ephemeris, and the mass of the planets, M_p , is bounded to less than $2 M_{\text{Jup}}$. We defined the reference coordinate system as described in Winn (2010),¹² therefore Kepler-9b has fixed Ω_b to 180° , effectively reducing the inclination vector of planet b to i_b . All other parameters have been fixed to the values determined in Section 4, e.g. stellar and planetary radii.

We used an updated version of TRADES that allows us to fit T_{0S} and T_{14S} simultaneously during the planetary orbit integration and perform a Bayesian analysis with the emcee package; we used the same form of the log-likelihood, $\ln \mathcal{L}$, introduced in Malavolta et al. (2017a). Although TRADES can also fit the observed RVs as well, we did not include the HARPS-N data at this stage of the analysis because we wanted to use those observations as an independent check of the TTV-derived solution. We ran a simulation with 200 walkers, 200 000 steps, and we discarded the first 50 000 steps as burn-in.¹³ The initial walkers have been generated from a neighbourhood of the solution from B14. As in the previous section, we checked for GR statistics and we obtained the final posterior distribution after the chains had been thinned with a factor of 100; the correlation plots of the posterior distributions of the fitted and derived parameters are shown in Figs 4 and 5, respectively. We computed the final best-fitting parameter set as the MLE (and HDI) of the posterior distribution; see the summary of the best-fitting parameters in Table 3 and how they reproduce the observed T_0 and T_{14} data for Kepler-9b and Kepler-9c in Figs 6 and 7, respectively. Fig. 8 shows how the new solution with the newly derived parameters fits the RV observations.

The final solution confirmed the masses and the orbital parameters found in B14.

We also tested a parameter search fixing the inclination to the value from the LC analysis and the longitude of node to 180° , for both planets. In this case, the best-fitting model found compatible masses but it did not reproduce the trend of the T_{14} data and the final Bayesian Information Criteria were higher.

6 COMPARISON OF LITERATURE SOLUTIONS WITH RVs

We propagated the solution of H10 and B14 to the epochs of our RVs using the dynamical integrator embedded in TRADES (Borsato et al. 2014). For consistency, we used the stellar masses of the two respective papers. For each solution, we generated 10 000 sets by varying each parameter of a quantity randomly extracted from a Gaussian distribution with variance equal to the errors listed in Table 3. We then computed the χ_r^2 of the RVs for each set of orbital parameters. In both cases, we have only one free parameter, i.e. the systemic RV of the target, γ , since all the other parameters have been determined independently of our RV data set. We can therefore use the χ_r^2 to select which solution better reproduces our measurements. We select 10 000 sets of parameters from the posterior distribution

¹²Astrosentric reference system, with the plane X – Y the sky plane and Z -axis pointing towards the observer; X -axis is aligned with line of nodes and Ω fixed to 180° .

¹³We visually checked the trace plot and we found that a few walkers reached the convergence just before 50 000 steps.




Figure 4. Correlation plot of the fitted parameters. The MLE solution has been shown as a dashed blue line.

computed in Section 5 and compute the χ_r^2 . The distributions of the samples from B14 and this paper overlap each other and they are centred at $\chi_r^2 = 1.71$, while the H10 distribution has a higher χ_r^2 of about 10.5 (see Fig. 9), i.e. the solutions obtained using the T_0 alone compare well with those from the HARPS-N RVs, and do not depend on the exact number of transits involved in the analysis.

In our analysis, we did not include the six Keck/HIRES observations gathered by H10, since their apparent discrepancy with the TTV solutions presented by B14 and following works was what motivated us obtaining new RVs with HARPS-N. While our RV data set is well described by the TTV solutions presented in B14 and in this work, we still do not have an explanation for the inconsistency with the previous RVs from H10. We note, however, that it is not the first time that a few, sparse Keck/HIRES RVs are

in disagreement with a larger RV data set obtained with HARPS-N (e.g. Sozzetti et al. 2015). Given the observational strategy we employed, the large number of RVs, and the inclusion of possible source of contamination from the Moon, we believe that our data set is more suitable to confirm or disprove the possible disagreement between TTV and RV methods.

7 DISCUSSION AND CONCLUSIONS

The apparent disagreement in the distribution of planets in the mass-radius diagram between masses obtained from TTV modelling and masses derived from RVs has been a long-standing problem in the exoplanet community (see Malavolta et al. 2017a, for a review of the most interesting cases). The Kepler-9 system is the first system characterized with TTV, and the already low densities of the planets




Figure 5. Same as in Fig. 4, but for derived parameters.

as determined in the discovery paper by Holman et al. (2010) have been further decreased in subsequent analysis by Borsato et al. (2014). Consequently, we selected this system as a proxy to compare RV and TTV mass measurements.

A differential comparison between the predicted RVs from H10 and B14 allowed us to set up the best observational strategy to be carried out with HARPS-N in order to independently confirm the true nature of the Kepler-9 planets and to overcome the fact that the faintness of the star would make a 5σ detection unfeasible. Additionally, we performed an independent analysis of the full *Kepler* light curve in order to compare the observed RVs with the prediction of the most precise orbital model that could be obtained from TTVs. In our analysis, we also included refined stellar parameters that take into account the precise parallax measurement provided by *Gaia* DR2, and both literature and newly derived photospheric parameters for the star.


The fit of the whole *Kepler* data set confirmed the masses, and hence the densities (see Table 3), originally found by B14. Our mass values are also well consistent with those in the recent work by Freudenthal et al. (2018), meaning that with a dynamical model we have been able to obtain masses with the same precision as the more computationally expensive photodynamical model, even when using space-borne photometry alone. Our analysis places Kepler-9b and c in the mass–radius region of super-Neptunes/inflated sub-Saturns together with other planets recently discovered that also have precise RV-derived masses, such as WASP-139b (Hellier et al. 2017), K2-24c (Dai et al. 2016), K2-39b (Van Eylen et al. 2016), and HATS-8b (Bayliss et al. 2015), as can be seen in Fig. 10.

The observed HARPS-N RVs noticeably agree with the model of this paper (and B14) with a χ_r^2 of 1.71, but are not consistent with the solution from H10 ($\chi_r^2 \sim 10.5$). The high value of the

Table 3. Table summarizing the dynamical fit solution. Parameter values as the MLE and HDI at 68.27 per cent equivalent. Dynamical parameters are computed at the epoch of reference $\text{BJD}_{\text{TDB}} 2455088.212$.


Parameter	Kepler-9b	Kepler-9c
<i>Fitted dynamical model</i>		
M_p/M_\star	$0.000128^{+0.000001}_{-0.000002}$	$0.000088^{+0.000001}_{-0.000001}$
P_p (day)	$19.23891^{+0.00006}_{-0.00006}$	$38.9853^{+0.0003}_{-0.0003}$
$\sqrt{e} \cos(\omega_p)$	$0.24651^{+0.0021}_{-0.0027}$	$-0.2526^{+0.0003}_{-0.0003}$
$\sqrt{e} \sin(\omega_p)$	$-0.014^{+0.002}_{-0.002}$	$0.0559^{+0.0005}_{-0.0005}$
i ($^\circ$)	$88.982^{+0.007}_{-0.005}$	–
$i \cos(\Omega_p)$	–	$-89.172^{+0.002}_{-0.005}$
$i \sin(\Omega_p)$	–	$1.7^{+0.2}_{-0.5}$
λ_p ($^\circ$) ^(a)	$179.49^{+0.15}_{-0.11}$	$293.9^{+0.3}_{-0.1}$
<i>Derived dynamical model</i>		
M_p (M_\oplus)	$43.4^{+1.6}_{-2.0}$	$29.9^{+1.1}_{-1.3}$
ρ_p (g cm^{-3})	$0.42^{+0.06}_{-0.09}$	$0.31^{+0.05}_{-0.06}$
e_p	$0.0609^{+0.0010}_{-0.0013}$	$0.06691^{+0.00010}_{-0.00012}$
ω_p ($^\circ$)	$357.0^{+0.5}_{-0.4}$	$167.5^{+0.1}_{-0.1}$
\mathcal{M}_p ($^\circ$)	$2.6^{+0.5}_{-0.6}$	$307.4^{+0.1}_{-0.1}$
i ($^\circ$)	–	$89.188^{+0.005}_{-0.006}$
Ω_p ($^\circ$)	180. (fixed)	$179.0^{+0.3}_{-0.1}$
dynamical model χ_r^2 (dof = 230)	1.16	–

Note. ^(a) λ_p is the mean longitude of the planet, defined as $\lambda_p = \Omega_p + \omega_p + \mathcal{M}_p$.

**Figure 6.** O–C (upper left panel) where O and C mean observed and calculated transit times, respectively, and duration (upper right panel) plots with residuals (lower panels) for the best-fitting solution for Kepler-9b. The grey lines represent 1000 realizations obtained by randomly selecting sets of parameters from the posterior distributions.

masses found by H10 is ascribable to the very short baseline of the photometric data and due to a possible underestimation of the HIRES RV uncertainties.

The *Kepler* mission has shown the power of the TTV method to determine the planetary nature of candidates in multiple systems and to characterize the orbital and physical parameters of the

**Figure 7.** Same as Fig. 6, but for Kepler-9c.

exoplanets. The lack of bright stars hosting multiple-planet systems showing TTV in the *Kepler* field did not allow for high-precision RV observation in order to confirm or rule out the mass determination discrepancy. With the advent of the *TESS* (Ricker et al. 2014) and *PLATO* (Rauer et al. 2014) missions, we will be able to measure the masses of many planets around bright stars with both the TTV and the RV methods, allowing us to further investigate the level of consistency in planet parameters between the two methods, or the presence of any bias in the solutions coming from the two techniques.




Figure 8. Kepler-9 RV plot (observed RVs as black-open circles) for best-fitting solution (blue-filled circles). The RV model of the best-fitting solution covering the full integration time has been plotted as a black-dashed line. We plotted 1000 realizations (as grey lines) of the model from the posterior distribution, but they all lie too close to the best-fitting RV model to be distinguishable from it.




Figure 9. Histograms of the χ_r^2 from the RV of the 10000 simulations for the three different solutions: **H10** from Holman et al. (2010), **B14** from Borsato et al. (2014), and from the posterior distribution of this work.

ACKNOWLEDGEMENTS

Based on observations made with the Italian Telescopio Nazionale Galileo (TNG) operated on the island of La Palma by the Fundacion Galileo Galilei of the INAF (Istituto Nazionale di Astrofisica) at the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofisica de Canarias. The HARPS-N project has been funded




Figure 10. Mass–radius plot for Kepler-9b and c. Results from **H10** are included for comparisons. Planets with mass determination with uncertainties greater than 20 per cent are shaded accordingly. Planets are colour coded according to their incident flux. Data obtained from exoplanet.eu on 2018 October.

by the Prodex Program of the Swiss Space Office (SSO), the Harvard University Origins of Life Initiative (HUOLI), the Scottish Universities Physics Alliance (SUPA), the University of Geneva, the Smithsonian Astrophysical Observatory (SAO), and the Italian National Astrophysical Institute (INAF), the University of St Andrews, Queen’s University Belfast, and the University of Edinburgh. The research leading to these results received funding from the European Union Seventh Framework Programme (FP7/2007–2013) under grant agreement number 313014 (ETA-EARTH) and support from Italian Space Agency (ASI) regulated by Accordo ASI-INAF no. 2013-016-R.0 del 9 luglio 2013 e integrazione del 9 luglio 2015. This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC; <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement. LM acknowledges the support by INAF/Frontiera through the ‘Progetti Premiali’ funding scheme of the Italian Ministry of Education, University, and Research. ACC acknowledges support from the Science & Technology Facilities Council (STFC) consolidated grant number ST/R000824/1. Some of this work has been carried out within the framework of the NCCR PlanetS, supported by the Swiss National Science Foundation. XD is grateful to The Branco Weiss Fellowship – Society in Science for its financial support. CAW acknowledges support from the STFC grant ST/P000312/1. DWL acknowledges partial support from the *Kepler* mission under NASA Cooperative Agreement NNX13AB58A with the Smithsonian Astrophysical Observatory. This material is based upon work supported by the National Aeronautics and Space Administration under grant nos. NNX15AC90G and NNX17AB59G issued through the Exoplanets Research Program. This research has made use of the Exoplanet Follow-up Observation Program, NASA’s Astrophysics Data System, and the NASA Exoplanet Archive, which are operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program. Some of the data presented in

this paper were obtained from the Mikulski Archive for Space Telescopes (MAST). STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. Support for MAST for non-*HST* data is provided by the NASA Office of Space Science via grant NNX13AC07G and by other grants and contracts.

REFERENCES

- Agol E., Steffen J., Sari R., Clarkson W., 2005, *MNRAS*, 359, 567
- Barros S. C. C. et al., 2015, *MNRAS*, 454, 4267
- Bayliss D. et al., 2015, *AJ*, 150, 49
- Becker J. C., Vanderburg A., Adams F. C., Rappaport S. A., Schwengeler H. M., 2015, *ApJ*, 812, L18
- Borsato L., Marzari F., Nascimbeni V., Piotto G., Granata V., Bedin L. R., Malavolta L., 2014, *A&A*, 571, A38 (B14)
- Borucki W. J. et al., 2011, *ApJ*, 736, 19
- Buchhave L. A. et al., 2012, *Nature*, 486, 375
- Buchhave L. A. et al., 2014, *Nature*, 509, 593
- Choi J., Dotter A., Conroy C., Cantiello M., Paxton B., Johnson B. D., 2016, *ApJ*, 823, 102
- Cochran W. D. et al., 2011, *ApJS*, 197, 7
- Cutri R. M. et al., 2003, *VizieR Online Data Catalog*, 2246
- Dai F. et al., 2016, *ApJ*, 823, 115
- Dotter A., 2016, *ApJS*, 222, 8
- Dotter A., Chaboyer B., Jevremović D., Kostov V., Baron E., Ferguson J. W., 2008, *ApJS*, 178, 89
- Eastman J., Gaudi B. S., Agol E., 2013, *PASP*, 125, 83
- Feroz F., Hobson M. P., 2008, *MNRAS*, 384, 449
- Feroz F., Hobson M. P., Bridges M., 2009, *MNRAS*, 398, 1601
- Feroz F., Hobson M. P., Cameron E., Pettitt A. N., 2013, preprint ([arXiv: e-prints](https://arxiv.org/abs/1307.1701))
- Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, *PASP*, 125, 306
- Freudenthal J. et al., 2018, *A&A*, 618, A41
- Gaia Collaboration, 2016, *A&A*, 595, A1
- Gaia Collaboration, 2018, *A&A*, 616, 22
- Gelman A., Rubin D. B., 1992, *Stat. Sci.*, 7, 457
- Goodman J., Weare J., 2010, *Commun. Appl. Math. Comput. Sci.*, 5, 65
- Hellier C. et al., 2017, *MNRAS*, 465, 3693
- Holman M. J. et al., 2010, *Science*, 330, 51 (H10)
- Holman M. J., Murray N. W., 2005, *Science*, 307, 1288
- Huber D. et al., 2014, *ApJS*, 211, 2
- Jontof-Hutter D. et al., 2016, *ApJ*, 820, 39
- Kipping D. M., 2010, *MNRAS*, 407, 301
- Kipping D. M., 2013, *MNRAS*, 435, 2152
- Kreidberg L., 2015, *PASP*, 127, 1161
- Lissauer J. J. et al., 2011, *Nature*, 470, 53
- Lovis C. et al., 2011, preprint ([arXiv:e-prints](https://arxiv.org/abs/1105.3456))
- Malavolta L. et al., 2017a, *AJ*, 153, 224
- Malavolta L., Lovis C., Pepe F., Sneden C., Udry S., 2017b, *MNRAS*, 469, 3965
- Malavolta L. et al., 2018, *AJ*, 155, 107
- Masuda K., Hirano T., Taruya A., Nagasawa M., Suto Y., 2013, *ApJ*, 778, 185
- Mills S. M., Mazeh T., 2017, *ApJ*, 839, L8
- Mortier A., Sousa S. G., Adibekyan V. Z., Brandão I. M., Santos N. C., 2014, *A&A*, 572, A95
- Morton T. D., 2015, *Isochrones: Stellar Model Grid Package*. Astrophysics Source Code Library, record ascl:1503.010
- Nespral D. et al., 2017, *A&A*, 601, A128
- Osborn H. P. et al., 2017, *A&A*, 604, A19
- Pace G., 2013, *A&A*, 551, L8
- Paxton B., Bildsten L., Dotter A., Herwig F., Lesaffre P., Timmes F., 2011, *ApJS*, 192, 3
- Petigura E. A. et al., 2017, *AJ*, 154, 107
- Rauer H. et al., 2014, *Exp. Astron.*, 38, 249
- Ricker G. R. et al., 2014, in Jacobus M. O., Jr, Mark C., Giovanni G. F., Howard A. M., eds, *Proc. SPIE Conf. Ser. Vol. 9143, Space Telescopes and Instrumentation 2014: Optical, Infrared, and Millimeter Wave*. SPIE, Bellingham, p. 914320
- Skrutskie M. F. et al., 2006, *AJ*, 131, 1163
- Sneden C., 1973, *ApJ*, 184, 839
- Sousa S. G., Santos N. C., Adibekyan V., Delgado-Mena E., Israelian G., 2015, *A&A*, 577, A67
- Sozzetti A. et al., 2015, *A&A*, 575, L15
- Stassun K. G., Torres G., 2018, *ApJ*, 862, 61
- Steffen J. H., 2016, *MNRAS*, 457, 4384
- Storn R., Price K., 1997, *J. Glob. Optim.*, 11, 341
- Torres G. et al., 2011, *ApJ*, 727, 24
- Van Eylen V. et al., 2016, *AJ*, 152, 143
- von Essen C. et al., 2018, *A&A*, 615, A79
- Wang S., Addison B., Fischer D. A., Brewer J. M., Isaacson H., Howard A. W., Laughlin G., 2018, *AJ*, 155, 70
- Weiss L. M. et al., 2013, *ApJ*, 768, 14
- Weiss L. M. et al., 2017, *AJ*, 153, 15
- Weiss L. M., Marcy G. W., 2014, *ApJ*, 783, L6
- Winn J. N., 2010, in Seager S., ed., *Exoplanets*. University of Arizona Press, Tucson, AZ, p. 55

SUPPORTING INFORMATION

Supplementary data are available at [MNRAS](https://www.mnras.org) online.

Borsato_Kepler-9_transit_times_durations_TableA1.dat.

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

APPENDIX A: TIME OF TRANSIT AND DURATIONS

Central time (T_0) and duration T_{14} with corresponding error obtained from the fit of each transit, as described in Section 5.

Table A1. Transit times and durations as described in Section 5. Full table available in an electronic form.

Transit number (a)	T_0 (KBJD ^(b) _{TDB})	σ_{T_0} (min)	T_{14} (min)	$\sigma_{T_{14}}$ (min)	SC/LC
<i>Kepler-9b</i>					
0	144.24992	1.6	254.3	5.0	LC
1	163.48362	1.1	249.4	3.6	LC
3	201.95379	1.2	258.5	3.6	LC
...
<i>Kepler-9c</i>					
0	136.30647	1.2	288.2	5.2	LC
1	175.33153	1.9	289.9	4.4	LC
2	214.33540	1.1	279.2	4.6	LC
...

Notes. ^(a)Transit number computed with respect to the linear ephemeris in Table 2.

^(b)BJD_{TDB} – 2454833.0.

APPENDIX B: RV MEASUREMENTS WITH HARPS-N

RV with corresponding errors obtained with HARPS-N facility.

Table B1. HARPS-N RVs.

Time (BJD _{TDB})	RV (m s ⁻¹)	σ_{RV} (m s ⁻¹)
2456783.60953	2353.88	12.88
2456783.63028	2348.14	10.00
2456798.63783	2338.88	8.42
2456798.66056	2356.35	8.23
2456801.65029	2344.83	9.25
2456801.66955	2333.04	10.26
2456813.53960	2340.04	7.38
2456813.56188	2335.65	7.33
2456815.54510	2331.71	9.40
2456815.56727	2334.02	8.52
2456829.53712	2337.37	10.28
2456829.55869	2326.08	10.58
2456831.58087	2325.38	9.56
2456831.60226	2336.21	11.00
2456834.50224	2311.57	15.00
2456845.67515	2319.74	10.48
2456845.69621	2315.50	10.85
2456848.44839	2342.51	15.55
2456848.46990	2318.14	16.26
2456851.45439	2345.14	9.33
2456851.47538	2326.24	9.37
2456853.45667	2323.64	16.36
2456853.47773	2335.83	18.50
2456922.38581	2314.59	13.09
2456922.40526	2315.97	16.20
2456925.37421	2314.17	19.65
2456934.36631	2338.37	11.78
2456934.38747	2337.96	11.48
2456936.46863	2365.09	10.49
2456936.49021	2348.56	10.31

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.