



Publication Year	2020
Acceptance in OA@INAF	2022-02-17T11:49:00Z
Title	ELT-HIRES, the high resolution spectrograph for the ELT: the Phase A study and the path to construction
Authors	Marconi, Alessandro; Abreu, M.; Adibekyan, V.; Aliverti, Matteo; Allende Prieto, C.; et al.
DOI	10.1117/12.2560489
Handle	http://hdl.handle.net/20.500.12386/31409
Series	PROCEEDINGS OF SPIE
Number	11447

PROCEEDINGS OF SPIE

[SPIDigitalLibrary.org/conference-proceedings-of-spie](https://spiedigitallibrary.org/conference-proceedings-of-spie)

ELT-HIRES, the high resolution spectrograph for the ELT: the Phase A study and the path to construction

Marconi , A., Abreu, M., Adibekyan, V., Aliverti, M., Allende Prieto, C., et al.

A. Marconi , M. Abreu, V. Adibekyan, M. Aliverti, C. Allende Prieto, P. Amado, M. Amate, E. Artigau, S. Augusto, S. Barros, S. Becerril, B. Benneke, E. Bergin, P. Berio, N. Bezawada, I. Boisse, X. Bonfils, F. Bouchy, C. Broeg, A. Cabral, R. Calvo-Ortega, B. L. Canto Martins, B. Chazelas, A. Chiavassa, L. Christensen, R. Cirami, I. Coretti, S. Covino, G. Cresci, S. Cristiani, V. Cunha Parro, G. Cupani, V. D'Odorico, I. de Castro Leão, J. R. de Medeiros, M. de Souza, P. Di Marcantonio, I. Di Varano, R. Doyon, H. Drass, P. Figueira, A. Fragoso, J. Fynbo, E. Gallo, M. Genoni, J. González Hernández, R. Gratton, M. Haehnelt, C. Hansen, J. Hlavacek-Larrondo, I. Hughes, P. Huke, A. Humphrey, H. Kjeldsen, A. Korn, D. Kouach, M. Landoni, J. Liske, C. Lovis, D. Lunney, R. Maiolino, L. Malo, T. Marquart, C. J. A. P. Martins, P. Maslowski, E. Mason, G. Micela, P. Molaro, J. Monnier, M. Monteiro, C. Mordasini, T. Morris, A. Mucciarelli, G. Murray, A. Niedzielski, E. Niemczura, B. Nisini, N. Nunes, E. Oliva, L. Origlia, E. Pallé, G. Pariani, P. Parr-Burman, L. Pasquini, J. Peñate, F. Pepe, G. Pietrzynski, E. Pinna, N. Piskunov, A. Pollo, J. Rasilla, R. Rebolo, P. Rees II, A. Reiners, M. Riva, D. Romano, S. Rousseau, N. Sanna, M. Sarajlic, T.-C. Shen, F. Sortino, D. Sosnowska, S. Sousa, E. Stempels, K. Strassmeier, F. Tenegi, A. Tozzi, S. Udry, L. Valenziano, L. Vanzì, M. Weber, M. Woche, M. Xompero, E. Zackrisson, M. R. Zapatero Osorio, "ELT-HIRES, the high resolution spectrograph for the ELT: the Phase A study and the path to construction," Proc. SPIE 11447, Ground-based and Airborne Instrumentation for Astronomy VIII, 1144726 (13 December 2020); doi: 10.1117/12.2560489

SPIE.

Event: SPIE Astronomical Telescopes + Instrumentation, 2020, Online Only

ELT-HIRES, the high resolution spectrograph for the ELT: Phase A study and path to construction

A. Marconi^{1,2}, M. Abreu^{3,4}, V. Adibekyan^{5,6}, M. Aliverti⁷, C. Allende Prieto^{8,9}, P. Amado¹⁰, M. Amate⁸, E. Artigau¹¹, S. Augusto¹², S. Barros^{5,6}, S. Becerril¹⁰, B. Benneke¹¹, E. Bergin¹³, P. Berio¹⁴, N. Bezawada¹⁵, I. Boisse¹⁶, X. Bonfils¹⁷, F. Bouchy^{18,16}, C. Broeg¹⁹, A. Cabral^{3,4}, R. Calvo-Ortega¹⁰, B. L. Canto Martins²⁰, B. Chazelas¹⁸, A. Chiavassa¹⁴, L. Christensen²¹, R. Cirami²², I. Coretti²², S. Covino⁷, G. Cresci², S. Cristiani²², V. Cunha Parro¹², G. Cupani²², V. D'Odorico²², I. de Castro Leão²⁰, J. R. de Medeiros²⁰, M. de Souza¹², P. Di Marcantonio²², I. Di Varano²³, R. Doyon¹¹, H. Drass^{24,25}, P. Figueira^{26,5}, A. Fragoso⁸, J. Fynbo²¹, E. Gallo¹³, M. Genoni⁷, J. González Hernández^{8,9}, R. Gratton²⁷, M. Haehnelt²⁸, C. Hansen²⁹, J. Hlavacek-Larrondo³⁰, I. Hughes¹⁸, P. Huke³¹, A. Humphrey⁵, H. Kjeldsen³², A. Korn³³, D. Kouach³⁴, M. Landoni⁷, J. Liske³⁵, C. Lovis¹⁸, D. Lunney³⁶, R. Maiolino³⁷, L. Malo¹¹, T. Marquart³³, C. J. A. P. Martins^{5,38}, P. Maslowski³⁹, E. Mason²², G. Micela⁴⁰, P. Molaro²², J. Monnier¹³, M. Monteiro⁵, C. Mordasini¹⁹, T. Morris⁴¹, A. Mucciarelli⁴², G. Murray⁴¹, A. Niedzielski⁴³, E. Niemczura⁴⁴, B. Nisini⁴⁵, N. Nunes^{3,4}, E. Oliva², L. Origlia⁴⁶, E. Pallé^{8,9}, G. Pariani⁷, P. Parr-Burman³⁶, L. Pasquini¹⁵, J. Peñate⁸, F. Pepe¹⁸, G. Pietrzynski⁴⁷, E. Pinna², N. Piskunov³³, A. Pollo⁴⁸, J. Rasilla⁸, R. Rebolo^{8,9}, P. Rees³⁶, A. Reiners³¹, M. Riva⁷, D. Romano⁴⁶, S. Rousseau¹⁴, N. Sanna², N. Santos^{5,6}, M. Sarajlic¹⁹, T.-C. Shen²⁴, F. Sortino⁴⁶, D. Sosnowska¹⁸, S. Sousa⁵, E. Stempels³³, K. Strassmeier²³, F. Tenegi⁸, A. Tozzi², S. Udry¹⁸, L. Valenziano⁴⁶, L. Vanzini²⁴, M. Weber²³, M. Woche²³, M. Xompero², E. Zackrisson³³, and M. R. Zapatero Osorio⁴⁹

¹Dip. di Fisica e Astronomia, Univ. di Firenze, via G. Sansone 1, I-50019, Sesto F.no (FI), Italy

²INAF-Osservatorio Astrofisico di Arcetri, Largo E. Fermi 2, I-50125, Firenze, Italy

³Instituto de Astrofísica e Ciências do Espaço, Universidade de Lisboa, Campo Grande 1749-016 Lisboa Portugal

⁴Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Campo Grande 1749-016 Lisboa Portugal

⁵Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, CAUP, Rua das Estrelas, PT4150-762 Porto, Portugal

⁶Departamento de Física e Astronomia, Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre, 4169-007 Porto, Portugal

⁷INAF-Osservatorio Astronomico di Brera, Via Bianchi 46, I-23807 Merate, Italy

⁸Instituto de Astrofísica de Canarias (IAC), C/ Via Lactea, s/n E-38205, La Laguna, Tenerife, Spain

⁹Universidad de La Laguna (ULL), Departamento de Astrofísica, E-38206 La Laguna, Tenerife, Spain

¹⁰Instituto de Astrofísica de Andalucía-CSIC Glorieta de la Astronomía s/n, 18008, Granada, Spain

¹¹Institut de Recherche sur les Exoplanètes and Observatoire du Mont-Mégantic, département de physique, Université de Montréal, CP 6128 Succ. Centre-ville, H3C 3J7, Montréal, QC, Canada

- ¹²IMT - Instituto Mauá de Tecnologia, Praça Mauá, 1 - Mauá, São Caetano do Sul - SP
Brazil, 09580-900
- ¹³Department of Astronomy, University of Michigan, 323 West Hall, 1085 S. University
Avenue, Ann Arbor, MI 48109, USA
- ¹⁴Laboratoire Lagrange, Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, Blvd
de L'Observatoire CS34229,06004 Nice Cedex 4, France
- ¹⁵European Southern Observatory, Karl-Schwarzschild-Straße 2, 85748 Garching bei München,
Germany
- ¹⁶Aix Marseille Univ, CNRS, CNES, LAM, Marseille, France
- ¹⁷Université Grenoble Alpes, CNRS, IPAG, 38000, Grenoble, France
- ¹⁸Département d'Astronomie, Université de Geneve, Chemin des Maillettes 51, Sauverny,
CH-1290 Versoix, Switzerland
- ¹⁹Physikalisches Institut, University of Bern, Gesellschaftsstrasse 6, CH-3012 Bern, Switzerland
- ²⁰Board of Observational Astronomy, Federal University of Rio Grande do Norte, Campus
Universitário 59078-970, Natal RN, Brasil
- ²¹Cosmic Dawn Center, Niels Bohr Institute, Copenhagen University, DK-2100 Copenhagen O,
Denmark
- ²²INAF Osservatorio Astronomico di Trieste, Via Giambattista Tiepolo 11, 34131 - Trieste
Italy
- ²³Leibniz Institute for Astrophysics Potsdam (AIP), An der Sternwarte 16, D-14482 Potsdam,
Germany
- ²⁴Centro de Astro Ingenieria, Pontificia Universidad Catolica de Chile, Avda. Libertador
Bernardo O'Higgins 340 - Santiago de Chile
- ²⁵Millennium Institute of Astrophysics, Santiago, Chile
- ²⁶European Southern Observatory, Alonso de Cordova 3107, Vitacura, Santiago, Chile
- ²⁷INAF-Osservatorio Astronomico di Padova, vicolo dell'Osservatorio 5, 35122 Padova, Italy
- ²⁸Kavli Institute for Cosmology and Institute of Astronomy, Madingley Road, Cambridge CB3
0HA, UK
- ²⁹Max Planck Institute for Astronomy, MPIA, Koenigstuhl 17, 69117 Heidelberg, Germany
- ³⁰Département de physique, Université de Montréal, CP 6128 Succ. Centre-ville, H3C 3J7,
Montréal, QC, Canada.
- ³¹Institute for Astrophysics University of Göttingen, Friedrich-Hund-Platz 1 37077 Göttingen,
Germany
- ³²Department of Physics and Astronomy, Ny Munkegade 120, building 1520, 527, 8000 Aarhus
C, Denmark
- ³³Division of Astronomy and Space Physics, Department of Physics and Astronomy, Uppsala
University, Box 516, S-75120 Uppsala, Sweden
- ³⁴CNRS, OMP, Université de Toulouse, 14 Avenue Belin, F-31400 Toulouse, France
- ³⁵Hamburger Sternwarte, Universität Hamburg, Gojenbergsweg 112, D-21029 Hamburg
- ³⁶UK Astronomy Technology Centre (part of the Science and Technology Facilities Council),
Blackford Hill, Edinburgh, EH9 3HJ, UK
- ³⁷Cavendish Laboratory, Univ. of Cambridge, JJ Thomson Avenue, Cambridge CB3 0HE, UK
- ³⁸Centro de Astrofísica da Universidade do Porto, Rua das Estrelas, PT4150-762 Porto,
Portugal
- ³⁹Institute of Physics, Nicolaus Copernicus University in Torun, Gagarina 11, 87-100 Torun,

Poland

⁴⁰INAF-Osservatorio Astronomico di Palermo, Piazza del Parlamento 1, 90134 Palermo, Italy

⁴¹Centre for Advanced Instrumentation, Department of Physics, Durham University, South Road, Durham, DH1 3LE, UK

⁴²Dipartimento di Fisica e Astronomia, Università degli Studi di Bologna, Via Gobetti 93/2, I-40129 Bologna, Italy

⁴³Institute of Astronomy, Nicolaus Copernicus University in Torun, Gagarina 11, 87-100 Torun, Poland

⁴⁴Astronomical Institute, University of Wrocław, ul. Kopernika 11, 51-622 Wrocław, Poland

⁴⁵INAF-Osservatorio Astronomico di Roma, via Frascati 33, 00078 Roma, Italy

⁴⁶INAF-Osservatorio di Astrofisica e Scienze dello Spazio di Bologna, via P. Gobetti 93/3, 40129 Bologna, Italy

⁴⁷Nicolaus Copernicus Astronomical Center of the Polish Academy of Sciences, ul. Bartycka 18, 00-716 Warszawa, Poland

⁴⁸Jagiellonian University and National Centre for Nuclear Research, ul. Hoza 69, 00-681 Warszawa, Poland

⁴⁹Centro de Astrobiología (CSIC-INTA), Carretera de Ajalvir km 4, 28850 Torrejón de Ardoz, Madrid, Spain

ABSTRACT

HIRES is the high-resolution spectrograph of the European Extremely Large Telescope at optical and near-infrared wavelengths. It consists of three fibre-fed spectrographs providing a wavelength coverage of 0.4-1.8 μm (goal 0.35-2.4 μm) at a spectral resolution of $\sim 100,000$. The fibre-feeding allows HIRES to have several, interchangeable observing modes including a SCAO module and a small diffraction-limited IFU in the NIR. Therefore, it will be able to operate both in seeing- and diffraction-limited modes. Its modularity will ensure that HIRES can be placed entirely on the Nasmyth platform, if enough mass and volume is available, or part on the Nasmyth and part in the Coudé room. ELT-HIRES has a wide range of science cases spanning nearly all areas of research in astrophysics and even fundamental physics. Among the top science cases there are the detection of biosignatures from exoplanet atmospheres, finding the fingerprints of the first generation of stars (PopIII), tests on the stability of Nature's fundamental couplings, and the direct detection of the cosmic acceleration. The HIRES consortium is composed of more than 30 institutes from 14 countries, forming a team of more than 200 scientists and engineers.

Keywords: EXTREMELY LARGE TELESCOPES, HIGH RESOLUTION SPECTROSCOPY, EXOPLANETS, STARS AND PLANETS FORMATION, PHYSICS AND EVOLUTION OF STARS, PHYSICS AND EVOLUTION OF GALAXIES, COSMOLOGY, FUNDAMENTAL PHYSICS

1. INTRODUCTION

The European Extremely Large Telescope (ELT) will be the largest ground-based telescope at visible and infrared wavelengths and its flagship science cases are the detection of life signatures in Earth-like exoplanets and the direct detection of the cosmic expansion re-acceleration: it is no coincidence that both science cases require observations with a high-resolution spectrograph.

High-resolution spectroscopy is a truly interdisciplinary tool and, during the past decades, has enabled some of the most extraordinary discoveries spanning all fields of Astrophysics, from Planetary Sciences to Cosmology. Indeed, high-resolution spectrometers have allowed astronomers to go beyond the classical domain of astrophysics and to address some of the fundamental questions of Physics. ESO has a long and successful tradition in high

Send correspondence to A. Marconi (alessandro.marconi@inaf.it)

resolution spectroscopy as demonstrated by the exquisite suite of medium-high resolution spectrographs offered to the community of Member States. UVES, FLAMES, CRIRES, Xshooter and HARPS have enabled European teams to lead in many areas of research. ESPRESSO has now joined this suite of very successful high-resolution spectrographs, fulfilling the promise of truly revolutionising some of these research areas. The scientific interest and high productivity of high-resolution spectroscopy is reflected by the fact that more than 30% of ESO publications are based on its high-resolution spectrographs.

However, it is becoming increasingly clear that, in most areas of research, high-resolution spectroscopy has reached the “photon-starved” regime with 8-10m class telescopes. Despite major progress on the instrumentation front, further major advances in these fields desperately require a larger photon collecting area. Due to its inherently “photon-starved” nature, amongst the various astronomical observing techniques, high-resolution spectroscopy requires the collecting area of Extremely Large Telescopes.

When defining the ELT instrumentation, ESO commissioned two phase-A studies for high-resolution spectrographs, CODEX¹ and SIMPLE² which were started in 2007 and completed in 2010. These studies demonstrated the importance of optical and near-IR high-resolution spectroscopy at the ELT and ESO thus decided to include a High-REsolution Spectrograph (HIRES) in the ELT instrumentation roadmap. Soon after conclusion of the respective phase A studies the CODEX and SIMPLE consortia realized the great scientific importance of covering the optical and near-infrared spectral ranges simultaneously. This marked the birth of the HIRES that started developing the concept of an X-Shooter-like spectrograph, but with higher resolution, capable of providing $R \sim 100,000$ over the full optical and near-infrared wavelengths range. Following a community workshop in September 2012 the HIRES Initiative has prepared a White Paper summarizing a wide range of science cases proposed by the community (Maiolino et al.³) and also prepared a Blue Book with a preliminary technical instrument concept.

With the start of construction of the ELT, the HIRES Initiative became a Consortium, recruiting additional institutes which expressed their interest in HIRES. ESO commissioned the consortium to perform a Phase A study which started in March 2016 and successfully concluded in May 2018. Since new Institutes from USA and Canada joined the HIRES consortium, and many activities in preparation of the start of construction were performed.

The HIRES Consortium is now composed of institutes from Brazil, Canada, Chile, Denmark, France, Germany, Italy, Poland, Portugal, Spain, Sweden, Switzerland, United Kingdom and USA. The full list of institutes is presented in table 1 together with the coordinator of each country.

2. SCIENCE GOALS

2.1 Exoplanets and protoplanetary disks

The study of exoplanet atmospheres for a wide range of planetary objects, from gas giants to rocky planets, and from hot to temperate planets, is a primary objective in the field for the next decade. In particular, the detection of components such as molecular oxygen, water and methane in Earth- or super-Earth sized planets is considered to be truly transformational, as they may be regarded as signature of habitability or even signatures of life. Simulations of HIRES observations have been performed by Snellen et al.^{4,5} and Hakwer & Parry.⁶

HIRES will be able to probe the atmospheres in transmission during the transit of an exoplanet in front of its host star. As an example, it will be possible to detect CO₂ absorption in Trappist-1 b with a S/N of 6 in 4 transits of the planet, while O₂ absorption at 0.75 μm can be detected in only 25 transits of the planet, i.e. less than 30 hours of observations. HIRES will also be able to directly probe exoplanets, by spatially resolving them from their host star, focussing on their reflected star light and taking advantage of the angular resolution of the ELT with AO-assisted observations. For example, it will be possible to detect the Proxima-Cen b planet in 4 nights of integration with a S/N of 8 with a relatively simple system of single conjugate adaptive optics (SCAO), similar to that used by other ELT first-light instruments. Figure 1, left, shows that HIRES will be able to detect O₂ from a Proxima-b like exoplanet in 70 h of integration.

Protoplanetary disks are a natural outcome of angular momentum conservation in star formation and are ubiquitous around young, forming stars. HIRES will be able to determine the properties of the gas in the inner




Figure 1. Hires science highlights. Left: Cross Correlation Signal indicating the clear detection of O2 in as Proxima-b like exoplanet in 70h of total integration (adapted from Fig. 4 of Hawker & Parry 2019). Right: Observations of a $z=9$ quasar with $H_{AB}=22$ and a total integration time of 10h showing Hires capability of distinguishing IGM enrichment by normal SNIi supernovae or by low mass and pair instability Supernovae from Pop III Stars (simulations by the Hires Science Advisory Team).

star-disk region, where different competing mechanisms of disk gas dispersal are at play. This will constrain on one side on the mechanisms through which the forming star acquires mass and removes the angular momentum, and on the other side the initial condition for planets formation.

2.2 Stars and Stellar Populations

The vast light-collecting power of the ELT will enable detailed high-resolution spectroscopy of individual stars, and in particular very faint red dwarfs and distant red giants in nearby galaxies, for which Hires will be able to provide tight constraints for the atmospheric parameters. These constraints will be extremely important to characterize the stellar hosts of exoplanets. Hires will also expand our horizon by measuring the heavy-elements abundances of the most primitive stars (low mass, low metallicity) in our galaxy and its satellites helping us to understand what is the lowest metallicity for which gas can collapse to form low-mass stars, and what are the nature and yields of the very first generation of stars and their supernovae. Last, but not least, the combination of very high spectral resolving power and diffraction-limited angular resolution makes the ELT a unique resource for deepening our understanding of the physics of stellar atmospheres and nucleosynthesis processes, by allowing to spectroscopically resolve the effects of surface convection and to measure isotopic abundances of atomic species.

2.3 Galaxy Formation and evolution and the intergalactic medium

The detection of the first generation of stars (so called Pop III stars) and the observational characterization of their properties is one of the main objectives of extragalactic astrophysics. Proto-galaxies hosting Pop III stars are expected to be too faint for direct detection, even with JWST. However, the signature of Pop III stars can be detected through their nucleosynthetic yields which can be potentially observed in the abundance patterns of very metal-poor absorption systems in the high-resolution, wide-range spectra of bright high-redshift sources provided by Hires in the NIR (Figure 1, right). The direct detection and characterization of the beginning of the reionization epoch is another very important goal in the study of galaxy formation. This process is believed to have been dominated by ultraviolet photons from the first generations of galaxies, most of which are too faint to be observed directly even with JWST. By targeting bright quasars at high redshift as background continuum sources, Hires will be able to study both transmission features in the Lyman- α forest and metal absorption lines associated with these reionization-epoch sources, constraining the patchiness of the reionization process, the properties of the ultraviolet background radiation and the chemical enrichment of the IGM in this epoch.

2.4 Cosmology and Fundamental Physics

The observational evidence for the acceleration of the expansion of the universe and the tensions that have been highlighted by different cosmological probes have shown that our canonical theories of cosmology of fundamental physics may be incomplete (and possibly incorrect), and that there might be unknown physics yet to

be discovered. HIRES will allow to search for, identify and ultimately characterize any new physics through several different but fundamentally inter-related observations which will enable a unique set of tests of the current cosmological paradigm. HIRES will be able to constrain the variation of fundamental physical constants like the fine-structure constant α and proton-electron mass ratio μ with the advantage, compared to laboratory measurements, of exploring variations over 12 Gyr timescales and 15 Gpc spatial scales. A detection of varying fundamental constants would be revolutionary: it would automatically prove that the Einstein Equivalence Principle is violated (i.e. gravity is not purely geometry), and that there is a fifth force. HIRES will enable a test of the CMB temperature-redshift relation, $T(z) = T_0(1+z)$, which is a robust prediction of standard cosmology but that must be directly verified by measurements. A departure from this relation can in turn reveal a violation of the hypothesis of local position invariance (and thus of the equivalence principle) or that the number of photons is not conserved. HIRES measurement will greatly improve on the existing constraints on $T(z)$ compared to existing data. The redshifts of cosmologically distant objects drift slowly with time (the so-called Sandage effect). A redshift drift measurement is fundamentally different from all other cosmological observations and can provide a direct detection of cosmic re-acceleration, thus undoubtedly confirming cosmic acceleration, the existence of dark energy and potentially provide evidence for new physics. HIRES will be capable of detecting the redshift drift in the Ly α forests of the brightest currently known QSOs (~ 6 cm/s/decade at $z = 4$ for a Planck-like standard cosmology). The ELT may thus become the first facility ever to watch the Universe change in "real time".

2.5 Science Priorities

These are just a few of the many science cases that can be addressed, a collection of many of these can be found in the community white paper.³ However, in order to define the instrument baseline design a prioritization of the science cases was performed by the HIRES Science Advisory Team following criteria of scientific impact (transformational versus incremental), feasibility and competitiveness. Then, if the TLR's of the top priority science cases were enabling other science cases, the latter were not considered any further in the subsequent prioritization, as considered accomplished together with the top priority science cases. The top science priorities and associated requirements are listed below:

1. Exoplanet atmospheres in transmission, requiring a spectral resolution of at least 100,000, a wavelength coverage of at least 0.50-1.80 μm and a wavelength calibration accuracy of 1 m/s. The implementation of the above TLRs would automatically enable the following science cases:
 - reionization of the universe,
 - the characterization of cool stars,
 - the detection and investigation of near pristine gas,
 - the study of Extragalactic transients.
2. Variation of the Fundamental Constants of Physics, requiring an extension to 0.37 μm in addition to the TLRs of priority 1. These extension towards the blue would also automatically enable to investigate:
 - the cosmic variation of the CMB temperature,
 - the determination of the deuterium abundance,
 - the investigation and characterization of primitive stars.

At $\lambda < 0.40\mu\text{m}$ the throughput of the ELT is expected to be low as a consequence of the planned coating. However, even in the range 0.37-0.40 μm the system is expected to outperform ESPRESSO at the VLT, and new coating is under study by ESO and may be available a few years after first light.

3. Detection of exoplanet atmospheres in reflection, requiring, on top of the TLRs of priority 1, the addition of an Adaptive Optics (SCAO) system and an Integral Field Unit. Reflected-light spectra allow tracing atmospheric emission from lower altitudes on the dayside of the exoplanet. These additional TLRs would automatically enable also the following cases:

- Planet formation in protoplanetary disks,
 - Characterization of stellar atmospheres,
 - Search of low mass Black Holes.
4. Sandage test. Its additional TLRs, are a wavelength range of 0.40-0.67 μm and a stability of 2 cm/s, enabling also:
- the mass determination of Earth-like exoplanets
 - radial velocity searches and mass determinations for exoplanets around M-dwarf stars.




Figure 2. HIRES architectural design, outlining the instrument subsystems: Front End (seeing-limited and AO assisted with SCAO unit), Fibre Link, Calibration Unit, VIS-Blue, VIS-Red and NIR (cold spectrograph).

3. INSTRUMENT CONCEPT

Following phase A and further studies before the start of construction, the HIRES baseline design is that of a modular instrument consisting of three fibre-fed cross dispersed echelle spectrographs VIS-BLUE (UBV), VIS-RED (RIZ) and NIR (YJH), providing a simultaneous spectral range of 0.4-1.8 μm (goal 0.35-2.4 μm) at a resolution of 100,000. The fibre feeding allows several, interchangeable, observing modes ensuring maximization of either accuracy, throughput or spatially resolved information. Together with the SCAO module, the proposed baseline design capable of fulfilling the requirements of the 4 top science cases.

The baseline design is summarized below but several alternatives have been evaluated during the Phase A study. Also, several add-ons made possible by the modular nature of the instrument have been considered (e.g. a polarimetric module in the intermediate focus, or a wavelength extension out to the K-band 2.0-2.4 μm). The overall concept is summarized in Figure 2: in the Front End the light from the telescope is split, via dichroics, into 3 wavelength channels. Each wavelength channel interfaces with several fibre bundles that feed the corresponding spectrograph module. Each fibre-bundle corresponds to an observing mode and all together they constitute the Fibre Link. All spectrographs, VIS-BLUE, VIS-RED and NIR, have a fixed configuration, i.e. no moving parts, allowing to fulfil the requirements on stability. They include a series of parallel entrance slits consisting of linear micro-lens arrays each glued to the fibre bundles. The split in wavelengths between the spectrographs is influenced, among other parameters by the optical throughput of the different types of fibres available on the market; therefore, the different modules can be positioned at different distances from the focal

plane of the telescope. The whole instrument should be placed on the Nasmyth platform, if enough volume and mass is available. If necessary, the fibre feeding allows the VIS-RED and NIR modules to be placed in the Coudé Room, which can also host the Calibration Unit.

The total estimated cost of the instrument has been estimated to be around 35 millions of Euros, with over 550 FTEs required for the duration of the project.




Figure 3. Hires limiting magnitudes obtained from the ETC for different S/N ratios (30 – top and 100 – bottom), compact and extended sources (left and right). Observations are in seeing-limited mode with $R = 100,000$ a total exposure time of 1800s.

4. PERFORMANCE

The Exposure Time Calculator, regularly updated to take into account modifications in the design, is maintained by INAF-Arcetri and can be run at the <http://hires.inaf.it/etc.html> web link. This ETC can compute the limiting magnitude achievable at a given wavelength, in a given exposure time and at a given signal to noise ratio or it can compute the signal to noise ratio achievable at a given wavelength, in a given exposure time and at a given magnitude. Hires expected performances computed with the ETC are summarized in Figure 3. Figure 4 uses simulations performed by the Science Team for the extragalactic science case to show the improvement of the combination of ELT and Hires with respect to existing, lower resolution instruments, like X-Shooter and UVES. Although Hires has a higher resolution and smaller spectral channels, it is able to reach much higher S/N in a much shorter time.

5. CONSORTIUM ORGANIZATION AND PATH TO CONSTRUCTION

The consortium is currently organized as shown in Figure 5. The Consortium is led by the PI, who is the point of contact with ESO, and by an Executive Board composed by the coordinators of each Country. The Project Office is composed of Project Scientist, Project Manager, Instrument Scientist and System Engineer.

The Project Scientist leads the Science team which is composed of a science team at large and of a core science team. The science team at large is composed of 4 working groups, each with a coordinator and a deputy:




Figure 4. Left: simulated observations using VLT+X-shooter (top) and ELT-HIRES (bottom) of a $z=7$ source $J_{AB} = 20.2$. Signal-to-noise ratios per spectral channel have been calculated assuming a 25 (5) hour integration with Xshooter (HIRES). The top spectra in both figures are the adopted models convolved with the instrument spectral resolution. Right: simulated observations using VLT+X-shooter (top), VLT+UVES (middle) and ELT-HIRES (bottom) of the region below Lyman- α for a $z \approx 6$ quasars at $m_{AB} \leq 21$ mag.

Exoplanets and Circumstellar Disks (Coordinator E. Palle, Deputy C. Lovis), Stars and Stellar Populations (C. Allende, A. Korn), Formation and Evolution of Galaxies and Intergalactic Medium (V. D’Odorico, E. Zackrisson), Cosmology and Fundamental Physics (J. Liske, C. Martins). Each working group is composed of about 15 members beyond coordinator and deputy. The PS, as chair, the coordinators and deputies of each working group constitute the core science team.

The Project Manager leads the Project Management team and is responsible for managing the project.

The System Engineer leads the System Team which includes the managers of each of the major sub-systems and the system architects.

Finally, figure 5 also shows the countries which are leading the effort for each of these major subsystems, as well as the major contributors for each work package.

The Consortium is currently awaiting the approval by ESO Council to start the construction phase, which is expected for the second half of 2021. With an expected duration of construction of about 8 years, the instrument should start commissioning at the telescope in 2029/2030.

6. CONCLUSIONS

The HIRES baseline design is that of three ultra-stable and modular fibre-fed cross dispersed echelle spectrographs providing a simultaneous spectral coverage of 0.4-1.8 μm (goal 0.35-1.8 μm) at a resolution of 100,000 with several, interchangeable, observing modes ensuring maximization of either accuracy, throughput or spatially resolved information. Overall, the studies conducted so far have shown that the HIRES baseline design can address the 4 top priority science cases, being able to provide ground-breaking science results with no obvious technical showstoppers.

The construction of HIRES includes the majority of the institutes in ESO member states with expertise in high resolution spectroscopy and will require an estimated 30 MEUR in hardware (excluding contingencies) and about 500 FTEs. Contingencies are expected to be low (5-10%) because the proposed baseline design is based on proven technical solutions and can benefit on heritage from HARPS and ESPRESSO and other previous high-resolution spectrographs, e.g. PEPSI at the 11.8m LBT, SPIRou and CARMENES. The construction will last about 8-10 years. Therefore, with Phase B starting in 2021, HIRES could be at the telescope as early as 2030.

Overall, HIRES is an instrument capable of addressing ground-breaking science cases while being almost (telescope) pupil independent, as it can operate both in seeing and diffraction limited modes; the modularity




Figure 5. Consortium organization and WBS pre-phase B

ensures flexibility during construction and the possibility to quickly adapt to new development in the technical as well as science landscape.

ACKNOWLEDGMENTS

The Italian effort for HIRES is supported by the Italian National Institute for Astrophysics (INAF). HIRES work in the UK is supported by the Science and Technology Facilities Council (STFC) at the UKATC, University of Cambridge (grants ST/S001387/1 and ST/N002873/1) and Heriot Watt University (grant ST/S001328/1), as part of the UK E-ELT Programme.

We acknowledge financial support from the Spanish Ministry of Science and Innovation (MICINN) projects AYA2017-86389-P, RYC-2013-14875, PGC2018-098153-B-C31, PID2019-109522GB-C51/52.

The German efforts for HIRES are funded by the Federal Ministry for Education and Research (BMBF). KGS thanks the BMBF-Verbundforschung for support through grants 05A17BAB and 05A2020.

This work was supported by FCT - Fundação para a Ciência e a Tecnologia through national funds and by FEDER through COMPETE2020 - Programa Operacional Competitividade e Internacionalização by these grants: UID/FIS/04434/2019; UIDB/04434/2020; UIDP/04434/2020; PTDC/FIS-AST/32113/2017 & POCI-01-0145-FEDER-032113; PTDC/FIS-AST/28953/2017 & POCI-01-0145-FEDER-028953; PTDC/FIS-AST/28987/2017 & POCI-01-0145-FEDER-028987.

Research activities of the observational astronomy board at the Federal University of Rio Grande do Norte are supported by continuous grants from the Brazilian funding agencies CNPq, FAPERN, and INCT-INEspaço. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

REFERENCES

- [1] Pasquini, L., Cristiani, S., García López, R., and et al., “Codex,” in [Ground-based and Airborne Instrumentation for Astronomy III], *Proc. SPIE* **7735**, 77352F (July 2010).

- [2] Origlia, L., Oliva, E., Maiolino, R., and et al., “SIMPLE: a high-resolution near-infrared spectrometer for the E-ELT,” in [*Ground-based and Airborne Instrumentation for Astronomy III*], *Proc. SPIE* **7735**, 77352B (July 2010).
- [3] Maiolino, R., Haehnelt, M., Murphy, M. T., and et al., “A Community Science Case for E-ELT HIRES,” *ArXiv e-prints (1310.3163)* (Oct. 2013).
- [4] Snellen, I. A. G., de Kok, R. J., le Poole, R., Brogi, M., and Birkby, J., “Finding Extraterrestrial Life Using Ground-based High-dispersion Spectroscopy,” *ApJ* **764**, 182 (Feb. 2013).
- [5] Snellen, I., de Kok, R., Birkby, J. L., Brandl, B., Brogi, M., Keller, C., Kenworthy, M., Schwarz, H., and Stuik, R., “Combining high-dispersion spectroscopy with high contrast imaging: Probing rocky planets around our nearest neighbors,” *A&A* **576**, A59 (Apr. 2015).
- [6] Hawker, G. A. and Parry, I. R., “High-resolution spectroscopy and high contrast imaging with the ELT: looking for O₂ in Proxima b,” *MNRAS* **484**, 4855–4864 (Apr. 2019).

Table 1. Consortium composition

Country	Coordinator	Consortium Members
Brazil	J. Renan de Medeiros	Núcleo de Astronomia Observacional, Universidade Federal do Rio Grande do Norte (CI); Instituto Mauá de Tecnologia
Canada	R. Doyon	Institut de Recherche sur les Exoplanètes and Observatoire du Mont-Mégantic, département de physique, Université de Montréal
Chile	L. Vanzi	Pontificia Universidad Catolica de Chile (CI); Centre of Astro Engineering, Universidad de Chile; Department of Astronomy, Universidad de Concepcion; Center of Astronomical Instrumentation, Universidad de Antofagasta
Denmark	J. Fynbo	Niels Bohr Institute, University of Copenhagen (CI); Department of Physics and Astronomy, Aarhus University
France	I. Boisse	Laboratoire d'Astrophysique de Marseille, CNRS, CNES, AMU (CI); Institut de Planétologie et d'Astrophysique de Grenoble, Université Grenoble Alpes; Laboratoire Lagrange, Observatoire de la Côte d'Azur; Observatoire de Haute Provence, CNRS, AMU, Institut Pythéas, Institut de Recherche en Astrophysique et Planetologie, Observatoire Midi-Pyrénées; Laboratoire Univers et Particules, Université de Montpellier
Germany	K: Strassmeier	Leibniz-Institut für Astrophysik Potsdam (CI); Institut für Astrophysik, Universität Göttingen; Zentrum für Astronomie Heidelberg, Landessternwarte; Thüringer Landessternwarte Tautenburg; Hamburger Sternwarte, Universität Hamburg
Italy	A. Marconi (PI)	INAF, Istituto Nazionale di Astrofisica (Lead Technical Institute)
Poland	A. Niedzielski	Faculty of Physics, Astronomy and Applied Informatics, Nicolaus Copernicus University in Torun
Portugal	N. Santos	Instituto de Astrofísica e Ciências do Espaço (IA) at Centro de Investigação em Astronomia/Astrofísica da Universidade do Porto (CI), Instituto de Astrofísica e Ciências do Espaço at Faculdade de Ciências da Universidade de Lisboa
Spain	R. Rebolo	Instituto de Astrofísica de Canarias (CI); Instituto de Astrofísica de Andalucía-CSIC; Centro de Astrobiología
Sweden	N. Piskunov	Dept. of Physics and Astronomy, Uppsala University
Switzerland	C. Lovis	Département d'Astronomie, Observatoire de Sauverny, Université de Genève (CI); Universität Bern, Physikalische Institut
United Kingdom	M. Haehnelt	Science and Technology Facilities Council (CI); Cavendish Laboratory & Institute of Astronomy, University of Cambridge; UK Astronomy Technology Centre; Institute of Photonics and Quantum Sciences, Heriot-Watt University
USA	E. Bergin	Department of Astronomy, University of Michigan