

Publication Year	2019
Acceptance in OA@INAF	2022-03-29T10:15:50Z
Title	Iqueye at the NTT : The best light curves of the optical pulsars in the Crab, LMC B0540-69 and Vela supernova remnants
Authors	Barbieri, Cesare; Barbieri, M.; Mignani, R.; Naletto, G.; ZAMPIERI, Luca
DOI	10.5281/zenodo.3245246
Handle	http://hdl.handle.net/20.500.12386/32005

Iqueye at the NTT: The best light curves of the optical pulsars in the Crab, LMC B0540-69 and Vela supernova remnants

Cesare Barbieri (University of Padova and INAF, Italy), Aleksandr Burtovoi (University of Padova and INAF) Roberto Mignani (INAF-IASF, Italy), Giampiero Naletto (University of Padova, Italy), Luca Zampieri (INAF-OAPd, Italy), Mauro Barbieri (University of Atacama, Chile),



QUANTUM OPTICS INSTRUMENTATION FOR ASTRONOMY

D. Dravins<sup>1</sup>, C. Barbieri<sup>2</sup>, V. Da Deppo<sup>3</sup>, D. Faria<sup>1</sup>, S. Fornasier<sup>2</sup>, R. A. E. Fosbury<sup>4</sup>, L. Lindegren<sup>1</sup>, G. Naletto<sup>3</sup>, R. Nilsson<sup>1</sup>, T. Occhipinti<sup>3</sup>, F. Tamburini<sup>2</sup>, H. Uthas<sup>1</sup>, L. Zampieri<sup>5</sup>

 (1) Lund Observatory, Box 43, SE-22100 Lund, Sweden
(2) Department of Astronomy, University of Padova, Vicolo dell'Osservatorio 2, IT-35122 Padova, Italy
(3) Dept. of Information Engineering, University of Padova, Via Gradenigo, 6/8, IT-35131 Padova, Italy
(4) Space Telescope-European Coordinating Facility & European Southern Observatory, Karl-Schwarzschild-Straße 2, DE-85748 Garching bei München, Germany
(5) INAF – Astronomical Observatory of Padova, Vicolo dell'Osservatorio 5, IT-35122 Padova, Italy



QuantEYE

In 2005, we completed the study of a novel instrument for the 100 m diameter Overwhelmingly Large (OWL) telescope (QuantEYE, the ESO Quantum Eye; Dravins et al. 2005). The main objective was to demonstrate the possibility to reach the picosecond time resolution (Heisenberg limit) needed to bring quantum optics concepts into the astronomical domain.

The study had two main scientific goals:

- To measure the *entropy of light* through the statistics of the photon arrival times.

- To perform optical *High Time Resolution Astrophysics* with unparalleled precision and accuracy.

## Development of AQUEYE and IQUEYE

Starting from QuantEYE, we realized two similar downscaled prototypes, Aqueye and Iqueye. Both instruments adopted the concept of *splitting the telescope entrance pupil* in four parts, each of them focused on a Single Photon Avalanche Detector (SPAD) operated in Geiger mode.



Both photometers are photon-counting nonimaging instruments, with a field of view of few arcsec, Filters can be inserted both in the primary beam before the pyramid and in each subbeam, thus providing multicolor simultaneous photometry.

## The two photometers



The second one was lqueye. designed for 4 m class telescopes, mounted at the 3.6 m ESO NTT in La Silla in 2009 and 2010. The first instrument was Aqueye (then upgraded to Aqueye+), mounted at the 1.8m Copernicus telescope in Asiago.



## **Detectors and Time Unit**



Single Photon Avalanche Photodiodes (SPADs) with 100 µm pixel size, 35 ps time resolution, ~100 dark count rate, 6-8 MHz maximum count rate, quantum efficiency in the visible up to 60%



The Time Unit tags the photon arrival times with sub-ns accuracy on a highly accurate time scale (UTC) provided by a **Rubidium clock** disciplined by a GPS unit.

## The overall system



Post-processing is done with any convenient selection of time bin (from ns to minutes), thus providing a dynamic range of 6 orders of magnitude.

## Aqueye and Iqueye are now both in Asiago



Having two identical photometers with the same accurate time frame provides another unique capability, namely the simultaneous observation of a celestial source with two distant, non optically coupled telescopes. In other words, a modern version of the Intensity Interferometry first performed by Hanbury Brown and Twiss at Narrabri more than 50 years ago. The first successful attempts have already been made.

March 26, 2019

The photometers have been used for studies on optical pulsars, lunar occultations, exo-planet transits and fast variable objects.

Optical pulsars indeed have been a most rewarding test bed for Aqueye and Iqueye. Here, we present the results obtained at the NTT on *three optical pulsars*. Their light curves, pulse shape and phase determination appear to be the best available in the literature.



# Iqueye at the NTT



March 26, 2019



# Pulsars in a nut shell -1

Hewish et al. 1968, Nature, 217, 709

## Rotating Neutron Stars as the Origin of the Pulsating Radio Sources

by

T. GOLD

Center for Radiophysics and Space Research, Cornell University, Ithaca, New York The constancy of frequency in the recently discovered pulsed radio sources can be accounted for by the rotation of a neutron star. Because of the strong magnetic fields and high rotation speeds, relativistic velocities will be set up in any plasma in the surrounding magnetosphere, leading to radiation in the pattern of a rotating beacon.

Pulsars are rapidly rotating magnetized neutron stars born from the collapse of massive stars, with rotational periods from 1 ms to 10s.

The mass of the pulsar is around 1.4 solar masses, the surface temperature is of the order of 10<sup>5</sup>-10<sup>6</sup> K. The density is of the order of 10<sup>14</sup> g cm<sup>-3</sup>. The inferred magnetic fields range from 10<sup>8</sup> to 10<sup>14</sup> G.

March 26, 2019

## Pulsars in a nut shell -2



Radiation is emitted in a beam that sweeps along our line of sight at every rotation. Timing means measuring the arrival time of these pulses of radiation, deriving the period and its derivatives. The characteristic age of the pulsar and strength of magnetic filed are derived from the following equations:

$$\tau_{\rm c} = \frac{1}{2} \frac{P}{\dot{P}}$$

$$B_{\rm sd} \approx 3.2 \times 10^{19} \sqrt{P\dot{P}} \, {\rm G}$$

The spin frequency  $\nu$  decreases with time:  $d\nu/dt = -k \nu^n$  with a **braking index n** which is =3 for a perfect dipole. Our data have allowed a precise determination of **n** for two optical pulsars.

## Pulsar in a nut shell - 3

To date, about 2500 pulsars have been identified in radio, both isolated and in binary systems. Around 250 are detected in  $\gamma$  rays by Fermi (all pulsating) Around 150 are detected in X rays (not all pulsating) Only a score are seen pulsating in UV, optical and IR.

Different emission mechanisms correspond to different emission regions in the magnetosphere and on the star surface. Therefore, pulsars are unique laboratories for studying:

- nuclear matter equation of state
- relativistic electrodynamics
- intense gravitational fields
- particle acceleration
- · interaction with their nebulae

#### Only multi-wavelength observations give the global picture.

# **Optical pulsars**

There are 5 isolated radio pulsars pulsating also in optical wavelengths, and we have studied 3 of them with Iqueye at the NTT:

1- the V = 16.5, 33 ms Crab Nebula pulsar, in the SN remnant exploded in year 1054. Thanks to its declination it is visible both from Asiago and La Silla

2 – the V=22.5, 50 ms pulsar in the Large Magellanic Cloud (designated as B0540-69), the only extragalactic optical pulsar known today

3 – the V=23.5, 89 ms Vela pulsar

March 26, 2019

La Silla 50th anniversary

## 1 - The Crab pulsar

The Crab pulsar was observed at the NTT in January 2009 and again in December 2009.

In the last occasion simultaneous data were obtained with Jodrell Bank. The radio telescope detected hundreds of Giant Radio Pulses during the Iqueye observations.



#### A series of individual pulses, raw data

## data binned at 3 microseconds = 10<sup>-4</sup> Period



# Pulsar slowing down



Phase of the pulses with respect to uniform rotation. Time on top axis is in MJD. Few days of Iqueye observations provide a robust estimation of the pulsar slowing down.

Combining Jan and Dec data we derive a braking index n = 2.44, similar to the radio one in the over the same period. The age is consistent with the explosion of AD 1054

# Accuracy of period and phase difference with Jodrell Bank

Iqueye and Jodrell Bank radio periods agree to the 1 ps level.



Iqueye data confirm the systematic optical-radio phase difference, with the optical pulse preceding the radio one by about 200 microseconds.

### Values of the radio – optical delay of the main peak



#### The error bars in the Aqueye and Iqueye data is **dominated by the** radio errors.

Our interpretation of the delay is that optical and radio beams are misaligned by 1.5–3 deg because at the position where electrons emit optical photons the magnetic field has a slightly different orientation. Regular monitoring observations

Regular monitoring observations are carried out in Asiago do not find significan evolution of the delay.

# **Optical signature of GRPs**

The night of Dec 14, 2009 Jodrell Bank detected 737 Giant Radio Pulses (GRPs) above a  $6.0-\sigma$  cutoff, of which 663 had concurrent Iqueye observations.



The Iqueye data confirm *a noticeable increase in optical flux* up to a 4-σ level in correspondence to the GRPs.

# The second brightest pulsar: B0540-69 in the Large Magellanic Cloud



This pulsar is approximately 100 times fainter than Crab's, therefore individual pulses could not be detected.

Our observations extended by 9 years the time span over which optical data were obtained.

The braking index n (measurable only in optical and in X rays) over 27 years of optical observations is n =2.087 +/- 0.013, decidedly lower than the magnetic dipole value.

# The UV light curve



Pulsations have been seen also in the UV with the HST. B0540-69 is one of the 4 pulsars that are seen to pulsate from the radio to the gamma-rays.

## 2015 The Fermi LAT detection



Figure 1: Sky maps of the LMC. (A) 0.2–200 GeV gamma-ray emission in a  $10^{\circ} \times 10^{\circ}$  region encompassing the LMC. The map was smoothed using a Gaussian kernel with  $\sigma = 0.2^{\circ}$ . Emission is strongest around 30 Doradus (approximately delimited by the blue box), but also fills much of the galaxy. Contours show the atomic gas distribution. (B) 2–200 GeV gamma-ray emission in a  $2^{\circ} \times 2^{\circ}$  region around 30 Doradus. The map was smoothed using a Gaussian kernel with  $\sigma = 0.1^{\circ}$ . Better angular resolution at higher energies resolves two components coincident with PSR J0540–6919 and PSR J0537–6910, whose locations are indicated as blue dots. Both maps are given in J2000 equatorial coordinates.

## The Gamma to Radio light curves





## The faintest pulsar: Vela



Figure A1. Leahy normalized power spectrum of Obs. 14 (see Table 1). The time bin is 0.01 s. The resolution in frequency is  $10^{-4}$  Hz.

Vela's pulsar is 10 times fainter than B0540-69. However, the periodic signal (period around 89 ms) is plainly evident from the Fourier transform.

# The Iqueye light curve



Two cycles are shown for clarity. The light curve is truly complex

## Phase drift



Figure 2. Phase drift of the Vela pulsar  $\psi(t)$  with respect to uniform rotation. Time t expressed in MJD.

Although we could not derive *n* from our data alone, for the first time we determined the relative time of arrival of the radio-opticalgamma ray peaks with an accuracy of a fraction of a millisecond.

A mosaic of light curves at increasing energies.

The morphology of thelight CURVES changes with increasing energies, different from Crab and B0540-69



## A detailed Iqueye vs Fermi-LAT

6 Spolon et al.



Figure 3. Two periods of the Vela pulsar pulsa profile from the Iqueye optical observations taken in 2009 (top panel) and the Ferni-LAT gamma-ray observations taken during 3 yrs of its operation (MJD 54682.66-65793.85, *bottom panel*). The blue line in each plot shows the best-fitting analytical function which reproduces the profile. Individual components of these functions are also shown with gray dashed lines. Phase 0 corresponds to the radio peak. The average for error bar is shown on the top left.

# Conclusions

IquEYE at the NTT and Aqueye at the Copernicus in Asiago have proven the validity of our design and realization. The instruments perform very well as extremely fast photon counters and multicolor photometers, both for very bright and very faint sources. The concept is very modular, and could be adapted to any telescope.

With telescopes of modest size (< 4 m diameter) it is not possible to achieve significant results on quantum effects.

However, several astrophysical problems requiring the highest time resolution like pulsars and gamma ray bursts, can be investigated better than with other photometers, thanks to the fair quantum efficiency, superior time tagging and time keeping capability and large dynamic range selectable at will.

# The examples here shown attest the validity of Iqueye and Aqueye for multi-wavelength astronomy.

## Main References (by date) - 1

Barbieri C et al. (2009) Very fast photon counting photometers for astronomical applications: IquEYE for the ESO 3.5m New Technology Telescope, Photon Counting Applications, Quantum Optics, and Quantum Information Transfer and Processing II. Edited by Prochazka, Ivan; Sobolewski, Roman; Dusek, Miloslav. Proceedings of the SPIE, Volume 7355

Naletto G et al. (2009) *Iqueye, a single photon-counting photometer applied to the ESO new technology telescope*, Astronomy & Astrophysics, Volume 508, Issue 1, 2009, pp.531-539

Naletto G et al. (2010) *Upgrade of Iqueye, a novel photon-counting photometer for the ESO New Technology Telescope*, Ground-based and Airborne Instrumentation for Astronomy III. Edited by McLean, Ian S.; Ramsay, Suzanne K.; Takami, Hideki. Proceedings of the SPIE, Volume 7735, pp. 773545-773545-12

Barbieri C et al. (2010) *Iqueye, a single photon counting very high speed photometer for the ESO 3.5m NTT*, Proceedings of SPIE - Volume 7681, 2010, Article number 768110 Advanced Photon Counting Techniques IV;Orlando, FL;7 April 2010 through 8 April 2010;Code80660

### Main References - 2

Gradari S et al. (2011). *The optical light curve of the LMC pulsar B0540-69 in 2009*, Monthly Notices of The Royal Astronomical Society, Volume 412, Issue 4, pp. 2689-2694

Barbieri C et al. (2012) Aqueye and Iqueye, Very-High-Time-Resolution Photon-Counting Photometers, New Horizons in Time-Domain Astronomy, Proceedings of the International Astronomical Union, IAU Symposium, Volume 285, p. 280-282

Collins S et al. (2012) *Crab Pulsar: Enhanced Optical Emission During Giant Radio Pulses*, New Horizons in Time-Domain Astronomy, Proceedings of the International Astronomical Union, IAU Symposium, Volume 285, p. 296-298

Zampieri L; Čadež A; Barbieri C; Naletto G; Calvani M; Barbieri M; Verroi E; Zoccarato P; Occhipinti T (2014) *Optical phase coherent timing of the Crab nebula pulsar with Iqueye at the ESO New Technology Telescope* Monthly Notices of the Royal Astronomical Society, 439, 2813

FERMI LAT Collaboration (2015) *An extremely bright gamma-ray pulsar in the Large Magellanic Cloud* Science, 350, 801

## Main References - 3

Čadež A; Zampieri L; Barbieri C; Calvani M; Naletto G; Barbieri M; Ponikvar D (2016) *What brakes the Crab pulsar?,* Astronomy & Astrophysics, Volume 587, id.A99, 11 pp.

Spolon A, Zampieri L, Burtovoi A, Naletto G, Barbieri C, Barbieri M, Patruno A, Verroi E (2019) *Timing analysis and pulse profile of the Vela pulsar in the optical band from Iqueye observations*, Monthly Notices of the Royal Astronomical Society, Volume 482, Issue 1, p.175-183, <u>https://doi.org/10.1093/mnras/sty2605</u>

Mignani, R. P. et al. (2019) *The First Ultraviolet Detection of the Large Magellanic Cloud Pulsar PSR B0540–69 and Its Multi-wavelength Properties*, The Astrophysical Journal, Volume 871, Issue 2, article id. 246, 15 pp. (2019)

## **THANK YOU!**