



Publication Year	2021
Acceptance in OA @INAF	2022-03-29T14:25:04Z
Title	HIRES, the High-resolution Spectrograph for the ELT
Authors	Marconi, Alessandro; Abreu, M.; Adibekyan, V.; Aliverti, Matteo; Allende Prieto, C.; et al.
DOI	10.18727/0722-6691/5219
Handle	http://hdl.handle.net/20.500.12386/32052
Journal	THE MESSENGER
Number	182

HIRES, the High-resolution Spectrograph for the ELT

- Alessandro Marconi^{1, 2}
 Manuel Abreu^{3, 4}
 Vardan Adibekyan^{5, 6}
 Matteo Aliverti⁷
 Carlos Allende Prieto^{8, 9}
 Pedro Amado¹⁰
 Manuel Amate⁸
 Etienne Artigau¹¹
 Sergio Augusto¹²
 Susana Barros^{5, 6}
 Santiago Becerril¹⁰
 Björn Benneke¹¹
 Edwin Bergin¹³
 Philippe Berio¹⁴
 Naidu Bezawada¹⁵
 Isabelle Boisse¹⁶
 Xavier Bonfils¹⁷
 François Bouchy^{18, 16}
 Christopher Broeg¹⁹
 Alexandre Cabral^{3, 4}
 Rocio Calvo-Ortega¹⁰
 Bruno Leonardo Canto Martins²⁰
 Bruno Chazelas¹⁸
 Andrea Chiavassa¹⁴
 Lise Christensen²¹
 Roberto Cirami²²
 Igor Coretti²²
 Stefano Covino⁷
 Giovanni Cresci²
 Stefano Cristiani²²
 Vanderlei Cunha Parro¹²
 Guido Cupani²²
 Izan de Castro Leão²⁰
 José Renan de Medeiros²⁰
 Marco Antonio Furlande Souza¹²
 Paolo Di Marcantonio²²
 Igor Di Varano²³
 Valentina D'Odorico²²
 René Doyon¹¹
 Holger Drass^{24, 25}
 Pedro Figueira^{15, 5}
 Ana Belen Fragoso⁸
 Johan Peter Uldall Fynbo²¹
 Elena Gallo¹³
 Matteo Genoni⁷
 Jonay González Hernández^{8, 9}
 Martin Haehnelt²⁶
 Julie Hlavacek-Larrondo²⁷
 Ian Hughes¹⁸
 Philipp Huke²⁸
 Andrew Humphrey⁵
 Hans Kjeldsen²⁹
 Andreas Korn³⁰
 Driss Kouach³¹
 Marco Landoni⁷
 Jochen Liske³²
 Christophe Lovis¹⁸
 David Lunney³³
 Roberto Maiolino³⁴
 Lison Malo¹¹
 Thomas Marquart³⁰
 Carlos Martins^{5, 35}
 Elena Mason²²
 Paolo Molaro²²
 John Monnier¹³
 Manuel Monteiro⁵
 Christoph Mordasini¹⁹
 Tim Morris³⁶
 Alessio Mucciarelli³⁷
 Graham Murray³⁶
 Andrzej Niedzielski³⁸
 Nelson Nunes^{3, 4}
 Ernesto Oliva²
 Livia Origlia³⁹
 Enric Pallé^{8, 9}
 Giorgio Pariani⁷
 Phil Parr-Burman³³
 José Peñate⁸
 Francesco Pepe¹⁸
 Enrico Pinna²
 Nikolai Piskunov³⁰
 José Luis Rasilla Piñero⁸
 Rafael Rebolo^{8, 9}
 Phil Rees³³
 Ansgar Reiners²⁸
 Marco Riva⁷
 Donatella Romano³⁹
 Sylvain Rousseau¹⁴
 Nicoletta Sanna²
 Nuno Santos^{5, 6}
 Mirsad Sarajlic¹⁹
 Tzu-Chiang Shen²⁴
 Francesca Sortino³⁹
 Danuta Sosnowska¹⁸
 Sérgio Sousa⁵
 Eric Stempels³⁰
 Klaus Strassmeier²³
 Fabio Tenegi⁸
 Andrea Tozzi²
 Stéphane Udry¹⁸
 Luca Valenziano³⁹
 Leonardo Vanzi²⁴
 Michael Weber²³
 Manfred Woche²³
 Marco Xompero²
 Erik Zackrisson³⁰
 María Rosa Zapatero Osorio⁴⁰
- ⁴ Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Portugal
⁵ Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, Portugal
⁶ Departamento de Física e Astronomia, Faculdade de Ciências, Universidade do Porto, Portugal
⁷ INAF – Osservatorio Astronomico di Brera, Italy
⁸ Instituto de Astrofísica de Canarias, La Laguna, Tenerife, Spain
⁹ Departamento de Astrofísica, Universidad de La Laguna, Tenerife, Spain
¹⁰ Instituto de Astrofísica de Andalucía-CSIC Glorieta de la Astronomía, Granada, Spain
¹¹ Institut de Recherche sur les Exoplanètes and Observatoire du Mont-Mégantic, Département de Physique, Université de Montréal, Canada
¹² Instituto Mauá de Tecnologia, São Caetano do Sul, Brazil
¹³ Department of Astronomy, University of Michigan, USA
¹⁴ Laboratoire Lagrange, Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, France
¹⁵ ESO
¹⁶ Aix Marseille University, CNRS, CNES, LAM, France
¹⁷ Université Grenoble Alpes, CNRS, IPAG, France
¹⁸ Département d'Astronomie, Université de Geneve, Switzerland
¹⁹ Physikalisches Institut, University of Bern, Switzerland
²⁰ Board of Observational Astronomy, Federal University of Rio Grande do Norte, Brazil
²¹ Cosmic Dawn Center, Niels Bohr Institute, Copenhagen University, Denmark
²² INAF – Osservatorio Astronomico di Trieste, Italy
²³ Leibniz Institute for Astrophysics Potsdam, Germany
²⁴ Centro de Astro Ingeniería, Pontificia Universidad Católica de Chile, Santiago, Chile
²⁵ Millennium Institute of Astrophysics, Santiago, Chile
²⁶ Kavli Institute for Cosmology and Institute of Astronomy, Cambridge, UK
²⁷ Département de physique, Université de Montréal, Canada
- ¹ Dipartimento di Fisica e Astronomia, Università di Firenze, Italy
² INAF – Osservatorio Astrofisico di Arcetri, Firenze, Italy
³ Instituto de Astrofísica e Ciências do Espaço, Universidade de Lisboa, Portugal

- ²⁸ Institute for Astrophysics, Georg-August-Universität, Göttingen, Germany
- ²⁹ Department of Physics and Astronomy, Aarhus University, Denmark
- ³⁰ Division of Astronomy and Space Physics, Department of Physics and Astronomy, Uppsala University, Sweden
- ³¹ CNRS, OMP, Université de Toulouse, France
- ³² Hamburger Sternwarte, Universität Hamburg, Germany
- ³³ UK Astronomy Technology Centre, Edinburgh, UK
- ³⁴ Cavendish Laboratory, University of Cambridge, UK
- ³⁵ Centro de Astrofísica da Universidade do Porto, Portugal
- ³⁶ Centre for Advanced Instrumentation, Department of Physics, Durham University, UK
- ³⁷ Dipartimento di Fisica e Astronomia, Università di Bologna, Italy
- ³⁸ Institute of Astronomy, Nicolaus Copernicus University in Torun, Poland
- ³⁹ INAF – Osservatorio di Astrofisica e Scienze dello Spazio di Bologna, Italy
- ⁴⁰ Centro de Astrobiología (CSIC-INTA), Madrid, Spain

HIRES will be the high-resolution spectrograph at optical and near-infrared (NIR) wavelengths for ESO's Extremely Large Telescope (ELT). It will consist of three fibre-fed spectrographs providing a wavelength coverage of 0.4–1.8 μm (with a goal of 0.35–1.8 μm) at a spectral resolution of $\sim 100\,000$. Fibre-feeding allows HIRES to have several interchangeable observing modes, including a single-conjugate adaptive optics (SCAO) module and a small diffraction-limited integral field unit in the NIR. It will therefore be able to operate in both seeing- and diffraction-limited modes. HIRES will address a wide range of science cases spanning nearly all areas of research in astrophysics and even fundamental physics. Some of the top science cases will be the detection of biosignatures from exoplanet atmospheres, finding the fingerprints of the first generation of stars (Pop III), tests on the stability of Nature's fundamental couplings, and the direct detection of the cosmic acceleration. The HIRES consortium is composed of more than 30 institutes

from 14 countries, forming a team of more than 200 scientists and engineers.

Introduction

At first light, the ELT will be the largest ground-based telescope at visible and infrared wavelengths. The flagship science cases supporting the successful ELT construction proposal were the detection of life signatures from Earth-like exoplanets and the direct detection of the cosmic expansion re-acceleration, and it is no coincidence that both science cases require observations with a high-resolution spectrograph.

Over the past few decades high-resolution spectroscopy has been a truly interdisciplinary tool, which has enabled some of the most extraordinary discoveries spanning all fields of astrophysics, from planetary sciences to cosmology. Astronomical high-resolution spectrometers have allowed scientists to go beyond the classical domain of astrophysics and to address some of the fundamental questions of physics. In the wide-ranging field of research exploiting high-resolution spectroscopy, ESO has a long and successful tradition, thanks to the exquisite suite of medium- and high-resolution spectrographs offered to the community of Member States. The Ultraviolet and Visual Echelle Spectrograph (UVES), the Fibre Large Array Multi Element Spectrograph (FLAMES), the Cryogenic high-resolution InfraRed Echelle Spectrograph (CRIRES), the medium-resolution spectrograph X-shooter and the High Accuracy Radial velocity Planet Searcher (HARPS) have enabled European teams to lead in many areas of research. The Echelle Spectrograph for Rocky Exoplanet and Stable Spectroscopic Observations (ESPRESSO), which is now joining this suite of very successful high-resolution spectrographs, is fulfilling its promise by truly revolutionising some of these research areas. The scientific interest and high productivity of high-resolution spectroscopy are reflected in the fact that more than 30% of ESO publications can be attributed to its high-resolution spectrographs.

However, it is becoming increasingly clear that in most areas of research high-resolution spectroscopy has

reached the “photon-starved” regime at 8–10-m-class telescopes. Despite major progress on the instrumentation front, further major advances in these fields desperately require a larger photon collecting area. Given its inherently “photon-starved” nature, amongst the various astronomical observing techniques high-resolution spectroscopy is most in need of the collecting area of Extremely Large Telescopes.

When defining the ELT instrumentation suite, ESO commissioned two Phase A studies for high-resolution spectrographs, one to work at visible wavelengths and known as CODEX (Pasquini et al., 2010), and SIMPLE (Origlia et al., 2010) to work in the NIR. Both studies were started in 2007 and completed in 2010. These studies demonstrated the importance of high-resolution optical and NIR spectroscopy at the ELT and ESO therefore decided to include a high-resolution spectrograph (HIRES) in the ELT instrumentation roadmap. Soon after conclusion of the respective Phase A studies the CODEX and SIMPLE consortia realised the great scientific importance of covering both the visible and NIR spectral ranges simultaneously. This marked the birth of the concept of an X-shooter-like spectrograph with higher resolution, capable of providing $R \sim 100\,000$ over the full visible and NIR wavelength range. Following a community workshop in September 2012 the HIRES Initiative prepared a White Paper summarising a wide range of science cases proposed by the community (Maiolino et al., 2013) and also prepared a preliminary technical instrument concept.

With the start of construction of the ELT, the HIRES Initiative decided to organise itself as the HIRES Consortium and recruited additional institutes that had expressed an interest in HIRES. The Consortium, strongly motivated by the unprecedented scientific achievements that the combination of such an instrument with the ELT would enable, was commissioned by ESO to perform a Phase A study. The Phase A study started in March 2016 and concluded successfully in May 2018. Following the conclusion of the Phase A study, other institutes in the USA and Canada joined the HIRES Consortium.

The HIRES Consortium^a is now composed of institutes from Brazil, Canada, Chile, Denmark, France, Germany, Italy, Poland, Portugal, Spain, Sweden, Switzerland, the United Kingdom and the USA. The Italian National Institute for Astrophysics (INAF) is the lead technical institute. See Marconi et al. (2018) for more details on the Consortium structure and organisation.


Science goals

During the Phase A study, the HIRES Science Advisory Team (SAT), chaired by the Project Scientist, defined the science priorities for HIRES and determined the corresponding top-level requirements. These science cases, briefly described below, were then prioritised in order to define the instrument baseline design. Many other science cases are possible with HIRES, but they will not be mentioned here, where we focus on a few representative science goals. A description of the prioritisation process can be found in, for example, Maiolino et al. (2013) and Marconi et al. (2018).

Exoplanets and protoplanetary discs

The study of exoplanet atmospheres for a wide range of objects, from gas giants to rocky planets, and from hot to temperate planets, is a primary objective in the field for the next decade. In particular, the detection of components such as molecular oxygen, water and methane in Earth- or super-Earth-sized planets would be truly transformational, as they may be regarded as signatures of habitability or even signatures of life. Simulations of HIRES observations have been performed by Snellen et al. (2013, 2015) and Hawker & Parry (2019).

HIRES will be able to probe the atmosphere of an exoplanet in transmission during a transit in front of its host star. As an example, it will be possible to detect CO₂ absorption in Trappist-1b with a signal-to-noise ratio (S/N) of 6 in 4 transits of the planet, while O₂ absorption at 0.75 μm would be detected in only 25 transits of the planet, i.e., less than 30 hours of observation. HIRES will also be able to probe exoplanets directly, by



spatially resolving them from their host star, focusing on their reflected starlight and taking advantage of the angular resolution of the ELT with AO-assisted observations. For example, it will be possible to detect the Proxima Centauri b planet in 4 nights of integration at a S/N of 8 with a relatively simple SCAO system similar to that used by other ELT first-light instruments. Figure 1, left, shows that HIRES will be able to detect O₂ from an exoplanet similar to Proxima b in 70 hours of integration.

Protoplanetary discs are a natural outcome of angular momentum conservation in star formation and are ubiquitous around young, forming stars. HIRES will be able to determine the properties of the gas in the inner star-disc region, where competing mechanisms of disc gas dispersal are at play. This will constrain, on the one hand, the mechanisms by which the forming star acquires mass and removes angular momentum, and, on the other hand, the initial conditions for planet formation.

Figure 1. HIRES science highlights. Top: Cross-correlation signal indicating the clear detection of O₂ in a Proxima-b-like exoplanet in 70 hours of total integration (adapted from Figure 4 of Hawker & Parry, 2019). Bottom: Observations of a z = 7 quasar with H_{AB} = 20 and a total integration time of 2 hours showing HIRES's ability to distinguish IGM enrichment by normal SNIi supernovae or by low mass and pair instability supernovae from Pop III stars (simulations by the HIRES Science Advisory Team).

Stars and stellar populations

The vast light-collecting power of the ELT will enable detailed high-resolution spectroscopy of individual stars, and in particular very faint red dwarfs and distant red giants in nearby galaxies, for which HIRES will be able to provide tight constraints on the atmospheric parameters. These constraints will be extremely important for characterising the stellar hosts of exoplanets.

HIRES will also expand our horizons by measuring the heavy-element abundances of the most primitive stars (with low mass and low metallicity) in our Galaxy and its satellites, helping us to understand what is the lowest metallicity




Figure 2. HIRES architectural design, outlining the instrument subsystems: Front-end (seeing-limited and AO assisted with SCAO unit), Fibre Link, Calibration Unit, VIS-Blue, VIS-Red and NIR (cold IR spectrograph).

at which gas can collapse to form low-mass stars, and what are the nature and yields of the very first generation of stars and their supernovae.

Last, but not least, the combination of very high spectral resolving power and diffraction-limited angular resolution makes the ELT a unique resource for deepening our understanding of the physics of stellar atmospheres and nucleosynthesis processes, by allowing us to spectroscopically resolve the effects of surface convection and to measure isotopic abundances of atomic species.

Galaxy formation and evolution, and the intergalactic medium

The detection of Population III stars and the observational characterisation of their properties are major objectives of extragalactic astrophysics. Protogalaxies hosting Pop III stars are expected to be too faint for direct detection, even with the JWST. However, the signature of Pop III stars can be detected through their nucleosynthetic yields, which can potentially be observed in the abundance patterns of very metal-poor absorption systems in the high-resolution, wide-range spectra of bright high-redshift

sources provided by HIRES in the NIR (Figure 1, bottom).

The direct detection and characterisation of the beginning of the reionisation epoch is another very important goal in the study of galaxy formation. This process is believed to have been dominated by ultraviolet photons from the first generations of galaxies, most of which are too faint to be observed directly even with the JWST. By targeting bright quasars at high redshift as background continuum sources, HIRES will be able to study both transmission features in the Lyman- α forest and metal absorption lines associated with these reionisation-epoch sources, constraining the patchiness of the reionisation process, the properties of the ultraviolet background radiation and the chemical enrichment of the intergalactic medium (IGM) in this epoch.

Cosmology and fundamental physics

The observational evidence for the acceleration of the expansion of the Universe and the tensions that have been highlighted by different cosmological probes have shown that our canonical theories of cosmology or fundamental physics may be incomplete (and possibly incorrect) and that there might be unknown physics yet to be discovered. HIRES will allow us to search for, identify and ultimately characterise any new physics through several different but fundamentally inter-related observations which will enable a unique

set of tests of the current cosmological paradigm.

HIRES will be able to constrain the variation of fundamental physical constants like the fine-structure constant α and the proton-electron mass ratio μ with the advantage, compared to laboratory measurements, of exploring variations over timescales of 12 Gyr and spatial scales of 15 Gpc. A detection of variation in the fundamental constants would be revolutionary: it would automatically prove that the Einstein Equivalence Principle is violated (i.e., gravity is not purely geometry), and that there is a fifth force.

HIRES will enable a test of the cosmic microwave background (CMB) temperature-redshift relation, $T(z) = T_0 (1 + z)$, which is a robust prediction of standard cosmology but one that must be directly verified by measurements. A departure from this relation can in turn reveal that the hypothesis of local position invariance (and thus the equivalence principle) is violated or that the number of photons is not conserved. HIRES measurements will greatly improve on the existing constraints on $T(z)$ compared to existing data.

The redshifts of cosmologically distant objects drift slowly with time — the so-called Sandage (or Sandage–Loeb) effect (see Liske et al., 2008). A redshift drift measurement is fundamentally different from all other cosmological observations; it can provide a direct detection of cosmic reacceleration, thus undoubtedly confirming cosmic acceleration and the existence of dark energy, and potentially providing evidence for new physics. HIRES will be capable of detecting the redshift drift in the Lyman- α forests of the brightest currently known QSOs ($\sim 6 \text{ cm s}^{-1} \text{ decade}^{-1}$ at $z = 4$ for a Planck-like standard cosmology). The ELT may thus become the first facility ever to watch the Universe change in “real time”.

Science priorities

These are just a few of the many science cases that can be addressed, a collection of which can be found in Maiolino et al. (2013). However, in order to define the instrument baseline design a prioritisation of the science cases was performed by

the HIRES Science Advisory Team following criteria of scientific impact (transformational versus incremental), feasibility and competitiveness.

Then, if the top level requirements (TLRs) of the top priority science cases were also enabling other science cases, the latter were not considered any further in the subsequent prioritisation, being considered as accomplished together with the top priority science cases. The top science priorities and associated requirements are:

1. **Exoplanet atmospheres in transmission**, requiring a spectral resolution of at least 100 000, a wavelength coverage of at least 0.50–1.80 μm and a wavelength calibration accuracy of 1 m s^{-1} . The implementation of the above TLRs would automatically enable the following science cases:
 - reionisation of the universe,
 - characterisation of cool stars,
 - detection and investigation of near-pristine gas,
 - the study of extragalactic transients.

2. **Variation of the fundamental constants of physics**, requiring an extension to 0.37 μm in addition to the TLRs of priority 1. At wavelengths less than 0.40 μm the throughput of the ELT is expected to be low as a consequence of the planned coating. However, even in the range 0.37–0.40 μm the system is expected to outperform ESPRESSO at the VLT, and a new coating is under study which may be available a few years after first light. This extension towards the blue would also automatically enable investigation of:
 - the cosmic variation of the CMB temperature,
 - the determination of the deuterium abundance,
 - the investigation and characterisation of primitive stars.

3. **Detection of exoplanet atmospheres in reflection**, requiring, on top of the TLRs of priority 1, the addition of an SCAO system and an integral field unit. Reflected-light spectra allow atmospheric emission to be traced from lower altitudes on the day side of the exoplanet. These additional TLRs

would automatically also enable the following science cases:

- planet formation in protoplanetary discs,
- characterisation of stellar atmospheres,
- searching for low-mass black holes.

4. **The Sandage test**, for which the additional TLRs are a wavelength range of 0.40–0.67 μm and a stability of 2 cm s^{-1} , also enabling:
 - the mass determination of Earth-like exoplanets,
 - radial velocity searches and mass determinations for exoplanets around M-dwarf stars.

Instrument concept

Following Phase A and further studies before the start of construction, the HIRES baseline design is for a modular instrument consisting of three fibre-fed cross-dispersed echelle spectrographs — VIS-BLUE (*UBV*), VIS-RED (*RIZ*) and NIR (*YJH*) — providing a simultaneous spectral range of 0.4–1.8 μm (with a goal of 0.35–1.8 μm) at a resolution of 100 000. Fibre-feeding allows several, interchangeable, observing modes, ensuring maximisation of either accuracy, throughput or spatially resolved information. Together with the

SCAO module, the proposed baseline design is capable of fulfilling the requirements of the 4 top science cases.

The baseline design is summarised below but several alternatives were also evaluated during the Phase A study. Also, several add-ons made possible by the modular nature of the instrument have been considered (for example, a polarimetric module at the intermediate focus, or a wavelength extension out to the *K* band (2.0–2.4 μm). The overall concept is summarised in Figure 2. In the front-end the light from the telescope is split, via dichroics, into 3 wavelength channels. Each wavelength channel interfaces with several fibre bundles that feed the corresponding spectrograph module. Each fibre bundle corresponds to an observing mode and together they constitute the Fibre Link. All three spectrographs, VIS-BLUE, VIS-RED and NIR, have a fixed configuration, i.e., there are no moving parts, thereby fulfilling the stability requirements. They include a series of parallel entrance slits consisting of linear micro-lens arrays each glued to the fibre bundles. The split in wavelengths between the spectrographs is influenced, along with other parameters, by the optical throughput of the different types of fibre available on the market; the different modules can therefore be positioned at




Figure 3. HIRES limiting magnitudes obtained from the Exposure Time Calculator for different S/N ratios (30 — top and 100 — bottom), and for compact and extended sources (left and right). Observations are in seeing-limited mode with $R = 100\,000$ and a total exposure time of 1800 seconds.

different distances from the focal plane of the telescope.

The whole instrument should be placed on the Nasmyth platform, if enough volume and mass is available. If necessary, the fibre-feeding allows the VIS-RED and NIR modules to be placed in the Coudé Room, which can also host the Calibration Unit.

Performance

The Exposure Time Calculator (ETC), regularly updated to take into account modifications to the design, is maintained by INAF-Arcetri and can be run online². The ETC can compute the limiting magnitude achievable at a given wavelength in a given exposure time and at a given S/N, or it can compute the S/N achievable at a given wavelength in a given exposure time and at a given magnitude. HIRES expected performances, computed with the ETC, are summarised in Figure 3.

Conclusions

The HIRES baseline design is for three ultra-stable and modular fibre-fed cross-dispersed echelle spectrographs providing a simultaneous spectral coverage of 0.4–1.8 μm (with a goal of 0.35–1.8 μm) at a resolution of 100 000 with several, interchangeable, observing modes ensuring maximisation of either accuracy, throughput or spatially resolved information. Overall, the studies conducted so far have shown that the HIRES baseline design can address the 4 top priority science cases, being able to provide ground-breaking science results with no obvious technical showstoppers.

The construction of HIRES involves the majority of the institutes in ESO Member States with expertise in high-resolution spectroscopy and will require an estimated 30 million euros for hardware (excluding contingencies) and about 500 full time equivalent personnel. Contingencies are expected to be low (5–10%) because the proposed baseline design is based on proven technical solutions and can benefit from heritage from

HARPS and ESPRESSO and other earlier high-resolution spectrographs, for example the Potsdam Echelle Polarimetric and Spectroscopic Instrument (PEPSI) at the 11.8-m Large Binocular Telescope, the SPectropolarimètre InfraRouge (SPIrou) at the 3.6-m Canada France Hawaii Telescope and the Calar Alto high-Resolution search for M dwarfs with Exoearths with Near-infrared and optical Échelle Spectrographs (CARMENES) instrument at the 3.5-m telescope of Calar Alto Observatory. Construction will take about 8–10 years, so with Phase B starting in 2021, HIRES could be at the telescope as early as 2030.

Overall, HIRES is an instrument capable of addressing ground-breaking science cases whilst being almost (telescope) pupil independent, as it can operate in both seeing- and diffraction-limited modes; the modularity ensures flexibility during construction and the possibility to adapt quickly to new developments in both the technical and the science landscapes.

Acknowledgements

The Italian effort for HIRES is supported by the Italian National Institute for Astrophysics (INAF). HIRES work in the UK is supported by the Science and Technology Facilities Council (STFC) at the UK Astronomy Technology Centre, the University of Cambridge (grants ST/S001387/1 and ST/N002873/1) and Heriot Watt University (grant ST/S001328/1), as part of the UK ELT Programme. We acknowledge financial support from the Spanish Ministry of Science and Innovation (MICINN) under projects AYA2017-86389-P, RYC-2013-14875, PGC2018-098153-B-C31, and PID2019-109522GB-C51/52. The German efforts for HIRES are funded by the Federal Ministry for Education and Research (BMBF). Klaus Strassmeier thanks the BMBF-Verbundforschung for support through grants 05A17BAB and 05A2020. This work was supported by FCT – Fundação para a Ciência e a Tecnologia through national funds and by FEDER through COMPETE2020 – Programa Operacional Competitividade e Internacionalização under these grants: UID/FIS/04434/2019; UIDB/04434/2020; UIDP/04434/2020; PTDC/FIS-AST/32113/2017 & POCI-01-0145-FEDER-032113; PTDC/FIS-AST/28953/2017 & POCI-01-0145-FEDER-028953; PTDC/FIS-AST/28987/2017 & POCI-01-0145-FEDER-028987. Research activities of the observational astronomy board at the Federal University of Rio Grande do Norte are supported by continuous grants from the Brazilian funding agencies CNPq, FAPERJ, and INCT-INESpaço. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brazil (CAPES), Finance Code 001.

References

- Hawker, G. A. & Parry, I. R. 2019, *MNRAS*, 484, 4855
 Liske, J. et al. 2008, *MNRAS*, 386, 1192
 Maiolino, R. et al. 2013, arXiv:1310.3163
 Marconi, A. et al. 2018, *Proc. SPIE*, 10702, 107021Y
 Origlia, L. et al. 2010, *Proc. SPIE*, 7735, 77352B
 Pasquini, L. et al. 2010, *Proc. SPIE*, 7735, 77352F
 Snellen, I. et al. 2013, *ApJ*, 764, 182
 Snellen, I. et al. 2015, *A&A*, 576, A59

Links

- ¹ Instrument Web Page <http://hires.inaf.it>
² Exposure time calculator <http://hires.inaf.it/etc.html>

Notes

- ^a Partners of the HIRES Consortium (CI = Coordinating Institute within a country)
 Brazil: Núcleo de Astronomia Observacional, Universidade Federal do Rio Grande do Norte (CI); Instituto Mauá de Tecnologia. Canada: Institut de Recherche sur les Exoplanètes and Observatoire du Mont-Mégantic, département de physique, Université de Montréal. Chile: Pontificia Universidad Católica de Chile (CI); Centre of Astro Engineering, Universidad de Chile; Department of Astronomy, Universidad de Concepcion; Center of Astronomical Instrumentation, Universidad de Antofagasta. Denmark: Niels Bohr Institute, University of Copenhagen (CI); Department of Physics and Astronomy, Aarhus University. France: Laboratoire d'Astrophysique de Marseille, CNRS, CNES, AMU (CI); Institut de Planétologie et d'Astrophysique de Grenoble, Université Grenoble Alpes; Laboratoire Lagrange, Observatoire de la Côte d'Azur; Observatoire de Haute Provence, CNRS, AMU, Institut Pythéas, Institut de Recherche en Astrophysique et Planétologie, Observatoire Midi-Pyrénées; Laboratoire Univers et Particules, Université de Montpellier. Germany: Leibniz-Institut für Astrophysik Potsdam (CI); Institut für Astrophysik, Universität Göttingen; Zentrum für Astronomie Heidelberg, Landessternwarte; Thüringer Landessternwarte Tautenburg; Hamburger Sternwarte, Universität Hamburg. Italy: INAF – Istituto Nazionale di Astrofisica (Lead Technical Institute). Poland: Faculty of Physics, Astronomy and Applied Informatics, Nicolaus Copernicus University in Torun. Portugal: Instituto de Astrofísica e Ciências do Espaço at Centro de Investigação em Astronomia/Astrofísica da Universidade do Porto (CI), Instituto de Astrofísica e Ciências do Espaço at Faculdade de Ciências da Universidade de Lisboa. Spain: Instituto de Astrofísica de Canarias (CI); Instituto de Astrofísica de Andalucía-CSIC; Centro de Astrobiología Sweden: Dept. of Physics and Astronomy, Uppsala University. Switzerland: Département d'Astronomie, Observatoire de Sauverny, Université de Genève (CI); Universität Bern, Physikalisches Institut. United Kingdom: Science and Technology Facilities Council (CI); Cavendish Laboratory & Institute of Astronomy, University of Cambridge; UK Astronomy Technology Centre; Institute of Photonics and Quantum Sciences, Heriot-Watt University. USA: Department of Astronomy, University of Michigan.