



Publication Year	2019
Acceptance in OA	2022-06-22T11:56:24Z
Title	ngVLA Key Science Goal 5 Understanding the Formation and Evolution of Black Holes in the Era of Multi-Messenger Astronomy
Authors	Lazio, T. Joseph W., Alatalo, Katherine, Blecha, Laura, Boizelle, Benjamin, Bower, Geoffrey C., Braatz, James, Bogdanovic, Tamara, Briskin, Walter, Burke-Spolaor, Sarah, Carbone, Dario, Chomiuk, Laura, Civano, Francesca M., Comerford, Julia, Condon, James, Coppejans, Deanne, Corsi, Alessandra, Darling, Jeremiah K., Davis, Timothy A., Frail, Dale A., Hall, Kirsten R., Hallinan, Gregg, Harwood, Jeremy, Kharb, Preeti, Kimball, Amy, Kirkpatrick, Allison, Kording, E. G., Lacy, Mark, Lazzati, Davide, Lister, Matthew L., Liu, Xin, Maccarone, Thomas J., Metzger, Brian, Miller-Jones, J. C. A., MUKHERJEE, DIPANJAN, Nyland, K. E., O'Shaughnessy, Richard, Owen, Benjamin, Patil, Pallavi, Pesce, Dominic, Plotkin, Richard M., PRANDONI, ISABELLA, Ravi, Vikram, Reid, Mark, Reines, Amy, Rujopakarn, Wiphu, Rupen, Michael P., Sand, David, Shen, Yue, Simon, Joseph, Sivakoff, Gregory R., Strader, Jay, Taylor, Greg B., Taylor, Stephen, van Velzen, Sjoert
Handle	http://hdl.handle.net/20.500.12386/32445
Volume	233

close enough to Sgr A* to have orbital timescales of ~years or faster. The conventional explanation for this is that strong scattering in the ISM broadens pulsed radio signals in time, to the point that they become undetectable; observing at higher frequencies reduces this effect, but the steep spectrum of most radio pulsars means the signals also become much fainter. The 2013 discovery of the magnetar J1745-2900 only 0.1 pc in projection from Sgr A* challenged this viewpoint, leading to some claims of a "missing pulsar" problem. However even with the reduced scattering strength inferred from this source, no existing radio telescopes have yet had the sensitivity to detect faint millisecond pulsars (MSPs) at the Galactic center. The proposed next-generation Very Large Array (ngVLA) will have nearly an order of magnitude more sensitivity than any current telescope in the ~5--30 GHz range that is expected to be the sweet spot for detection of pulsars -- including MSPs -- at the Galactic center. In this presentation we will outline the motivations, current knowledge, and predictions for a pulsar search in the Galactic center using the ngVLA, one of the key science goals for the instrument.

Author(s): Michael Kramer, T. Joseph W Lazio, Norbert Wex, Scott M. Ransom, Julia Deneva, James Cordes, Shami Chatterjee, Jason Dexter, Geoffrey C. Bower, Paul Demorest, Robert Wharton, Lijing Shao

Institution(s): National Radio Astronomy Observatory, Cornell University, ASIAA, Max Planck Institute for Extraterrestrial Physics, George Mason University, JPL, Caltech, Max Planck Institute for Radio Astronomy, National Radio Astronomy Observatory

361.25 - ngVLA Key Science Goal 5

Understanding the Formation and Evolution of Black Holes in the Era of Multi-Messenger Astronomy (T. Joseph W Lazio)

The next-generation Very Large Array (ngVLA) will be a powerful telescope for finding and studying black holes across the entire mass range. High-resolution imaging abilities will allow the separation of low-luminosity black holes in the local Universe from background sources, thereby providing critical constraints on the mass function, formation, and growth of black holes. Its combination of sensitivity and angular resolution will provide new constraints on the physics of black hole accretion and jet formation. Combined with facilities across the spectrum and gravitational wave observatories, the ngVLA will provide crucial constraints on the interaction of black holes with their environments, with specific implications for the relationship between evolution of galaxies and the emission of gravitational waves from in-spiraling supermassive black holes and potential implications for stellar mass and intermediate mass black holes. The ngVLA will identify the radio counterparts to transient sources discovered by electromagnetic, gravitational wave, and neutrino observatories, and its high-resolution, fast-mapping capabilities will make it the preferred instrument to pinpoint electromagnetic counterparts to events such as supermassive

black hole mergers. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. Part of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration.

Author(s): E G Kording, Joseph Simon, Sarah Burke-Spolaor, Jeremiah K Darling, Mark Reid, Laura Chomiuk, Dario Carbone, Gregory R Sivakoff, Brian Metzger, Davide Lazzati, Xin Liu, J. C. A. Miller-Jones, Pallavi Patil, Vikram Ravi, Katherine Alatalo, Preeti Kharb,

Institution(s): International Centre for Radio Astronomy Research-Curtin University, Rochester Institute of Technology, Univ degli Studi di Torino, INAF-Istituto di Radioastronomia, Oregon State University, Purdue University, University of Illinois Urbana-Champaign,

361.26 - VLBA Observations of the Structure and Motions of the Inner Parsec of the M87 Jet (Robert Craig Walker)

Walker et al (2018, Ap.J. 855, 128) reported results from Very Long Baseline Array (VLBA) observations of M87 at 43 GHz which include intensive monitoring in 2007 and 2008 plus 17 years of roughly annual observations, for a total of 50 individual observations. The results from that study are reviewed in this poster. The central radio source in M87 provides the best opportunity to study jet formation because it has a large angular size for the gravitational radius of the black hole and has a bright jet that is well resolved by VLBI observations. The 43 GHz VLBA observations have a resolution of about 0.21 x 0.43 milli-arcseconds (mas) which is about 30 by 60 Schwarzschild radii (R_s) for $D = 16.7$ Mpc and $M_{bh} = 6.1 \times 10^9 M_{sun}$. Our high-dynamic-range images clearly show the wide-opening-angle structure of the jet and show the counter-jet. The jet and counter-jet are nearly symmetric in the inner 1.5 milli-arcseconds (0.12 pc in projection) with both being edge brightened. Both show deviations from parabolic shape in the form of an initial rapid expansion in width and subsequent contraction followed by further rapid expansion and, beyond the visible counter-jet, subsequent collimation. Proper motions and counter-jet/jet intensity ratios both indicate acceleration from apparent speeds of less than about 0.5c to greater than about 2c in the inner about 2 mas (0.16 pc or 240 R_s in projection) and suggest a helical flow. The jet displays a sideways shift with an approximately 8 to 10 year quasi-periodicity. The shift propagates outwards non-ballistically and significantly more slowly than the flow speed revealed by the fastest moving components. Polarization data show a systematic structure with magnetic field vectors that suggest a toroidal field close to the core.

Author(s): Frederick B. Davies, Chun Ly, William Junor, Robert Craig Walker, Philip E. Hardee

Institution(s): National Radio Astronomy Observatory, Steward Observatory, University of Arizona, University of Alabama, Los Alamos National Laboratory, University of