



Publication Year	2019
Acceptance in OA@INAF	2022-06-22T12:25:35Z
Title	Erratum: Radiolysis of N ₂ O:CO ₂ Ice by Heavy Ions: Simulation of Cosmic Ray Effects
Authors	Pereira, R C; de Barros, A L F; Fulvio, Daniele; Boduch, P; Rothard, H; et al.
DOI	10.1093/mnras/sty3207
Handle	http://hdl.handle.net/20.500.12386/32450
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
Number	483

Erratum: Radiolysis of N₂O:CO₂ Ice by Heavy Ions: Simulation of Cosmic Ray Effects

by R. C. Pereira,¹ A. L. F. de Barros,^{1,2*} D. Fulvio,^{3★} P. Boduch,⁴ H. Rothard⁴ and E. F. da Silveira³

¹Programa de Pós-graduação em Instrumentação Óptica, Centro Federal de Educação Tecnológica Celso Suckow da Fonseca, Av. Maracanã 229, 20271-110, Rio de Janeiro, RJ, Brazil

²Departamento de Física, Centro Federal de Educação Tecnológica Celso Suckow da Fonseca, Av. Maracanã 229, 20271-110, Rio de Janeiro, RJ, Brazil

³Departamento de Física, Pontifícia Universidade Católica do Rio de Janeiro, Rua Marquês de São Vicente 225, 22451-900, Rio de Janeiro, RJ, Brazil

⁴Centre de Recherche sur les Ions, les Matériaux et la Photonique Normandie Université, ENSICAEN, UNICAEN, CEA, CNRS, CIMAP, 14000 Caen, France

Key words: errata, addenda – astrochemistry – molecular data – methods: laboratory: molecular – interplanetary medium – cosmic rays – ISM: molecules.

In our recent paper ‘Radiolysis of N₂O:CO₂ Ice by Heavy Ions: Simulation of Cosmic Ray Effects’ published in Mon. Not. R. Astron. Soc. 478, 4939 – 4951 (2018), some errors came to our attention after the publication online, such as two missing spectra in Fig. 1, and some minor errors in Table 3.

In this Erratum, errors have been identified and corrected, as follows:

1 - Column density unit, throughout the paper, is always molecule cm⁻².

2 - A-values, throughout the paper, always refer to band strength.

3 - Fig. 1: the missing spectra in panels (a) and (b) are now shown (see on right column).

4 - In Fig. 2, panel (a), the correct label over the peak at about 3600 cm⁻¹ is CO₂.

5 - Page 2 (Experimental Setup), column 2, third paragraph, the correct value for the density of pure CO₂ is 0.98 g cm⁻³ (Luna R. et al. (2012)). As a consequence, the estimated value for the ice thickness is about 0.38 μm. Since the penetration depth of the 90 MeV ¹³⁶Xe²³⁺ ions used in this work is equal to about 18 μm (>>0.38 μm), the new value of ice thickness does not change the current results and conclusions.

6 - Page 3 (Results), column 2: 4th paragraph has a typo: Tables 1 and 2 instead of Tables1tbl2.tab.

7 - Table 1: the footnote reference Yamada & Person (1964) is (d).

8 - Table 2: the correct reference for footnote (b) refers to de Barros et al. (2017).

9 - In Table 3, References^fJamieson, Mebel & Kaiser (2006) and^hGerakines et al. (2001) are not relevant for this work. The correct Table 3 is presented below.

10 - Fig. 6: the band at about 1980 cm⁻¹ is an artifact.

11 - Table 4, the correct caption is: Characteristics of the N₂O and CO₂ products formed in the N₂O:CO₂ radiolysis.

12 - Page 9, Section 4.2.2: typing error: 216-cm⁻¹ is 2164 cm⁻¹.

13 - Fig. 10 (a): the label of the band at about 2280 cm⁻¹ in the bottom left corner refers to ¹³CO₂.

14 - In the reference list there are two missing references:




Figure 1. Infrared spectra showing radiolysis effects on N₂O:CO₂ ice: (a) spectrum of the non-irradiated ice mixture from 4000 to 600 cm⁻¹; (b) spectrum after 1.0×10^{12} ions cm⁻²; and (c) zoom of the previous spectrum, in the 2170 to 900 cm⁻¹ region. The N₂O₃ band peaked at 1305 cm⁻¹ appears as a shoulder band after de-convolution of the N₂O fundamental band (1296 cm⁻¹).

* E-mail: ana.barros@cefet-rj.br (ALFDB); dfulvio@puc-rio.br (DF)

Table 3. IR band characteristics for the observed product species: assignment, wavenumber position, wavelengths, and band strength (A-value).

Molecule	Assignment	Current wavenumber (cm $^{-1}$)	Wavelengths (μm)	Wavenumber literature Value (cm $^{-1}$)	Band strength (10 $^{-18}$ cm molecule $^{-1}$)
CO	v_1	2138	4.67	2139 ^a	11.0 ^a
CO_3	v_3	2045	4.89	2045 ^b	89.0 ^b
NO	v_1	1848 → 1869*	5.41 → 5.35	1869 ^d	4.5 ^c
N_2O_4	v_9	1741	5.74	1741 ^e	5.0 ^e
N_2O_2/N_2O_4	v_5	1765	5.67	1764 ^d	—
N_2O_4	v_2	1621	6.17	1628 ^f	59.5 ^f
NO_3	v_1	1460	6.85	1471 ^f	7.6 ^f
N_2O_3	v_1/v_3	1305	7.66	1303 ^f	46.3 ^f
N_2O_4	v_{11}	1260	7.94	1262 ^f	85.0 ^f
O ₃	v_1	1037	9.64	1040 ^f	14.0 ^f

Note. ^aPalumbo & Strazzulla (1993), ^bBennett et al. (2004), ^cSicilia et al. (2012), ^dFateley, Bent & Crawford (1959), ^eFulvio et al. (2009), ^fJamieson et al. (2005). *The arrow means blue shift of the NO band along the irradiation.

REFERENCES

- Bennett C. J., Jamieson C., Mebel A. M., Kaiser R. I., 2004, *Physical Chemistry Chemical Physics*, 6, 735
de Barros A. L. F., da Silveira E. F., Fulvio D., Rothard H., Boduch P., 2017,
MNRAS, 465, 3281

Luna R., Satorre M. A., Domingo M., Millán C., Santonja C., 2012, *Icarus*, 221, 186

This paper has been typeset from a TeX/LaTeX file prepared by the author.