

Publication Year	2007
Acceptance in OA@INAF	2022-12-27T14:40:05Z
Title	Massive star-formation in G24.78+0.08 studied by means of maser VLBI and thermal interferometric observations
Authors	MOSCADELLI, Luca; Goddi, C.; CESARONI, Riccardo; BELTRAN SOROLLA, MARIA TERESA
DOI	10.1017/S1743921307012690
Handle	http://hdl.handle.net/20.500.12386/32805
Series	PROCEEDINGS OF THE INTERNATIONAL ASTRONOMICAL UNION
Number	vol. 3, S242

Massive star-formation in G24.78+0.08 studied by means of maser VLBI and thermal interferometric observations

L. Moscadelli¹, C. Goddi², R. Cesaroni¹ and M. T. Beltràn³

¹INAF, Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy email: mosca@arcetri.astro.it

²Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

³Departament d'Astronomia i Meteorologia, Universitat de Barcelona, Av. Diagonal 647, 08028, Barcelona, Catalunya, Spain

Abstract. This work presents the results of VLBI observations of 6.7 GHz methanol and 22.2 GHz water masers towards the mm core A in the massive star-forming region G24.78+0.08. Comparing the maser with previous millimeter interferometric and recent continuum VLA observations, the physical properties and the gas kinematics of the G24 A core on linear scales from ~100 AU to ~0.1 pc are determined.

Keywords. Masers, stars: formation, HII regions, techniques: interferometric

1. Introduction

The star forming region G24.78+0.08 is located at a distance of 7.7 kpc and has a large bolometric luminosity ($\leq 7 \times 10^4 L_{\odot}$). Furuya *et al.* (2002) first detected three molecular cores (in continuum emission at 2 mm) and two HII regions (one of these embedded in one of the cores) spread over ~ 0.4 pc. The region contains at least four distinct centers of star formation, named A, B, C, and D, which Furuya *et al.* (2002) speculate to be in different evolutionary states. Both the molecular core A and the source B have associated HII regions (Codella *et al.* 1997) and, hence, are already ZAMS stars, whereas the other two appear to be less evolved. Cores A and C power bipolar molecular outflows (Furuya *et al.* 2002).

1.4 mm Plateau de Bure Interferometer (PdBI) one-arcsecond angular resolution observations by Beltrán *et al.* (2004) have resolved the structure of the G24 A core into two distinct subcores: G24 A1, towards the southeast (SE), and G24 A2, towards the northwest (NW). G24 A1 is associated with the most compact HII region, while G24 A2 shows strong pointlike emission from 3.6 to 8 μ m (GLIMPSE) but weak radio 22 GHz continuum emission. Both G24 A1 and A2 are hot molecular cores (HMCs) (Beltrán *et al.* 2004), with a V_{LSR} gradient along a northeast-southwest (NE-SW) direction, approximately perpendicular to the axis of the ¹²CO molecular outflow. This fact suggests that the two cores might be rotating around the outflow axis (Beltrán *et al.* 2004). Towards G24 A1 and A2 several species of molecular masers have been detected: 6.7 GHz CH₃OH, observed with the Australian Compact Array (ATCA) by Walsh *et al.* (1998), and 22.2 GHz H₂O and 1.6 GHz OH masers observed with the VLA by Forster & Caswell (1989).



Figure 1. a) Map of the CH₃CN ($12 \rightarrow 11$) line peak velocity towards G24 A, with the contour levels in kilometers per seconds indicated in the wedge to the right of the panel (Beltrán *et al.* 2004); overlaid on the CH₃CN velocity map, is shown the contour map of the 1.4 mm continuum emission (*solid lines*) and the positions of the CH₃OH 6.7 GHz masers (*filled spheres*); the straight black line represents the axis of the molecular outflow observed with PdBI in ¹²CO(1–0) by Furuya *et al.* (2002). b) and c) Positions and line-of-sight velocities of the 6.7 GHz maser features observed towards the subcore A2 and A1, respectively, with the colors (see online version) denoting different line-of-sight velocities, according to the scale on the right-hand side of the panels.

2. Observational results

2.1. Class II CH₃ OH masers

Figure 1 shows a map of the CH_3CN (12–11) line line-of-sight velocity overlaid on the contour map of the 1.4 mm continuum emission towards G24 A1 and G24 A2 (Beltrán *et al.* 2004). The absolute positions (accurate to within 50 mas) of the 6.7 GHz methanol masers observed with the EVN are plotted as filled black circles (Fig. 1, panel a). The 6.7 GHz maser emission occurs from two distinct groups of features, each associated with one of the mm subcores (A1 or A2). Both groups of maser features are elongated along a direction that agrees well with the axis of the velocity gradient measured in the CH_3CN 1.4 mm line.

Panels c and b of Fig. 1 show the line-of-sight velocities of the 6.7 GHz maser features, associated respectively with the A1 and A2 subcores. Across each maser group, the variation of velocities agrees with the velocity gradient observed using the CH₃CN (12–11) line in the harbouring subcore, with redshifted velocities to the NE and blueshifted velocities to the SW. The methanol maser features are distributed over regions ~three times smaller (0.023 and 0.02 pc) and cover an observed velocity range ~three times larger (9.1 and 5.7 km s⁻¹) than the emission of the CH₃CN thermal line in both subcores.

2.2. $H_2 O$ masers

By observing G24.78+0.08 at 22.2 GHz with the VLBA in phase-reference mode at four different epochs, absolute positions and proper motions of the water maser features have been derived. Figure 2 shows the water maser distribution in the subcore A1, which contains the strongest water maser features of the entire G24.78+0.08 region, with



Figure 2. Comparison of the distribution of the VLBA 22.2 GHz water (*triangles*) and EVN 6.7 GHz methanol (*dots*) maser features in the mm subcore G24 A1. The colors (see online version) denote different velocities, according to the scale on the right-hand side of the panel. Black arrows show the measured absolute proper motions of water maser features, using dotted arrows to denote the most uncertain measurements; the amplitude scale for proper motions is reported at the bottom of the panel. The ellipse shows the position and the deconvolved FWHM size of the G24 A1 continuum source observed recently in K-band with the VLA by Beltran *et al.* (in preparation).

intensities ranging from ≈ 0.05 to ≈ 100 Jy beam⁻¹. Maser emission arises from two distinct regions found in the NE and SW corner of the plotted area. The strongest features of the NE maser cluster (called "A1N") are distributed along an EW line with a length of ≈ 100 mas, and along an arc-shaped structure of diameter ≈ 50 mas. With a few exceptions, most of the maser features have velocities within ± 5 km s⁻¹ of the LSR systemic velocity (111 km s⁻¹) of the region. The measured absolute proper motions, with amplitudes in the range 20–60 km s⁻¹, have directions roughly perpendicular to the linear and arcuate spatial distributions of the maser features.

3. Rotation and expansion in the mm core G24 A

We now compare the VLBI CH₃OH and H₂O maser observations towards the core G24 A with previous millimeter PdBI and recent VLA observations, and discuss a model for explaining the gas kinematics from small (100 AU) to large (0.1 pc) scale.

3.1. Evidence for rotation

Figure 1 shows that, in both cores G24 A1 and A2, the velocity distribution of the methanol 6.7 GHz maser spots agrees well with the velocity gradient measured in the thermal CH₃CN (12–11) line. If, as suggested by Beltrán *et al.* (2004), the CH₃CN velocity gradient is due to rotation, the three times larger spread of velocities of the methanol masers across a three times smaller region, might be explained in terms of conservation of angular momentum from large to small scales inside the cores. Assuming that inside the region (of diameter ≈ 0.02 pc) traced by the 6.7 GHz maser emission in both cores, the gas is in rotational equilibrium, a dynamical mass of 55 and 19 M_{\odot}

for G24 A1 and G24 A2, is derived respectively. From the free-free continuum emission of the ultra-compact HII (UCHII) region in G24 A1, Codella *et al.* (1997) estimate a spectral type of O9.5 for the exciting star, corresponding to a stellar mass of $\approx 20 M_{\odot}$. This must be added to the mass of the gas within 0.02 pc from the YSO, which is \approx $30 M_{\odot}$ for a mean density of 10^8 cm^{-3} . The total mass (star plus gas) is consistent with the mass of 55 M_{\odot} computed from the methanol maser velocity range in G24 A1.

3.2. Evidence for expansion

The water maser proper motions measured in the A1N cluster are oriented approximately perpendicular to the geometrical distribution of the maser features, which are mainly concentrated in two structures, a line oriented along the EW direction and an arc (see Fig. 2). The separation from the LSR systemic velocity of the line-of-sight velocities of most maser features of the A1N cluster is quite small ($\leq 5 \text{ km s}^{-1}$) compared with the sky-projected velocity components (20–60 km s⁻¹), implying that the velocity vectors of the maser features are close to the plane of the sky. The observed velocity distribution suggests that the A1N water masers are tracing gas *expanding* towards N-NE.

Figure 2 shows that the UCHII region falls close to the A1N maser cluster. Considering that the estimated size of the HII region is likely an upper limit, most of the water maser features appear to be distributed along the border of the ionized gas. Furthermore, the elongation of the continuum source (P.A. = 66°) agrees well with the average orientation of the water maser proper motions. Thus, the position and the geometry of the continuum source suggests that the water maser motions are driven by the expansion of the ionized gas. Freely expanding ionized gas should move at a maximum velocity comparable to the sound speed in the ionized gas, $\approx 10 \text{ km s}^{-1}$ assuming a kinetic temperature of 10^4 K . The observed velocities of the water masers are significantly higher than 10 km s^{-1} , indicating that the ionized gas is *not* freely expanding, but that the expansion must be driven by some additional mechanism than thermal pressure.

Stellar winds are commonly observed towards early-type OB main-sequence stars and observational evidence is being accumulated that such winds play an important role during the early expansion phase of an UCHII region (Hoare *et al.* 2007). A detailed study by Lamers & Leitherer (1993) indicates wind velocities $V_{\rm w} = 1500 - 2500$ km s⁻¹ and mass loss rates $\dot{M}_{\rm w} = 1-5 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$, as appropriate for an O9.5 star. Such a powerful stellar wind might be already emitted during the early ZAMS evolution of a massive star. We have considered the case that the expansion of the water maser shell in G24 A1 is driven by a strong wind. Our VLBA water maser and VLA continuum radio observations allow us to set a value for the shell radius and velocity with $R_0 \approx 500$ AU, and $V_0 \approx 40$ km s⁻¹ respectively. To model the dynamics of the wind-driven shell, we have used equations given in Castor *et al.* (1975) and Shull (1980). A detailed report of our calculations is given in Moscadelli *et al.* 2007 (in preparation). We find that, over the

139

explored parameter space for the stellar wind properties, and fixing the wind mechanical luminosity, gives a unique value of ambient density for which the wind-driven shell can attain the observed values of radius and velocity. For the likely values of wind luminosities for a O9.5 star, the wind-driven shell models require the hydrogen particle ambient density to lie in the range 6×10^6 cm⁻³ $\leq n_{\rm H} \leq 2.5 \times 10^7$ cm⁻³. The shell age, t_0 , is predicted to be in the narrow interval $t_0 = 30$ -35 yr.

3.3. Rotation or expansion in the G24 A1 core?

On the basis of the discussed observational evidences, we propose two alternative interpretations to explain the gas kinematics in the G24 A1 core on scales from 100 AU to 0.1 pc.

The expansion detected in the 22.2 GHz water masers on scales of ≈ 100 AU (see Fig. 2) might be the dominant motion across the whole molecular core. Thus, the regular variation of line-of-sight velocity observed in the thermal CH₃CN (12–11) line as well as in the CH₃OH 6.7 GHz maser emission (see Fig. 1)on linear scales of 0.05 pc, might be due to expansion rather than to rotation, as proposed by Beltrán *et al.* (2004). In particular, the regular velocity gradient can be accounted for only by a *collimated* flow. In this interpretation, the 22.2 GHz water masers would mark the inner portion of a molecular outflow, closer to the YSO and not yet collimated, while the CH₃OH 6.7 GHz maser and the CH₃CN thermal emissions would trace the same outflow at larger scale, where collimation is already effective.

Interpreting the motion of the gas in the G24 A1 core in terms of a collimated outflow oriented along the NE–SW direction, apparently contrasts with the PdBI ¹²CO(1–0) observations by (Furuya et al. 2002), indicating the presence of a bipolar outflow on scale ≥ 0.1 pc oriented SE–NW. We propose an alternative scenario, where expansion and rotation can coexist. Figure 2 shows that the detected CH_3OH 6.7 GHz maser spots are on average more distant from the center of the continuum source than the water maser spots and located beyond the border of the UCHII region. We speculate that methanol masers are tracing the pre-shock ambient gas radiatively heated by the YSO radiation. The value of pre-shock ambient gas density indicated by the wind-driven shell model fits well with the observations of both H_2O and CH_3OH masers around the G24 A1 continuum source, since a pre-shock gas density $n_{\rm H} \sim 10^7 \ {\rm cm}^{-3}$ is predicted in the excitation models of both water and methanol masers. The kinematics of methanol masers is expected to differ from that of water masers as the former are located near the G24 A1 core and are unaffected by expansion in the HII region. The water masers should mark the border of the dynamical interaction between the YSO and the surrounding ambient medium: gas moving beyond the water maser shell is not vet perturbed by the YSO evolution (and might be still rotating and slightly contracting), whereas gas inside the maser shell is rapidly expanding due to the YSO wind pressure.

References

Beltrán, M. T., Cesaroni, R., Neri, R., et al. 2004, ApJL, 601, L187
Castor, J., McCray, R., & Weaver, R. 1975, ApJL, 200, L107
Codella, C., Testi, L., & Cesaroni, R. 1997, A&A, 325, 282
Forster, J. R. & Caswell, J. L. 1989, A&A, 213, 339
Furuya, R. S., Cesaroni, R., Codella, C., et al. 2002, A&A, 390, L1
Hoare, M. G., Kurtz, S. E., Lizano, S., Keto, E., & Hofner, P. 2007, in Protostars and Planets V, ed. B. Reipurth, D. Jewitt, & K. Keil, 181–196
Lamers, H. J. G. L. M. & Leitherer, C. 1993, ApJ, 412, 771
Shull, J. M. 1980, ApJ, 238, 860
Walsh, A. J., Burton, M. G., Hyland, A. R., & Robinson, G. 1998, MNRAS, 301, 640