



<b>Publication Year</b>	2020
<b>Acceptance in OA @INAF</b>	2024-01-19T09:56:14Z
<b>Title</b>	Candidate Population III stellar complex at $z = 6.629$ in the MUSE Deep Lensed Field
<b>Authors</b>	VANZELLA, Eros; MENEGHETTI, MASSIMO; Caminha, G. B.; CASTELLANO, Marco; CALURA, Francesco; et al.
<b>DOI</b>	10.1093/mnrasl/slaa041
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/34558">http://hdl.handle.net/20.500.12386/34558</a>
<b>Journal</b>	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY. LETTERS
<b>Number</b>	494

# Candidate Population III stellar complex at $z = 6.629$ in the MUSE Deep Lensed Field

E. Vanzella,<sup>1★</sup> M. Meneghetti<sup>1★</sup>, G. B. Caminha,<sup>2</sup> M. Castellano,<sup>3</sup> F. Calura<sup>1★</sup>,  
P. Rosati,<sup>1,4</sup> C. Grillo,<sup>5</sup> M. Dijkstra, M. Gronke<sup>6</sup>, E. Sani,<sup>7</sup> A. Mercurio,<sup>8</sup> P. Tozzi,<sup>9</sup>  
M. Nonino,<sup>10</sup> S. Cristiani,<sup>10</sup> M. Mignoli,<sup>1</sup> L. Pentericci,<sup>3</sup> R. Gilli,<sup>1</sup> T. Treu,<sup>11</sup>  
K. Caputi,<sup>2</sup> G. Cupani,<sup>10</sup> A. Fontana,<sup>3</sup> A. Grazian<sup>12</sup> and I. Balestra<sup>10,13</sup>

<sup>1</sup>INAF – Osservatorio di Astrofisica e Scienza dello Spazio, via Gobetti 93/3, I-40129 Bologna, Italy

<sup>2</sup>Kapteyn Astronomical Institute, University of Groningen, Postbus 800, NL-9700 AV Groningen, the Netherlands

<sup>3</sup>INAF – Osservatorio Astronomico di Roma, Via Frascati 33, I-00078 Monte Porzio Catone (RM), Italy

<sup>4</sup>Dipartimento di Fisica e Scienze della Terra, Università degli Studi di Ferrara, via Saragat 1, I-44122 Ferrara, Italy

<sup>5</sup>Dipartimento di Fisica, Università degli Studi di Milano, via Celoria 16, I-20133 Milano, Italy

<sup>6</sup>Department of Physics, University of California, Santa Barbara, CA 93106, USA

<sup>7</sup>European Southern Observatory, Alonso de Cordova 3107, Casilla 19, Santiago 19001, Chile

<sup>8</sup>INAF – Osservatorio Astronomico di Capodimonte, Via Moiariello 16, I-80131 Napoli, Italy

<sup>9</sup>INAF – Osservatorio Astrofisico di Arcetri, Largo E. Fermi, I-50125 Firenze, Italy

<sup>10</sup>INAF – Osservatorio Astronomico di Trieste, via G. B. Tiepolo 11, I-34143 Trieste, Italy

<sup>11</sup>Department of Physics and Astronomy, University of California, Los Angeles, CA 90095, USA

<sup>12</sup>INAF – Osservatorio Astronomico di Padova, Vicolo Osservatorio 5, I-35122 Padova, Italy

<sup>13</sup>OmegaLambdaTec GmbH, Lichtenbergstrasse 8, D-85748 Garching bei Munchen, Germany

Accepted 2020 March 4. Received 2020 February 26; in original form 2020 January 9

## ABSTRACT

We discovered a strongly lensed ( $\mu \gtrsim 40$ ) Ly $\alpha$  emission at  $z = 6.629$  ( $S/N \simeq 18$ ) in the MUSE Deep Lensed Field (MDLF) targeting the Hubble Frontier Field (HFF) galaxy cluster MACS J0416. Dedicated lensing simulations imply that the Ly $\alpha$  emitting region necessarily crosses the caustic. The arc-like shape of the Ly $\alpha$  extends 3 arcsec on the observed plane and is the result of two merged multiple images, each one with a de-lensed Ly $\alpha$  luminosity  $L \lesssim 2.8 \times 10^{40}$  erg s $^{-1}$  arising from a confined region ( $\lesssim 150$  pc effective radius). A spatially unresolved *Hubble Space Telescope* (*HST*) counterpart is barely detected at  $S/N \simeq 2$  after stacking the near-infrared bands, corresponding to an observed (intrinsic) magnitude  $m_{1500} \gtrsim 30.8$  ( $\gtrsim 35.0$ ). The inferred rest-frame Ly $\alpha$  equivalent width is  $EW_0 > 1120$  Å if the IGM transmission is  $T_{\text{IGM}} < 0.5$ . The low luminosities and the extremely large Ly $\alpha$   $EW_0$  match the case of a Population III (Pop III) star complex made of several dozens stars ( $\sim 10^4 M_\odot$ ) that irradiate an H II region crossing the caustic. While the Ly $\alpha$  and stellar continuum are among the faintest ever observed at this redshift, the continuum and the Ly $\alpha$  emissions could be affected by differential magnification, possibly biasing the  $EW_0$  estimate. The aforementioned tentative *HST* detection tends to favour a large  $EW_0$ , making such a faint Pop III candidate a key target for the *James Webb Space Telescope* and Extremely Large Telescopes.

**Key words:** gravitational lensing: strong – galaxies: formation – galaxies: starburst.

## 1 INTRODUCTION

Finding and characterizing the first galaxies are the next frontier in observational astronomy. It is thought that the Universe was initially

metal-enriched by the first generation of Population III (Pop III) stars, which could also have played a key role in cosmic reionization before the formation of primeval galaxies (e.g. Dayal & Ferrara 2018; Wise 2019; Zackrisson & Vikaeus 2019, and references therein). Late ( $z < 7$ ) Pop III star formation might also have occurred in pristine regions due to inhomogeneous metal enrichment of the first galaxies (Tornatore, Ferrara & Schneider 2007; Salvaterra, Ferrara & Dayal 2011; Visbal, Haiman & Bryan 2016). Given

\* E-mail: [eros.vanzella@inaf.it](mailto:eros.vanzella@inaf.it) (EV); [massimo.meneghetti@inaf.it](mailto:massimo.meneghetti@inaf.it) (MM); [francesco.calura@inaf.it](mailto:francesco.calura@inaf.it) (FC)


the exceptionally high-effective temperatures of Pop III stars in the zero-age main sequence, they emit a large fraction of their luminosity in the Lyman continuum and have a much harder ionizing spectrum than stars with higher metallicity. The main characteristics of their predicted spectral energy distribution (SED) are the presence of a prominent rest-frame Ly $\alpha$  (Lyman-alpha) emission line due to the strong ionizing flux up to  $\sim 1000\text{--}4000$  Å rest-frame equivalent width (denoted as  $EW_0$ , hereafter) and significant He recombination line (especially He II $\lambda 1640$ , with  $EW_0$  up to  $15\text{--}40$  Å) due to spectral hardness, while a clear deficit of all the metal lines is expected. In particular, Inoue (2011) suggested the following criteria for the identification of extremely metal poor or Pop III galaxies:  $EW_0(\text{Ly}\alpha) > 230$  Å,  $EW_0([\text{O III}]\lambda 5007) < 20$  Å, and  $EW_0(\text{He II}\lambda 1640) > 1$  Å, and prominent Balmer lines like  $EW_0(\text{H}\alpha) > 1900$  Å, while showing an extremely blue ultraviolet spectral slope [ $\beta \sim -3$ ,  $F(\lambda) \sim \lambda^\beta$ ].

Observations have yielded candidates for Pop III stellar populations at high redshift (e.g. Kashikawa et al. 2012; Sobral et al. 2015, and references therein), yet without any definitive detection. These include a controversial  $z = 6.6$  galaxy dubbed CR7 that displays He II $\lambda 1640$  emission (Shibuya et al. 2018; Sobral et al. 2019). Thus, to date, there has not been a confirmed observation of a galaxy dominated by the flux of Pop III stars. The possibility of observing signatures from very metal poor or Pop III star clusters through gravitational lensing has also been discussed, e.g. Zackrisson et al. (2015) (see also Hernán-Caballero et al. 2017), including the detection of single Pop III stars with fluxes temporarily magnified to extreme values (with the magnification parameter  $\mu \simeq 10^3\text{--}10^5$ ) during their transit across the caustic of a galaxy cluster. Such single-star-transit events can boost the flux of the star by  $7\text{--}12$  mag (Windhorst et al. 2018), making such objects visible for a limited amount of time even down to intrinsic magnitudes of  $35\text{--}38$ . Examples of such events detecting single normal stars at  $z < 2$  have been reported recently by Rodney et al. (2018).

Very low luminosity emission-line galaxies have been identified in Hubble Ultra Deep Field, down to magnitude  $30\text{--}32$  ( $M_{1500} = -15$ ) and  $S/N \sim 1\text{--}5$  (Maseda et al. 2018). Strong gravitational lensing allowed us to shed further light on similar low-luminosity objects, providing higher  $S/N \sim 20$  for individual cases (e.g. Vanzella et al. 2017, 2019). In this letter, we present an object at  $z = 6.629$  showing (1) the faintest Ly $\alpha$  emission ever detected at  $z > 6$  crossing the caustic of the Hubble Frontier Field (HFF) galaxy cluster MACS J0416 (Lotz et al. 2017) and (2) a large Ly $\alpha$   $EW_0$ , potentially implying that extreme stellar populations are present. We assume a flat cosmology with  $\Omega_M = 0.3$ ,  $\Omega_\Lambda = 0.7$ , and  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ .

## 2 THE MUSE DEEP LENSED FIELD: MDLF

VLT/MUSE (Bacon et al. 2010) deep spectroscopic observations of 17.1 h of integration time in a single pointing have been obtained on the HFF galaxy cluster MACS J0416 (Prog. 0100.A-0763 (A), PI Vanzella). The data reduction follows the technique described in Caminha et al. (2017), eventually achieving a point spread function (PSF) with full width at half-maximum (FWHM) of 0.6 arcsec in the final data cube. A more detailed description of the observational campaign of the MUSE Deep Lensed Field (MDLF) and of the data reduction will be presented elsewhere. A refined lens model of HFF J0416 using a new set of confirmed multiple images from the MDLF, will also be presented in a forthcoming paper.




**Figure 1.** A  $30 \text{ arcsec} \times 30 \text{ arcsec}$  region extracted from the MDLF at the  $z = 6.629$  Ly $\alpha$  wavelength and averaged over  $dv = 200 \text{ km s}^{-1}$ . The indicated Ly $\alpha$  arclet straddles the critical line (marked with the blue dotted line), close to a highly magnified system already confirmed at  $z = 6.145$  and well constrained by the three giant Ly $\alpha$  arcs (black contours). The one-dimensional spectrum of the Ly $\alpha$  line at  $z = 6.629$  is shown in the inset, where the asymmetry towards the red side is evident (a Gaussian with  $\text{FWHM} = 100 \text{ km s}^{-1}$  is superimposed with a blue line).

### 2.1 An Ly $\alpha$ arc at $z = 6.629$ and the faint *HST* counterpart

Fig. 1 shows the extended (3 arcsec) arc from the continuum subtracted narrow band image extracted from the MUSE data cube and the one-dimensional profile of the emission line at  $\lambda = 9270.7$  Å, in a region free from OH sky emission lines. The arc is detected at  $S/N = 18$  with flux  $4.4 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2}$  calculated within an elliptical aperture with major and minor axes of 4 and 1.5 arcsec, respectively, and shows an asymmetric profile having an instrumental corrected FWHM of  $98 (\pm 7) \text{ km s}^{-1}$ . We identify this line as Ly $\alpha$  at  $z = 6.629$  for the following reasons: (1) the weighted skewness  $S_W$  (as defined by Shimasaku et al. 2006) is  $3.4 \pm 0.7$ , in line with the typical values observed for asymmetric Ly $\alpha$  emissions at high- $z$  (it is zero for symmetric shapes, Fig. 1); (2) if it is identified to other typical lines like [O II] $\lambda 3727$ , 3729, [O III] $\lambda 4959$ , [O III] $\lambda 5007$ , H $\beta$ , or H $\alpha$ , each of them would imply the presence and detection of additional lines in the same spectrum; (3) the MUSE spectral resolution at  $\lambda > 9000$  Å is  $R \simeq 3500$ , high enough to resolve the single components of the doublets like C IV $\lambda 1548$ , 1550, O III] $\lambda 1661$ , 1666, C III] $\lambda \lambda 1907$ , 1909, and [O II] $\lambda 3727$ , 3729 further excluding these lines for identification. As Fig. 2 shows, there is no clear detection in the F105W (Y), F125W (J), F140W (JH), and F160W (H) bands in the HFF images, probing the ultraviolet stellar continuum down to the nominal depth of the HFFs (mag  $\simeq 29$ , at  $5\sigma$  limit; Lotz et al. 2017). We therefore computed the  $Y + J + JH + H$  weighted-mean-stacked images (probing  $\lambda \sim 1500$  Å), reaching a  $1\sigma$  limit of 31.6 within circular apertures of diameter 0.4 arcsec. Such a limit has been derived by inserting 30 apertures in free regions surrounding the source position and computing the standard deviation among them (the A-PHOT tool has been used for this task, Merlin et al. 2019). As discussed in the next section, there is a configuration in that we expect the presence of two very close multiple images near the





**Figure 2.** From left to right: the MUSE Ly $\alpha$  emission averaged over  $dv = 160 \text{ km s}^{-1}$  (the  $2\sigma$  and  $4\sigma$  contours are shown, with the dotted line marking the critical line); the Y-band *HST* image; the stacked Y + J + JH + H and J + JH + H images with indicated the positions of the two possible high- $z$  counterparts (solid circles with diameter 0.4 arcsec); the colour stacked image; the differential stacked *HST* images highlighting the two possible counterparts (A–C) and (B–C) after a Gaussian smoothing with  $\sigma = 1$  pix; the B + V + I stacked image (C).


Ly $\alpha$  arc. Indeed, there is a possible detection at  $S/N \sim 2$  lying within the arclet indicated as 1 in Fig. 2, with  $m_{1500} \simeq 31$  and showing a photometric drop in the F435W + F606W + F814W image. The same test has been performed adopting an elliptical aperture oriented along the arc, with semi-axis 0.7 and 0.2 arcsec, and no signal has been detected down to  $m_{1500} \simeq 30.85$  at  $1\sigma$ . We expect a second nearby image with similar magnification that, however, is contaminated by a foreground object clearly detected in the blue bands. While image 1 could be the *HST* counterpart, a tentative second image detection marked as 2 is shown in Fig. 2.

### 3 THE Ly $\alpha$ EMISSION IS ON THE CAUSTIC

The Ly $\alpha$  arc lies in a well-known region of the galaxy cluster where Vanzella et al. (2017, 2019) already discussed another star-forming complex system at  $z = 6.145$  showing several multiple images identified in deep *HST* data, producing three clear Ly $\alpha$  arcs in the MUSE observations (Fig. 1). The presence of such a system adds valuable constraints for the case studied in this work. In fact, any detection at  $z > 6$  in the region where the  $z = 6.629$  arc lies, would produce multiple images as in the case observed at  $z = 6.145$ , unless such images are so close to merge into a single spatially unresolved mildly elongated arc. It is exactly the case for the  $z = 6.629$  arclet discussed here: the absence of two distinct images (Figs 1 and 2) implies that the Ly $\alpha$  arc straddles the critical line and is indeed the result of two spatially unresolved Ly $\alpha$  images, generated by an Ly $\alpha$  emitting region lying on the corresponding caustic.

#### 3.1 Simulating the caustic crossing

In order to perform a quantitative estimate of the magnification of the Ly $\alpha$  emission, we use dedicated simulations with a customized version of the software SKYLENS (e.g. Meneghetti et al. 2010; Plazas et al. 2019). The method will be extensively described in a future paper. In short, we perform the following steps. We start from the assumption that the source of the Ly $\alpha$  emission can be described by a single, circularly symmetric Sérsic surface brightness profile. The profile is characterized by the Sérsic index  $n$  and by the effective radius  $R_e$ . We ray trace  $2000 \times 2000$  light rays through a squared region containing the Ly $\alpha$  emission (cyan square in Fig. 3, whose size is  $\sim 6.2 \text{ arcsec} \times 6.2 \text{ arcsec}$ ) and propagate them towards the source plane at redshift  $z_s = 6.629$ , accounting for the deflections induced by the lensing cluster. In this work, we use the strong lensing model described in Caminha et al. (2017).<sup>1</sup> On the source



**Figure 3.** The tangential caustic (orange) and the critical line (white) are superimposed to the (false-colour) image of the galaxy cluster members used for the strong lensing model. The observed Ly $\alpha$  arc is well reproduced when the source is very close to the caustic (black arrow). Three multiple images are predicted (marked as a, b, and c): images b and c merge into a single small arc, whereas image a is not detected (bottom left inset). In the top panels, the observed, simulated, and residuals Ly $\alpha$  images are shown for the best-fitting values of  $n$ ,  $R_e$ , and  $\mu = \mu(a) + \mu(b)$ .

plane, the arrival positions of the light rays are used to sample the brightness distribution of the source, which is then mapped on to the image plane. By performing this operation, we reconstruct the lensed image, which is subsequently convolved with a Gaussian PSF with an FWHM of 0.6 arcsec. The resulting convolved image is first rebinned at the same resolution of the MUSE image ( $0.2 \text{ arcsec pixel}^{-1}$ ) and then compared to the observed Ly $\alpha$  arclet. We use the python package LMFIT to perform a Non-linear Least Square Minimization of a cost function defined as the squared difference between the simulated and the observed images. As shown in the upper panels of Fig. 3, the source model, despite its simplicity, reproduces very well the observed arclet morphology. The two parameters  $n$  and  $R_e$  are quite degenerate. For a Sérsic index in the range  $n = 1-4$ , the best-fitting effective radius  $R_e$  varies between

<sup>1</sup>The Lenstool model is published in the HFF lens model format at <http://www.fe.infn.it/astro/lensing/>.

**Table 1.** Properties of the Ly  $\alpha$  emitter in the source plane.

Ly $\alpha$ (erg s <sup>-1</sup> cm <sup>-2</sup> )	$5.5 \times 10^{-20}$
Ly $\alpha$ (erg s <sup>-1</sup> )	$2.8 \times 10^{40}$
EW <sub>0</sub> (Ly $\alpha$ ) (Å) ( $T_{\text{IGM}} < 0.5$ )	$> 1120$
$M_{1500}(m_{1500})$ ( $2\sigma$ )	$\gtrsim -11.9$ ( $\gtrsim 35$ )
$R_e$ Ly $\alpha$ region (pc)	$< 150$
Magnification [ $\mu(b) = \mu(c)$ ]	$\mu(b) + \mu(c) \gtrsim 80$ ; $\mu(a) \simeq 4.5$

0.012 and 0.027 arcsec. The best-fitting solution corresponds to  $n = 3.66$  and  $R_e = 0.026$  arcsec. The latest is equivalent to 140 pc on the source plane. In all cases, the model suggests that the Ly  $\alpha$  emitting region partially overlaps with the caustic, meaning that only part of the source is reproduced twice in the arclet.

Having obtained a model for the Ly  $\alpha$  source, we can estimate the magnification of the arclet by comparing the lensed and the intrinsic fluxes of the source after integrating the surface brightness over the region used for ray tracing. We estimate that the total magnification of the Ly  $\alpha$  arclet (images b and c) is  $\mu(b + c) = \mu(b) + \mu(c) \simeq 80$ , implying that the de-lensed Ly  $\alpha$  flux of the arclet is  $\sim 5.5 \times 10^{-20}$  erg s<sup>-1</sup> cm<sup>-2</sup>. Keeping  $n \sim 1$  (exponential profile) in our fitting procedure the magnification increases to  $\mu(b + c) \sim 110$ .

It is well known that such magnification estimate is prone to systematic errors due to the uncertainties of the lens model in regions where the magnification gradients are very strong (e.g. Meneghetti et al. 2017). In order to circumvent this limitation, we can make use of the counter image a, which is predicted to form much farther from the critical lines (Fig. 3). Using our best-fitting source determined above, we estimate that  $\mu(a) \sim 4.5$  with an uncertainty smaller than 20 per cent. We also find that, because of the much shallower magnification gradient in this region, the estimate is quite insensitive to the properties of the source. Indeed,  $\mu(a)$  would change by  $\sim 5$  per cent [ $\mu(a) \sim 4.3$ ] by adopting a point source approximation. The image a is not detected at the depth of the MDLF down to  $1\sigma$  Ly  $\alpha$  flux limit  $f_{a,\text{lim}}$  ( $\sim 2.4 \times 10^{-19}$  erg s<sup>-1</sup> cm<sup>-2</sup>, for a point-like source, consistently to Inami et al. 2017). This sets a lower limit  $\mu(b + c) \gtrsim \mu(a)f_{bc}/f_{a,\text{lim}}$ , where the observed ratio is  $f_{bc}/f_{a,\text{lim}} \simeq 18$ . Therefore, we can conclude that  $\mu(b + c) \gtrsim 80$ , in keeping with the fitting procedure ( $\mu = 80\text{--}110$ ).

### 3.2 A large Ly $\alpha$ EW

The computation of the EW of the Ly  $\alpha$  line ( $L$ ) requires an estimate of the underlying stellar continuum ( $S$ ), taking into account that magnifications associated with  $S$  ( $\mu_S$ ) and  $L$  ( $\mu_L$ ) might differ. A general expression for EW<sub>0</sub> is

$$\text{EW}_0 = \frac{1}{(1+z)} \frac{\mu_S}{\mu_L} \frac{f(\text{Ly } \alpha)}{F_\lambda(\text{UV}) \times T_{\text{IGM}}(\text{Ly } \alpha)}, \quad (1)$$

where  $f(\text{Ly } \alpha)$  is the Ly  $\alpha$  flux ( $5.5 \times 10^{-20}$  erg s<sup>-1</sup> cm<sup>-2</sup>),  $F_\lambda(\text{UV})$  is the ultraviolet continuum at the Ly  $\alpha$  wavelength for that we assume a value  $< 1.27 \times 10^{-23}$  erg s<sup>-1</sup> cm<sup>-2</sup> Å<sup>-1</sup>, corresponding to  $m_{1500} \gtrsim 35$  [ $= 31 + 2.5 \log_{10}(\mu)$ ,  $\mu \gtrsim 40$ ]. Given the large uncertainties on the *HST* detection ( $S/N \sim 2$ ,  $\sigma_m \sim 0.5$ ), any assumption on the ultraviolet slope  $\beta$  would not be significant. Indeed, a slope  $\beta = -2.5$  ( $-3$ ) would imply a magnitude difference of  $m_{1216} - m_{1500} = -0.11$  ( $-0.23$ ).  $T_{\text{IGM}}(\text{Ly } \alpha)$  is the transmission of the intergalactic medium for Ly  $\alpha$  photons (see below, and Table 1). We identify two scenarios

(i) if we assume that  $L$  and  $S$  have the same intrinsic size and brightness profile, then  $\mu_S \simeq \mu_L$  and the EW<sub>0</sub> is independent on

the lens model. Thus, under this assumption, a lower limit on EW<sub>0</sub> can be found by using the very low significance detection (if not the non-detection) of the UV continuum,  $m_{1500} \gtrsim 35$ . Combined with the de-lensed Ly  $\alpha$  flux of  $5.5 \times 10^{-20}$  erg s<sup>-1</sup> cm<sup>-2</sup> leads to the result that EW<sub>0</sub>  $> 564$  Å, in the case  $T_{\text{IGM}}(\text{Ly } \alpha) = 1.0$ . The low significance of the *HST* detection prevents us from verifying whether  $S$  is extended as the arclet.

(ii) if, in contrast, we assume that the size of  $S$  is smaller than that of  $L$ , as it might be the case when  $S$  is embedded and generates the H II region (e.g. Steidel et al. 2011), then we expect that  $\mu_S > \mu_L$  and consequently EW<sub>0</sub> would be even larger than in the previous case.

There is still the possibility that  $S$  is located outside the lens caustic. In this scenario, no continuum flux is expected near the Ly  $\alpha$  arclet. We could only use the non-detection in the *HST* data of image a ( $m > 31.6$  at  $1\sigma$ ), where  $S$  is certainly present, to set an upper limit of  $F_\lambda(\text{UV})$ . Using the fact that  $\mu(a) \sim 4.5$ , the de-lensed magnitude limit is  $m > 33.2$ . Combined with the lower limit of the magnification of  $L$  ( $\mu_L \gtrsim 80$ ), we obtain that EW<sub>0</sub>  $\gtrsim 110$  Å in the case  $T_{\text{IGM}}(\text{Ly } \alpha) = 1.0$ . Note, however, that the marginal detection of image 1 in the stacked *HST* images (see Fig. 2), which could be the image of  $S$ , seems to disfavour this scenario.

It is now worth discussing the IGM transmission  $T_{\text{IGM}}(\text{Ly } \alpha)$  that depends on both the ‘intrinsic’ (pre-IGM) Ly  $\alpha$  spectrum emerging and the IGM properties. Due to the resonant absorption of the neutral or partially neutral IGM combined with cosmological inflow, the Ly  $\alpha$  spectrum blueward of  $v_{\text{cut-off}} \lesssim 200$  km s<sup>-1</sup> is absorbed (Dijkstra & Wyithe 2007). Thus, the more asymmetric the intrinsic Ly  $\alpha$  is towards the blue or the lower the red peak offset from systemic, the lower  $T_{\text{IGM}}$ . All this implies that  $T_{\text{IGM}}$  is highly uncertain and estimates reach from tens of per cent (e.g.  $T_{\text{IGM}} = 0.20^{+0.12}_{-0.15}$  at  $z = 6.6$ , Laursen, Sommer-Larsen & Razoumov 2011) to values approaching unity for an intrinsic single read peak with offset  $> 300$  km s<sup>-1</sup>. However, since the Ly  $\alpha$  spectrum presented here is very narrow and asymmetric (cf. Section 2.1) it is likely that either a significant part of the red peak has been removed from the IGM (in case of an intrinsic spectrum with a large offset, and thus, large width), or an intrinsic blue component existed (in the case of a small intrinsic offset). Both cases would imply a significant absorption of the IGM, and hence,  $T_{\text{IGM}} < 1$ . Knowledge of the systemic redshift, through, e.g. H  $\alpha$  information would be helpful in reconstructing the intrinsic Ly  $\alpha$  line, and thus, to constrain  $T_{\text{IGM}}$  more qualitatively. This could, furthermore, rule out the radiative transfer effects as an origin of the large EW – that we already deem unlikely due to the asymmetry of the observed line.

We conclude that, even assuming  $T_{\text{IGM}}(\text{Ly } \alpha) = 1$ , a still quite extreme EW<sub>0</sub>(Ly  $\alpha$ )  $> 550$  Å emerges from a region crossing the caustic, that can easily approach (or exceed) 1000 Å assuming a more plausible  $T_{\text{IGM}}(\text{Ly } \alpha) < 1.0$ . Though not totally excluded, we do not consider in this work the possibilities that the large EW(Ly  $\alpha$ ) originates from a very faint AGN (e.g. with BH mass of  $10^{2-3}$  M<sub>⊙</sub>, Fan 2012) or a multiphase scattering medium (e.g. Neufeld 1991).

### 4 CANDIDATE Pop III STARS

The predicted EW<sub>0</sub>(Ly  $\alpha$ ) for metal-free stellar populations exceeds 400 Å and it goes up to a few thousands rest frame (Inoue 2011; Schaerer 2013), and is observable if neighbouring sources (either Pop III or Pop II stars) have already contributed towards ionizing a local bubble (Sobral et al. 2015). The large EW<sub>0</sub>(Ly  $\alpha$ ) value



reported in this work opens for a possible dominant contribution by extremely metal-poor stars. It is interesting to calculate how many Pop III stars are needed to reproduce both the observed  $M_{1500}$  and  $L(\text{Ly}\alpha)$ :

(i) *UV continuum*: The apparent magnitudes at 1500 Å rest frame at  $z = 7$  for Pop III star at ZAMS with masses 1–1000  $M_{\odot}$  are reported by Windhorst et al. (2018). In particular, stars with masses of 100, 300, and 1000  $M_{\odot}$  have magnitudes  $m_{1500} = 40.08, 38.64,$  and  $37.44$ , respectively, neglecting dust attenuation. Adopting  $m_{1500} \gtrsim 35$  ( $M_{\text{UV}} \gtrsim -11.9$ , Section 3.2) and assuming for simplicity the same masses for all stars, the number of Pop III stars required to reproduce the intrinsic UV flux corresponding to  $M_{\text{UV}}$  amounts to  $N(M_{\text{UV}}) = 10, 30,$  and  $110$  for stellar masses of 1000, 300, and 100  $M_{\odot}$ .

(ii) *Ly $\alpha$  emission*: Mas-Ribas, Dijkstra & Forero-Romero (2016) provide the photon flux  $Q[\text{H I}]$  ( $\text{s}^{-1}$ ) for different Pop III ZAMS stars and the conversion to  $L(\text{Ly}\alpha)$  luminosity considering case-B departure, stochastic sampling of the Salpeter, and Top-Heavy initial mass functions (IMFs) and zero escape fraction of the Lyman continuum radiation (note, however, that if a fraction of the ionizing radiation escapes, the emerging Ly $\alpha$  would be dimmed linearly by the same factor, e.g. Schaerer 2013). We perform the calculation as above, assuming again the same mass for all stars (no boosting from the stochastic sampling of the IMF is considered). The Ly $\alpha$  luminosity emerging from Pop-III stars of mass 1000, 300, and 100  $M_{\odot}$  are  $3.20 \times 10^{40}, 8.11 \times 10^{39},$  and  $1.72 \times 10^{39} \text{ erg s}^{-1}$ . Under the assumption that  $N(\text{Ly}\alpha) = N(M_{\text{UV}})$  (being Ly $\alpha$  and  $M_{\text{UV}}$  referring to the same star complex), the resulting  $T_{\text{IGM}}(\text{Ly}\alpha)$  are 0.09, 0.11, and 0.16, respectively. These values double if the case B is assumed, i.e. the predicted  $L(\text{Ly}\alpha)$  is about a factor two fainter (Mas-Ribas et al. 2016). With such values of  $T_{\text{IGM}}(\text{Ly}\alpha)$  the resulting  $\text{EW}_0(\text{Ly}\alpha)$  ranges between 4000–1500 Å, for the three classes of Pop III stellar masses.

Future facilities are necessary to make a significant step forward. First, only the *James Webb Space Telescope* will access the optical rest frame looking for the possible deficit of metals and the expected enormous Balmer emissions (e.g. Inoue 2011), eventually gaining in depth with respect *HST* imaging. The next generation of Extremely Large Telescopes will also investigate the currently vague stellar component  $S$  by performing very deep imaging, while spectroscopy will address the deficiency of high-ionization metal lines and the possible key He II $\lambda$ 1640 emission. The intrinsic He II $\lambda$ 1640/Ly $\alpha$  line ratio predicted for Pop III spans the range 0.01–0.10 (e.g. Schaerer 2013; Mas-Ribas et al. 2016), implying the expected flux of He II $\lambda$ 1640 would be  $1.1 \times (10^{-21} - 10^{-20})/T_{\text{IGM}}(\text{Ly}\alpha) \text{ erg s}^{-1} \text{ cm}^{-2}$ , clearly requiring an ELT-like telescope or an 8–10 m class telescope in the most optimistic cases [ $T_{\text{IGM}}(\text{Ly}\alpha) \ll 1$ ].

## ACKNOWLEDGEMENTS

We thank the anonymous referee for a constructive report. We thank L.I. Mas-Ribas and D. Schaerer for very stimulating discussions.

This work is supported by PRIN-MIUR 2017 WSCC32. We acknowledge funding from the INAF main-stream (1.05.01.86.31). KC and GBC acknowledge funding from the ERC through the award of the Consolidator Grant ID 681627-BUILDUP. This work was supported in part by the NSF grant: COLLABORATIVE RESEARCH: The Final Frontier: Spectroscopic Probes of Galaxies at the Epoch of Reionization (AST-1815458, AST-1810822). MG acknowledges funding through HST-HF2-51409.

## REFERENCES

- Bacon R. et al., 2010, in McLean I. S., Ramsay S. K., Takami H., eds, Proc. SPIE Conf. Ser. Vol. 7735, Ground-based and Airborne Instrumentation for Astronomy III. SPIE, Bellingham, p. 773508
- Caminha G. B. et al., 2017, *A&A*, 600, A90
- Dayal P., Ferrara A., 2018, Phys. Rep., 780, 1
- Dijkstra M., Wyithe J. S. B., 2007, *MNRAS*, 379, 1589
- Fan X., 2012, *Res. Astron. Astrophys.*, 12, 865
- Hernán-Caballero A. et al., 2017, *ApJ*, 849, 82
- Inami H. et al., 2017, *A&A*, 608, A2
- Inoue A. K., 2011, *MNRAS*, 415, 2920
- Kashikawa N. et al., 2012, *ApJ*, 761, 85
- Laursen P., Sommer-Larsen J., Razoumov A. O., 2011, *ApJ*, 728, 52
- Lotz J. M. et al., 2017, *ApJ*, 837, 97
- Mas-Ribas L., Dijkstra M., Forero-Romero J. E., 2016, *ApJ*, 833, 65
- Maseda M. V. et al., 2018, *ApJ*, 865, L1
- Meneghetti M., Rasia E., Merten J., Bellagamba F., Ettori S., Mazzotta P., Dolag K., Marri S., 2010, *A&A*, 514, A93
- Meneghetti M. et al., 2017, *MNRAS*, 472, 3177
- Merlin E., Pilo S., Fontana A., Castellano M., Paris D., Roscani V., Santini P., Torelli M., 2019, *A&A*, 622, A169
- Neufeld D. A., 1991, *ApJ*, 370, L85
- Plazas A. A., Meneghetti M., Maturi M., Rhodes J., 2019, *MNRAS*, 482, 2823
- Rodney S. A. et al., 2018, *Nat. Astron.*, 2, 324
- Salvaterra R., Ferrara A., Dayal P., 2011, *MNRAS*, 414, 847
- Schaerer D., 2013, in Wiklind T., Mobasher B., Bromm V., eds, *Astrophysics and Space Science Library*, Vol. 396, The First Galaxies. Springer-Verlag, Berlin, p. 345
- Shibuya T. et al., 2018, *PASJ*, 70, S15
- Shimasaku K. et al., 2006, *PASJ*, 58, 313
- Sobral D., Matthee J., Darvish B., Schaerer D., Mobasher B., Röttgering H. J. A., Santos S., Hemmati S., 2015, *ApJ*, 808, 139
- Sobral D. et al., 2019, *MNRAS*, 482, 2422
- Steidel C. C., Bogosavljević M., Shapley A. E., Kollmeier J. A., Reddy N. A., Erb D. K., Pettini M., 2011, *ApJ*, 736, 160
- Tornatore L., Ferrara A., Schneider R., 2007, *MNRAS*, 382, 945
- Vanzella E. et al., 2017, *MNRAS*, 467, 4304
- Vanzella E. et al., 2019, *MNRAS*, 483, 3618
- Visbal E., Haiman Z., Bryan G. L., 2016, *MNRAS*, 460, L59
- Windhorst R. A. et al., 2018, *ApJS*, 234, 41
- Wise J. H., 2019, *Contemp. Phys.*, 60, 145
- Zackrisson E., Vikaeus A., 2019, preprint (arXiv:1903.12555)
- Zackrisson E., González J., Eriksson S., Asadi S., Safranek-Shrader C., Trenti M., Inoue A. K., 2015, *MNRAS*, 449, 3057

This paper has been typeset from a  $\text{\LaTeX}$  file prepared by the author.