



Publication Year	2007
Acceptance in OA @INAF	2024-02-12T15:06:00Z
Title	Prominent activity of the blazar OJ 287 in 2005. XMM-Newton and multiwavelength observations
Authors	Ciprini, S.; RAITERI, Claudia Maria; Rizzi, N.; Agudo, I.; FOSCHINI, LUIGI; et al.
Handle	http://hdl.handle.net/20.500.12386/34745
Journal	MEMORIE DELLA SOCIETA ASTRONOMICA ITALIANA



Prominent activity of the blazar OJ 287 in 2005. XMM-Newton and multiwavelength observations

S. Ciprini^{1,2},

C.M. Raiteri³, N. Rizzi^{4,1}, I. Agudo^{5,6}, L. Foschini⁷, M. Fiorucci⁸, L.O. Takalo², M. Villata³,
L. Ostorero^{9,10}, A. Sillanpää², M. Valtonen², G. Tosti¹, S.J. Wagner¹⁰, H. Aller¹¹, M.F. Aller¹¹, A. Arai¹²,
A.A. Arkharov¹³, U. Bach^{6,3}, V. Bakis¹⁴, M. Bagaglia⁸, M. Böttcher¹⁵, C. Buemi¹⁶, D. Carosati¹⁷,
W.P. Chen¹⁸, Y. Efimov¹⁹, D. Emmanoulopoulos¹⁰, A. Erdem¹⁴, L. Fuhrmann⁶, A. Frasca³¹, M. Fullhart¹⁵,
A. Goyal²⁰, J. Heidt¹⁰, T. Hovatta²¹, F. Hroch²², M.A. Ibrahimov²³, L. Jilková²², M. Joshi¹⁵, M. Kamada¹²,
M. Katsuura¹², D. Kinoshita¹⁸, A. Kostov²⁴, D. Kotaka¹², Y.Y. Kovalev²⁵, T. Krejčová²², T. Krichbaum⁶,
G. Krishna²⁶, M. Kurosaki¹², O. Kurtanidze²⁷, A. Lahteenmaki²¹, L. Lanteri³, V. Larionov²⁸, C.-U. Lee²⁹,
P. Leto¹⁶, J. Li³⁰, E. Lindfors², F. Münz²², E. Marilli³¹, Y. Matsubara¹², S. Mizoguchi¹², S. Mondal¹⁸,
K. Nakamura¹², E. Nieppola²¹, K. Nilsson², S. Nishiyama¹², G. Nucciarelli⁸, A. Ogino¹², J. Ohlert³²,
A. Oksanen³³, E. Ovcharov³⁴, S. Pak²⁹, M. Pasanen², C. Pullen³⁵, T. Pursimo³⁶, J.A. Ros³⁷, K. Sadakane¹²,
A.C. Sadun³⁸, R. Sagar²⁰, B.-W. Sohn²⁹, N. Sumitomo¹², K. Tanaka¹², C. Trigilio¹⁶, I. Tornaiainen²¹,
M. Tornikoski²¹, G. Umana¹⁶, H. Ungerechts³⁹, E. Valtaoja², A. Volvach⁴⁰, J.R. Webb⁴¹, J. Wu⁴²,
H.-S. Yim²⁹, Y. Zhang³⁰

Abstract. Two guest-observer XMM-Newton pointings of the blazar OJ 287 in 2005 are introduced, along with part of the radio, mm, near-IR, and optical data obtained during a coordinated and intensive WEBT campaign, during longer-term monitoring observations performed by teams of the ENIGMA network, and during other independent observing programs (like VLBA observations). In that year OJ 287 showed an interesting variable behavior in the optical band. An optical outburst, well matched by our WEBT observations, is claimed in the period Oct.-Nov. 2005, and the XMM-Newton X-ray observations are performed in correspondence with two active optical states (an intermediate flare and such outburst). X-ray data indicates different flux levels, spectral slopes, and emission components, and VLBA radio maps are consistent with a jet precession model. This appreciable observing effort is still ongoing (a further XMM-Newton pointing is planned in 2008), joined with further parallel/multi-monitoring observing programmes devoted to this interesting object.

Key words. BL Lacertae objects: general – BL Lacertae objects: individual: OJ 287 – quasars: individual: OJ 287 – X-rays: galaxies – radio continuum: galaxies – galaxies: jets – galaxies: photometry – radiation mechanisms: non-thermal

Send offprint requests to: S. Ciprini
e-mail: stefano.ciprini@pg.infn.it

1) Physics Dept., University of Perugia, & INFN Perugia, via Pascoli, I-06123 Perugia, Italy. 2) Tuorla Observatory, University of Turku, Väisälantie 20, 21500 Piikkiö, Finland. 3) INAF, Osservatorio Astronomico di Torino, I-10025 Pino Torinese, Torino, Italy. 4) Osservatorio Astronomico Sirio, Grotte di Castellana, Piazzale Anelli, I-70013, Bari, Italy. 5) Instituto de Astrofísica de Andalucía, CSIC, E-18008 Granada, Spain. 6) Max-Planck-Institut fuer Radioastronomie, Auf dem Huegel 69, D-53121, Bonn, Germany. 7) INAF, Istituto di Astrofisica Spaziale e Fisica Cosmica, I-40129 Bologna, Italy. 8) Physics Dept.,

University of Perugia, via Pascoli, I-06123 Perugia, Italy. 9) Dipartimento di Fisica Generale, University of Torino, I-10125 Torino, Italy. 10) Landessternwarte Heidelberg-Königstuhl, D-69117, Heidelberg, Germany. 11) Department of Astronomy, Dennison Building, University of Michigan, Ann Arbor, MI 48109, USA. 12) Astronomical Institute, Osaka Kyoiku University, Kashiwara, Osaka, Japan. 13) Pulkovo Astronomical Observatory, Russian Academy of St. Petersburg, Russia. 14) Çanakkale Onsekiz Mart University Observatory, Terzioğlu Kampüsü, 17040 Çanakkale, Turkey. 15) Department of Physics and Astronomy, Clippinger Hall, 251B Ohio University Athens, OH 45701-2979, USA. 16) INAF, Istituto di Radioastronomia, Sezione di Noto, 96017 Noto, Siracusa, Italy. 17) Armenzano Astronomical




Fig. 1. Optical (R -band magnitude) light curve obtained during the coordinated, intensive and short MW campaign by the Whole Earth Blazar Telescope (WEBT) consortium, and during longer-term monitoring observations (ENIGMA network observatories, plus other independent programmes and observatories), requested or triggered by the 2 XMM-Newton pointings in 2005. This light curve covers two optical observing seasons for the blazar (Oct.2004 - Mar.2006). The XMM-Newton space observatory pointed OJ 287 twice (Apr.12 and Nov.3-4, 2005, vertical lines in the figs.) following our granted GO AO-4 proposal. X-ray observations were coincident with two main active stages in the optical band (a mild flare in Apr. and an outburst in Oct.-Nov.). The second X-ray observation was performed during an optical enduring (~ 20 days) and time-structured (3 major symmetric wiggles) outburst ($R_{max} \sim 13.3$). The whole dataset of radio, mm, near-IR, and multiband optical observations has been collected by more than 30 ground-based observatories. The intensive coordinated WEBT campaign covers a period of about 1 week around the first XMM-Newton pointing in Apr.2005, and about 1 month around the second pointing of Nov.2005.

1. The OJ 287 campaign in 2005

Observatory, Assisi, Perugia, Italy. 18) Institute of Astronomy and Dept. of Physics, National Central University, Jungli City 320-54, Taoyuan, Taiwan, ROC. 19) Crimean Astrophysical Observatory, Nauchny, 98409 Crimea, Ukraine. 20) Aryabhata Research Institute of Observational Sciences, Manora Peak, Nainital 263 129, India. 21) Metsähovi Radio Observatory, Helsinki Univ. of Technology, 02540 Kylmäla, Finland. 22) Institute of Theoretical Physics and Astrophysics, Masaryk University, 611 37 Brno, Czech Republic. 23) Ulugh Beg Astronomical Inst., Academy of Sciences of Uzbekistan, Tashkent 700052, Uzbekistan. 24) Institute of Astronomy, Bulgarian Academy of Sciences, 1784 Sofia, Bulgaria. 25) National Radio Astronomy Observatory, Green Bank, WV 24944, USA. 26) National Centre for Radio Astronomy, TIFR, Ganeshkhind, Pune 411007, India. 27) Abastumani Observatory, 383762 Abastumani, Georgia. 28) Astronomical Institute, St. Petersburg State University, Universitetsky pr. 28, Petrodvorets, 198504 St. Petersburg, Russia. 29) Korea Astronomy and Space Science Institute, Daejeon 305-348, Republic of Korea. 30) Department of Physics, Tsinghua University, Beijing 100084, China. 31) INAF, Osservatorio Astrofisico di Catania, 95125 Catania, Italy. 32) Michael Adrian Observatory, 65468 Trebur, Germany. 33) Nyrölä Observatory, Jyväskylän Sirius ry, 40950 Jyväskylä, Finland. 34) Department of Astronomy, Faculty of Physics, Sofia University, 1164, Sofia, Bulgaria. 35) Coyote Hill Observatory, Wilton, Sacramento, CA 95693, USA. 36) Nordic Optical Telescope, 38700 Santa Cruz de La Palma, Spain. 37) Agrupació Astronòmica de Sabadell, 08200 Sabadell, Spain. 38) Department of Physics, University of Colorado at Denver and Health Sciences Center, Denver, CO 80217-3364, USA. 39) IRAM, Avd. Div. Pastora 7NC, 18012 Granada, Spain. 40) Laboratory of Radioastronomy of Crimean Astrophysical Observatory RT-22, Katsively, Yalta, 334247 Crimea, Ukraine. 41) SARA Observatory, Florida International Univ., Miami, FL 33199, USA. 42) National Astronomical Observatories, Datun Road, Chaoyang District, Beijing, China.

The high-energy and variable flux, the broad-band synchrotron and inverse Compton (possibly connected) emission components, make blazars ideal targets for multiwavelength (MW) campaigns, involving both space-borne observatories and ground-based facilities. In particular the blazar OJ 287 (PG 0851+202, 3EG J0853+1941, $z = 0.306$) is an optically highly-variable (> 3 mag variations) and low/intermediate-energy peaked BL Lac object (LBL/IBL), interesting mainly for a twofold reason. First, this source is historically among the best observed AGN in the optical/radio bands, having a very good database of observations (see Pursimo et al. 2000; Takalo 1994, for a review of earlier observations), hence allowing a more significant statistical investigation, and a wider study of the parameter space of variability (Wagner 2007). Secondly it is one of the very few extragalactic sources, where a major periodical or quasi-periodical signature is claimed (e.g., Valtonen 2007; Nilsson et al. 2006), and the search for supermassive binary black holes should become a major research field in the next years.




Fig. 3. 22 GHz VLBA image of OJ 287 obtained in Apr.02, 2005. Contours represent the total intensity, the colour scale the polarized intensity, and the superimposed sticks the orientation of the polarization electric vectors. The positions of the fitted Gaussian components are indicated by the crosses, whereas the circles (of radius equal to the FWHM of each Gaussian) symbolize their size. The 22 GHz image convolving beam is $(1.05 \times 0.41) \text{ mas}^2$, with major axis position angle at -12° .

in Oct.-Nov. 2005; (ii) a broken power-law X-ray spectrum (i.e. a two-component, synchrotron + IC spectrum, typical signature of intermediate blazars, or a thermal tail end) discovered by the EPIC instrument on Nov.3-4; (iii) a clear frequency dependence of the mean structural position angle of the radio-jet in VLBA maps, consistent with a ballistic jet precession model, and a polarization structure mostly concentrated on the emission core. A more detailed analysis will be available in (Ciprini et al. 2007).

Acknowledgements. Based on observations obtained with XMM-Newton, an ESA science mission with instruments and contributions directly funded by ESA Member States and NASA. The teams belonged to the former Research Training Network ENIGMA, acknowledge the funding by European




Fig. 4. XMM-Newton combined EPIC (pn, MOS1 and MOS2 detectors) X-ray spectra of OJ 287, belonging to the two observations performed on Apr.12, and Nov.3-4, 2005. Upper panel: the Apr.12 spectrum can be described by a simple power-law + galactic absorption, $N_H = 3.09 \times 10^{20} \text{ cm}^{-2}$, $\Gamma = 1.63 \pm 0.02$, $\chi_r^2 = 1.03$ *d.o.f.* = 367, $F_{2-10\text{KeV}} = (2.5 \pm 0.8) \times 10^{-12} \text{ erg s}^{-1} \text{ cm}^{-2}$. Lower panel: the Nov.3-4 spectrum can be described by a broken power-law + galactic absorption, $N_H = 3.09 \times 10^{20} \text{ cm}^{-2}$, $\Gamma_1 = 2.65^{+0.12}_{-0.07}$, $\Gamma_2 = 1.79 \pm 0.02$, $E_{br} = 0.69^{+0.04}_{-0.05} \text{ keV}$, $\chi_r^2 = 1.03$ *d.o.f.* = 927, $F_{2-10\text{keV}} = (1.82 \pm 0.07) \times 10^{-12} \text{ erg s}^{-1} \text{ cm}^{-2}$.

Community's Human Potential Programme under contract HPRN-CT-2002-00321.

References

- Ciprini S., Raiteri C.M., Rizzi N., et al., in prep.
 Fuhrmann L., Ciprini S., Marchili N., et al., in prep
 Isobe N., Tashiro M., Sugiho M., & Makishima K. 2001, PASJ, 53, 79




Fig. 5. The overall spectral energy distribution (SED) of OJ 287 quasi-simultaneous to the first XMM-Newton observation of Apr.12, 2005 (OM and EPIC-pn spectral data reported), assembled with radio, mm, near-IR and optical data collected around the satellite pointing date (filled red/orange symbols). A synchrotron self-Compton (SSC) model for this epoch is reported (black line). The SSC fit attempt reported is a first approximation model (one-zone, homogeneous region with injection of a power law energy distribution of accelerated electrons, characterized by cooling and escape times), taking into account also the radio emission, and providing the following parameters: $\gamma_{min} = 135$, $\gamma_{max} = 6.2 \times 10^4$, $p = 1.8$, $k = 0.05$, $B = 0.4\text{Gauss}$, $R = 9.5 \times 10^{16}\text{cm}$, $\mathcal{D} = 6$, $z = 0.306$. In this date XMM-Newton data suggests a pure inverse-Compton origin for the X-ray emission. On the other hand data referring to the Nov.3-4, 2005 pointing (not reported here), suggests a different situation. More details can be found in Ciprini et al. (2007).

- Massaro E., Giommi P., Perri M. et al. 2003, A&A, 399,
 Nilsson K., Takalo L.O., Sillanpää A., & Ciprini S. 2006, ASP Conf. Ser. 350, 47
 Pursimo, T., Takalo, L.O., Sillanpää A., et al. 2000, A&AS, 146, 141
 Takalo, L.O., 1994, Vistas in Astron., 38, 77
 Valtonen, M. J. 2007, ApJ, 659, 1074
 Wagner, S. J., 2007, AIP Conf. Proc., in press