



Publication Year	2018
Acceptance in OA	2024-02-16T09:09:07Z
Title	THESEUS: A key space mission concept for Multi-Messenger Astrophysics
Authors	Stratta, G., CIOLFI, RICCARDO, AMATI, LORENZO, Bozzo, E., GHIRLANDA, Giancarlo, MAIORANO, Elisabetta, NICASTRO, LUCIANO, ROSSI, Andrea, Vinciguerra, S., Frontera, F., Götz, D., Guidorzi, C., O'Brien, P., Osborne, J. P., Tanvir, N., Branchesi, M., BROCATO, Enzo, Dainotti, M. G., De Pasquale, M., GRADO, ANIELLO, Greiner, J., Longo, F., Maio, U., Mereghetti, D., MIGNANI, Roberto, PIRANOMONTE, Silvia, Rezzolla, L., SALVATERRA, Ruben, Starling, R., Willingale, R., Böer, M., BULGARELLI, ANDREA, Caruana, J., Colafrancesco, S., Colpi, M., COVINO, Stefano, D'AVANZO, Paolo, D'Elia, V., Drago, A., FUSCHINO, FABIO, Gendre, B., Hudec, R., Jonker, P., LABANTI, CLAUDIO, Malesani, D., Mundell, C. G., PALAZZI, ELIANA, Patricelli, B., Razzano, M., CAMPANA, RICCARDO, Rosati, P., Rodic, T., Szécsi, D., STAMERRA, Antonio, van Putten, M., Vergani, S., Zhang, B., BERNARDINI, Maria Grazia
Publisher's version (DOI)	10.1016/j.asr.2018.04.013
Handle	http://hdl.handle.net/20.500.12386/34758
Journal	ADVANCES IN SPACE RESEARCH
Volume	62

THESEUS: a key space mission for Multi-Messenger Astrophysics

G. Stratta^a, R. Ciolfi^{b,c}, L. Amati^d, E. Bozzo^e, G. Ghirlanda^f, E. Maiorano^d, L. Nicastro^d, A. Rossi^d, S. Vinciguerra^g, F. Frontera^{h,d}, D. Gotzⁱ, C. Guidorzi^h, P. O'Brien^j, J. P. Osborne^j, N. Tanvir^k, M. Branchesi^{l,m}, E. Brocatoⁿ, M. G. Dainotti^o, M. De Pasquale^p, A. Grado^q, J. Greiner^r, F. Longo^{s,t}, U. Maio^{u,v}, R. Mignani^{w,x}, S. Piranomonte^y, L. Rezzolla^{z,aa}, R. Salvaterra^w, R. Starling^j, M. Boer^{ab}, A. Bulgarelli^d, J. Caruana^{ac}, S. Colafrancesco^{ad}, M. Colpi^{ae}, S. Covino^f, P. D'Avanzo^f, V. D'Elia^{af,y}, A. Drago^{ag}, F. Fuschino^d, B. Gendre^{ah,ai}, R. Hudec^{aj,ak}, P. Jonker^{al,am}, C. Labanti^d, D. Malesani^{an}, C. G. Mundell^{ao}, E. Palazzi^d, B. Patricelli^{ap}, M. Razzano^{ap}, C. Riccardo^d, P. Rosati^h, T. Rodic^{av}, D. Szécsi^{ar,as}, A. Stamerra^{ap}, M. van Putten^{ai}, S. Vergani^{au,f}, B. Zhang^{av}

^aUrbino University, via S. Chiara 27, 60129, Urbino (PU, Italy)

^bINAF, Osservatorio Astronomico di Padova, Vicolo dell' Osservatorio 5, I-35122 Padova, Italy

^cINFN-TIFPA, Trento Institute for Fundamental Physics and Applications, via Sommarive 14, I-38123 Trento, Italy

^dINAF-IASF Bologna, via P. Gobetti, 101, I-40129 Bologna, Italy

^eDepartment of Astronomy, University of Geneva, ch. d'Écogia 16, CH-1290 Versoix, Switzerland

^fINAF - Osservatorio astronomico di Brera, Via E. Bianchi 46, Merate (LC), I-23807, Italy

^gInstitute of Gravitational Wave Astronomy & School of Physics and Astronomy, University of Birmingham, Birmingham, B15 2TT, United Kingdom

^hDepartment of Physics and Earth Sciences, University of Ferrara, Via Saragat 1, I-44122 Ferrara, Italy

ⁱIRFU/Département d'Astrophysique, CEA, Université Paris-Saclay, F-91191, Gif-sur-Yvette, France

^jDepartment of Physics and Astronomy, University of Leicester, Leicester LE1 7RH, United Kingdom

^kUniversity of Leicester, Department of Physics and Astronomy and Leicester Institute of Space & Earth Observation, University Road, Leicester, LE1 7RH, United Kingdom

^lUniversità degli Studi di Urbino Carlo Bo, via A. Saffi 2, 61029, Urbino

^mINFN, Sezione di Firenze, via G. Sansone 1, 50019, Sesto Fiorentino, Italy

ⁿINAF - Astronomico di Teramo, Mentore Maggini s.n.c., 64100 Teramo, Italy

^oDepartment of Physics & Astronomy, Stanford University, Via Pueblo Mall 382, Stanford CA, 94305-4060, USA

^pDepartment of Astronomy and Space Sciences, Istanbul University, Beyazit, 34119, Istanbul, Turkey

^qINAF - Capodimonte Astronomical observatory Naples, Via Moiariello 16 I-80131, Naples, Italy

^rMax Planck Institute for Astrophysics, Karl-Schwarzschild-Str. 1, 85741 Garching, Germany

^sDepartment of Physics, University of Trieste, via Valerio 2, Trieste, Italy

^tINFN Trieste, via Valerio 2, Trieste, Italy

^uLeibniz Institut for Astrophysics, An der Sternwarte 16, 14482 Potsdam, Germany

^vINAF-Osservatorio Astronomico di Trieste, via G. Tiepolo 11, 34131 Trieste, Italy

^wINAF - Istituto di Astrofisica Spaziale e Fisica Cosmica Milano, via E. Bassini 15, 20133, Milano, Italy

^xJanusz Gil Institute of Astronomy, University of Zielona Góra, Lubuska 2, 65-265, Zielona Góra, Poland

^yINAF-Osservatorio Astronomico di Roma, Via Frascati 33, I-00040 Monte Porzio Catone, Italy

^zInstitut für Theoretische Physik, Johann Wolfgang Goethe-Universität, Max-von-Laue-Straße 1, 60438 Frankfurt, Germany

^{aa}Frankfurt Institute for Advanced Studies, Ruth-Moufang-Straße 1, 60438 Frankfurt, Germany

^{ab}ARTEMIS, CNRS UMR 5270, Université Côte d'Azur, Observatoire de la Côte d'Azur, boulevard de l'Observatoire, CS 34229, F-06304 Nice Cedex 04, France

^{ac}Department of Physics & Institute of Space Sciences & Astronomy, University of Malta, Msida MSD 2080, Malta

^{ad}School of Physics, University of Witwatersrand, Private Bag 3, Wits-2050, Johannesburg, South Africa

^{ae}Dipartimento di Fisica G. Occhialini, Università degli Studi di Milano Bicocca & INFN, Sezione di Milano-Bicocca, Piazza della Scienza 3, 20126 Milano, Italy

^{af}Space Science Data Center (SSDC), Agenzia Spaziale Italiana, via del Politecnico, s.n.c., I-00133, Roma, Italy

^{ag}INFN, Via Enrico Fermi 40, Frascati, Italy

^{ah}University of the Virgin Islands, 2 John Brewer's Bay, 00802 St Thomas, US Virgin Islands

^{ai}Etelman Observatory, Bonne Resolution, St Thomas, US Virgin Islands

^{aj}Czech Technical University, Faculty of Electrical Engineering, Prague 16627, Czech Republic

^{ak}Kazan Federal University, Kazan 420008, Russian Federations

^{al}SRON, Netherlands Institute for Space Research, Sorbonnelaan 2, NL-3584 CA Utrecht, The Netherlands

^{am}Department of Astrophysics/IMAPP, Radboud University, P.O. Box 9010, NL-6500 GL Nijmegen, The Netherlands

^{an}Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, Juliane Maries Vej 30, DK-2100 Copenhagen, Denmark

^{ao}Department of Physics, University of Bath, Claverton Down, Bath BA2 7AY, United Kingdom

^{ap}Scuola Normale Superiore, Piazza dei Cavalieri 7, I-56126 Pisa, Italy

^{aq}SPACE-SI, Slovenian Centre of Excellence for Space Sciences and Technologies, Ljubljana, Slovenia

^{ar}Astronomical Institute of the Czech Academy of Sciences, Fričova 298, 25165 Ondřejov, Czech Republic

^{as}School of Physics and Astronomy and Institute of Gravitational Wave Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, United Kingdom

^{at}Sejong University, 98 Gunja-Dong Gwangjin-gu, Seoul 143-747, Korea

^{au}GEPI, Observatoire de Paris, PSL Research University, CNRS, Place Jules Janssen, 92190 Meudon

Abstract

The recent discovery of the electromagnetic counterpart of the gravitational wave source GW170817, has demonstrated the huge informative power of multi-messenger observations. During the next decade the nascent field of multi-messenger astronomy will mature significantly. In 2030s, third generation gravitational wave detectors will be roughly ten times more sensitive than the current ones. At the same time, neutrino detectors currently upgrading to multi km³ telescopes, will include a 10 km³ facility in the Southern hemisphere that is expected to be operational during the thirties. In this review, we describe the most promising high frequency gravitational wave and neutrino sources that will be detected in the next two decades. In this context, we show the important role of the *Transient High Energy Sky and Early Universe Surveyor* (THESEUS), a mission concept proposed to ESA by a large international collaboration in response to the call for the Cosmic Vision Programme M5 missions. THESEUS aims at providing a substantial advancement in early Universe science as well as playing a fundamental role in multi-messenger and time-domain astrophysics, operating in strong synergy with future gravitational wave and neutrino detectors as well as major ground- and space-based telescopes. This review is an extension of the THESEUS white paper (Amati et al., 2017), also in light of the discovery of GW170817/GRB170817A that was announced on October 16th, 2017.

Keywords: X-ray sources; X-ray bursts; gamma-ray sources; gamma-ray bursts; Astronomical and space-research instrumentation

1. Introduction

With the first detection in 2015 of gravitational waves (GWs) from black hole binary systems during their coalescing phase (Abbott et al., 2016a,b), a new observational window on the Universe has been opened. Stellar-mass black hole coalescences, together with binary neutron star (NS-NS), NS-black hole (BH) mergers, and burst sources as core-collapsing massive stars and possibly NS instability episodes, are among the main targets of ground-based GW detectors, an ensemble of Michelson-type interferometers sensitive to the high frequency range, from few Hz to few thousand Hz. Some of these sources are also expected to produce neutrinos and electromagnetic (EM) signals over the entire spectrum, from radio to gamma-rays.

These expectations were astonishingly satisfied for the first time on August 17th, 2017, when a GW signal consistent with a binary neutron star merger system (Abbott et al., 2017e) was found shortly preceding the short gamma-ray burst GRB170817A (Abbott et al., 2017d). The GW170817 90% confidence sky area obtained with the LIGO and Virgo network was fully contained within the GRB error box. In addition, a “kilonova” or “macronova” emission (AT2017gfo), theoretically predicted from such systems (e.g. Li and Paczyński, 1998a), has been found within the GW-GRB error-box and positionally consistent with NGC4993, a lenticular galaxy at a distance compatible with the GW signal (Abbott et al., 2017f; Smartt et al., 2017; Tanvir et al., 2017; Pian et al., 2017; Coulter et al., 2017).

By the end of the twenties, the sky will be routinely monitored by the second-generation GW detector network, composed by the two Advanced LIGO (aLIGO) detectors in the US (Harry and LIGO Scientific Collaboration, 2010), Advanced Virgo (aVirgo) in Italy (Acernese et al., 2015), ILIGO in India (e.g. Abbott et al., 2016c) and KAGRA in Japan (Somiya, 2012). Then, in the 2030s, more sensitive third generation GW detectors, such as the Einstein Telescope (ET, e.g. Punturo et al., 2010) and LIGO Cosmic Explorer (LIGO-CE, e.g. Abbott et al., 2017b), are planned to be operational and to provide an increase of roughly one order of magnitude in sensitivity. In parallel to these advancements, IceCube and KM3nNeT and the advent of 10 km³ detectors (e.g. IceCube-Gen2, IceCube-Gen2 Collaboration: Aartsen et al., 2014, and references therein) will enable to gain high-statistics samples of astrophysical neutrinos.

The end of the 2020s will therefore coincide with a golden era of multi-messenger astronomy. By that time, the ESA M5 approved missions for space-based astronomy will be launched. THESEUS (*Transient High Energy Sky*




Figure 1: THESEUS within the multi-messenger Astrophysics context of 2020-2030. Green and orange labels are for presently operating and future planned or under construction instruments (Figure credit: S. Schanne).




Figure 2: THESEUS Satellite Baseline Configuration and Instrument suite accommodation.

and Early Universe Surveyor¹) is a space mission concept (Fig. 2), developed by a large International collaboration currently under evaluation by ESA within the selection process for next M5 mission of the Cosmic Vision Programme (Amati et al., 2017). If selected, the launch of THESEUS (2029) will provide a very strong contribution to multi-messenger astronomy. Its instrumental capabilities will ensure fast reaction to GW and neutrino triggers and high

¹<http://www.isdc.unige.ch/theseus>

Table 1: THESEUS instruments

SXI		XGIS				IRT	
Energy range	0.3-6 keV	2 -30 keV	30-150 keV	> 150 keV	ZYJH (0.7 – 1.8 μ m)		
Field of View	1 sr	Half sens.: Total:	50 \times 50 deg ² 64 \times 64 deg ²	50 \times 50 deg ² 85 \times 85 deg ²	2π sr	imaging low res high res	10' \times 10' 10' \times 10' 5' \times 5'
Source location accuracy	< 10'' (best) 105'' (worse)	5' (for > 6 σ source)				< 1''	
Sensitivity	erg(ph) cm ⁻² s ⁻¹ 2×10^{-8} (10) (1s) 2×10^{-11} (0.01) (10 ks)	ph cm ⁻² s ⁻¹ 1 (1s) 0.02 (1ks)	0.15 (1s) 0.004 (1ks)	0.22 (1s) 0.008 (1ks)	H (AB mag) imaging 20.6 (300 s) low res. 18.5 (300 s) high res. 17.5 (1800 s)		

energy (from soft X-rays to gamma-rays) and near infrared (NIR) coverage in the electromagnetic (EM) spectrum. Its wide field of view (FOV) will ensure autonomous triggers of a large number of transient X-ray and gamma-ray sources (Fig. 4). This will enable independent trigger of the EM counterpart of several GW/neutrino sources, as it was the case for GRB170817A triggered by Fermi/GBM. THESEUS will also provide a much better localization of the transient, with uncertainty of the order of 5 arcmin with the hard-X and gamma-ray imager and spectrometer (XGIS), less than 1 arcmin with the X-ray imager (SXI) as sketched in Figure 5, and about 1 arcsecond with the infrared telescope (IRT, see Tab. 1). At the same time, in response to THESEUS triggers, GW and neutrino archival data analysis will enable to search for simultaneous events at the time of the trigger (e.g., due to GRBs or supernovae), since these type of detectors record all their data almost continuously. This strategy has been already pursued by the LIGO-Virgo collaboration for a number of GRBs (e.g. Abbott et al., 2005, 2008, 2017a).

THESEUS will observe in synergy with several telescopes operating at different wavelengths, as illustrated in Figure 1, among which it is worth mentioning: 1) the space-based telescopes James Webb Space Telescope (JWST), ATHENA and WFIRST; 2) the ground-based telescopes with large FOV like zPTF and LSST; 3) the 30-m class telescopes GMT, TMT and ELT; 4) the Square Kilometer Array (SKA) in the radio; 5) the very high-energy (GeV-TeV) Cherenkov Telescope Array (CTA). We note that the main differences between THESEUS and the other large X-ray telescope facility operational in the 2030's, ATHENA, are the much larger field of views of the X-ray and gamma-ray detectors on board THESEUS that will make it a “surveyor” instrument, and the presence of an infrared telescope with both imaging and spectroscopic capabilities.

In the following sections, after a short review of the main THESEUS characteristics (§2; see Amati et al. 2017 for a more exhaustive description of the mission concept), we describe the main role of THESEUS is the Multi-Messenger Astrophysics (MMA) and the main properties of the most promising GW (§4) and neutrino (§5) sources that THESEUS will observe, together with their expected joint GW+EM and neutrino+EM detection rates taking into account the facilities planned to be operational by the end of the twenties.

2. The THESEUS Mission

The THESEUS mission aims at exploiting GRBs for investigating the early Universe and at providing a substantial advancement in multi-messenger and time-domain astrophysics (see Amati et al. 2017 for a detailed review).

The instrumentation foreseen on board THESEUS, illustrated in Figure 2, includes:

- Soft X-ray Imager (SXI, 0.3-6 keV): a set of 4 lobster-eye telescopes units, covering a total FOV of \sim 1 sr with source location accuracy < 1 arcmin;
- X-Gamma ray Imaging Spectrometer (XGIS, 2 keV-20 MeV): a set of coded-mask cameras using monolithic X-gamma ray detectors based on bars of Silicon Drift Diodes coupled with CsI crystal scintillator, granting an




Figure 3: The yearly cumulative distribution of GRBs with redshift determination as a function of the redshift for Swift and THESEUS (Amati et al., 2017). The THESEUS expected improvement in the detection and identification of GRBs at very high redshift w/r to present situation is impressive (more than 100–150 GRBs at $z>6$ and several tens at $z>8$ in a few years) and will allow the mission to shade light on main open issues early Universe science (star formation rate evolution, re-ionization, pop III stars, metallicity evolution of first galaxies, etc.).

unprecedentedly broad energy band, a FOV up to ~ 4 sr, a source location accuracy of ~ 5 arcmin, and an energy resolution of ~ 200 – 300 eV in 2–30 keV;

- InfraRed Telescope (IRT, 0.7 – $1.8 \mu\text{m}$): a 0.7 m class IR telescope with 10×10 arcmin FOV, for fast response, with both imaging and spectroscopy capabilities.

The main characteristics and sensitivities of these instruments are summarized in Table 1. The mission profile includes fast slewing capability, allowing to point the IRT to the position of GRBs and of other transient sources detected and localised by the SXI and/or the XGIS within a few minutes at most, and the possibility of promptly transmitting to ground trigger time, position, and redshift of these events (as evaluated on-board by means of IRT photometry and spectroscopy), thus enabling quick follow-up with large ground- and space-based multi-wavelength observatories. As shown in Figure 3 and detailed in citetAmati17, this unique combination of scientific instruments and mission profile will allow THESEUS to make a giant leap in the use of Gamma-Ray Bursts for shading light on the main open questions on the early Universe (star formation rate evolution up to the end of "dark ages", cosmic re-ionization, metallicity evolution of the early galaxies, pop III stars, ...).

If compared to current generation X-ray facilities, such as for instance the X-ray telescope on board Swift, THESEUS/SXI has a grasp (i.e. $\text{FoV} \times \text{Effective Area}$) of ~ 150 times higher. The large grasp of the SXI, joined with the broad energy band, large effective area and few arcmin source location accuracy of the XGIS, will enable the discovery and study of a wealth of transient sources, both Galactic and extra-galactic (Fig. 4). THESEUS will be also used as a flexible infrared observatory complementary to other facilities, as it was the case for the Swift mission. Rapid follow-up photometric and spectroscopic observations will allow IRT to measure the redshift for a substantial fraction of these events, also refining their localisation down to 0.5 – 1 arcsec (see Amati et al. 2017).




Figure 4: Sensitivity of the SXI (black curves) and XGIS (red) vs. integration time (Amati et al., 2017). The solid curves assume a source column density of $5 \times 10^{20} \text{ cm}^{-2}$ (i.e., well out of the Galactic plane and very little intrinsic absorption). The dotted curves assume a source column density of 10^{22} cm^{-2} (significant intrinsic absorption). The black dots are the peak fluxes for Swift BAT GRBs plotted against T90/2 (where T90 is defined as the time interval over which 90% of the total background-subtracted counts are observed, with the interval starting when 5% of the total counts have been observed, Koshut et al., 1995). The flux in the soft band 0.3–10 keV was estimated using the T90 BAT spectral fit including the absorption from the XRT spectral fit. The red dots are those GRBs for which T90/2 is less than 1 s. The green dots are the initial fluxes and times since trigger at the start of the Swift XRT GRB light-curves. The horizontal lines indicate the duration of the first time bin in the XRT light-curve. The various shaded regions illustrate variability and flux regions for different types of transients and variable sources.

3. The role of THESEUS in the Multi-Messenger Astronomy

The detection of EM counterparts of GW and neutrino signals will enable a multitude of science programmes (see, e.g., Bloom et al., 2009; Phinney, 2009) by allowing for parameter constraints that the GW or neutrino observations alone cannot fully provide. GW detectors have relatively poor sky localisation capabilities, mainly based on triangulation methods, that on average will not be better than few dozens of square degrees. For GW sources at distances larger than the horizon of second-generation detectors (200 Mpc), therefore accessible only by the third-generation ones in the 2030s (e.g. Einstein Telescope and Cosmic Explorer, Punturo et al. 2010; Abbott et al. 2017d), sky localization may even worsen if the new generation network will be composed by only one or two detectors, with possible values of the order of few hundred square degrees or more (e.g. Zhao and Wen, 2017). Neutrino detectors can localise to an accuracy of better than a few squares degrees (see, e.g., Santander, 2016, and references therein). In order to maximise the science return of the multi-messenger investigation it is essential to have a facility that (i) can detect and disseminate an EM signal independently to the GW/neutrino event and (ii) can rapidly search with good sensitivity in the large error boxes provided by the GW and neutrino facilities.

These combined requirements are uniquely fulfilled by THESEUS. Specifically, THESEUS will trigger and localize transient sources within the uncertain GW and/or neutrino error boxes with the hard XGIS and/or with SXI. A very large fraction of the error boxes of poorly localised GW sources can be covered with SXI FoV within one orbit due to the large grasp of the instrument (see Amati et al. 2017). In response to an SXI/XGIS trigger, if an optical counterpart is present, the source sky localization can be refined down to few arcseconds with IRT observations. Precise localizations will be disseminated within minutes to the astronomical community, thus enabling large ground-based telescopes to observe and deeply characterise the transient nature.




Figure 5: The plot shows the THESEUS/SXI field of view ($\sim 110 \times 30 \text{ deg}^2$, pink rectangle) superimposed on the probability skymap of GW 170817 obtained with the two Advanced LIGO only (cyan) and with the addition of Advanced Virgo (green) (Abbott et al., 2017e). THESEUS not only will cover a large fraction of the skymap (even those obtained with only two GW-detectors, e.g. cyan area), but will also localize the counterpart with uncertainty of the order of 5 arcmin with the XGIS and to less than 1 arcmin with SXI. *The THESEUS location accuracy of GW events produced by NS-NS mergers can be as good as 1 arcsec in case of detection of the kilonova emission by the IRT.* By the end of the 2020s, if ET will be a single detector, almost no directional information will be available for GW sources (e.g. $> 1000 \text{ deg}^2$ for BNS at $z > 0.3$, Zhao and Wen e.g. 2017), and a GRB-localising satellite will be essential to discover EM counterparts.

As we will discuss in more details in the next sections, several multi-messenger sources are among the main targets of THESEUS, as for example GRBs, flaring magnetars, core-collapse supernovae (CCSNe) and AGNs. We here briefly recall the main THESEUS capabilities in GRB detections while we address the reader to Amati et al. 2017 for the detection capabilities of the other mentioned sources.

The combination of SXI and XGIS, makes THESEUS a unique machine to explore both the populations of long/high redshift and hard/short GRBs. Figure 6 shows the density contours of the population of short GRBs (dashed contours) in the peak energy - peak flux plane (Ghirlanda et al., 2016). The density contours of the short GRB population detectable by THESEUS is shown by the shaded contours. Due to their harder spectrum, short GRBs are better triggered by XGIS than SXI (see Fig. 8). Compared to the detection thresholds of BATSE and Fermi/GBM, THESEUS will slightly extend the detected population leftwards of these thresholds (cyan and yellow lines in Figure 6). The star symbol in Figure 6 shows the position of GRB170817A as revealed by GBM (Goldstein et al., 2017). THESEUS will be able to fully access similar events and explore their nature. Although the XGIS sensitivity threshold improves over GBM, its smaller (by a factor of 2) field of view compensates this gain reaching a detection rate of short GRBs which is comparable to that of GBM. What makes THESEUS XGIS unique, with respect to GBM, is the possibility to locate, thanks to the soft (2 keV-30 keV) coded mask detectors of the XGIS, most of the detected short GRBs with an expected accuracy of 5 arcmin (to be compared with the average $>$ few degrees of GBM GRBs). THESEUS will ensure short GRB detection with a rate of 15-35 per year (Fig. 8). Since short GRBs are expected to be the EM counterpart of compact binary coalescences that are known to emit high-frequency GWs, and the case of GRB170817A has definitively confirmed this expectation, the short GRB detection capabilities highlight the crucial relevance of the role of THESEUS for multi-messenger astronomy in an epoch where almost all short GRBs will be accompanied by a GW signal detected by the third-generation interferometers (e.g. ET or LIGO-CE).




Figure 6: Density contours (dashed lines) corresponding to 1, 2, 3 σ levels of the synthetic population of Short GRBs (from Ghirlanda et al. 2016). Shaded coloured regions show the density contours of the population detectable by THESEUS. The yellow and cyan lines show the trigger threshold of Fermi/GBM and GCRO/Batse (from Nava et al. 2011). The flux is integrated over the 10-1000 keV energy range. The star symbol shows the short GRB170817A (Goldstein et al., 2017).

Figure 7 shows the density contours (dashed lines) of synthetic population of long GRBs (Ghirlanda et al., 2015c) in the observer-frame plane representing the peak energy E_{peak} versus the (10-1000 keV) peak flux. Density contours of long GRBs detected by THESEUS are shown by the shaded coloured regions. The trigger thresholds of Fermi/GBM and CGRO/BATSE are also shown for comparison (adapted from Nava et al. 2011) by the yellow and cyan lines, respectively. Bursts located at the right of these lines are detectable by the Fermi/GBM and BATSE. THESEUS will access a region of the E_{peak} -Peak flux plane totally unexplored by past and current instruments. A large fraction of its population will be constituted by soft low flux events. Among these there will be (i) low redshift/low luminosity events (with a E_{peak} due to the correlation between these two observables; Yonetoku et al., 2004) which are of paramount importance to constrain the faint end of the GRB luminosity function (Pescalli et al., 2015) and (ii) long GRBs at high redshifts which, used as beacons, will allow us to explore the high redshift Universe and its evolution. Given the association of long GRB to CCSNe and the expected GW radiation as well as neutrino emission of these events, THESEUS detection capabilities will make this mission crucial for multi-messenger joint observational campaigns related to long GRB sources.

As explained in the next sections, besides the expected collimated GRB “prompt” emission, softer X-ray emission is also expected from the side and/or afterglow emission from the GRB jet, with a much lower degree of collimation. For short GRB sources and in particular NS-NS mergers, an additional nearly isotropic soft X-ray emission is possibly expected when the merger remnant is a long-lived NS or magnetar, where the corresponding transient is powered by the spindown of the latter (see Section 4.2).

GRB afterglows can be monitored also at the NIR wavelengths with THESEUS/IRT. Figure 9 shows the optical afterglow fluxes of on-axis GRBs compared with the sensitivity of IRT in both imaging and spectroscopic mode.

3.1. Science return from joint GW+EM detections with THESEUS

Each individual joint observation of an EM source and its GW and/or neutrino counterpart, provides an enormous science return. We mention just few examples in the case of compact binary coalescences: *i*) the determination of the GW polarization ratio would constrain the binary orbit inclination and hence, when combined with an EM signal, the




Figure 7: Density contours (dashed lines) corresponding to 1, 2, 3 σ levels of the synthetic population of Long GRBs (from Ghirlanda et al. 2015c). Shaded coloured regions show the density contours of the population detectable by THESEUS. The yellow and cyan lines show the trigger threshold of Fermi/GBM and GCRO/Batse (from Nava et al. 2011). The flux is integrated over the 10-1000 keV energy range. As can be seen, *THESEUS* will carry on-board the ideal instruments suite for detecting all classes of GRBs (classical long GRBs, short/hard GRBs, sub-energetic GRBs, and very high-redshift GRBs, which, in this plane, populate the region of weak/soft events), providing a redshift estimate for most of them ? detecting




Figure 8: Cumulative distribution of the rate of short GRBs as a function of redshift that THESEUS will detect (yellow stripe filled region). The fraction of the population that will be detected by the soft coded mask instruments of XGIS (2-30 keV) is shown by the green stripe. The cumulative distribution of the fewer short GRBs also detected by SXI is shown by the red stripe. The vertical width of the stripes account for the uncertainties of the model parameters of the short GRB population adopted (Ghirlanda et al., 2016).

jet geometry and source energetics; *ii*) a better understanding of the NS equation of state can follow from combined GW and X-ray emission signals (see §4.2) (see, e.g., Bauswein and Janka, 2012; Takami et al., 2014; Lasky et al., 2014; Ciolfi and Siegel, 2015b,a; Messenger et al., 2015; Rezzolla and Takami, 2016; Drago et al., 2016); *iii*) an estimate of the amount of matter expelled during a NS-NS or a NS-BH merger (e.g. Fernández and Metzger, 2016, and references therein); *iv*) a better understanding of the physical mechanisms underlying the core-collapse phase of massive stars; *v*) whether the magnetar scenario and the current interpretation of X-ray giant flares are correct; *vi*) tracing the history of heavy-metal enrichment of the Universe; *vii*) redshift measurements of a large sample of short GRBs combined with the absolute source luminosity distance provided by the CBC-GW signals can deliver precise measurements of the Hubble constant (Schutz, 1986), helping to break the degeneracies in determining other cosmological parameters via CMB, SNIa and BAO surveys (see, e.g., Dalal et al., 2006).

The last point on the Hubble constant measure is of particular relevance for THESEUS especially during the third-generation GW detector era, when, as we will show in the next section, almost all short GRB detected will have a GW counterpart. A first attempt of Hubble constant measurement has been explored with GW170817 for which the recession velocity v_r of the optical transient AT2017gfo host galaxy NGC4993, was combined with the luminosity distance D_L measured directly from the waveform of GW170817. For small distances, as in the case of GW170817 (~ 40 Mpc), the Hubble constant depends only on these two variables as $H_0 = v_r/D_L$. Despite the large uncertainties on this first measurement, the results are very encouraging (Abbott et al., 2017c). The value obtained, $H_0 = 70.0^{+12.0}_{-8.0}$ km s $^{-1}$ Mpc $^{-1}$, lies in between the measurements obtained from SNIa from SHoES (73.24 ± 1.74 km s $^{-1}$ Mpc $^{-1}$, (Riess et al., 2016) and CMB from Planck (67.74 ± 0.46 km s $^{-1}$ Mpc $^{-1}$, (Planck Collaboration et al., 2016). Furthermore, combining the observing angle vs. GW amplitude degeneracy measured by LIGO-Virgo interferometers with independent information on the observing angle derived from modelling the associated broadband afterglow, a further reduction by $\sim 5\%$ on the uncertainty interval of H_0 could be obtained (Guidorzi et al., 2017).

3.2. THESEUS and GRB170817A

In this section we explore THESEUS capabilities in the detection and characterization of the short GRB170817A associated with GW170817 (Abbott et al., 2017e; Goldstein et al., 2017; Savchenko et al., 2017), after a brief summary of the main observational properties of this event.

The two events were found consistent with being originated from a common source with high confidence ($\sim 10^{-7}$ probability of being independent, Abbott et al. 2017e). In addition, a bright optical transient was observed in NGC4993, which was then identified as the host galaxy (Smartt et al., 2017; Tanvir et al., 2017; Pian et al., 2017;




Figure 9: R-band light curves of long (grey lines) and short (black dots) GRBs (adapted from Kann et al. 2011). The limiting magnitudes achievable with THESEUS/IRT with 300 s of exposure (blue lines), 8-m class telescopes (red lines) and ELT/MICADO (green lines) are also shown. Dashed horizontal line for the spectroscopy, solid line for the imaging. The magnitudes are rescaled from H-band to R-band assuming achromatic behaviour and a spectral index $\beta=0.7$. A tentative observation strategy could consist of the following steps: first starting the follow up with IRT, then activating the observations with 8-m class telescopes after few hours, finally, according to the brightness of the afterglow and thanks to the very high sensitivity of ELT, performing late observations for weeks.

Abbott et al., 2017d; Coulter et al., 2017). This galaxy has a distance estimate of ~ 40 Mpc, which was consistent with the luminosity distance measured from the gravitational wave signal. This was by far the closest short GRB yet observed. The gamma-ray peak photon flux (3.7 ± 0.9 ph $\text{cm}^{-2} \text{s}^{-1}$ in the 10-1000 keV band, Goldstein et al. 2017) implies an extremely low isotropic luminosity short GRB if compared with typical values ($\sim 10^{51} - 10^{53}$ erg s^{-1} , see e.g. Ghirlanda et al. 2015b), with 1.7×10^{47} erg s^{-1} (e.g. Zhang et al. 2017). Since various indications point at a binary merger seen with viewing angle $\sim 20 - 40$ deg away from the normal direction to the orbital plane, a possible explanation of the low luminosity is that the event was a short GRB with a structured jet observed off-axis (e.g. Troja et al., 2017; Alexander et al., 2017; Margutti et al., 2017; Haggard et al., 2017; Hallinan et al., 2017; Lazzati et al., 2017c). Within the latter scenario, GRB170817A suggests a significant extension of the observed short GRB population to include a larger fraction of dimmer events (e.g. Burgess et al., 2017), which can strongly enhance the coincident short GRB/GW detection rate up to relatively small distances (i.e. < 100 Mpc) with sensible instruments such as THESEUS. The late-time X-ray and radio counterparts detected respectively ~ 9 and ~ 16 days after merger (Troja et al., 2017; Margutti et al., 2017; Alexander et al., 2017; Hallinan et al., 2017; Haggard et al., 2017) and the additional X-ray and optical detections ~ 109 and ~ 112 days after trigger, respectively (GCNs 22201, GCN 22203, 22206, GCN 22207), are consistent with both the expectation of an afterglow emission from a structured jet (Rossi et al., 2002; Zhang and Mészáros, 2002; Kathirgamaraju et al., 2017; Lazzati et al., 2017a,b; Gottlieb et al., 2017, 2018; Salafia et al., 2017a; Lazzati et al., 2017c) and from the deceleration of an isotropic mildly relativistic outflow (Mooley et al., 2017; Salafia et al., 2017b). Both scenarios can account reasonably well for the low luminosity of the prompt emission and the late-time rise up of the X-ray and radio emission, although the slow late-time radio increase might require some modifications of the simplest assumptions (Mooley et al. 2017; but see Lazzati et al. 2017c).




Figure 10: The Fermi/GBM peak photon flux of the short GW/GRB 170817 (red star, [Goldstein et al. 2017](#)) rescaled with the distance (blue line) and compared with THESEUS/XGIS 1-s sensitivity in the 50-300 keV energy range. Off-axis short GRB similar to GRB 170817 could had been detected with THESEUS/XGIS up to $\sim 70 - 80$ Mpc.

In Figure 10 the measured Fermi/GBM flux in the 50-300 keV of $0.8 \pm 0.3 \text{ ph cm}^{-2} \text{ s}^{-1}$ is extrapolated at distances larger than the distance of GRB170817A (~ 40 Mpc) and compared with the THESEUS/XGIS sensitivity in the 30-150 keV band ([Amati et al., 2017](#), see Fig. 36) rescaled to the 50-300 keV. From this plot, we can see that not only this source could had been clearly detected with XGIS but it could have been detected up to $\sim 70 - 80$ Mpc, that is nearly twice the actual distance of the source. On the other hand, the faint X-ray emission detected 9 days after the trigger with a flux of the order of a few times $10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$ ([Troja et al., 2017](#)), could not had been detected with the SXI.

Figure 11 shows how IRT could had clearly detected the NIR counterpart of GW170817 recognized to be the expected "kilonova" (or "macronova", see next section). In particular, the observed emission was bright enough during the first 2 days to have allowed IRT low-resolution spectroscopy with unprecedented coverage.

More in general, Figure 12 shows how THESEUS/IRT will be able to promptly follow-up the kilonova/macronova IR emission and thus enable a detailed characterization of these key astrophysical phenomena associated with compact binary coalescences and short GRBs. Specifically, the plot shows the expected macronova apparent magnitudes for a source between 50 and 200 Mpc ([Barnes et al., 2016](#)) plotted against the IRT sensitivity in imaging and spectroscopic mode. If bright enough, spectroscopic observations could be performed on-board, thus providing redshift estimates and information on chemical composition of circum-burst medium. Alternatively, if the optical/NIR counterpart is not bright enough for onboard spectroscopy, precise IRT sky coordinates will be disseminated to ground based telescopes to perform spectroscopic observations and redshift measurements.

4. Gravitational wave sources

4.1. NS-NS / NS-BH mergers: Collimated emission from Short GRBs

Compact binary coalescences (CBCs) involving neutron stars (NS) and stellar mass black holes (BH) are among the sources of GWs that will be likely detected in spades in the next decade. These systems radiate GWs within the




Figure 11: Light curve of the kilonova associated to the gravitational wave/short GRB event GW170817/GRB170817A in the IRT filters (from Tanvir et al. 2017; Pian et al. 2017; Arcavi et al. 2017; Cowperthwaite et al. 2017; Drout et al. 2017; Kasliwal et al. 2017; Smartt et al. 2017; Troja et al. 2017). The continuous and dashed red lines indicate the THESEUS/IRT limiting H magnitudes for imaging and prism spectroscopy, respectively, with 300s of exposure (see Amati et al., 2017). Credit: A. Melandri

most sensitive frequency range of ground-based GW detectors (1-2000 Hz), with large GW energy output, of the order of $10^{-2} M_{\odot}c^2$, and gravitational waveforms well predicted by General Relativity (see, e.g., Baiotti and Rezzolla 2017 for a review). From the merger of two stellar-mass black holes the current consensus is that no significant EM counterpart emission is expected, with exceptions where a short GRB-like EM signal may be produced from those BH-BH systems merging in very high density environments (as for example in an AGN disk, Bartos 2016), or in other exotic conditions (e.g. Perna et al., 2016; Seto and Muto, 2011; Loeb, 2016). On the other hand, EM emission is expected on robust theoretical ground for merging NS-NS or NS-BH. For more than a decade, mounting indirect evidence supported the long-standing hypothesis that short GRB progenitors are associated with CBC systems with at least one neutron star (e.g. Paczynski, 1986; Eichler et al., 1989; Narayan et al., 1992; Barthelmy et al., 2005; Fox et al., 2005; Gehrels et al., 2005). The most compelling pieces of evidence include (i) the observation of short GRBs in both elliptical and late-type star forming galaxies, with a preference for the most massive ones (ii) the relatively large projected offsets of these events with respect to the center of their host galaxies, and (iii) the lack of supernova associations, as opposed to the case of long GRBs (e.g., Berger 2014 and refs therein). The most recent numerical simulations also provided supporting (though not conclusive) evidence that such merging systems might act as short GRB central engines (e.g. Rezzolla et al., 2011; Paschalidis et al., 2015; Ruiz et al., 2016; Kawamura et al., 2016; Rosswog et al., 2013). Short GRBs have been historically distinguished from long GRBs as those with gamma-ray burst (prompt emission) duration less than 2 s (Kouveliotou et al., 1993). However, there are similarities between the two classes of GRBs as far as their prompt emission (e.g. Ghirlanda et al., 2015a, 2009, 2011) and their afterglows (e.g. D’Avanzo, 2015). It is likely that the two populations overlaps and contaminate each other, especially when selection is based solely on the observed duration (Bromberg et al. 2013, but see Zhang et al. 2012). Therefore, GW signals from short GRBs might be an additional parameter to be considered to firmly distinguish core-collapse (long) from compact merger (short) progenitors.

On August 17th, 2017 the association of short GRBs and CBCs involving NSs found direct confirmation, with the first detection of a gravitational wave signal consistent with a binary NS merger named GW170817 and the




Figure 12: Theoretical H -band lightcurves of kilonova based on models (from Barnes et al., 2016). The lightcurves are in observer frame for a source between 50 and 200 Mpc. Gray model is for the most optimistic case of a kilonova with $0.1 M_{\odot}$ ejected mass with speed of $0.3 c$. Green model is for a weaker emission, corresponding to $0.01 M_{\odot}$ ejected mass with speed of $0.1 c$. The continuous and dashed red lines indicate the THESEUS/IRT limiting H magnitudes for imaging and prism spectroscopy, respectively, with 300 s of exposure (see Amati et al., 2017).

observation, less than two seconds later, of GRB170817A (Abbott et al., 2017e; Goldstein et al., 2017; Savchenko et al., 2017). Since GW events enable to individuate faint short GRBs as GRB170817A, future joint multi-messenger detections of such sources will shed light on several still debated topics as: 1) the jet intrinsic structure and their properties, and ultimately the crucial issue of short GRB energetics; 2) the physics regulating the on- and off-axis emission; 3) the late-time (e.g. after 1-2 weeks) component origin. THESEUS will significantly contribute to all these topics.

Table 2 shows the expected rate of THESEUS/XGIS short GRB detections with a GW counterpart from merging NS-NS systems (i.e. within the GW detector horizon). The quoted numbers are obtained by correcting the realistic estimate of NS-NS merger rate, $\sim 40 \text{ yr}^{-1}$ (Abadie et al. 2010, see also Belczynski et al. 2017) for the fraction of the sky covered by the XGIS FoV, that is $\sim 50\%$, and the short GRB jet collimation factor, by assuming a jet half-opening angle range of $10\text{--}40$ deg, taking into account the possibility to observe off-axis short GRBs up to 5 times a jet half-opening angle of 10 degrees² and 2 times a jet of 40 degrees (Kathirgamaraju et al., 2017; Pescalli et al., 2016). We are here assuming that every BNS merger produces a jetted short GRB, which is still an open issue. Results show that, during the 2020's, the GW+EM detection rate of short GRBs with THESEUS is found to be of the order of 5–15 per year. By the time of the launch of THESEUS, gravitational radiation from NS-NS and NS-BH mergers will be detectable by third-generation detectors such as the Einstein Telescope (ET) up to redshifts $z \sim 2$ or larger (see, e.g., Sathyaprakash et al., 2012; Punturo et al., 2010), thus dramatically increasing the GW+EM on-axis short GRB detection rate. The important implication is that almost all THESEUS short GRBs will have a detectable GW emission. Indeed, it is likely that at the typical distances at which ET detects GW events, the only EM counterparts that could feasibly be detected are short GRBs and their afterglows, making the role of THESEUS crucial for multi-messenger astronomy by that time.

²The 5-times factor was obtained by considering the THESEUS/XGIS 1 sec photon flux sensitivity $\sim 0.2 \text{ ph cm}^{-2} \text{ s}^{-1}$ (see also Amati et al. 2017)

Table 2: Number of NS-NS (BNS) mergers expected to be detected in the next years by second- (2020+) and third- (2030+) generation GW detectors and the expected detection number of electromagnetic counterparts as short GRBs (collimated) and X-ray isotropic emitting counterparts (see §3.1 and 3.2) with THESEUS SXI and XGIS (see text for more details). BNS rate is a realistic estimate from [Abadie et al. 2010](#) and [Sathyaprakash et al. 2012](#) and the BNS range indicates the sky- and orbital inclination-averaged distance up to which GW detectors can detect a BNS with $S/N = 8$.

GW observations		THESEUS XGIS/SXI joint GW+EM observations			
Epoch	GW detector	BNS range	BNS rate (yr ⁻¹)	XGIS/sGRB rate (yr ⁻¹)	SXI/X-ray isotropic counterpart rate (yr ⁻¹)
2020+	Second-generation (advanced LIGO, Advanced Virgo, India-LIGO, KAGRA)	~200 Mpc	~40*	~5-15	~1-3 (simultaneous) ~6-12 (+follow-up)
2030+	Second + Third-generation (e.g. ET, Cosmic Explorer)	~15-20 Gpc	>10000	~15-35	≥100

* from Abadie et al. 2010a

4.2. NS-NS / NS-BH mergers: non-collimated soft X-ray and optical/NIR emission

GW emission from CBCs depends only weakly on the inclination angle of the inspiral orbit and therefore these events are in general observable at any viewing angle. As a consequence, most of the GW-detected mergers are expected to be observed off-axis (i.e. with a large angular distance of the observer from the orbital axis). This makes the non collimated, nearly isotropic EM components extremely relevant for the multi-messenger investigation of CBCs.

A potentially powerful nearly-isotropic emission is expected if a NS-NS merger produces a long-lived millisecond magnetar. In this case, soft X-ray to optical transients can be powered by the magnetar spin-down emission reprocessed by the baryon-polluted environment surrounding the merger site (mostly due to isotropic matter ejection in the early post-merger phase), with time scales of minutes to days and luminosities in the range 10^{43} – 10^{48} erg s⁻¹ (e.g. [Yu et al., 2013](#); [Metzger and Piro, 2014](#); [Siegel and Ciolfi, 2016a,b](#)). In particular, in soft X-rays (at ~keV photon energies) these transients can last from minutes to hours and, for the most optimistic models, reach luminosities as high as 10^{48} erg s⁻¹ ([Siegel and Ciolfi, 2016a,b](#)). According to alternative models, X-ray emission may also be generated via direct dissipation of magnetar winds (see, e.g., [Zhang, 2013](#); [Rezzolla and Kumar, 2015](#)). Furthermore, the high pressure of the magnetar wind can in some cases accelerate the expansion of previously ejected matter into the interstellar medium up to relativistic velocities, causing a front shock which in turn produces synchrotron radiation in the X-ray band (with a high beaming factor of ~0.8; see, e.g., [Gao et al. 2013](#)).

Figure 13 shows predictions for magnetar-powered X-ray emission following a NS-NS merger according to a number of different models. Overall, typical time scales for these transients are comparable to magnetar spin-down time scales of $\sim 10^3$ – 10^5 s and the predicted luminosities span a wide range that goes from 10^{41} to 10^{48} erg s⁻¹. Joint GW+EM detection rates with THESEUS/SXI are discussed below. These rates depend not only on the rate of NS-NS mergers, but also on the (essentially unknown) fraction of mergers forming a long-lived NS remnant, which is necessary to produce spindown-powered transients. The observation of this type of emission after a NS-NS merger would indeed indicate that the remnant is long-lived, allowing for significant constraints on the equation of state of the remnant itself (e.g. [Piro et al., 2017](#); [Drago and Pagliara, 2017](#)).

In the case of GW170817/GRB170817A, no evidence for this type of emission was found in the soft X-ray band. However, the first deep pointed observations at ~keV photon energies only started ~15 hours after merger with Swift/XRT ([Evans et al., 2017](#)). Possible constraints could be provided by the MAXI (2-10 keV) observations taken at 4.5 hours after the trigger, with a flux limit of $\sim 10^{-8}$ erg cm⁻² s⁻¹. We note that for GW170817 the nature of the remnant (BH vs. long-lived NS) was not established for this event, thus making it difficult to put constraints on theoretical expectations. For future observations, being able to catch the soft X-ray emission (or to firmly assess its absence) within the relevant time scale after a GW trigger will require a monitoring (wide-field) instrument sensitive to ~keV energies. THESEUS/SXI will perfectly respond to this need.

The expected detection rate of the isotropic X-ray emission from NS-NS mergers is quoted in Table 2 where, from the realistic rate of NS-NS mergers that will be detected with GW observatories in 2020s and 2030s, we have accounted for: 1) the fraction of the sky covered by the SXI FoV, that is ~ 8%, for serendipitous discoveries, and 2) the fraction of NS-NS systems that can produce X-ray emission (i.e. that do not form immediately a BH), that we assumed




Figure 13: Expected X-ray fluxes at peak luminosity from two different luminosity distances ($z=0.05$ on the left panel, and $z=1$ on the right panel) and from different models of magnetar-powered X-ray emission from long-lived NS-NS merger remnants. Predictions from each model are represented by a coloured region and/or by single dots that are indicative of fiducial cases (see the legend on the right). Grey solid lines in the left panel show typical GRB X-ray afterglows observed with Swift/XRT. Finally, the black curves show the SXI sensitivity vs. exposure time, assuming a source column density of $5 \times 10^{20} \text{ cm}^{-2}$ (i.e., well out of the Galactic plane and very little intrinsic absorption, solid line) and 10^{22} cm^{-2} (significant intrinsic absorption, dashed line).

to be within [30-60]% (Gao et al., 2013; Piro et al., 2017). Moreover, we consider the fraction of BNS sources that could be followed-up with SXI after a GW alert, estimated to be of the order of $\sim 40\%$. From these computations, we find that during the 2020s the joint GW+EM detection rate with THESEUS of these X-ray counterparts of NS-NS mergers is ~ 6 -12 per year. During 2030s, with the third-generation GW detectors, isotropic X-ray emission from NS-NS mergers as predicted by some models (e.g. Siegel and Ciolfi, 2016b) could be detected up to ~ 10 times larger distances, with an improved joint GW+EM detection rate of few hundreds per year (depending on the largely uncertain intrinsic luminosity of such X-ray component, see Table 2 and Fig. 13). With such statistics, THESEUS will provide a unique contribution to characterize this X-ray emission from NS-NS systems.

Another well known type of nearly-isotropic emission expected from CBCs involving NSs is the so-called “kilonova” or “macronova” (e.g., Li and Paczyński, 1998b; Metzger et al., 2010). NS-NS or NS-BH mergers can eject a substantial amount of matter (up $10^{-2} M_{\odot}$ or more) which becomes unbound and leaves the system. This material can be expelled both during merger (dynamical ejecta) and in the post-merger phase, in the form of baryon-loaded winds from the accretion disk surrounding the merger remnant (or from the remnant NS itself, for NS-NS mergers without prompt collapse to BH). Due to the unique conditions of high neutron density and temperature, r-process nucleosynthesis of very heavy elements takes place in the ejected matter and days after merger the radioactive decay of such elements heats up the material producing a thermal transient signal peaking in the optical/NIR bands and with typical luminosities of $\sim 10^{40}$ – $10^{41} \text{ erg s}^{-1}$ (see, e.g., Fernández and Metzger, 2016; Metzger, 2017). The temporal and spectral properties of these signals encode crucial information on the nature of the merger progenitor (e.g., NS-NS or NS-BH), the equation of state of neutron stars, and the heavy element chemical enrichment of the Universe.

Before the GW170817 event, observational and photometric-only evidence of kilonova/macronova transients relied only on a few candidates observed during short GRB follow-up campaigns (e.g. Tanvir et al., 2013; Jin et al., 2015; Berger et al., 2013). The recent discovery of an optical/IR transient associated with GW170817 (counterpart named AT2017gfo) has now provided the first compelling evidence, both photometric and spectroscopic, of the existence of kilonovae/macronovae (Pian et al. 2017, see also Abbott et al. 2017d; Tanvir et al. 2017; Nicholl et al. 2017; Smartt et al. 2017; Tanaka et al. 2017; Chornock et al. 2017). The observations of GW 170817 consolidated the presence of a strong IR emission component (reaching its maximum 1.5 days after merger, with 17.2 and 17.5 mag in the J and K bands, respectively (e.g. Tanvir et al., 2017). This provides a very strong science case for the IR instrument on-board THESEUS (IRT) as shown in Figure 12.

Both serendipitous discoveries within the large THESEUS/SXI FoV and re-pointing of THESEUS in response to a GW trigger will allow to study off-axis X-ray emission expected from NS-NS systems (see §4.2). With THESEUS/SXI in combination with the second-generation detector network, almost all predicted non-collimated X-ray

counterparts of GW events from NS-NS merging systems will be easily detected simultaneously with the GW trigger and/or with rapid follow-up of the GW-individuated sky region. Among the open questions that THESEUS will help to address there are: 1) does the NS-NS merger create a NS or a BH, and how fast?; 2) how much matter is expelled in the NS mergers? At which speeds?; 3) what is the amount of asymmetry in the NS-NS merger ejecta and the corresponding optical emission?

4.3. Core-collapse of massive stars: supernovae and long GRBs

Core-collapse supernovae (CCSNe) represent another type of GW sources that are of great interest for the involved community. Their expected GW emission is highly uncertain as it strongly depends on the rather unknown SN explosion mechanism (e.g. [Logue et al., 2012](#); [Powell et al., 2016](#)). Not only the signal morphology (waveform), but also the expected energy output are still under debate. Thus, depending on the assumed model as well as dedicated data analysis techniques, during the second-generation GW detector network era the detection of GW from CCSNe is predicted up to few Mpc or less (e.g. [Ott, 2010](#)), or up to ~ 100 Mpc (e.g. [van Putten et al., 2017](#)). While this makes it difficult to predict the GW signal and its detectability, it also represents a unique opportunity to probe the CCSN inner dynamics that cannot be explored via the sole observation of EM signals.

The firm association of nearby long GRBs with temporally and spatially coincident CCSNe (e.g. [Woosley and Bloom, 2006](#); [Galama et al., 1998](#); [Stanek et al., 2003](#))³ implies that any long GRB, if close enough, should be associated with a detectable GW emission and thus offers a very interesting potential synergy between gamma-ray and GW detectors. At present, this synergy is severely limited by the rather small GW detection horizon of these events, which implies rates that can be as low as a few events per century, but third generation GW detectors such as the Einstein Telescope will offer much better prospects. The first joined GW/GRB/SN observations, possibly combined also with neutrino detections (Sect. 5), will prove crucial to unravel the nature of these sources and their explosion mechanism.

Wolf-Rayet stars as well as red and blue supergiants are expected to exhibit bright shock breakout soon after their core collapse, with X-ray bursts lasting 10-1000 s and with luminosities expected in the range $10^{43} - 10^{46}$ erg s⁻¹. These progenitors are likely responsible for Type Ibc and most Type II SNe, which occur at rates of 2.6×10^{-5} and 4.5×10^{-5} Mpc⁻³ yr⁻¹, respectively ([Li et al., 2011](#)). THESEUS/SXI and XGIS can detect these events up to ~ 50 Mpc leading to a rate of a few per year. Third generation interferometers may thus detect CCSNe up to ~ 1 Gpc. We thus expect up to few shock breakout events per year that can be detected with THESEUS/SXI simultaneously with their GW counterpart during the 2030s. SBO components are temporally closer to the possibly associated GW events than the optical CCSNe counterpart, thus their detection can mark with more precision the start time of the gravitational radiation emission and can be used in the challenging signal search processes ([Andreoni et al., 2016](#)).

Observed long GRB rate density is ~ 1 Gpc⁻³ yr⁻¹ (e.g. [Le and Dermer, 2007](#)) and simultaneous GW+EM detection rate of more than 1 event per year is expected only with the third-generation GW interferometers. Off-axis X-ray afterglow detections (“orphan afterglows”) (e.g. [Granot et al., 2002](#); [Ghirlanda et al., 2013, 2015b](#)) can potentially increase the simultaneous GW+EM detection rate for nearby long GRBs by a factor that strongly depends on the jet opening angle and the observer viewing angle. THESEUS may also observe the appearance of a NIR orphan afterglow few days after the reception of a GW signal due to a collapsing massive star. In addition, the possible large number of low luminosity GRBs (LLGRBs, e.g. [Toma et al., 2007](#); [Virgili et al., 2009](#)) in the nearby Universe, expected to be up to 1000 times more numerous than long GRBs, will provide clear signatures in the GW detectors because of their much smaller distances with respect to long GRBs.

4.4. Magnetars

Fractures of the solid crust on the surface of highly magnetized neutrons stars and/or dramatic magnetic field readjustments represent the most widely accepted explanations to interpret the magnetar bursting activity and in particular the rare giant flares observed in X-rays from three different soft gamma repeaters (SGRs; see, e.g., [Thompson](#)

³For GRB 060614 ([Della Valle et al., 2006](#); [Fynbo et al., 2006](#)), in spite of marginal evidence for an associated kilonova, which would make it a short GRB, this event, along with GRB 060505, leaves the possibility of long SN-less GRBs, see also [Xu et al. 2009](#).

and Duncan 1995; Guidorzi et al. 2004; Mereghetti et al. 2015). The above events will inevitably excite non-radial oscillation modes that may produce detectable GWs (see, e.g., Corsi and Owen, 2011; Ciolfi et al., 2011). The most recent estimates for the energy reservoir available in a giant flare are between 10^{45} erg (about the same as the total EM emission) and 10^{47} erg. The efficiency of conversion of this energy into GWs was estimated in numerical relativity simulations and it was found to be likely too small to be within the sensitivity range of present GW detectors (Ciolfi and Rezzolla, 2012; Lasky et al., 2012). However, at the typical dominant (i.e. f-mode) oscillation frequencies in NSs (\sim kHz), ET will be sensitive to much lower GW energies (Punturo et al., 2010). Therefore, a relatively close giant flare event might lead to a detectable GW emission.

5. Neutrino sources

Several gamma-ray and X-ray sources that THESEUS will observe as GRBs, CCSNe and AGNs, are also expected to originate neutrinos. Due to their low interaction cross-section, neutrinos can probe the innermost regions similarly to gravitational waves but, in addition, neutrino detectors can provide a more refined sky localisation than GW interferometers, with an uncertainty that goes from few degrees down to a fraction of a degree. Current neutrino deep-water-based detectors include DUMAND, Lake Baikal, and ANTARES. These Northern hemisphere detectors complement the South Pole based IceCube, the first km-scale neutrino observatory, completed and in full operation since 2010. Two major upgrades for the near and far future are planned with the construction of Km3Net in the Northern hemisphere, started in 2015, and IceCubeGen2, an upgrade to a 10 km^3 detector of IceCube (e.g. IceCube-Gen2 Collaboration: Aartsen et al., 2014, and references therein). These are prevalently high-energy neutrino detectors but IceCube and Km3Net can detect also MeV neutrinos due to the capabilities to suppress background rate, together with other liquid scintillators and liquid Argon Time-Projection Chamber detectors (e.g. see reviews by Scholberg, 2012; Gil-Botella, 2016, and references therein).

Pulses of low energy neutrinos ($<10 \text{ MeV}$) are expected to be released during CCSNe with an energy release up to 10^{53} erg. Indeed, MeV neutrinos have been detected so far only from one CCSN, namely SN1987A, in the Large Magellanic Cloud at 50 kpc distance (e.g. Gaisser et al., 1995). Comparison of the SN1987A neutrino signal with theoretical predictions showed that the general features of CCSNe are compatible with the observations (e.g. Giunti and Chung, 2007). However, significant uncertainties are still affecting cCCSNe modelings and, more in general, the core-collapse processes of massive stars. Great advances are expected from GW and further neutrino detections that will be achieved with the next generation detectors. Both GW and neutrinos can provide important information as the degree of asymmetry in the matter distribution, as well as the rotation rate and the strength of the magnetic fields, that can be used as priors in numerical simulations (see, e.g., Chassande-Mottin et al., 2010, and reference therein).

Significant evidence of high-energy (TeV-PeV) cosmic neutrinos has recently been obtained from an extensive IceCube fourth-year data analysis (e.g. Aartsen et al., 2014). The lack of significant anisotropy in the data sky direction distribution is consistent with an (at least partially) extragalactic origin of the neutrino sources. High-energy neutrinos originate from hadrons acceleration, for example in jets, where after interacting with high energy photons produce charged pions decaying as high energy neutrinos ($>10^5 \text{ GeV}$; see, e.g., Waxman and Bahcall, 1997). Among the candidates that have been proposed to be responsible for the observed high-energy neutrino flux there are GRBs, AGN and blazars that are part of the main THESEUS targets in the context of the Time-domain Universe (Amati et al., 2017).

GRB are historically addressed among the best candidates of Ultra High Energy Cosmic Rays (UHECR) (e.g. Ghisellini et al., 2008), together with AGNs. Recent results from the Pierre Auger Observatory found evidence for dipolar anisotropy in CR at $E > 8 \times 10^{18} \text{ eV}$ towards a given direction in the sky, which is compatible with an extra-galactic origin, with possible suggestion that they are due to Large Scale Structures, with relatively nearby sources within 300 Mpc (e.g. Globus and Piran, 2017). As UHECR sources, GRBs are therefore addressed as promising high-energy neutrinos source candidates together with AGN. However, searches for neutrino events in coincidence with GRBs have not provided any confirmed association so far, possibly because of the average large distances of GRBs and/or a low neutrino production efficiency in bright GRBs. On the other hand, GRBs sources are particularly interesting since they could potentially emit also GWs. Possible detection could be achieved with the next generation of neutrino detectors. For long GRB, neutrinos emitted along the jet direction give the highest chances of detection. The expected rate of on-axis GRB that can be detected with IceCube has been estimated to be of the order of ~ 0.3 per year. (e.g.

Xiao et al., 2017). The lack of neutrinos from the very nearby short GRB associated with the GW170817 source has been interpreted to be due to the off-axis viewing angle of our line of sight with respect to the jet direction (Albert et al., 2017).

The feasibility of future joint EM and GW/neutrino observations are supported by theoretical background. In particular, for the case of short GRBs, according to the most recent studies (Kimura et al., 2017), high-energy neutrinos are thought to be most efficiently produced during the so called "Extended Emission" (EE), a softer, prolonged emission lasting few tens up to hundreds of seconds, that follows the initial count rate spike that characterises some short GRBs (Norris and Bonnell, 2006; Kaneko et al., 2015). It has been observed that about $\sim 25\%$ of short GRBs are accompanied by an EE (Sakamoto et al., 2011). This fraction is likely biased by the lack of X-ray survey instruments that could detect this component and likely more short GRBs are accompanied by EE (Nakamura et al., 2014), possibly up to 50% (Kimura et al., 2017). According to the neutrino detection probability estimates as a function of the short GRB with EE distance computed by Kimura et al. (2017), we expect that THESEUS/neutrino counterpart detection rate of on-axis short GRBs with EE within the horizon of IceCube and IceCubeGen2 is of the order of 0.02-0.25 and 0.1-0.5 per year, respectively. By considering the possibility to observe neutrinos also from short GRB with EE viewed off-axis, the THESEUS/neutrino counterpart detection rate may increase up to 0.2-4, and 0.5-7 per year, respectively. Future multi-messenger campaigns with deeper detector sensitivities will likely further constrain GRB progenitor models, clarifying the presence of a jet and its composition, and the relative neutrino/EM energy budgets and the role of GRBs as sources of UHCRs (Abbasi et al., 2012).

THESEUS/SXI and XGIS can detect SN shock breakout events up to ~ 50 Mpc (see previous section), thus leading to a potential joint neutrino detections of a few events per year with new generation neutrino detectors as Km3Net or IceCubeGen2. Blazars have been considered among the possible source candidates for the recently detected IceCube cosmic high-energy neutrino flux. THESEUS/blazars detection rate is estimated to be of hundreds per year (see Tab.2 in Amati et al. 2017).

6. Summary

The first detection of the electromagnetic counterparts of a GW source has confirmed a number of theoretical expectations and boosted the nascent multi-messenger astronomy. In this review we have discussed several classes of sources, including compact binary coalescences, core-collapsing massive stars, and instability episodes on NSs that are expected to originate simultaneously high-frequency GWs, neutrinos and EM emission across the entire EM spectrum, including in particular high energy emission (in X-rays and gamma-rays). We have shown that the mission concept THESEUS has the potential to play a crucial role in the multimessenger investigation of these sources. THESEUS, if approved, will have the capability to detect a very large number of transient sources in the X-ray and gamma-ray sky due to its wide field of view, and to automatically follow-up any high energy detection in the near infrared. In addition, it will be able to localize the sources down to arcminute (in gamma and X-rays) or to arcsecond (in NIR). As we have shown in this paper, the instrumental characteristics of THESEUS are ideal to operate in synergy with the facilities that will be available by the time of the mission: several new generation ground- and space-based telescopes, second- and third-generation GW detector networks and 10 km^3 neutrino detectors. This makes THESEUS perfectly suited for the coming golden era of multi-messenger astronomy and astrophysics.

References

- Aartsen, M.G., Ackermann, M., Adams, J., Aguilar, J.A., Ahlers, M., Ahrens, M., Altmann, D., Anderson, T., Argüelles, C., Arlen, T.C., et al., 2014. Observation of High-Energy Astrophysical Neutrinos in Three Years of IceCube Data. *Physical Review Letters* 113, 101101. doi:[10.1103/PhysRevLett.113.101101](https://doi.org/10.1103/PhysRevLett.113.101101), [arXiv:1405.5303](https://arxiv.org/abs/1405.5303).
- Abadie, J., Abbott, B.P., Abbott, R., Abernathy, M., Accadia, T., Acernese, F., Adams, C., Adhikari, R., Ajith, P., Allen, B., et al., 2010. TOPICAL REVIEW: Predictions for the rates of compact binary coalescences observable by ground-based gravitational-wave detectors. *Classical and Quantum Gravity* 27, 173001. doi:[10.1088/0264-9381/27/17/173001](https://doi.org/10.1088/0264-9381/27/17/173001), [arXiv:1003.2480](https://arxiv.org/abs/1003.2480).
- Abbasi, R., Abdou, Y., Abu-Zayyad, T., Ackermann, M., Adams, J., Aguilar, J.A., Ahlers, M., Altmann, D., Andeen, K., Auffenberg, J., et al., 2012. An absence of neutrinos associated with cosmic-ray acceleration in gamma-ray bursts. *Nature* 484, 351–354. doi:[10.1038/nature11068](https://doi.org/10.1038/nature11068), [arXiv:1204.4219](https://arxiv.org/abs/1204.4219).
- Abbott, B., Abbott, R., Adhikari, R., Ageev, A., Allen, B., Amin, R., Anderson, S.B., Anderson, W.G., Araya, M., Armandula, H., et al., 2005. Search for gravitational waves associated with the gamma ray burst GRB030329 using the LIGO detectors. *Phys.RevD* 72, 042002. doi:[10.1103/PhysRevD.72.042002](https://doi.org/10.1103/PhysRevD.72.042002), [arXiv:gr-qc/0501068](https://arxiv.org/abs/gr-qc/0501068).

- Abbott, B., Abbott, R., Adhikari, R., Agresti, J., Ajith, P., Allen, B., Amin, R., Anderson, S.B., Anderson, W.G., Arain, M., et al., 2008. Implications for the Origin of GRB 070201 from LIGO Observations. *Astroph. J.* 681, 1419–1430. doi:[10.1086/587954](https://doi.org/10.1086/587954), [arXiv:0711.1163](https://arxiv.org/abs/0711.1163).
- Abbott, B.P., Abbott, R., Abbott, T.D., Abernathy, M.R., Acernese, F., Ackley, K., Adams, C., Adams, T., Addesso, P., Adhikari, R.X., et al., 2016a. GW151226: Observation of Gravitational Waves from a 22-Solar-Mass Binary Black Hole Coalescence. *Physical Review Letters* 116, 241103. doi:[10.1103/PhysRevLett.116.241103](https://doi.org/10.1103/PhysRevLett.116.241103), [arXiv:1606.04855](https://arxiv.org/abs/1606.04855).
- Abbott, B.P., Abbott, R., Abbott, T.D., Abernathy, M.R., Acernese, F., Ackley, K., Adams, C., Adams, T., Addesso, P., Adhikari, R.X., et al., 2016b. Properties of the Binary Black Hole Merger GW150914. *Physical Review Letters* 116, 241102. doi:[10.1103/PhysRevLett.116.241102](https://doi.org/10.1103/PhysRevLett.116.241102), [arXiv:1602.03840](https://arxiv.org/abs/1602.03840).
- Abbott, B.P., Abbott, R., Abbott, T.D., Abernathy, M.R., Acernese, F., Ackley, K., Adams, C., Adams, T., Addesso, P., Adhikari, R.X., et al., 2016c. Prospects for Observing and Localizing Gravitational-Wave Transients with Advanced LIGO and Advanced Virgo. *Living Reviews in Relativity* 19, 1. doi:[10.1007/lrr-2016-1](https://doi.org/10.1007/lrr-2016-1), [arXiv:1304.0670](https://arxiv.org/abs/1304.0670).
- Abbott, B.P., Abbott, R., Abbott, T.D., Abernathy, M.R., Acernese, F., Ackley, K., Adams, C., Adams, T., Addesso, P., Adhikari, R.X., et al., 2017a. Search for Gravitational Waves Associated with Gamma-Ray Bursts during the First Advanced LIGO Observing Run and Implications for the Origin of GRB 150906B. *Astroph. J.* 841, 89. doi:[10.3847/1538-4357/aa6c47](https://doi.org/10.3847/1538-4357/aa6c47), [arXiv:1611.07947](https://arxiv.org/abs/1611.07947).
- Abbott, B.P., Abbott, R., Abbott, T.D., Abernathy, M.R., Ackley, K., Adams, C., Addesso, P., Adhikari, R.X., Adya, V.B., Affeldt, C., et al., 2017b. Exploring the sensitivity of next generation gravitational wave detectors. *Classical and Quantum Gravity* 34, 044001. doi:[10.1088/1361-6382/aa51f4](https://doi.org/10.1088/1361-6382/aa51f4), [arXiv:1607.08697](https://arxiv.org/abs/1607.08697).
- Abbott, B.P., Abbott, R., Abbott, T.D., Acernese, F., Ackley, K., Adams, C., Adams, T., Addesso, P., Adhikari, R.X., Adya, V.B., et al., 2017c. A gravitational-wave standard siren measurement of the Hubble constant. *Nature* 551, 85–88. doi:[10.1038/nature24471](https://doi.org/10.1038/nature24471), [arXiv:1710.05835](https://arxiv.org/abs/1710.05835).
- Abbott, B.P., Abbott, R., Abbott, T.D., Acernese, F., Ackley, K., Adams, C., Adams, T., Addesso, P., Adhikari, R.X., Adya, V.B., et al., 2017d. Gravitational Waves and Gamma-Rays from a Binary Neutron Star Merger: GW170817 and GRB 170817A. *Astroph. J.* 848, L13. doi:[10.3847/2041-8213/aa920c](https://doi.org/10.3847/2041-8213/aa920c), [arXiv:1710.05834](https://arxiv.org/abs/1710.05834).
- Abbott, B.P., Abbott, R., Abbott, T.D., Acernese, F., Ackley, K., Adams, C., Adams, T., Addesso, P., Adhikari, R.X., Adya, V.B., et al., 2017e. GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral. *Physical Review Letters* 119, 161101. doi:[10.1103/PhysRevLett.119.161101](https://doi.org/10.1103/PhysRevLett.119.161101), [arXiv:1710.05832](https://arxiv.org/abs/1710.05832).
- Abbott, B.P., Abbott, R., Abbott, T.D., Acernese, F., Ackley, K., Adams, C., Adams, T., Addesso, P., Adhikari, R.X., Adya, V.B., et al., 2017f. Multi-messenger Observations of a Binary Neutron Star Merger. *Astroph. J.* 848, L12. doi:[10.3847/2041-8213/aa91c9](https://doi.org/10.3847/2041-8213/aa91c9), [arXiv:1710.05833](https://arxiv.org/abs/1710.05833).
- Acernese, F., Agathos, M., Agatsuma, K., Aisa, D., Allemandou, N., Allocca, A., Amarni, J., Astone, P., Balestri, G., Ballardini, G., et al., 2015. Advanced Virgo: a second-generation interferometric gravitational wave detector. *Classical and Quantum Gravity* 32, 024001. doi:[10.1088/0264-9381/32/2/024001](https://doi.org/10.1088/0264-9381/32/2/024001), [arXiv:1408.3978](https://arxiv.org/abs/1408.3978).
- Albert, A., Andre, M., Anghinolfi, M., Ardid, M., Aubert, J.J., Aublin, J., Avgitas, T., Baret, B., Barrios-Marti, J., Basa, S., et al., 2017. Search for High-energy Neutrinos from Binary Neutron Star Merger GW170817 with ANTARES, IceCube, and the Pierre Auger Observatory. *ArXiv e-prints* [arXiv:1710.05839](https://arxiv.org/abs/1710.05839).
- Alexander, K.D., Berger, E., Fong, W., Williams, P.K.G., Guidorzi, C., Margutti, R., Metzger, B.D., Annis, J., Blanchard, P.K., Brout, D., Brown, D.A., Chen, H.Y., Chornock, R., Cowperthwaite, P.S., Drout, M., Eftekhari, T., Frieman, J., Holz, D.E., Nicholl, M., Rest, A., Sako, M., Soares-Santos, M., Villar, V.A., 2017. The Electromagnetic Counterpart of the Binary Neutron Star Merger LIGO/Virgo GW170817. VI. Radio Constraints on a Relativistic Jet and Predictions for Late-time Emission from the Kilonova Ejecta. *Astroph. J.* 848, L21. doi:[10.3847/2041-8213/aa905d](https://doi.org/10.3847/2041-8213/aa905d), [arXiv:1710.05457](https://arxiv.org/abs/1710.05457).
- Amati, L., O'Brien, P., Goetz, D., Bozzo, E., Tenzer, C., Frontera, F., Ghirlanda, G., Labanti, C., Osborne, J.P., Stratta, G., Tanvir, N., Willingale, R., Attina, P., Campana, R., Castro-Tirado, A.J., Contini, C., Fuschino, F., Gomboc, A., Hudec, R., Orleanski, P., Renotte, E., Rodic, T., Bagoly, Z., Blain, A., Callanan, P., Covino, S., Ferrara, A., Le Floch, E., Marisaldi, M., Mereghetti, S., Rosati, P., Vacchi, A., D'Avanzo, P., Giommi, P., Gomboc, A., Piranomonte, S., Piro, L., Reglero, V., Rossi, A., Santangelo, A., Salvaterra, R., Tagliaferri, G., Vergani, S., Vinciguerra, S., Briggs, M., Campolongo, E., Ciolfi, R., Connaughton, V., Cordier, B., Morelli, B., Orlandini, M., Adami, C., Argan, A., Atteia, J.L., Auricchio, N., Balazs, L., Balazsi, G., Basa, S., Basak, R., Bellutti, P., Bernardini, M.G., Bertuccio, G., Braga, J., Branchesi, M., Brandt, S., Brocato, E., Budtz-Jorgensen, C., Bulgarelli, A., Burderi, L., Camp, J., Capozziello, S., Caruana, J., Casella, P., Cenko, B., Chardonnet, P., Ciardi, B., Colafrancesco, S., Dainotti, M.G., D'Elia, V., De Martino, D., De Pasquale, M., Del Monte, E., Della Valle, M., Drago, A., Evangelista, Y., Feroci, M., Finelli, F., Fiorini, M., Fynbo, J., Gal-Yam, A., Gendre, B., Ghisellini, G., Grado, A., Guidorzi, C., Hafizi, M., Hanlon, L., Hjorth, J., Izzo, L., Kiss, L., Kumar, P., Kuvvetli, I., Lavagna, M., Li, T., Longo, F., Lyutikov, M., Maio, U., Maiorano, E., Malcovati, P., Malesani, D., Margutti, R., Martin-Carrillo, A., Masetti, N., McBreen, S., Mignani, R., Morgante, G., Mundell, C., Nargaard-Nielsen, H.U., Nicastro, L., Palazzi, E., Paltani, S., Panessa, F., Pareschi, G., Pe'er, A., Penacchioni, A.V., Pian, E., Piedipalumbo, E., Piran, T., Rauw, G., Razzano, M., Read, A., Rezzolla, L., Romano, P., Ruffini, R., Savaglio, S., Sguera, V., Schady, P., Skidmore, W., Song, L., Stanway, E., Starling, R., Topinka, M., Troja, E., van Putten, M., Vanzella, E., Vercellone, S., Wilson-Hodge, C., Yonetoku, D., Zampa, G., Zampa, N., Zhang, B., Zhang, B.B., Zhang, S., Zhang, S.N., Antonelli, A., Bianco, F., Boci, S., Boer, M., Botticella, M.T., Boulade, O., Butler, C., Campana, S., Capitanio, F., Celotti, A., Chen, Y., Colpi, M., Comastri, A., Cuby, J.G., Dadina, M., De Luca, A., Dong, Y.W., Etori, S., Gandhi, P., Geza, E., Greiner, J., Guiriec, S., Harms, J., Hernanz, M., Hornstrup, A., Hutchinson, I., Israel, G., Jonker, P., Kaneko, Y., Kawai, N., Wiersema, K., Korpela, S., Lebrun, V., Lu, F., MacFadyen, A., Malaguti, G., Maraschi, L., Melandri, A., Modjaz, M., Morris, D., Omodei, N., Paizis, A., Pata, P., Petrosian, V., Rachevski, A., Rhoads, J., Ryde, F., Sabau-Graziati, L., Shigehiro, N., Sims, M., Soomin, J., Szecsi, D., Urata, Y., Uslenghi, M., Valenziano, L., Vianello, G., Vojtech, S., Watson, D., Zicha, J., 2017. The Transient High Energy Sky and Early Universe Surveyor (THESEUS). *ArXiv e-prints* [arXiv:1710.04638](https://arxiv.org/abs/1710.04638).
- Andreoni, I., D'Avanzo, P., Campana, S., Branchesi, M., Bernardini, M.G., Della Valle, M., Mannucci, F., Melandri, A., Tagliaferri, G., 2016. A time domain experiment with Swift: monitoring of seven nearby galaxies. *Astron. & Astroph.* 587, A147. doi:[10.1051/0004-6361/201527167](https://doi.org/10.1051/0004-6361/201527167), [arXiv:1601.03739](https://arxiv.org/abs/1601.03739).
- Arcavi, I., Hosseinzadeh, G., Howell, D.A., McCully, C., Poznanski, D., Kasen, D., Barnes, J., Zaltzman, M., Vasylyev, S., Maoz, D., Valenti, S., 2017. Optical emission from a kilonova following a gravitational-wave-detected neutron-star merger. *Nature* 551, 64–66. doi:[10.1038/nature24291](https://doi.org/10.1038/nature24291), [arXiv:1710.05843](https://arxiv.org/abs/1710.05843).

- Baiotti, L., Rezzolla, L., 2017. Binary neutron star mergers: a review of Einstein richest laboratory. *Reports on Progress in Physics* 80, 096901. doi:[10.1088/1361-6633/aa67bb](https://doi.org/10.1088/1361-6633/aa67bb), [arXiv:1607.03540](https://arxiv.org/abs/1607.03540).
- Barnes, J., Kasen, D., Wu, M.R., Martínez-Pinedo, G., 2016. Radioactivity and Thermalization in the Ejecta of Compact Object Mergers and Their Impact on Kilonova Light Curves. *Astroph. J.* 829, 110. doi:[10.3847/0004-637X/829/2/110](https://doi.org/10.3847/0004-637X/829/2/110), [arXiv:1605.07218](https://arxiv.org/abs/1605.07218).
- Barthelmy, S.D., Chincarini, G., Burrows, D.N., Gehrels, N., Covino, S., Moretti, A., Romano, P., O'Brien, P.T., Sarazin, C.L., Kouveliotou, C., Goad, M., Vaughan, S., Tagliaferri, G., Zhang, B., Antonelli, L.A., Campana, S., Cummings, J.R., D'Avanzo, P., Davies, M.B., Giommi, P., Grupe, D., Kaneko, Y., Kennea, J.A., King, A., Kobayashi, S., Melandri, A., Meszaros, P., Nousek, J.A., Patel, S., Sakamoto, T., Wijers, R.A.M.J., 2005. An origin for short γ -ray bursts unassociated with current star formation. *Nature* 438, 994–996. doi:[10.1038/nature04392](https://doi.org/10.1038/nature04392), [arXiv:astro-ph/0511579](https://arxiv.org/abs/astro-ph/0511579).
- Bartos, I., 2016. Rapid and Bright Stellar-mass Binary Black Hole Mergers in Active Galactic Nuclei, in: *American Astronomical Society Meeting Abstracts*, p. 208.03.
- Bauswein, A., Janka, H.T., 2012. Measuring Neutron-Star Properties via Gravitational Waves from Neutron-Star Mergers. *Physical Review Letters* 108, 011101. doi:[10.1103/PhysRevLett.108.011101](https://doi.org/10.1103/PhysRevLett.108.011101), [arXiv:1106.1616](https://arxiv.org/abs/1106.1616).
- Belczynski, K., Ryu, T., Perna, R., Berti, E., Tanaka, T.L., Bulik, T., 2017. On the likelihood of detecting gravitational waves from Population III compact object binaries. *Mon. Not. R. Astron. Soc.* 471, 4702–4721. doi:[10.1093/mnras/stx1759](https://doi.org/10.1093/mnras/stx1759), [arXiv:1612.01524](https://arxiv.org/abs/1612.01524).
- Berger, E., 2014. Short-Duration Gamma-Ray Bursts. *Annu. Rev. Astro. Astroph.* 52, 43–105. doi:[10.1146/annurev-astro-081913-035926](https://doi.org/10.1146/annurev-astro-081913-035926), [arXiv:1311.2603](https://arxiv.org/abs/1311.2603).
- Berger, E., Fong, W., Chornock, R., 2013. An r-process Kilonova Associated with the Short-hard GRB 130603B. *Astroph. J.* 774, L23. doi:[10.1088/2041-8205/774/2/L23](https://doi.org/10.1088/2041-8205/774/2/L23), [arXiv:1306.3960](https://arxiv.org/abs/1306.3960).
- Bloom, J.S., Holz, D.E., Hughes, S.A., Menou, K., Adams, A., Anderson, S.F., Becker, A., Bower, G.C., Brandt, N., Cobb, B., Cook, K., Corsi, A., Covino, S., Fox, D., Fruchter, A., Fryer, C., Grindlay, J., Hartmann, D., Haiman, Z., Kocsis, B., Jones, L., Loeb, A., Marka, S., Metzger, B., Nakar, E., Nissanke, S., Perley, D.A., Piran, T., Poznanski, D., Prince, T., Schnittman, J., Soderberg, A., Strauss, M., Shawhan, P.S., Shoemaker, D.H., Sievers, J., Stubbs, C., Tagliaferri, G., Ubertini, P., Wozniak, P., 2009. Astro2010 Decadal Survey Whitepaper: Coordinated Science in the Gravitational and Electromagnetic Skies. *ArXiv e-prints* [arXiv:0902.1527](https://arxiv.org/abs/0902.1527).
- Bromberg, O., Nakar, E., Piran, T., Sari, R., 2013. Short versus Long and Collapsars versus Non-collapsars: A Quantitative Classification of Gamma-Ray Bursts. *Astroph. J.* 764, 179. doi:[10.1088/0004-637X/764/2/179](https://doi.org/10.1088/0004-637X/764/2/179), [arXiv:1210.0068](https://arxiv.org/abs/1210.0068).
- Burgess, J.M., Greiner, J., Begue, D., Giannios, D., Berlato, F., Lipunov, V.M., 2017. Viewing short Gamma-ray Bursts from a different angle. *ArXiv e-prints* [arXiv:1710.05823](https://arxiv.org/abs/1710.05823).
- Chassande-Mottin, E., LIGO Scientific Collaboration, Virgo Collaboration, 2010. Joint searches for gravitational waves and high-energy neutrinos, in: *Journal of Physics Conference Series*, p. 012002. doi:[10.1088/1742-6596/243/1/012002](https://doi.org/10.1088/1742-6596/243/1/012002).
- Chornock, R., Berger, E., Kasen, D., Cowperthwaite, P.S., Nicholl, M., Villar, V.A., Alexander, K.D., Blanchard, P.K., Eftekhari, T., Fong, W., Margutti, R., Williams, P.K.G., Annis, J., Brout, D., Brown, D.A., Chen, H.Y., Drout, M.R., Farr, B., Foley, R.J., Frieman, J.A., Fryer, C.L., Herner, K., Holz, D.E., Kessler, R., Matheson, T., Metzger, B.D., Quataert, E., Rest, A., Sako, M., Scolnic, D.M., Smith, N., Soares-Santos, M., 2017. The Electromagnetic Counterpart of the Binary Neutron Star Merger LIGO/Virgo GW170817. IV. Detection of Near-infrared Signatures of r-process Nucleosynthesis with Gemini-South. *Astroph. J.* 848, L19. doi:[10.3847/2041-8213/aa905c](https://doi.org/10.3847/2041-8213/aa905c), [arXiv:1710.05454](https://arxiv.org/abs/1710.05454).
- Cioffi, R., Lander, S.K., Manca, G.M., Rezzolla, L., 2011. Instability-driven Evolution of Poloidal Magnetic Fields in Relativistic Stars. *Astroph. J.* 736, L6. doi:[10.1088/2041-8205/736/1/L6](https://doi.org/10.1088/2041-8205/736/1/L6), [arXiv:1105.3971](https://arxiv.org/abs/1105.3971).
- Cioffi, R., Rezzolla, L., 2012. Poloidal-field Instability in Magnetized Relativistic Stars. *Astroph. J.* 760, 1. doi:[10.1088/0004-637X/760/1/1](https://doi.org/10.1088/0004-637X/760/1/1), [arXiv:1206.6604](https://arxiv.org/abs/1206.6604).
- Cioffi, R., Siegel, D.M., 2015a. Short gamma-ray bursts from binary neutron star mergers: the time-reversal scenario. *ArXiv e-prints* [arXiv:1505.01420](https://arxiv.org/abs/1505.01420).
- Cioffi, R., Siegel, D.M., 2015b. Short Gamma-Ray Bursts in the “Time-reversal” Scenario. *Astroph. J.* 798, L36. doi:[10.1088/2041-8205/798/2/L36](https://doi.org/10.1088/2041-8205/798/2/L36), [arXiv:1411.2015](https://arxiv.org/abs/1411.2015).
- Corsi, A., Owen, B.J., 2011. Maximum gravitational-wave energy emissible in magnetar flares. *Phys.RevD* 83, 104014. doi:[10.1103/PhysRevD.83.104014](https://doi.org/10.1103/PhysRevD.83.104014), [arXiv:1102.3421](https://arxiv.org/abs/1102.3421).
- Coulter, D.A., Foley, R.J., Kilpatrick, C.D., Drout, M.R., Piro, A.L., Shappee, B.J., Siebert, M.R., Simon, J.D., Ulloa, N., Kasen, D., Madore, B.F., Murguía-Berthier, A., Pan, Y.C., Prochaska, J.X., Ramirez-Ruiz, E., Rest, A., Rojas-Bravo, C., 2017. Swope Supernova Survey 2017a (SSS17a), the Optical Counterpart to a Gravitational Wave Source. *ArXiv e-prints* [arXiv:1710.05452](https://arxiv.org/abs/1710.05452).
- Cowperthwaite, P.S., Berger, E., Villar, V.A., Metzger, B.D., Nicholl, M., Chornock, R., Blanchard, P.K., Fong, W., Margutti, R., Soares-Santos, M., Alexander, K.D., Allam, S., Annis, J., Brout, D., Brown, D.A., Butler, R.E., Chen, H.Y., Diehl, H.T., Doctor, Z., Drout, M.R., Eftekhari, T., Farr, B., Finley, D.A., Foley, R.J., Frieman, J.A., Fryer, C.L., García-Bellido, J., Gill, M.S.S., Guillochon, J., Herner, K., Holz, D.E., Kasen, D., Kessler, R., Marriner, J., Matheson, T., Neilsen, Jr., E.H., Quataert, E., Palmese, A., Rest, A., Sako, M., Scolnic, D.M., Smith, N., Tucker, D.L., Williams, P.K.G., Balbinot, E., Carlin, J.L., Cook, E.R., Durret, F., Li, T.S., Lopes, P.A.A., Lourenço, A.C.C., Marshall, J.L., Medina, G.E., Muir, J., Muñoz, R.R., Sauseda, M., Schlegel, D.J., Secco, L.F., Vivas, A.K., Wester, W., Zenteno, A., Zhang, Y., Abbott, T.M.C., Banerji, M., Bechtol, K., Benoit-Lévy, A., Bertin, E., Buckley-Geer, E., Burke, D.L., Capozzi, D., Carrero Rosell, A., Carrasco Kind, M., Castander, F.J., Croce, M., Cunha, C.E., D'Andrea, C.B., da Costa, L.N., Davis, C., DePoy, D.L., Desai, S., Dietrich, J.P., Drlica-Wagner, A., Eifler, T.F., Evrard, A.E., Fernandez, E., Flaugher, B., Fosalba, P., Gaztanaga, E., Gerdes, D.W., Giannantonio, T., Goldstein, D.A., Gruen, D., Gruendl, R.A., Gutierrez, G., Honscheid, K., Jain, B., James, D.J., Jeltama, T., Johnson, M.W.G., Johnson, M.D., Kent, S., Krause, E., Kron, R., Kuehn, K., Nuropatkin, N., Lahav, O., Lima, M., Lin, H., Maia, M.A.G., March, M., Martini, P., McMahon, R.G., Menanteau, F., Miller, C.J., Miquel, R., Mohr, J.J., Neilsen, E., Nichol, R.C., Ogando, R.L.C., Plazas, A.A., Roe, N., Romer, A.K., Roodman, A., Rykoff, E.S., Sanchez, E., Scarpine, V., Schindler, R., Schubnell, M., Sevilla-Noarbe, I., Smith, M., Smith, R.C., Sobreira, F., Suchyta, E., Swanson, M.E.C., Tarle, G., Thomas, D., Thomas, R.C., Troxel, M.A., Vikram, V., Walker, A.R., Wechsler, R.H., Weller, J., Yanny, B., Zuntz, J., 2017. The Electromagnetic Counterpart of the Binary Neutron Star Merger LIGO/Virgo GW170817. II. UV, Optical, and Near-infrared Light Curves and Comparison to Kilonova Models. *Astroph. J.* 848, L17. doi:[10.3847/2041-8213/aa8fc7](https://doi.org/10.3847/2041-8213/aa8fc7), [arXiv:1710.05840](https://arxiv.org/abs/1710.05840).
- Dalal, N., Holz, D.E., Hughes, S.A., Jain, B., 2006. Short GRB and binary black hole standard sirens as a probe of dark energy. *Phys.RevD* 74,

063006. doi:[10.1103/PhysRevD.74.063006](https://doi.org/10.1103/PhysRevD.74.063006), [arXiv:astro-ph/0601275](https://arxiv.org/abs/astro-ph/0601275).
- D'Avanzo, P., 2015. Short gamma-ray bursts: A review. *Journal of High Energy Astrophysics* 7, 73–80. doi:[10.1016/j.jheap.2015.07.002](https://doi.org/10.1016/j.jheap.2015.07.002).
- Della Valle, M., Chincarini, G., Panagia, N., Tagliaferri, G., Malesani, D., Testa, V., Fugazza, D., Campana, S., Covino, S., Mangano, V., Antonelli, L.A., D'Avanzo, P., Hurley, K., Mirabel, I.F., Pellizza, L.J., Piranomonte, S., Stella, L., 2006. An enigmatic long-lasting γ -ray burst not accompanied by a bright supernova. *Nature* 444, 1050–1052. doi:[10.1038/nature05374](https://doi.org/10.1038/nature05374), [arXiv:astro-ph/0608322](https://arxiv.org/abs/astro-ph/0608322).
- Drago, A., Lavagno, A., Metzger, B.D., Pagliara, G., 2016. Quark deconfinement and the duration of short gamma-ray bursts. *Phys.RevD* 93, 103001. doi:[10.1103/PhysRevD.93.103001](https://doi.org/10.1103/PhysRevD.93.103001), [arXiv:1510.05581](https://arxiv.org/abs/1510.05581).
- Drago, A., Pagliara, G., 2017. Merger of two neutron stars: predictions from the two-families scenario. *ArXiv e-prints* [arXiv:1710.02003](https://arxiv.org/abs/1710.02003).
- Drout, M.R., Piro, A.L., Shappee, B.J., Kilpatrick, C.D., Simon, J.D., Contreras, C., Coulter, D.A., Foley, R.J., Siebert, M.R., Morrell, N., Boutsias, K., Di Mille, F., Holoien, T.W.S., Kasen, D., Kollmeier, J.A., Madore, B.F., Monson, A.J., Murguía-Berthier, A., Pan, Y.C., Prochaska, J.X., Ramirez-Ruiz, E., Rest, A., Adams, C., Alatalo, K., Bañados, E., Baughman, J., Beers, T.C., Bernstein, R.A., Bitsakis, T., Campillay, A., Hansen, T.T., Higgs, C.R., Ji, A.P., Maravelias, G., Marshall, J.L., Moni Bidin, C., Prieto, J.L., Rasmussen, K.C., Rojas-Bravo, C., Strom, A.L., Ulloa, N., Vargas-González, J., Wan, Z., Whitten, D.D., 2017. Light Curves of the Neutron Star Merger GW170817/SSS17a: Implications for R-Process Nucleosynthesis. *ArXiv e-prints* [arXiv:1710.05443](https://arxiv.org/abs/1710.05443).
- Eichler, D., Livio, M., Piran, T., Schramm, D.N., 1989. Nucleosynthesis, neutrino bursts and gamma-rays from coalescing neutron stars. *Nature* 340, 126–128. doi:[10.1038/340126a0](https://doi.org/10.1038/340126a0).
- Evans, P.A., Cenko, S.B., Kennea, J.A., Emery, S.W.K., Kuin, N.P.M., Korobkin, O., Wollaeger, R.T., Fryer, C.L., Madsen, K.K., Harrison, F.A., Xu, Y., Nakar, E., Hotokezaka, K., Lien, A., Campana, S., Oates, S.R., Troja, E., Breeveld, A.A., Marshall, F.E., Barthelmy, S.D., Beardmore, A.P., Burrows, D.N., Cusumano, G., D'Ai, A., D'Avanzo, P., D'Elia, V., de Pasquale, M., Even, W.P., Fontes, C.J., Forster, K., García, J., Giommi, P., Grefenstette, B., Gronwall, C., Hartmann, D.H., Heida, M., Hungerford, A.L., Kasliwal, M.M., Krimm, H.A., Levan, A.J., Malesani, D., Melandri, A., Miyasaka, H., Nousek, J.A., O'Brien, P.T., Osborne, J.P., Pagani, C., Page, K.L., Palmer, D.M., Perri, M., Pike, S., Racusin, J.L., Rosswog, S., Siegel, M.H., Sakamoto, T., Sbarufatti, B., Tagliaferri, G., Tanvir, N.R., Tohuvaohu, A., 2017. Swift and NuSTAR observations of GW170817: detection of a blue kilonova. *ArXiv e-prints* [arXiv:1710.05437](https://arxiv.org/abs/1710.05437).
- Fernández, R., Metzger, B.D., 2016. Electromagnetic Signatures of Neutron Star Mergers in the Advanced LIGO Era. *Annual Review of Nuclear and Particle Science* 66, 23–45. doi:[10.1146/annurev-nucl-102115-044819](https://doi.org/10.1146/annurev-nucl-102115-044819), [arXiv:1512.05435](https://arxiv.org/abs/1512.05435).
- Fox, D.B., Frail, D.A., Price, P.A., Kulkarni, S.R., Berger, E., Piran, T., Soderberg, A.M., Cenko, S.B., Cameron, P.B., Gal-Yam, A., Kasliwal, M.M., Moon, D.S., Harrison, F.A., Nakar, E., Schmidt, B.P., Penprase, B., Chevalier, R.A., Kumar, P., Roth, K., Watson, D., Lee, B.L., Shtetman, S., Phillips, M.M., Roth, M., McCarthy, P.J., Rauch, M., Cowie, L., Peterson, B.A., Rich, J., Kawai, N., Aoki, K., Kosugi, G., Totani, T., Park, H.S., MacFadyen, A., Hurley, K.C., 2005. The afterglow of GRB 050709 and the nature of the short-hard γ -ray bursts. *Nature* 437, 845–850. doi:[10.1038/nature04189](https://doi.org/10.1038/nature04189), [arXiv:astro-ph/0510110](https://arxiv.org/abs/astro-ph/0510110).
- Fynbo, J.P.U., Watson, D., Thöne, C.C., Sollerman, J., Bloom, J.S., Davis, T.M., Hjorth, J., Jakobsson, P., Jørgensen, U.G., Graham, J.F., Fruchter, A.S., Bersier, D., Kewley, L., Cassan, A., Castro Cerón, J.M., Foley, S., Gorosabel, J., Hinse, T.C., Horne, K.D., Jensen, B.L., Klose, S., Kocevski, D., Marquette, J.B., Perley, D., Ramirez-Ruiz, E., Stritzinger, M.D., Vreeswijk, P.M., Wijers, R.A.M., Woller, K.G., Xu, D., Zub, M., 2006. No supernovae associated with two long-duration γ -ray bursts. *Nature* 444, 1047–1049. doi:[10.1038/nature05375](https://doi.org/10.1038/nature05375), [arXiv:astro-ph/0608313](https://arxiv.org/abs/astro-ph/0608313).
- Gaisser, T.K., Halzen, F., Stanev, T., 1995. Particle astrophysics with high energy neutrinos. *Physics Reports* 258, 173–236. doi:[10.1016/0370-1573\(95\)00003-Y](https://doi.org/10.1016/0370-1573(95)00003-Y), [arXiv:hep-ph/9410384](https://arxiv.org/abs/hep-ph/9410384).
- Galama, T.J., Vreeswijk, P.M., van Paradijs, J., Kouveliotou, C., Augusteijn, T., Bönhardt, H., Brewer, J.P., Doublier, V., Gonzalez, J.F., Leibundgut, B., Lidman, C., Hainaut, O.R., Patat, F., Heise, J., in't Zand, J., Hurley, K., Groot, P.J., Strom, R.G., Mazzali, P.A., Iwamoto, K., Nomoto, K., Umeda, H., Nakamura, T., Young, T.R., Suzuki, T., Shigeyama, T., Koshut, T., Kippen, M., Robinson, C., de Wildt, P., Wijers, R.A.M.J., Tanvir, N., Greiner, J., Pian, E., Palazzi, E., Frontera, F., Masetti, N., Nicastro, L., Feroci, M., Costa, E., Piro, L., Peterson, B.A., Tinney, C., Boyle, B., Cannon, R., Stathakis, R., Sadler, E., Begam, M.C., Ianna, P., 1998. An unusual supernova in the error box of the γ -ray burst of 25 April 1998. *Nature* 395, 670–672. doi:[10.1038/27150](https://doi.org/10.1038/27150), [arXiv:astro-ph/9806175](https://arxiv.org/abs/astro-ph/9806175).
- Gao, H., Ding, X., Wu, X.F., Zhang, B., Dai, Z.G., 2013. Bright Broadband Afterglows of Gravitational Wave Bursts from Mergers of Binary Neutron Stars. *Astroph. J.* 771, 86. doi:[10.1088/0004-637X/771/2/86](https://doi.org/10.1088/0004-637X/771/2/86), [arXiv:1301.0439](https://arxiv.org/abs/1301.0439).
- Gehrels, N., Sarazin, C.L., O'Brien, P.T., Zhang, B., Barbier, L., Barthelmy, S.D., Blustin, A., Burrows, D.N., Cannizzo, J., Cummings, J.R., Goad, M., Holland, S.T., Hurkett, C.P., Kennea, J.A., Levan, A., Markwardt, C.B., Mason, K.O., Meszaros, P., Page, M., Palmer, D.M., Rol, E., Sakamoto, T., Willingale, R., Angelini, L., Beardmore, A., Boyd, P.T., Breeveld, A., Campana, S., Chester, M.M., Chincarini, G., Cominsky, L.R., Cusumano, G., de Pasquale, M., Fenimore, E.E., Giommi, P., Gronwall, C., Grupe, D., Hill, J.E., Hinshaw, D., Hjorth, J., Hullinger, D., Hurley, K.C., Klose, S., Kobayashi, S., Kouveliotou, C., Krimm, H.A., Mangano, V., Marshall, F.E., McGowan, K., Moretti, A., Mushotzky, R.F., Nakazawa, K., Norris, J.P., Nousek, J.A., Osborne, J.P., Page, K., Parsons, A.M., Patel, S., Perri, M., Poole, T., Romano, P., Roming, P.W.A., Rosen, S., Sato, G., Schady, P., Smale, A.P., Sollerman, J., Starling, R., Still, M., Suzuki, M., Tagliaferri, G., Takahashi, T., Tashiro, M., Tueller, J., Wells, A.A., White, N.E., Wijers, R.A.M.J., 2005. A short γ -ray burst apparently associated with an elliptical galaxy at redshift $z = 0.225$. *Nature* 437, 851–854. doi:[10.1038/nature04142](https://doi.org/10.1038/nature04142), [arXiv:astro-ph/0505630](https://arxiv.org/abs/astro-ph/0505630).
- Ghirlanda, G., Bernardini, M.G., Calderone, G., D'Avanzo, P., 2015a. Are short Gamma Ray Bursts similar to long ones? *Journal of High Energy Astrophysics* 7, 81–89. doi:[10.1016/j.jheap.2015.04.002](https://doi.org/10.1016/j.jheap.2015.04.002).
- Ghirlanda, G., Ghisellini, G., Nava, L., 2011. Short and long gamma-ray bursts: same emission mechanism? *Mon. Not. R. Astron. Soc.* 418, L109–L113. doi:[10.1111/j.1745-3933.2011.01154.x](https://doi.org/10.1111/j.1745-3933.2011.01154.x), [arXiv:1109.1833](https://arxiv.org/abs/1109.1833).
- Ghirlanda, G., Nava, L., Ghisellini, G., Celotti, A., Firmani, C., 2009. Short versus long gamma-ray bursts: spectra, energetics, and luminosities. *Astron. & Astroph.* 496, 585–595. doi:[10.1051/0004-6361/200811209](https://doi.org/10.1051/0004-6361/200811209), [arXiv:0902.0983](https://arxiv.org/abs/0902.0983).
- Ghirlanda, G., Salafia, O.S., Pescalli, A., Ghisellini, G., Salvaterra, R., Chassande-Mottin, E., Colpi, M., Nappo, F., D'Avanzo, P., Melandri, A., Bernardini, M.G., Branchesi, M., Campana, S., Ciolfi, R., Covino, S., Götz, D., Vergani, S.D., Zennaro, M., Tagliaferri, G., 2016. Short gamma-ray bursts at the dawn of the gravitational wave era. *Astron. & Astroph.* 594, A84. doi:[10.1051/0004-6361/201628993](https://doi.org/10.1051/0004-6361/201628993), [arXiv:1607.07875](https://arxiv.org/abs/1607.07875).
- Ghirlanda, G., Salvaterra, R., Burlon, D., Campana, S., Melandri, A., Bernardini, M.G., Covino, S., D'Avanzo, P., D'Elia, V., Ghisellini, G.,

- Nava, L., Prandoni, I., Sironi, L., Tagliaferri, G., Vergani, S.D., Wolter, A., 2013. Radio afterglows of a complete sample of bright Swift GRBs: predictions from present days to the SKA era. *Mon. Not. R. Astron. Soc.* 435, 2543–2551. doi:[10.1093/mnras/stt1466](https://doi.org/10.1093/mnras/stt1466), [arXiv:1307.7704](https://arxiv.org/abs/1307.7704).
- Ghirlanda, G., Salvaterra, R., Campana, S., Vergani, S.D., Japelj, J., Bernardini, M.G., Burlon, D., D’Avanzo, P., Melandri, A., Gomboc, A., Nappo, F., Paladini, R., Pescalli, A., Salafia, O.S., Tagliaferri, G., 2015b. Unveiling the population of orphan γ -ray bursts. *Astron. & Astroph.* 578, A71. doi:[10.1051/0004-6361/201526112](https://doi.org/10.1051/0004-6361/201526112), [arXiv:1504.02096](https://arxiv.org/abs/1504.02096).
- Ghirlanda, G., Salvaterra, R., Ghisellini, G., Mereghetti, S., Tagliaferri, G., Campana, S., Osborne, J.P., O’Brien, P., Tanvir, N., Willingale, D., Amati, L., Basa, S., Bernardini, M.G., Burlon, D., Covino, S., D’Avanzo, P., Frontera, F., Götz, D., Melandri, A., Nava, L., Piro, L., Vergani, S.D., 2015c. Accessing the population of high-redshift Gamma Ray Bursts. *Mon. Not. R. Astron. Soc.* 448, 2514–2524. doi:[10.1093/mnras/stv183](https://doi.org/10.1093/mnras/stv183), [arXiv:1502.02676](https://arxiv.org/abs/1502.02676).
- Ghisellini, G., Ghirlanda, G., Tavecchio, F., Fraternali, F., Pareschi, G., 2008. Ultra-high energy cosmic rays, spiral galaxies and magnetars. *Mon. Not. R. Astron. Soc.* 390, L88–L92. doi:[10.1111/j.1745-3933.2008.00547.x](https://doi.org/10.1111/j.1745-3933.2008.00547.x), [arXiv:0806.2393](https://arxiv.org/abs/0806.2393).
- Gil-Boella, I., 2016. Detection of Supernova Neutrinos. ArXiv e-prints [arXiv:1605.02204](https://arxiv.org/abs/1605.02204).
- Giunti, C., Chung, W.K., 2007. *Fundamentals of Neutrino Physics and Astrophysics*. Oxford University Press.
- Globus, N., Piran, T., 2017. The Extragalactic Ultra-high-energy Cosmic-Ray Dipole. *Astroph. J.* 850, L25. doi:[10.3847/2041-8213/aa991b](https://doi.org/10.3847/2041-8213/aa991b), [arXiv:1709.10110](https://arxiv.org/abs/1709.10110).
- Goldstein, A., Veres, P., Burns, E., Briggs, M.S., Hamburg, R., Kocevski, D., Wilson-Hodge, C.A., Preece, R.D., Poolakkil, S., Roberts, O.J., Hui, C.M., Connaughton, V., Racusin, J., von Kienlin, A., Dal Canton, T., Christensen, N., Littenberg, T., Siellez, K., Blackburn, L., Broida, J., Bissaldi, E., Cleveland, W.H., Gibby, M.H., Giles, M.M., Kippen, R.M., McBreen, S., McEnery, J., Meegan, C.A., Paciesas, W.S., Stanbro, M., 2017. An Ordinary Short Gamma-Ray Burst with Extraordinary Implications: Fermi-GBM Detection of GRB 170817A. *Astroph. J.* 848, L14. doi:[10.3847/2041-8213/aa8f41](https://doi.org/10.3847/2041-8213/aa8f41), [arXiv:1710.05446](https://arxiv.org/abs/1710.05446).
- Gottlieb, O., Nakar, E., Piran, T., 2018. The cocoon emission - an electromagnetic counterpart to gravitational waves from neutron star mergers. *Mon. Not. R. Astron. Soc.* 473, 576–584. doi:[10.1093/mnras/stx2357](https://doi.org/10.1093/mnras/stx2357), [arXiv:1705.10797](https://arxiv.org/abs/1705.10797).
- Gottlieb, O., Nakar, E., Piran, T., Hotokezaka, K., 2017. A cocoon shock breakout as the origin of the γ -ray emission in GW170817. ArXiv e-prints [arXiv:1710.05896](https://arxiv.org/abs/1710.05896).
- Granot, J., Panaitescu, A., Kumar, P., Woosley, S.E., 2002. Off-Axis Afterglow Emission from Jetted Gamma-Ray Bursts. *Astroph. J.* 570, L61–L64. doi:[10.1086/340991](https://doi.org/10.1086/340991), [arXiv:astro-ph/0201322](https://arxiv.org/abs/astro-ph/0201322).
- Guidorzi, C., Frontera, F., Montanari, E., 2004. The two large flares from SGR1900+14 observed with the BeppoSAX Gamma-Ray Burst Monitor: new results. *Nuclear Physics B Proceedings Supplements* 132, 536–541. doi:[10.1016/j.nuclphysbps.2004.04.090](https://doi.org/10.1016/j.nuclphysbps.2004.04.090).
- Guidorzi, C., Margutti, R., Brout, D., Scolnic, D., Fong, W., Alexander, K.D., Cowperthwaite, P.S., Annis, J., Berger, E., Blanchard, P.K., Chornock, R., Coppejans, D.L., Eftekhari, T., Frieman, J.A., Huterer, D., Nicholl, M., Soares-Santos, M., Terreran, G., Villar, V.A., Williams, P.K.G., 2017. Improved constraints on H_0 from a combined analysis of gravitational-wave and electromagnetic emission from GW170817. ArXiv e-prints [arXiv:1710.06426](https://arxiv.org/abs/1710.06426).
- Haggard, D., Nynka, M., Ruan, J.J., Kalogera, V., Cenko, S.B., Evans, P., Kennea, J.A., 2017. A Deep Chandra X-Ray Study of Neutron Star Coalescence GW170817. *Astroph. J.* 848, L25. doi:[10.3847/2041-8213/aa8ede](https://doi.org/10.3847/2041-8213/aa8ede), [arXiv:1710.05852](https://arxiv.org/abs/1710.05852).
- Hallinan, G., Corsi, A., Mooley, K.P., Hotokezaka, K., Nakar, E., Kasliwal, M.M., Kaplan, D.L., Frail, D.A., Myers, S.T., Murphy, T., De, K., Dobie, D., Allison, J.R., Bannister, K.W., Bhalerao, V., Chandra, P., Clarke, T.E., Giacintucci, S., Ho, A.Y.Q., Horesh, A., Kassim, N.E., Kulkarni, S.R., Lenc, E., Lockman, F.J., Lynch, C., Nichols, D., Nissanke, S., Palliyaguru, N., Peters, W.M., Piran, T., Rana, J., Sadler, E.M., Singer, L.P., 2017. A Radio Counterpart to a Neutron Star Merger. ArXiv e-prints [arXiv:1710.05435](https://arxiv.org/abs/1710.05435).
- Harry, G.M., LIGO Scientific Collaboration, 2010. Advanced LIGO: the next generation of gravitational wave detectors. *Classical and Quantum Gravity* 27, 084006. doi:[10.1088/0264-9381/27/8/084006](https://doi.org/10.1088/0264-9381/27/8/084006).
- IceCube-Gen2 Collaboration: Aartsen, M.G., Ackermann, M., Adams, J., Aguilar, J.A., Ahlers, M., Ahrens, M., Altmann, D., Anderson, T., et al., 2014. IceCube-Gen2: A Vision for the Future of Neutrino Astronomy in Antarctica. ArXiv e-prints [arXiv:1412.5106](https://arxiv.org/abs/1412.5106).
- Jin, Z.P., Li, X., Cano, Z., Covino, S., Fan, Y.Z., Wei, D.M., 2015. The Light Curve of the Macronova Associated with the Long-Short Burst GRB 060614. *Astroph. J.* 811, L22. doi:[10.1088/2041-8205/811/2/L22](https://doi.org/10.1088/2041-8205/811/2/L22), [arXiv:1507.07206](https://arxiv.org/abs/1507.07206).
- Kaneko, Y., Bostanci, Z.F., Göğüş, E., Lin, L., 2015. Short gamma-ray bursts with extended emission observed with Swift/BAT and Fermi/GBM. *Mon. Not. R. Astron. Soc.* 452, 824–837. doi:[10.1093/mnras/stv1286](https://doi.org/10.1093/mnras/stv1286), [arXiv:1506.05899](https://arxiv.org/abs/1506.05899).
- Kasliwal, M.M., Korobkin, O., Lau, R.M., Wollaeger, R., Fryer, C.L., 2017. Infrared Emission from Kilonovae: The Case of the Nearby Short Hard Burst GRB 160821B. *Astroph. J.* 843, L34. doi:[10.3847/2041-8213/aa799d](https://doi.org/10.3847/2041-8213/aa799d), [arXiv:1706.04647](https://arxiv.org/abs/1706.04647).
- Kathirgamaraju, A., Barniol Duran, R., Giannios, D., 2017. Off-axis short GRBs from structured jets as counterparts to GW events. ArXiv e-prints [arXiv:1708.07488](https://arxiv.org/abs/1708.07488).
- Kawamura, T., Giacomazzo, B., Kastaun, W., Ciolfi, R., Endrizzi, A., Baiotti, L., Perna, R., 2016. Binary neutron star mergers and short gamma-ray bursts: Effects of magnetic field orientation, equation of state, and mass ratio. *Phys.RevD* 94, 064012. doi:[10.1103/PhysRevD.94.064012](https://doi.org/10.1103/PhysRevD.94.064012), [arXiv:1607.01791](https://arxiv.org/abs/1607.01791).
- Kimura, S.S., Murase, K., Mészáros, P., Kiuchi, K., 2017. High-energy Neutrino Emission from Short Gamma-Ray Bursts: Prospects for Coincident Detection with Gravitational Waves. *Astroph. J.* 848, L4. doi:[10.3847/2041-8213/aa8d14](https://doi.org/10.3847/2041-8213/aa8d14), [arXiv:1708.07075](https://arxiv.org/abs/1708.07075).
- Koshut, T.M., Paciesas, W.S., Kouveliotou, C., van Paradijs, J., Pendleton, G.N., Fishman, G.J., Meegan, C.A., 1995. T_{90} as a Measurement of the Duration of GRBs, in: *American Astronomical Society Meeting Abstracts* #186, p. 886.
- Kouveliotou, C., Meegan, C.A., Fishman, G.J., Bhat, N.P., Briggs, M.S., Koshut, T.M., Paciesas, W.S., Pendleton, G.N., 1993. Identification of two classes of gamma-ray bursts. *Astroph. J.* 413, L101–L104. doi:[10.1086/186969](https://doi.org/10.1086/186969).
- Lasky, P.D., Haskell, B., Ravi, V., Howell, E.J., Coward, D.M., 2014. Nuclear equation of state from observations of short gamma-ray burst remnants. *Phys.RevD* 89, 047302. doi:[10.1103/PhysRevD.89.047302](https://doi.org/10.1103/PhysRevD.89.047302), [arXiv:1311.1352](https://arxiv.org/abs/1311.1352).
- Lasky, P.D., Zink, B., Kokkotas, K.D., 2012. Gravitational Waves and Hydromagnetic Instabilities in Rotating Magnetized Neutron Stars. ArXiv e-prints [arXiv:1203.3590](https://arxiv.org/abs/1203.3590).
- Lazzati, D., Deich, A., Morsony, B.J., Workman, J.C., 2017a. Off-axis emission of short γ -ray bursts and the detectability of electromagnetic counterparts of gravitational-wave-detected binary mergers. *Mon. Not. R. Astron. Soc.* 471, 1652–1661. doi:[10.1093/mnras/stx1683](https://doi.org/10.1093/mnras/stx1683),

- [arXiv:1610.01157](#).
- Lazzati, D., López-Cámara, D., Cantiello, M., Morsony, B.J., Perna, R., Workman, J.C., 2017b. Off-axis Prompt X-Ray Transients from the Cocoon of Short Gamma-Ray Bursts. *Astroph. J.* 848, L6. doi:[10.3847/2041-8213/aa8f3d](#), [arXiv:1709.01468](#).
- Lazzati, D., Perna, R., Morsony, B.J., López-Cámara, D., Cantiello, M., Ciolfi, R., Giacomazzo, B., Workman, J.C., 2017c. Late time afterglow observations reveal a collimated relativistic jet in the ejecta of the binary neutron star merger GW170817. ArXiv e-prints [arXiv:1712.03237](#).
- Le, T., Dermer, C.D., 2007. On the Redshift Distribution of Gamma-Ray Bursts in the Swift Era. *Astroph. J.* 661, 394–415. doi:[10.1086/513460](#), [arXiv:astro-ph/0610043](#).
- Li, L.X., Paczyński, B., 1998a. Transient Events from Neutron Star Mergers. *Astroph. J.* 507, L59–L62. doi:[10.1086/311680](#), [arXiv:astro-ph/9807272](#).
- Li, L.X., Paczyński, B., 1998b. Transient Events from Neutron Star Mergers. *Astroph. J.* 507, L59–L62. doi:[10.1086/311680](#), [arXiv:astro-ph/9807272](#).
- Li, W., Chornock, R., Leaman, J., Filippenko, A.V., Poznanski, D., Wang, X., Ganeshalingam, M., Mannucci, F., 2011. Nearby supernova rates from the Lick Observatory Supernova Search - III. The rate-size relation, and the rates as a function of galaxy Hubble type and colour. *Mon. Not. R. Astron. Soc.* 412, 1473–1507. doi:[10.1111/j.1365-2966.2011.18162.x](#), [arXiv:1006.4613](#).
- Loeb, A., 2016. Electromagnetic Counterparts to Black Hole Mergers Detected by LIGO. *Astroph. J.* 819, L21. doi:[10.3847/2041-8205/819/2/L21](#), [arXiv:1602.04735](#).
- Logue, J., Ott, C.D., Heng, I.S., Kalmus, P., Scargill, J.H.C., 2012. Inferring core-collapse supernova physics with gravitational waves. *Phys.RevD* 86, 044023. doi:[10.1103/PhysRevD.86.044023](#), [arXiv:1202.3256](#).
- Margutti, R., Berger, E., Fong, W., Guidorzi, C., Alexander, K.D., Metzger, B.D., Blanchard, P.K., Cowperthwaite, P.S., Chornock, R., Eftekhari, T., Nicholl, M., Villar, V.A., Williams, P.K.G., Annis, J., Brown, D.A., Chen, H., Doctor, Z., Frieman, J.A., Holz, D.E., Sako, M., Soares-Santos, M., 2017. The Electromagnetic Counterpart of the Binary Neutron Star Merger LIGO/Virgo GW170817. V. Rising X-Ray Emission from an Off-axis Jet. *Astroph. J.* 848, L20. doi:[10.3847/2041-8213/aa9057](#), [arXiv:1710.05431](#).
- Mereghetti, S., Pons, J.A., Melatos, A., 2015. Magnetars: Properties, Origin and Evolution. *Space Sci. Mod. Rev.* 191, 315–338. doi:[10.1007/s11214-015-0146-y](#), [arXiv:1503.06313](#).
- Messenger, C., Bulten, H.J., Crowder, S.G., Dergachev, V., Galloway, D.K., Goetz, E., Jonker, R.J.G., Lasky, P.D., Meadors, G.D., Melatos, A., Premachandra, S., Riles, K., Sammut, L., Thrane, E.H., Whelan, J.T., Zhang, Y., 2015. Gravitational waves from Scorpius X-1: A comparison of search methods and prospects for detection with advanced detectors. *Phys.RevD* 92, 023006. doi:[10.1103/PhysRevD.92.023006](#), [arXiv:1504.05889](#).
- Metzger, B.D., 2017. Kilonovae. *Living Reviews in Relativity* 20, 3. doi:[10.1007/s41114-017-0006-z](#), [arXiv:1610.09381](#).
- Metzger, B.D., Martínez-Pinedo, G., Darbha, S., Quataert, E., Arcones, A., Kasen, D., Thomas, R., Nugent, P., Panov, I.V., Zinner, N.T., 2010. Electromagnetic counterparts of compact object mergers powered by the radioactive decay of r-process nuclei. *Mon. Not. R. Astron. Soc.* 406, 2650–2662. doi:[10.1111/j.1365-2966.2010.16864.x](#), [arXiv:1001.5029](#).
- Metzger, B.D., Piro, A.L., 2014. Optical and X-ray emission from stable millisecond magnetars formed from the merger of binary neutron stars. *Mon. Not. R. Astron. Soc.* 439, 3916–3930. doi:[10.1093/mnras/stu247](#), [arXiv:1311.1519](#).
- Mooley, K.P., Nakar, E., Hotokezaka, K., Hallinan, G., Corsi, A., Frail, D.A., Horesh, A., Murphy, T., Lenc, E., Kaplan, D.L., De, K., Dobie, D., Chandra, P., Deller, A., Gottlieb, O., Kasliwal, M.M., Kulkarni, S.R., Myers, S.T., Nissanke, S., Piran, T., Lynch, C., Bhalerao, V., Bourke, S., Bannister, K.W., Singer, L.P., 2017. A mildly relativistic wide-angle outflow in the neutron star merger GW170817. ArXiv e-prints [arXiv:1711.11573](#).
- Nakamura, T., Kashiyama, K., Nakauchi, D., Suwa, Y., Sakamoto, T., Kawai, N., 2014. Soft X-Ray Extended Emissions of Short Gamma-Ray Bursts as Electromagnetic Counterparts of Compact Binary Mergers: Possible Origin and Detectability. *Astroph. J.* 796, 13. doi:[10.1088/0004-637X/796/1/13](#), [arXiv:1312.0297](#).
- Narayan, R., Paczynski, B., Piran, T., 1992. Gamma-ray bursts as the death throes of massive binary stars. *Astroph. J.* 395, L83–L86. doi:[10.1086/186493](#), [arXiv:astro-ph/9204001](#).
- Nava, L., Ghirlanda, G., Ghisellini, G., Celotti, G., 2011. Fermi/GBM and BATSE gamma-ray bursts: comparison of the spectral properties. *Mon. Not. R. Astron. Soc.* 415, 3153–3162. doi:[10.1111/j.1365-2966.2011.18928.x](#), [arXiv:1012.3968](#).
- Nicholl, M., Berger, E., Kasen, D., Metzger, B.D., Elias, J., Briceño, C., Alexander, K.D., Blanchard, P.K., Chornock, R., Cowperthwaite, P.S., Eftekhari, T., Fong, W., Margutti, R., Villar, V.A., Williams, P.K.G., Brown, W., Annis, J., Bahramian, A., Brout, D., Brown, D.A., Chen, H.Y., Clemens, J.C., Dennihy, E., Dunlap, B., Holz, D.E., Marchesini, E., Massaro, F., Moskowitz, N., Pelisoli, I., Rest, A., Ricci, F., Sako, M., Soares-Santos, M., Strader, J., 2017. The Electromagnetic Counterpart of the Binary Neutron Star Merger LIGO/Virgo GW170817. III. Optical and UV Spectra of a Blue Kilonova from Fast Polar Ejecta. *Astroph. J.* 848, L18. doi:[10.3847/2041-8213/aa9029](#), [arXiv:1710.05456](#).
- Norris, J.P., Bonnell, J.T., 2006. Short Gamma-Ray Bursts with Extended Emission. *Astroph. J.* 643, 266–275. doi:[10.1086/502796](#), [arXiv:astro-ph/0601190](#).
- Ott, C.D., 2010. Computational Models of Stellar Collapse, Core-Collapse Supernovae, and Black Hole Formation, in: APS April Meeting Abstracts.
- Paczynski, B., 1986. Gamma-ray bursters at cosmological distances. *Astroph. J.* 308, L43–L46. doi:[10.1086/184740](#).
- Paschalidis, V., Ruiz, M., Shapiro, S.L., 2015. Relativistic Simulations of Black Hole-Neutron Star Coalescence: The Jet Emerges. *Astroph. J.* 806, L14. doi:[10.1088/2041-8205/806/1/L14](#), [arXiv:1410.7392](#).
- Perna, R., Lazzati, D., Giacomazzo, B., 2016. Short Gamma-Ray Bursts from the Merger of Two Black Holes. *Astroph. J.* 821, L18. doi:[10.3847/2041-8205/821/1/L18](#), [arXiv:1602.05140](#).
- Pescalli, A., Ghirlanda, G., Salafia, O.S., Ghisellini, G., Nappo, F., Salvaterra, R., 2015. Luminosity function and jet structure of Gamma-Ray Burst. *Mon. Not. R. Astron. Soc.* 447, 1911–1921. doi:[10.1093/mnras/stu2482](#), [arXiv:1409.1213](#).
- Pescalli, A., Ghirlanda, G., Salvaterra, R., Ghisellini, G., Vergani, S.D., Nappo, F., Salafia, O.S., Melandri, A., Covino, S., Götz, D., 2016. The rate and luminosity function of long gamma ray bursts. *Astron. & Astroph.* 587, A40. doi:[10.1051/0004-6361/201526760](#), [arXiv:1506.05463](#).
- Phinney, E.S., 2009. Finding and Using Electromagnetic Counterparts of Gravitational Wave Sources, in: *astro2010: The Astronomy and Astrophysics Decadal Survey*. [arXiv:0903.0098](#).

- Pian, E., D'Avanzo, P., Benetti, S., Branchesi, M., Brocato, E., Campana, S., Cappellaro, E., Covino, S., D'Elia, V., Fynbo, J.P.U., Getman, F., Ghirlanda, G., Ghisellini, G., Grado, A., Greco, G., Hjorth, J., Kouveliotou, C., Levan, A., Limatola, L., Malesani, D., Mazzali, P.A., Melandri, A., Møller, P., Nicastro, L., Palazzi, E., Piranomonte, S., Rossi, A., Salafia, O.S., Selsing, J., Stratta, G., Tanaka, M., Tanvir, N.R., Tomasella, L., Watson, D., Yang, S., Amati, L., Antonelli, L.A., Ascenzi, S., Bernardini, M.G., Boër, M., Bufano, F., Bulgarelli, A., Capaccioli, M., Casella, P., Castro-Tirado, A.J., Chassande-Mottin, E., Ciolfi, R., Copperwheat, C.M., Dadina, M., De Cesare, G., di Paola, A., Fan, Y.Z., Gendre, B., Giuffrida, G., Giunta, A., Hunt, L.K., Israel, G.L., Jin, Z.P., Kasliwal, M.M., Klose, S., Lisi, M., Longo, F., Maiorano, E., Mapelli, M., Masetti, N., Nava, L., Patricelli, B., Perley, D., Pescalli, A., Piran, T., Possenti, A., Pulone, L., Razzano, M., Salvaterra, R., Schipani, P., Spera, M., Stammer, A., Stella, L., Tagliaferri, G., Testa, V., Troja, E., Turatto, M., Vergani, S.D., Vergani, D., 2017. Spectroscopic identification of r-process nucleosynthesis in a double neutron-star merger. *Nature* 551, 67–70. doi:[10.1038/nature24298](https://doi.org/10.1038/nature24298), [arXiv:1710.05858](https://arxiv.org/abs/1710.05858).
- Piro, A.L., Giacomazzo, B., Perna, R., 2017. The Fate of Neutron Star Binary Mergers. *Astroph. J.* 844, L19. doi:[10.3847/2041-8213/aa7f2f](https://doi.org/10.3847/2041-8213/aa7f2f), [arXiv:1704.08697](https://arxiv.org/abs/1704.08697).
- Planck Collaboration, Ade, P.A.R., Aghanim, N., Arnaud, M., Ashdown, M., Aumont, J., Baccigalupi, C., Banday, A.J., Barreiro, R.B., Bartlett, J.G., et al., 2016. Planck 2015 results. XIII. Cosmological parameters. *Astron. & Astroph.* 594, A13. doi:[10.1051/0004-6361/201525830](https://doi.org/10.1051/0004-6361/201525830), [arXiv:1502.01589](https://arxiv.org/abs/1502.01589).
- Powell, J., Gossan, S.E., Logue, J., Heng, I.S., 2016. Inferring the core-collapse supernova explosion mechanism with gravitational waves. *Phys.RevD* 94, 123012. doi:[10.1103/PhysRevD.94.123012](https://doi.org/10.1103/PhysRevD.94.123012), [arXiv:1610.05573](https://arxiv.org/abs/1610.05573).
- Punturo, M., Abernathy, M., Acernese, F., Allen, B., Andersson, N., Arun, K., Barone, F., Barr, B., Barsuglia, M., Beker, M., Beveridge, N., Birindelli, S., Bose, S., Bosi, L., Braccini, S., Bradaschia, C., Bulik, T., Calloni, E., Cella, G., Chassande Mottin, E., Chelkowski, S., Chincarini, A., Clark, J., Coccia, E., Colacino, C., Colas, J., Cumming, A., Cunningham, L., Cuoco, E., Danilishin, S., Danzmann, K., De Luca, G., De Salvo, R., Dent, T., De Rosa, R., Di Fiore, L., Di Virgilio, A., Doets, M., Fafone, V., Falferi, P., Flaminio, R., Franc, J., Frasconi, F., Freise, A., Fulda, P., Gair, J., Gemme, G., Gennai, A., Giazotto, A., Glampedakis, K., Granata, M., Grote, H., Guidi, G., Hammond, G., Hannam, M., Harms, J., Heiwerdt, D., Hendry, M., Heng, I., Hennes, E., Hild, S., Hough, J., Husa, S., Huttner, S., Jones, G., Khalili, F., Kokeyama, K., Kokkotas, K., Krishnan, B., Lorenzini, M., Lück, H., Majorana, E., Mandel, I., Mandic, V., Martin, I., Michel, C., Minenkov, Y., Morgado, N., Mosca, S., Mours, B., Müller-Ebhardt, H., Murray, P., Nawrodt, R., Nelson, J., Oshaughnessy, R., Ott, C.D., Palomba, C., Paoli, A., Parguez, G., Pasqualetti, A., Passaquieti, R., Passuello, D., Pinard, L., Poggiani, R., Popolizio, P., Prato, M., Puppo, P., Rabeling, D., Rapagnani, P., Read, J., Regimbau, T., Rehbein, H., Reid, S., Rezzolla, L., Ricci, F., Richard, F., Rocchi, A., Rowan, S., Rüdiger, A., Sassolas, B., Sathyaprakash, B., Schnabel, R., Schwarz, C., Seidel, P., Sintes, A., Somiya, K., Speirits, F., Strain, K., Strigin, S., Sutton, P., Tarabrin, S., Thüring, A., van den Brand, J., van Leewen, C., van Veggel, M., van den Broeck, C., Vecchio, A., Veitch, J., Vetrano, F., Vicere, A., Müller Vyatchanin, S., Willke, B., Woan, G., Wolfango, P., Yamamoto, K., 2010. The Einstein Telescope: a third-generation gravitational wave observatory. *Classical and Quantum Gravity* 27, 194002. doi:[10.1088/0264-9381/27/19/194002](https://doi.org/10.1088/0264-9381/27/19/194002).
- Rezzolla, L., Giacomazzo, B., Baiotti, L., Granot, J., Kouveliotou, C., Aloy, M.A., 2011. The Missing Link: Merging Neutron Stars Naturally Produce Jet-like Structures and Can Power Short Gamma-ray Bursts. *Astroph. J.* 732, L6. doi:[10.1088/2041-8205/732/1/L6](https://doi.org/10.1088/2041-8205/732/1/L6), [arXiv:1101.4298](https://arxiv.org/abs/1101.4298).
- Rezzolla, L., Kumar, P., 2015. A Novel Paradigm for Short Gamma-Ray Bursts With Extended X-Ray Emission. *Astroph. J.* 802, 95. doi:[10.1088/0004-637X/802/2/95](https://doi.org/10.1088/0004-637X/802/2/95), [arXiv:1410.8560](https://arxiv.org/abs/1410.8560).
- Rezzolla, L., Takami, K., 2016. Gravitational-wave signal from binary neutron stars: A systematic analysis of the spectral properties. *Phys.RevD* 93, 124051. doi:[10.1103/PhysRevD.93.124051](https://doi.org/10.1103/PhysRevD.93.124051), [arXiv:1604.00246](https://arxiv.org/abs/1604.00246).
- Riess, A.G., Macri, L.M., Hoffmann, S.L., Scolnic, D., Casertano, S., Filippenko, A.V., Tucker, B.E., Reid, M.J., Jones, D.O., Silverman, J.M., Chornock, R., Challis, P., Yuan, W., Brown, P.J., Foley, R.J., 2016. A 2.4% Determination of the Local Value of the Hubble Constant. *Astroph. J.* 826, 56. doi:[10.3847/0004-637X/826/1/56](https://doi.org/10.3847/0004-637X/826/1/56), [arXiv:1604.01424](https://arxiv.org/abs/1604.01424).
- Rossi, E., Lazzati, D., Rees, M.J., 2002. Afterglow light curves, viewing angle and the jet structure of γ -ray bursts. *Mon. Not. R. Astron. Soc.* 332, 945–950. doi:[10.1046/j.1365-8711.2002.05363.x](https://doi.org/10.1046/j.1365-8711.2002.05363.x), [arXiv:astro-ph/0112083](https://arxiv.org/abs/astro-ph/0112083).
- Rosswog, S., Piran, T., Nakar, E., 2013. The multimessenger picture of compact object encounters: binary mergers versus dynamical collisions. *Mon. Not. R. Astron. Soc.* 430, 2585–2604. doi:[10.1093/mnras/sts708](https://doi.org/10.1093/mnras/sts708), [arXiv:1204.6240](https://arxiv.org/abs/1204.6240).
- Ruiz, M., Lang, R.N., Paschalidis, V., Shapiro, S.L., 2016. Binary Neutron Star Mergers: A Jet Engine for Short Gamma-Ray Bursts. *Astroph. J.* 824, L6. doi:[10.3847/2041-8205/824/1/L6](https://doi.org/10.3847/2041-8205/824/1/L6), [arXiv:1604.02455](https://arxiv.org/abs/1604.02455).
- Sakamoto, T., Barthelmy, S.D., Baumgartner, W.H., Cummings, J.R., Fenimore, E.E., Gehrels, N., Krimm, H.A., Markwardt, C.B., Palmer, D.M., Parsons, A.M., Sato, G., Stamatikos, M., Tueller, J., Ukwatta, T.N., Zhang, B., 2011. The Second Swift Burst Alert Telescope Gamma-Ray Burst Catalog. *Astroph. J. Suppl.* 195, 2. doi:[10.1088/0067-0049/195/1/2](https://doi.org/10.1088/0067-0049/195/1/2), [arXiv:1104.4689](https://arxiv.org/abs/1104.4689).
- Salafia, O.S., Ghisellini, G., Ghirlanda, G., 2017a. Jet-driven and jet-less fireballs from compact binary mergers. *ArXiv e-prints* [arXiv:1710.05859](https://arxiv.org/abs/1710.05859).
- Salafia, O.S., Ghisellini, G., Ghirlanda, G., Colpi, M., 2017b. GRB170817A: a giant flare from a jet-less double neutron-star merger? *ArXiv e-prints* [arXiv:1711.03112](https://arxiv.org/abs/1711.03112).
- Santander, M., 2016. The Dawn of Multi-Messenger Astronomy. *ArXiv e-prints* [arXiv:1606.09335](https://arxiv.org/abs/1606.09335).
- Sathyaprakash, B., Abernathy, M., Acernese, F., Ajith, P., Allen, B., Amaro-Seoane, P., Andersson, N., Aoudia, S., Arun, K., Astone, P., et al., 2012. Scientific objectives of Einstein Telescope. *Classical and Quantum Gravity* 29, 124013. doi:[10.1088/0264-9381/29/12/124013](https://doi.org/10.1088/0264-9381/29/12/124013), [arXiv:1206.0331](https://arxiv.org/abs/1206.0331).
- Savchenko, V., Ferrigno, C., Kuulkers, E., Bazzano, A., Bozzo, E., Brandt, S., Chenevez, J., Courvoisier, T.J.L., Diehl, R., Domingo, A., Hanlon, L., Jourdain, E., von Kienlin, A., Laurent, P., Lebrun, F., Lutovinov, A., Martin-Carrillo, A., Mereghetti, S., Natalucci, L., Rodi, J., Roques, J.P., Sunyaev, R., Ubertini, P., 2017. INTEGRAL Detection of the First Prompt Gamma-Ray Signal Coincident with the Gravitational-wave Event GW170817. *Astroph. J.* 848, L15. doi:[10.3847/2041-8213/aa8f94](https://doi.org/10.3847/2041-8213/aa8f94), [arXiv:1710.05449](https://arxiv.org/abs/1710.05449).
- Scholberg, K., 2012. Supernova Neutrino Detection. *Annual Review of Nuclear and Particle Science* 62, 81–103. doi:[10.1146/annurev-nucl-102711-095006](https://doi.org/10.1146/annurev-nucl-102711-095006), [arXiv:1205.6003](https://arxiv.org/abs/1205.6003).
- Schutz, B.F., 1986. Determining the Hubble constant from gravitational wave observations. *Nature* 323, 310. doi:[10.1038/323310a0](https://doi.org/10.1038/323310a0).
- Seto, N., Muto, T., 2011. Resonant trapping of stars by merging massive black hole binaries. *Mon. Not. R. Astron. Soc.* 415, 3824–3830.

- doi:[10.1111/j.1365-2966.2011.18988.x](https://doi.org/10.1111/j.1365-2966.2011.18988.x), [arXiv:1105.1845](https://arxiv.org/abs/1105.1845).
- Siegel, D.M., Ciolfi, R., 2016a. Electromagnetic Emission from Long-lived Binary Neutron Star Merger Remnants. I. Formulation of the Problem. *Astroph. J.* 819, 14. doi:[10.3847/0004-637X/819/1/14](https://doi.org/10.3847/0004-637X/819/1/14), [arXiv:1508.07911](https://arxiv.org/abs/1508.07911).
- Siegel, D.M., Ciolfi, R., 2016b. Electromagnetic Emission from Long-lived Binary Neutron Star Merger Remnants. II. Lightcurves and Spectra. *Astroph. J.* 819, 15. doi:[10.3847/0004-637X/819/1/15](https://doi.org/10.3847/0004-637X/819/1/15), [arXiv:1508.07939](https://arxiv.org/abs/1508.07939).
- Smartt, S.J., Chen, T.W., Jerkstrand, A., Coughlin, M., Kankare, E., Sim, S.A., Fraser, M., Inerra, C., Maguire, K., Chambers, K.C., Huber, M.E., Krühler, T., Leloudas, G., Magee, M., Shingles, L.J., Smith, K.W., Young, D.R., Tonry, J., Kotak, R., Gal-Yam, A., Lyman, J.D., Homan, D.S., Agliozzo, C., Anderson, J.P., Angus, C.R., Ashall, C., Barbarino, C., Bauer, F.E., Berton, M., Botticella, M.T., Bulla, M., Bulger, J., Cannizzaro, G., Cano, Z., Cartier, R., Cikota, A., Clark, P., De Cia, A., Della Valle, M., Denneau, L., Dennefeld, M., Dessart, L., Dimitriadis, G., Elias-Rosa, N., Firth, R.E., Flewelling, H., Flörs, A., Franckowiak, A., Frohmaier, C., Galbany, L., González-Gaitán, S., Greiner, J., Gromadzki, M., Guelbenzu, A.N., Gutiérrez, C.P., Hamanowicz, A., Hanlon, L., Harmanen, J., Heintz, K.E., Heinze, A., Hernandez, M.S., Hodgkin, S.T., Hook, I.M., Izzo, L., James, P.A., Jonker, P.G., Kerzendorf, W.E., Klose, S., Kostrzewa-Rutkowska, Z., Kowalski, M., Kromer, M., Kuncarayakti, H., Lawrence, A., Lowe, T.B., Magnier, E.A., Manulis, I., Martin-Carrillo, A., Matilla, S., McBrien, O., Müller, A., Nordin, J., O’Neill, D., Onori, F., Palmerio, J.T., Pastorello, A., Patat, F., Pignata, G., Podsiadlowski, P., Pumo, M.L., Prentice, S.J., Rau, A., Razza, A., Rest, A., Reynolds, T., Roy, R., Rüter, A.J., Rybicki, K.A., Salmon, L., Schady, P., Schultz, A.S.B., Schweyer, T., Seitzenzahl, I.R., Smith, M., Sollerman, J., Stalder, B., Stubbs, C.W., Sullivan, M., Szegedi, H., Taddia, F., Taubenberger, S., Terreran, G., van Soelen, B., Vos, J., Wainscoat, R.J., Walton, N.A., Waters, C., Weiland, H., Willman, M., Wiseman, P., Wright, D.E., Wyrzykowski, Ł., Yaron, O., 2017. A kilonova as the electromagnetic counterpart to a gravitational-wave source. *Nature* 551, 75–79. doi:[10.1038/nature24303](https://doi.org/10.1038/nature24303), [arXiv:1710.05841](https://arxiv.org/abs/1710.05841).
- Somiya, K., 2012. Detector configuration of KAGRA-the Japanese cryogenic gravitational-wave detector. *Classical and Quantum Gravity* 29, 124007. doi:[10.1088/0264-9381/29/12/124007](https://doi.org/10.1088/0264-9381/29/12/124007), [arXiv:1111.7185](https://arxiv.org/abs/1111.7185).
- Stanek, K.Z., Matheson, T., Garnavich, P.M., Martini, P., Berlind, P., Caldwell, N., Challis, P., Brown, W.R., Schild, R., Krisciunas, K., Calkins, M.L., Lee, J.C., Hathi, N., Jansen, R.A., Windhorst, R., Echevarria, L., Eisenstein, D.J., Pindor, B., Olszewski, E.W., Harding, P., Holland, S.T., Bersier, D., 2003. Spectroscopic Discovery of the Supernova 2003dh Associated with GRB 030329. *Astroph. J.* 591, L17–L20. doi:[10.1086/376976](https://doi.org/10.1086/376976), [arXiv:astro-ph/0304173](https://arxiv.org/abs/astro-ph/0304173).
- Takami, K., Rezzolla, L., Baiotti, L., 2014. Constraining the Equation of State of Neutron Stars from Binary Mergers. *Physical Review Letters* 113, 091104. doi:[10.1103/PhysRevLett.113.091104](https://doi.org/10.1103/PhysRevLett.113.091104), [arXiv:1403.5672](https://arxiv.org/abs/1403.5672).
- Tanaka, M., Utsumi, Y., Mazzali, P.A., Tominaga, N., Yoshida, M., Sekiguchi, Y., Morokuma, T., Motohara, K., Ohta, K., Kawabata, K.S., Abe, F., Aoki, K., Asakura, Y., Baar, S., Barway, S., Bond, I.A., Doi, M., Fujiyoshi, T., Furusawa, H., Honda, S., Itoh, Y., Kawabata, M., Kawai, N., Kim, J.H., Lee, C.H., Miyazaki, S., Morihana, K., Nagashima, H., Nagayama, T., Nakaoka, T., Nakata, F., Ohsawa, R., Ohshima, T., Okita, H., Saito, T., Sumi, T., Tajitsu, A., Takahashi, J., Takayama, M., Tamura, Y., Tanaka, I., Terai, T., Tristram, P.J., Yasuda, N., Zenko, T., 2017. Kilonova from post-merger ejecta as an optical and near-infrared counterpart of GW170817. *ArXiv e-prints* [arXiv:1710.05850](https://arxiv.org/abs/1710.05850).
- Tanvir, N.R., Levan, A.J., Fruchter, A.S., Hjorth, J., Hounsell, R.A., Wiersema, K., Tunnicliffe, R.L., 2013. A ‘kilonova’ associated with the short-duration γ -ray burst GRB 130603B. *Nature* 500, 547–549. doi:[10.1038/nature12505](https://doi.org/10.1038/nature12505), [arXiv:1306.4971](https://arxiv.org/abs/1306.4971).
- Tanvir, N.R., Levan, A.J., González-Fernández, C., Korobkin, O., Mandel, I., Rosswog, S., Hjorth, J., D’Avanzo, P., Fruchter, A.S., Fryer, C.L., Kangas, T., Milvang-Jensen, B., Rosetti, S., Steeghs, D., Wollaeger, R.T., Cano, Z., Copperwheat, C.M., Covino, S., D’Elia, V., de Ugarte Postigo, A., Evans, P.A., Even, W.P., Fairhurst, S., Figuera Jaimes, R., Fontes, C.J., Fujii, Y.I., Fynbo, J.P.U., Gompertz, B.P., Greiner, J., Hodosan, G., Irwin, M.J., Jakobsson, P., Jørgensen, U.G., Kann, D.A., Lyman, J.D., Malesani, D., McMahon, R.G., Melandri, A., O’Brien, P.T., Osborne, J.P., Palazzi, E., Perley, D.A., Pian, E., Piranomonte, S., Rabus, M., Rol, E., Rowlinson, A., Schulze, S., Sutton, P., Thöne, C.C., Ulaczyk, K., Watson, D., Wiersema, K., Wijers, R.A.M.J., 2017. The Emergence of a Lanthanide-rich Kilonova Following the Merger of Two Neutron Stars. *Astroph. J.* 848, L27. doi:[10.3847/2041-8213/aa90b6](https://doi.org/10.3847/2041-8213/aa90b6), [arXiv:1710.05455](https://arxiv.org/abs/1710.05455).
- Thompson, C., Duncan, R.C., 1995. The soft gamma repeaters as very strongly magnetized neutron stars - I. Radiative mechanism for outbursts. *Mon. Not. R. Astron. Soc.* 275, 255–300. doi:[10.1093/mnras/275.2.255](https://doi.org/10.1093/mnras/275.2.255).
- Toma, K., Ioka, K., Sakamoto, T., Nakamura, T., 2007. Low-Luminosity GRB 060218: A Collapsar Jet from a Neutron Star, Leaving a Magnetar as a Remnant? *Astroph. J.* 659, 1420–1430. doi:[10.1086/512481](https://doi.org/10.1086/512481), [arXiv:astro-ph/0610867](https://arxiv.org/abs/astro-ph/0610867).
- Troja, E., Lipunov, V.M., Mundell, C.G., Butler, N.R., Watson, A.M., Kobayashi, S., Cenko, S.B., Marshall, F.E., Ricci, R., Fruchter, A., Wieringa, M.H., Gorbovskoy, E.S., Kornilov, V., Kutyrev, A., Lee, W.H., Toy, V., Tyurina, N.V., Budnev, N.M., Buckley, D.A.H., Gonzalez, J., Gress, O., Horesh, A., Panasyuk, M.I., Prochaska, J.X., Ramirez-Ruiz, E., Rebolo Lopez, R., Richer, M.G., Roman-Zuniga, C., Serra-Ricart, M., Yurkov, V., Gehrels, N., 2017. Significant and variable linear polarization during the prompt optical flash of GRB 160625B. *Nature* 547, 425–427.
- van Putten, M.H.P.M., Levinson, A., Frontera, F., Guidorzi, C., Amati, L., Della Valle, M., 2017. GPU-searches for broadband extended emission in gravitational waves in nearby energetic core-collapse supernovae. *ArXiv e-prints* [arXiv:1709.04455](https://arxiv.org/abs/1709.04455).
- Virgili, F.J., Liang, E.W., Zhang, B., 2009. Low-luminosity gamma-ray bursts as a distinct GRB population: a firmer case from multiple criteria constraints. *Mon. Not. R. Astron. Soc.* 392, 91–103. doi:[10.1111/j.1365-2966.2008.14063.x](https://doi.org/10.1111/j.1365-2966.2008.14063.x), [arXiv:0801.4751](https://arxiv.org/abs/0801.4751).
- Waxman, E., Bahcall, J., 1997. High Energy Neutrinos from Cosmological Gamma-Ray Burst Fireballs. *Physical Review Letters* 78, 2292–2295. doi:[10.1103/PhysRevLett.78.2292](https://doi.org/10.1103/PhysRevLett.78.2292), [arXiv:astro-ph/9701231](https://arxiv.org/abs/astro-ph/9701231).
- Woosley, S.E., Bloom, J.S., 2006. The Supernova Gamma-Ray Burst Connection. *Annu. Rev. Astro. Astroph.* 44, 507–556. doi:[10.1146/annurev.astro.43.072103.150558](https://doi.org/10.1146/annurev.astro.43.072103.150558), [arXiv:astro-ph/0609142](https://arxiv.org/abs/astro-ph/0609142).
- Xiao, D., Liu, L.D., Dai, Z.G., Wu, X.F., 2017. Afterglows and Kilonovae Associated with Nearby Low-Luminosity Short-Duration Gamma-Ray Bursts: Application to GW170817/GRB170817A. *ArXiv e-prints* [arXiv:1710.05910](https://arxiv.org/abs/1710.05910).
- Xu, D., Starling, R.L.C., Fynbo, J.P.U., Sollerman, J., Yost, S., Watson, D., Foley, S., O’Brien, P.T., Hjorth, J., 2009. In Search of Progenitors for Supernovaeless Gamma-Ray Bursts 060505 and 060614: Re-examination of Their Afterglows. *Astroph. J.* 696, 971–979. doi:[10.1088/0004-637X/696/1/971](https://doi.org/10.1088/0004-637X/696/1/971), [arXiv:0812.0979](https://arxiv.org/abs/0812.0979).
- Yonetoku, D., Murakami, T., Nakamura, T., Yamazaki, R., Inoue, A.K., Ioka, K., 2004. Gamma-Ray Burst Formation Rate Inferred from the Spectral Peak Energy-Peak Luminosity Relation. *Astroph. J.* 609, 935–951. doi:[10.1086/421285](https://doi.org/10.1086/421285), [arXiv:astro-ph/0309217](https://arxiv.org/abs/astro-ph/0309217).
- Yu, Y.W., Zhang, B., Gao, H., 2013. Bright “Merger-nova” from the Remnant of a Neutron Star Binary Merger: A Signature of a Newly Born, Massive, Millisecond Magnetar. *Astroph. J.* 776, L40. doi:[10.1088/2041-8205/776/2/L40](https://doi.org/10.1088/2041-8205/776/2/L40), [arXiv:1308.0876](https://arxiv.org/abs/1308.0876).

- Zhang, B., 2013. Early X-Ray and Optical Afterglow of Gravitational Wave Bursts from Mergers of Binary Neutron Stars. *Astroph. J.* 763, L22. doi:[10.1088/2041-8205/763/1/L22](https://doi.org/10.1088/2041-8205/763/1/L22), [arXiv:1212.0773](https://arxiv.org/abs/1212.0773).
- Zhang, B., Mészáros, P., 2002. An Analysis of Gamma-Ray Burst Spectral Break Models. *Astroph. J.* 581, 1236–1247. doi:[10.1086/344338](https://doi.org/10.1086/344338), [arXiv:astro-ph/0206158](https://arxiv.org/abs/astro-ph/0206158).
- Zhang, F.W., Shao, L., Yan, J.Z., Wei, D.M., 2012. Revisiting the Long/Soft-Short/Hard Classification of Gamma-Ray Bursts in the Fermi Era. *Astroph. J.* 750, 88. doi:[10.1088/0004-637X/750/2/88](https://doi.org/10.1088/0004-637X/750/2/88), [arXiv:1201.1549](https://arxiv.org/abs/1201.1549).
- Zhao, W., Wen, L., 2017. Localization accuracy of compact binary coalescences detected by the third-generation gravitational-wave detectors and implication for cosmology. ArXiv e-prints [arXiv:1710.05325](https://arxiv.org/abs/1710.05325).