

Publication Year	2005
Acceptance in OA@INAF	2024-03-14T14:23:50Z
Title	Testing GR with the Double Pulsar: Recent Results
Authors	Kramer, M.; Lorimer, D. R.; Lyne, A. G.; McLaughlin, M.; BURGAY, MARTA; et al.
DOI	10.48550/arXiv.astro-ph/0503386
Handle	http://hdl.handle.net/20.500.12386/34964

Testing GR with the Double Pulsar: Recent Results

M. Kramer, D.R. Lorimer, A.G. Lyne, M. McLaughlin
University of Manchester, Jodrell Bank Observatory, UK

M. Burgay, N. D'Amico, A. Possenti
INAF - Osservatorio Astronomica di Cagliari, Italy

F. Camilo
Columbia Astrophysics Laboratory, Columbia University, U SA

P.C.C. Freire
NAIC, Arecibo Observatory, USA

B.C. Joshi
NCRA, Pune, India

R.N. Manchester, J. Reynolds, J. Sarkissian
Australia Telescope National Facility, CSIRO, Australia

I.H. Stairs, R. D. Ferdman
Department of Physics and Astronomy, University of British Columbia, Canada

This first ever double pulsar system consists of two pulsars orbiting the common center of mass in a slightly eccentric orbit of only 2.4-hr duration. The pair of pulsars with pulse periods of 22 ms and 2.8 sec, respectively, confirms the long-proposed recycling theory for millisecond pulsars and provides an exciting opportunity to study the works of pulsar magnetospheres by a very fortunate geometrical alignment of the orbit relative to our line-of-sight. In particular, this binary system represents a truly unique laboratory for relativistic gravitational physics. This contribution serves as an update on the currently obtained results and their consequences for the test of general relativity in the strong-field regime. A complete and more up-to-date report of the timing results will be presented elsewhere shortly.

I. INTRODUCTION

In this year 2005 we celebrate the work of Albert Einstein, remembering his enormous contribution to our understanding of nature and the Universe. One of the best ways for honoring his work is to point out that still today, a hundred of years later, hundreds of scientists around the world are deeply involved in searching for the limits up to which his centennial theory of general relativity (GR) can be applied. To date GR has passed all observational tests with flying colours. Still, it is the continued aim of many physicists to achieve more stringent tests by either increasing the precision of the tests or by testing different aspects. Some of the most stringent tests are obtained by satellite experiments in the solar system. One must not forget, however, that these solar-system experiments are all made in the gravitational weak-field regime and that they will never be able to provide tests in the strong-field limit where deviations from GR may appear more clearly or even for the first time (see e.g. [1]). This strong-field regime is best explored using radio pulsars.

Pulsars, highly magnetized rotating neutron stars, are unique and versatile objects which can be used to

study an extremely wide range of physical and astrophysical problems. Beside testing theories of gravity one can study the Galaxy and the interstellar medium, stars, binary systems and their evolution, plasma physics and solid state physics under extreme conditions. In these proceedings, we will present such applications for gravitational physics made possible by the first ever discovered double pulsar [2, 3]. We will demonstrate that this rare binary system represents a truly unique laboratory for relativistic gravity. We will present an update on the currently obtained timing results and their consequences for tests of GR. A complete and more up-to-date report of the timing results will be presented elsewhere shortly.

II. THE DOUBLE PULSAR

Our team discovered the 22.8-ms pulsar J0737-3039 in April 2003 [2] in an extension to the hugely successful Parkes Multi-beam survey [4]. It was soon found to be a member of the most extreme relativistic binary system ever discovered: its short orbital period ($P_b = 2:4$ hrs) is combined with a remarkably high value of periastron advance

($\dot{\Omega} = 16.9 \text{ deg yr}^{-1}$, i.e. four times larger than for PSR B1913+16!) that was measurable after only a few days of observations. The system parameters predict that the two members of the binary system will coalesce on a short time scale of only 85 Myr. This boosts the hopes for detecting a merger of two neutron stars with first-generation ground-based gravitational wave detectors by a factor of several compared to previous estimates based on only the double neutron stars B1534+12 and B1913+16 [2, 5].

In October 2003, we detected radio pulses from the second neutron star when data sets covering the full orbital period were analysed [3]. The reason why signals from the 2.8-s pulsar companion (now called PSR J0737-3039B, hereafter "B") to the millisecond pulsar (now called PSR J0737-3039A, hereafter "A") had not been found earlier, became clear when it was realized that B was only bright for two short parts of the orbit. For the remainder of the orbit, the pulsar B is extremely weak and only detectable with the most sensitive equipment. The detection of a young companion B around an old millisecond pulsar A confirms the evolution scenario proposed for recycled pulsars (e.g. [6, 7]) and made this already exciting system sensational, providing a truly unique testbed for relativistic gravity.

III. STRONG-FIELD TESTS OF GENERAL RELATIVITY

Since neutron stars are very compact massive objects, the double pulsar (and other double neutron

star systems) can be considered as almost ideal point sources for testing theories of gravity in the strong-gravitational-field limit. Tests can be performed when a number of relativistic corrections to the Keplerian description of an orbit, the so-called "post-Keplerian" (PK) parameters, can be measured. For point masses with negligible spin contributions, the PK parameters in each theory should only be functions of the a priori unknown neutron star masses and the well measurable Keplerian parameters.

With the two masses as the only free parameters, the measurement of three or more PK parameters over-constrains the system, and thereby provides a test ground for theories of gravity. In a theory that describes a binary system correctly, the PK parameters produce theory-dependent lines in a mass-mass diagram that all intersect in a single point.

As A has the faster pulse period (and is bright throughout the entire orbit apart from a 27-s eclipse at superior conjunction), we can time A much more accurately than B and measure precise PK parameters for A's orbit. In GR, the five most important PK parameters are given to first post-Newtonian (1PN, or $O(v^2=c^2)$) order by [8]:

$$\dot{\Omega} = 3T^{2=3} \frac{P_b}{2}^{5=3} \frac{1}{1-e^2} (M_A + M_B)^{2=3}; \quad (1)$$

$$= T^{2=3} \frac{P_b}{2}^{1=3} e \frac{M_B (M_A + 2M_B)}{(M_A + M_B)^{4=3}}; \quad (2)$$

$$P_b = \frac{192}{5} T^{5=3} \frac{P_b}{2}^{5=3} \frac{1 + \frac{73}{24}e^2 + \frac{37}{96}e^4}{(1-e^2)^{7=2}} \frac{M_A M_B}{(M_A + M_B)^{1=3}}; \quad (3)$$

$$r = T M_B; \quad (4)$$

$$s = T^{1=3} \frac{P_b}{2}^{2=3} \times \frac{(M_A + M_B)^{2=3}}{M_B}; \quad (5)$$

where P_b is the period and e the eccentricity and x the semi-major axis measured in light-s of the binary orbit. The masses M_A and M_B of A and B, respectively, are expressed in solar masses (M_\odot). We define the constant $T = GM_\odot/c^3 = 4.925490947 \text{ s}$ where G denotes the Newtonian constant of gravity and c the speed of light. The first PK parameter, $\dot{\Omega}$, is the eas-

iest to measure and describes the relativistic advance of periastron. According to Eqn. 1 it provides an immediate measurement of the total mass of the system, $(M_A + M_B)$. The parameter s denotes the amplitude of delays in arrival times caused by the varying effects of the gravitational redshift and time dilation (second order Doppler) as the pulsar moves in its elliptical or-

bit at varying distances from the companion and with varying speeds. The decay of the orbit due to gravitational wave damping is expressed by the change in orbital period, P_b . The other two parameters, r and s , are related to the Shapiro delay caused by the gravitational field of the companion. These parameters are only measurable, depending on timing precision, if the orbit is seen nearly edge-on. For pulsar A, all these quantities have indeed been measured, providing a large number of available tests. In fact, in addition to tests with these PK parameters, the possibility to measure the orbit of both A and B opens up opportunities that go well beyond what is possible with previously known double neutron stars, as we will describe now.

With a measurement of the projected semi-major axes of the orbits of both A and B, we obtain a precise measurement of the mass ratio $R(M_A; M_B)$, from Kepler's third law,

$$R(M_A; M_B) = M_A/M_B = x_B/x_A. \quad (6)$$

For every realistic theory of gravity, we can expect the mass ratio, R , to follow this simple relation [9], at least to 1PN order. Most importantly, the R value is not only theory-independent, but also independent of strong-field (self-field) effects which is not the case for PK-parameters. This provides a stringent and new constraint for tests of gravitational theories as any combination of masses derived from the PK-parameters must be consistent with the mass ratio derived from Kepler's 3rd law. With five PK parameters already available, this additional constraint makes the double pulsar the most overdetermined system to date where the most relativistic effects can be studied in the strong-field limit.

IV. TIMING OF THE DOUBLE PULSAR

Our observations already provide measurements for all five PK parameters listed above. This includes a measurement of an orbital decay of the binary orbit due to gravitational wave emission with a rate of 7mm/day. As indicated earlier, we can use these results to test GR in a very elegant way [9]. The unique relationship between the two masses of the system predicted by GR (or any other theory) for each PK parameter can be plotted in a diagram showing the mass of A on one axis and B on the other. We expect all curves, including that of the mass ratio R , to intersect in a single point if the chosen theory (here GR) is a valid description of the nature of this system. Such tests have been possible to date in PSR B1913+16 (e.g. [10]) and for PSR B1534+12 (e.g. [11]). However, in neither of these systems were so many curves available as for the double pulsar system for which we derive a $M_A - M_B$ plot as shown in Fig. 1.

It turns out that, as another stroke of luck, we are observing the system almost completely edge-on which allows us to determine a Shapiro delay to very high precision. It also means, however, that we can probe a pulsar magnetosphere for the very first time using a background beacon. Results of the magnetospheric interactions between A and B [12] and the eclipse of A's signal at its superior conjunction [3, 13, 14] have been presented elsewhere and must be considered in search for a possible contamination of the timing data and hence a violation of our assumption that we deal with a "clean" system of point sources.

While all studies so far confirm the cleanness of the system, we also have to consider that the times-of-arrival (TOAs) are obtained with a standard "template matching" procedure that involves a cross-correlation of the observed pulse profile with high signal-to-noise ratio template (e.g. [15]). Any change in the pulse profile could therefore lead to systematic variations in the measured TOAs. For this reason it was necessary to perform detailed studies of the profiles of A and B and to investigate any possible profile evolution with time. Indeed, profile changes on secular time scales are expected for the double pulsar for the following reason.

In GR, the proper reference frame of a freely falling object suffers a precession with respect to a distant observer, called geodetic precession. In a binary pulsar system this geodetic precession leads to a relativistic spin-orbit coupling, analogous of spin-orbit coupling in atomic physics [16]. As a consequence, the pulsar spins precess about the total angular momentum, changing the relative orientation of the pulsars to one another and toward Earth. Since the orbital angular momentum is much larger than the pulsars' angular momenta, the total angular momentum is practically represented by the orbital angular momentum. The precession rate [17] depends on the period and the eccentricity of the orbit as well as the masses of A and B. With the orbital parameters of the double pulsar, GR predicts precession periods of only 75 yr for A and 71 yr for B.

Geodetic precession should have a direct effect on the timing as it causes the polar angles of the spins and hence the effects of aberration to change with time [9]. These changes modify the observed orbital parameters, like projected semi-major axis and eccentricity, which differ from the intrinsic values by an aberration dependent term, potentially allowing us to infer the system geometry [18]. Other consequences of geodetic precession can be expected to be detected much sooner and are directly relevant for the timing of A and B. These arise from variations in the pulse shape due to changing cuts through the emission beam as the pulsar spin axes precess. Moreover, geodetic precession also leads to a change in the relative alignment of the pulsar magnetospheres, so that the visibility pat-

TABLE I: Observed and derived parameters of PSRs J0737 3039A and B. Standard errors are given in parentheses after the values and are in units of the least significant digit(s).

Pulsar	PSR J0737 3039A	PSR J0737 3039B
Pulse period P (ms)	22.699378556138(2)	2773.4607474(4)
Period derivative \dot{P}	$1.7596(2) \times 10^{-18}$	$0.88(13) \times 10^{-15}$
Epoch of period (MJD)	52870.0	
Right ascension (J2000)	$07^{\text{h}}37^{\text{m}}51^{\text{s}}.24795(2)$	
Declination (J2000)	$30^{\circ}39'40''.7247(6)$	
Orbital period P_b (day)	0.1022515628(2)	
Eccentricity e	0.087778(2)	
Epoch of periastron T_0 (MJD)	52870.0120588(3)	
Advance of periastron $\dot{\omega}$ (deg yr $^{-1}$)	16.900(2)	
Longitude of periastron ω (deg)	73.805(1)	$73.805 + 180.0$
Projected semi-major axis $x = asini=c$ (sec)	1.415032(2)	1.513(4)
Gravitational redshift parameter (ms)	0.39(2)	
Shapiro delay parameter $s = \sin i$	0.9995(4)	
Shapiro delay parameter r (s)	6.2(6)	
Orbital decay \dot{P}_b (10^{-12})	1:20(8)	
Mass ratio $R = M_A/M_B$		1.071(1)

tern or even the profile of B may vary due to these changes as well.

The possibility to observe such phenomena is exciting but requires a careful analysis to exclude any impact onto the timing of the two pulsars. Indeed, the necessity to check for these effects has delayed the publication of the currently final timing results of the double pulsar. The analysis is now almost complete and results will be published shortly. A preliminary update is given in Table 1 and discussed below.

V. PRESENT RESULTS

The study of the profile evolution of A [19] did not lead to the detection of any profile change over a period of 15 months. This present non-detection greatly simplifies the timing of A but does not exclude the possibility that changes may not happen in the future. While the effects of geodetic precession could be small due to a near alignment of pulsar A's spin and the orbital momentum vector, the results could also be explained by observing the system at a particular precession phase. While this case appears to be relatively unlikely, it must not be excluded as such a situation had indeed occurred for PSR B1913+16 [20]. Indeed, a modelling of the results suggests that this present non-detection of profile changes is consistent with a rather wide range of possible system geometries. One conclusion that can be drawn, however, is that the observations are inconsistent with the large profile changes that had been predicted by some models [21].

In contrast to the results for A, similar studies of

the profile evolution of B [22] reveal a clear evolution of B's emission on orbital and secular time-scales. The profile of B is changing with time, while also the light-curves of B (i.e. the visibility of B versus orbital phase) undergo clear changes. These phenomena may be caused by a changing magnetospheric interaction due to geometry variations resulting from geodetic precession. In any case, these changes require sophisticated timing analysis techniques and the preliminary results obtained with this techniques are listed in Table 1, while final results will be published shortly.

The present timing results already indicate that the proper motion of this system is surprisingly small. While a significant measurement of a proper motion vector via pulsar timing will be available shortly, the present limit suggests a systemic velocity of less than 30 km/s for a dispersion measure distance of 600 pc [23].[30] While such a small velocity may be indicative of a small kick imparted onto B during its supernova explosion [25], other studies find this limit still to be consistent with a kick of average magnitude [26]. In any case, such a small velocity is good news for tests of GR. Usually, the observed value of \dot{P}_b is positively biased by an effect known as secular acceleration arising from a relative motion and acceleration of the system (e.g. [27]). Computing the magnitude of this observational bias using the obtained limit on the proper motion, however, suggests that the contribution is much less than 1%, so that the orbital decay measurement will be available for another precise GR test.

Finally, scintillation measurements have recently suggested an orbital inclination angle that is extremely close to 90 deg (i.e. within (0.29 ± 0.14) deg) [24]. This measurement appears to be inconsistent



