

Publication Year	2018
Acceptance in OA@INAF	2024-03-14T16:04:26Z
Title	IR spectroscopy of ammoniated phyllosilicates at low pressure/high temperature conditions
Authors	DE ANGELIS, Simone; FERRARI, MARCO; DE SANCTIS, MARIA CRISTINA; Ammannito, Eleonora
Handle	http://hdl.handle.net/20.500.12386/34968



# IR spectroscopy of ammoniated phyllosilicates at low pressure/high temperature conditions

S. De Angelis (1), M. Ferrari (1), M.C. De Sanctis (1), E. Ammannito (2) (1) Institute for Space Astrophysics and Planetology, IAPS-INAF, Rome Italy, (<u>simone.deangelis@iaps.inaf.it</u>) (2) Italian Space Agency – ASI, Rome, Italy

## Introduction

Ammonium phyllosilicates are thought to be among the constituents of dwarf planet (1) Ceres surface, based on ground-based telescopic [1] and VIR-Dawn spectral data [2]. Following these findings, several works are currently trying to reproduce in the laboratory the Ceres surface composition, in terms of multi-component mineral mixtures [3,4]. Thus it is of interest to investigate the behavior and stability of ammonium compounds, when measured at pressuretemperature conditions that are different from standard laboratory values. Here we describe Visible-Infrared spectroscopic measurements of an ensemble of ammonium phyllosilicates: spectra have been acquired at various pressure-temperature conditions, by means of a P-T cell realized at INAF-IAPS laboratory.

#### 1. Samples and experimental setup

Ammoniated phyllosilicates were produced in the laboratory starting from natural samples, following a procedure described in a series of works [e.g. 6].

Infrared spectra of five ammoniated samples (montmorillonite, SCa-3, two nontronites, NAu-1 and NAu-2, illite-smectite, ISCz-1 and hectorite, SHCa-1) were then acquired in the spectral range 0.35-2.5  $\mu$ m, by using an ASD FieldSpec Pro 4 spectro-photometer equipped with a QTH lamp. The instrument is characterized by a spectral resolution of about 3-10 nm in the whole range; the spatial resolution of the setup was about 5 mm on the sample. All samples were analyzed in the form of powder, with grain size d<36  $\mu$ m.

In order to acquire reflectance spectra at varying conditions, the samples were placed inside a P-T environmental cell, developed at INAF-IAPS [5]. The measurements strategy was the following: (i) acquisition at room P-T, (ii) sequence of acquisitions

at room T during pumping; in this stage the pumping was first performed with only primary diaphragm pump (down to a limit of 3-4 mbar) and then also with turbo-molecular pump (down to vacuum pressure of  $10^{-4}$ - $10^{-5}$  mbar); (iii) acquisitions in vacuum at higher temperatures.

#### 2. IR Spectral measurements.

Here we report, as an example, on the analyses performed on one sample, nontronite (NAu1). The spectra of NH<sub>4</sub>-Nontronite are shown in fig.1. Data were acquired in three stages. The first spectrum (fig.1, A) is at room pressure and temperature. In panel B spectra were acquired at room T during primary pumping from ambient to  $10^{-4}$  mbar. Finally in panel C spectra acquired in vacuum at different temperatures (50-240°C) are shown.

### 3. Results and Conclusions

Spectra of nontronite are characterized by  $Fe^{2+}$ - $Fe^{3+}$ bands at 0.7-1 µm, and by OH<sup>-</sup>/H<sub>2</sub>O bands at 1.4 and 1.9 µm [7]. The feature at 2.3 µm is Fe-OH [7]. NH<sub>4</sub><sup>+</sup> absorption is visible at 2.12 µm in the room P-T spectrum. After pumping the adsorbed water is removed and NH<sub>4</sub><sup>+</sup> features become evident also at 1.55 and 2.01 µm: at room P-T these two features are shoulders in the 1.4 and 1.9 µm bands. After heating up to 240°C all NH<sub>4</sub><sup>+</sup> bands are visible and quite separate from hydration bands.

We can see that  $NH_4^+$  bands remain quite unaltered both by the process of pumping, at least down to a vacuum of about  $10^{-4}$  mbar, and by the process of heating up to 240°C. For higher temperatures the nontronite sample is subject to structural changes: all the water is removed, then ammonia and finally dehydroxylation occurs (fig. 1C).



Fig.1.  $NH_4$ -Nontronite (NAu-1). A): ambient pressure and temperature; B): ambient temperature, during pumping; C): higher temperatures, in vacuum. Vertical lines indicate  $NH_4^+$ absorptions.

# Acknowledgements

The experiment is funded by ASI.

#### References

- [1] King T.V.V. et al., Science, vol.255, 1551-1553, 1992
- [2] De Sanctis M.C., et al., Nature, vol.528, 241-244, 2015. [3] De Angelis S. et al., EPSC abstract n.830, vol.11, 2017
- [4] Ehlmann B.L. et al., MAPS, 10.1111/maps.13103, 2018
  [5] De Angelis S. et al., 49<sup>th</sup> LPSC, abstract n.1428, 2018
  [6] Ferrari M. et al., 49<sup>th</sup> LPSC, abstract n.2413, 2018
  [7] Clark R.N. et al., JGR, vol.95, B8, 12,653-12,680, 1990