



<b>Publication Year</b>	2018
<b>Acceptance in OA</b>	2020-11-24T14:44:10Z
<b>Title</b>	Extreme quasars at high redshift
<b>Authors</b>	Martínez-Aldama, M. L., del Olmo, A., MARZIANI, Paola, Sulentic, J. W., Negrete, C. A., Dultzin, D., D'Onofrio, M., Perea, J.
<b>Publisher's version (DOI)</b>	10.1051/0004-6361/201833541
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/28510">http://hdl.handle.net/20.500.12386/28510</a>
<b>Journal</b>	ASTRONOMY & ASTROPHYSICS
<b>Volume</b>	618

# Extreme quasars at high redshift

M. L. Martínez-Aldama<sup>1</sup>, A. del Olmo<sup>1</sup>, P. Marziani<sup>2</sup>, J. W. Sulentic<sup>1</sup>, C.A. Negrete<sup>3</sup>, D. Dultzin<sup>4</sup>, M. D’Onofrio<sup>5</sup>, and J. Perea<sup>1</sup>

<sup>1</sup> Instituto de Astrofísica de Andalucía, IAA-CSIC, Glorieta de la Astronomía s/n, E-18008 Granada, Spain

<sup>2</sup> Istituto Nazionale d’Astrofisica (INAF), Osservatorio Astronomico di Padova, IT 35122, Padova, Italy

<sup>3</sup> CONACYT Research Fellow, Instituto de Astronomía, UNAM, México D.F. 04510, México

<sup>4</sup> Instituto de Astronomía, UNAM, México D.F. 04510, México

<sup>5</sup> Dipartimento di Fisica & Astronomia “Galileo Galilei”, Università di Padova, Padova, Italy

Preprint online version: October 1, 2018

## ABSTRACT

**Context.** Quasars radiating at extreme Eddington ratios (hereafter xA quasars) are likely a prime mover of galactic evolution and have been hailed as potential distance indicators. Their properties are still scarcely known.

**Aims.** We test the effectiveness of the selection criteria defined on the “4D Eigenvector 1” (4DE1) for identifying xA sources. We provide a quantitative description of their rest-frame UV spectra (1300 – 2200 Å) in the redshift range  $2 \lesssim z \lesssim 2.9$ , with a focus on major emission features.

**Methods.** Nineteen extreme quasar candidates were identified using 4DE1 selection criteria applied to SDSS spectra:  $\text{AlIII}\lambda 1860/\text{SiIII}\lambda 1892 \gtrsim 0.5$  and  $\text{CIII}\lambda 1909/\text{SiIII}\lambda 1892 \lesssim 1$ . The emission line spectra was studied using multicomponent fits of deep spectroscopic observations ( $S/N \gtrsim 40 - 50$ ; spectral resolution  $\approx 250 \text{ km s}^{-1}$ ) obtained with the OSIRIS at Gran Telescopio Canarias (GTC).

**Results.** GTC spectra confirm that almost all of these quasars are xA sources with very similar properties. We provide spectrophotometric and line profile measurements for the  $\text{SiIV}\lambda 1397 + \text{OIV}\lambda 1402$ ,  $\text{CIV}\lambda 1549 + \text{HeII}\lambda 1640$ , and the 1900Å blend. This last feature is found to be predominantly composed of  $\text{AlIII}\lambda 1860$ ,  $\text{SiIII}\lambda 1892$  and  $\text{FeII}$  emission features, with weak  $\text{CIII}\lambda 1909$ . The spectra can be characterized as very low ionization (ionization parameter,  $\log U \sim -3$ ), a condition that explains the significant  $\text{FeII}$  emission observed in the spectra. xA quasars show extreme properties in terms of  $\text{CIV}\lambda 1549$  equivalent width and blueshift amplitudes.  $\text{CIV}\lambda 1549$  shows low equivalent width, with a median value of 15Å ( $\lesssim 30 \text{ Å}$  for the most sources), and high or extreme blueshift amplitudes ( $-5000 \lesssim c(\frac{1}{2}) \lesssim -1000 \text{ km s}^{-1}$ ). Weak-lined quasars appear as extreme xA quasars and not as an independent class. The  $\text{CIV}\lambda 1549$  high amplitude blueshifts coexists in all cases save one with symmetric and narrower  $\text{AlIII}\lambda 1860$  and  $\text{SiIII}\lambda 1892$  profiles. Estimates of the Eddington ratio using the  $\text{AlIII}\lambda 1860$  FWHM as a virial broadening estimator are consistent with the ones of a previous xA sample.

**Conclusions.** xA quasars show distinguishing properties that make them easily identifiable in large surveys and potential “standard candles” for cosmological applications. It is now feasible to assemble large samples of xA quasars from the latest data releases of the SDSS. We provide evidence that  $\text{AlIII}\lambda 1860$  could be associated with a low-ionization virialized sub-system, supporting previous suggestions that  $\text{AlIII}$  is a reliable virial broadening estimator.

**Key words.** quasars: general – quasars: emission lines – quasars: supermassive black holes

Manuscript accepted for publication in A&A, July 2018

## 1. Introduction

Quasars are found over an enormous range of distances ( $z \sim 0 - 7.5$ ) in the Universe. For this reason they have occasionally been cited as the ultimate possible standard candles for use in cosmology (See for example the Chapter by in D’Onofrio & Burigana (2009) and the recent reviews by Sulentic et al. (2014b) and Czerny et al. (2018)). The problem with such a use has been the lack of a clear definition of “quasar” and a contextualization of their diversity. Since 2000 a clearer idea of their nature and diversity has emerged using the 4D Eigenvector 1 (4DE1) formalism. We are now able to identify a quasar main sequence (See Marziani et al. 2018, for a recent review), and recognize an extreme accretor (xA) quasar population at the end of this sequence (e.g., Marziani & Sulentic 2014, hereafter MS14). This xA population radiating at  $L/L_{\text{Edd}} \sim 1$  offer the best opportunity to use quasars for cosmology (MS14, Wang et al. 2014). This

paper describes a search for extreme quasars at  $z \sim 2.3$  built upon an extension of low-redshift 4DE1 studies.

Quasar spectra show diverse properties in measures of line intensity ratios and line profiles. These measures offer multifold diagnostics of emitting region structure and physical conditions (Marziani et al. 2018). Organizing the diversity of quasar properties has been an ongoing effort for many years. Perhaps the first successful attempt was carried out by Boroson & Green (1992). They proposed an Eigenvector 1 scheme based on a principal component analysis of the Palomar-Green quasar sample (See Gaskell et al. 1999; Sulentic et al. 2000a, for reviews up to the late 1990s). Trends between measures of  $[\text{O III}]\lambda\lambda 4959, 5007$ , optical  $\text{FeII}$  emission and full width at half maximum (FWHM)  $H\beta$  were found, and appreciation of their importance has grown with time (Sulentic & Marziani 2015, and references therein). Sulentic et al. (2000b) expanded upon this work and defined a new scheme called 4D Eigenvector 1 (4DE1) with the addition of

two new parameters. Principal 4DE1 measures for low- $z$  quasars involve: 1) FWHM of broad line H $\beta$  (excluding any narrow emission component)<sup>1</sup>; 2) the strength of the optical FeII blend at 4570 Å normalized by the intensity of H $\beta$ :  $R_{\text{FeII}}=I(\text{FeII})/I(\text{H}\beta)$ ; 3) the velocity shift at half maximum ( $c(\frac{1}{2})$ ) of the high-ionization line (HIL) C $\text{IV}\lambda 1549$  profile relative to a rest-frame (usually defined by measures of the [O III] $\lambda 5007$  and/or narrow H $\beta$  centroid) and 4) the soft X-ray photon index ( $\Gamma_{\text{soft}}$ ).

The 4DE1 optical plane (OP), defined by the measures of the FWHM of H $\beta$  and  $R_{\text{FeII}}$ , shows a reasonably well-defined sequence (a quasar “main sequence”, MS, Sulentic et al. 2000a; Marziani et al. 2001). The nomenclature is motivated by an analogy with the role of the stellar main sequence in the H-R diagram which connects observational measures to physical properties (Sulentic et al. 2008). The stellar MS is driven by stellar mass, while the quasar sequence is thought to be driven by Eddington ratio ( $\propto M$ , Marziani et al. 2001, 2003b). Sulentic et al. (2000a) noted a change in all 4DE1 measures near FWHM H $\beta$  = 4000 km s<sup>-1</sup> in low- $z$  quasar samples ( $L \lesssim 10^{47}$  erg s<sup>-1</sup>). This change motivated an empirical designation of two quasar populations: Population A with FWHM H $\beta$   $\leq$  4000 km s<sup>-1</sup>,  $R_{\text{FeII}} > 0.5$ , frequent C $\text{IV}\lambda 1549$  profile blueshifts and sources with a soft X-ray excess, and Population B with FWHM H $\beta$   $>$  4000 km s<sup>-1</sup>,  $R_{\text{FeII}} < 0.5$ , absence of C $\text{IV}\lambda 1549$  blueshift and little or no soft X-ray excess (Bensch et al. 2015). Radio-loud (RL) quasars are strongly concentrated in the Population B domain along with 30% of radio quiet (RQ) sources (e.g., Sulentic et al. 2003; Zamfir et al. 2008, and references therein).

Low-ionization emission lines (LILs - H $\beta$  best studied feature) in quasars frequently show asymmetric profiles. The H $\beta$  broad component (H $\beta_{\text{BC}}$ ) in Population B sources is usually well described by a double Gaussian profile with one of the components centered on the rest-frame and the second one redshifted by  $\approx$  1000 – 3000 km s<sup>-1</sup> (Zamfir et al. 2010), the very broad component (VBC). Population A sources rarely show the redshifted component. High-ionization line (C $\text{IV}\lambda 1549$  best studied prototype) profiles usually show blueward shifts/asymmetries in Population A sources. Pop. B objects also show weak or moderate strength LILs like FeII and the CaII IR triplet (Sulentic et al. 2006b; Martínez-Aldama et al. 2015). Usually Population B quasars do not show any strong soft X-ray excess (Sulentic et al. 2000a; Bensch et al. 2015, and references therein). Largely radio-quiet (RQ) population A sources usually show symmetric profiles well-modeled with a Lorentz function (Sulentic et al. 2002; Zamfir et al. 2010; Cracco et al. 2016). Population A sources with the narrowest H $\beta$  profiles ( $<$ 2000 km/s) are often called narrow line Seyfert 1 sources (NLSy1), but in no sense represent a distinct class of quasars (Zamfir et al. 2008; Sulentic et al. 2015).

All of the above description involves quasars with  $z < 1.0$  where moderate to high S/N ground-based spectra exist for significant numbers of sources H $\beta$  (ground based) and C $\text{IV}\lambda 1549$  (HST FOS archival data). Within each quasar population systematic trends are revealed in composite spectra of H $\beta$  (Sulentic et al. 2002; Zamfir et al. 2010) and C $\text{IV}$  (Bachev et al. 2004). Sulentic et al. (2002) defined Pop. A subclasses A1, A2, A3 and A4 in order of increasing intervals of  $0.5R_{\text{FeII}}$  have been defined. Extreme Population A sources in bins A3 and A4 (xA) involve quasars with very strong  $R_{\text{FeII}} \geq 1.0$ . This criterion is applicable only to sources at low redshift. This study involves searching for higher redshift analogues of such extreme quasars. At high

redshift ( $z > 2$ ) the spectral range used to define sources in the 4DE1 (FWHM H $\beta$ ,  $R_{\text{FeII}}$ ) context are lost unless NIR spectra are available for H $\beta$  (Marziani et al. 2009). Many C $\text{IV}$  spectra in the intermediate/high- $z$  range are available in the SDSS/BOSS archives although low S/N often precludes detailed analysis.

The 4DE1 provides a consistent picture of quasar observational properties in low- $z$  samples: beginning with the low- $R_{\text{FeII}}$  and broad FWHM H $\beta$ , involving high black hole mass “disk-dominated” quasars. As we move along the sequence we encounter sources whose spectra show narrower LIL profiles, lower ionization spectra, and blueshifted C $\text{IV}\lambda 1549$  profiles providing evidence of strong HIL emitting outflows: “wind-dominated” quasars (Richards et al. 2011). Eddington ratio (convolved with the effects of orientation) appears to be the physical parameter driving systematic changes of observational properties along the quasar MS (Sulentic et al. 2000a; Marziani et al. 2001; Boroson 2002). The main sequence observational trends can be interpreted as driven by small  $M_{\text{BH}}$ , higher  $L/L_{\text{Edd}}$  (young? high accretors) towards the high  $R_{\text{FeII}}$  end of the sequence. The xA sources cluster around  $L/L_{\text{Edd}} \approx 1^2$ . This would imply that the low- $z$  xA sources are the “youngest” (less massive than Pop. B) quasar population radiating at the highest Eddington ratios (Fraix-Burnet et al. 2017).

There is a growing consensus that sources at the high  $R_{\text{FeII}}$  end of the MS are accreting at the highest rates, and are expected to be close to the radiative limit per unit black hole mass (Sun & Shen 2015; Du et al. 2016a,b; Sniegowska et al. 2017). If this is the case, xA quasars acquire a special meaning. While Population A HILs are dominated by blue shifted emission associated with outflows, the presence of almost symmetric and unshifted LILs (Balmer lines, but also Paschen lines, La Franca et al. 2014) indicate the coexistence of a LIL emitting region that is virialized. Since  $L/L_{\text{Edd}}$  tends toward a constant limiting value (Mineshige et al. 2000), xA quasars can be considered as “Eddington standard candles.” If so that, if  $M_{\text{BH}}$  can be retrieved under the virial assumption, an estimate of the luminosity becomes possible since  $L/L_{\text{Edd}} \propto L/M_{\text{BH}}$  (Wang et al. 2013; La Franca et al. 2014, MS14). This approach is conceptually analogous to the use of the link between the velocity dispersion in virialized systems (i.e. the rotational velocity of HI disks in spiral galaxies Tully & Fisher 1977). Initial computations for samples of 100-200 low- $z$  quasars ( $\lesssim 1$ ) confirm the conceptual validity of the “virial luminosity” estimates (Negrete et al. 2017; Negrete et al. 2018, submitted), although scatter in the distance modulus is still too large to draw meaningful inferences for cosmology.

4DE1 trends can also be helpful for interpreting high- $L$ , high- $z$  quasars, although there are two caveats. At high redshift,  $z \sim 2$ , the majority of sources show large FWHM due to a bias in luminosity (Sulentic et al. 2014a, 2017): quasars with luminosities comparable to the low- $z$  low- $L$  sources are still too faint to be efficiently discovered. More fundamentally, there is a minimum possible FWHM H $\beta$  at fixed luminosity, if the line emitting region is virialized and its size follows a scaling law with luminosity. In practice this means that at  $\log L \gtrsim 47$ , all lines have to be broader than FWHM 2000 km s<sup>-1</sup>. By the same token the FWHM limit for Population A becomes luminosity dependent. The limit established at 4000 km s<sup>-1</sup> is valid only for low- $z$ , relatively low- $L$  quasars. Sources with larger FWHM and emission line properties similar to the ones of the low- $z$  xA quasars have

<sup>1</sup> In the following we understand for broad profile of a line the total broad profile excluding the narrow component

<sup>2</sup> The precise  $L/L_{\text{Edd}}$  values depend on the black hole mass  $M_{\text{BH}}$  scaling law and on the bolometric corrections. Following the 4DE1 based assumptions described above, the highest values along the MS are  $L/L_{\text{Edd}} \approx 1 - 2$

**Table 1.** Source identification and basic properties of the GTC-xA quasars.

SDSS identification (1)	$z$ (2)	$\Delta z$ (3)	Line (4)	$m_v$ (5)	$M_B$ (6)	$g - r$ (7)	Comments (8)
SDSSJ000807.27-103942.7	2.4660	0.0010	iii	19.15	-26.2	0.08	
SDSSJ004241.95+002213.9	2.0560	0.0053	iii	19.05	-25.6	0.24	
SDSSJ021606.41+011509.5	2.2236	0.0008	i,ii	19.36	-25.2	0.49	BAL
SDSSJ024154.42-004757.5	2.3919	0.0015	iii	19.24	-26.0	0.24	
SDSSJ084036.16+235524.7	2.1879	0.0012	iii	19.46	-25.3	0.40	
SDSSJ101822.96+203558.6	2.2502	0.0035	iii	19.27	-25.5	0.33	
SDSSJ103527.40+445435.6	2.2639	0.0060	iii	19.34	-25.6	0.20	
SDSSJ105806.16+600826.9	2.9406	0.0001	i,ii	19.29	-26.7	0.16	
SDSSJ110022.53+484012.6	2.0884	0.0028	iii	18.90	-25.9	0.13	
SDSSJ125659.79-033813.8	2.9801	0.0004	ii	19.27	-26.7	0.26	BAL
SDSSJ131132.92+052751.2	2.1234	0.0009	iii	19.06	-25.4	0.72	BAL
SDSSJ143525.31+400112.2	2.2615	0.0006	iii	18.30	-26.6	0.17	
SDSSJ144412.37+582636.9	2.3455	0.0018	iii	19.21	-25.8	0.37	Mini-BAL
SDSSJ151258.36+352533.2	2.2382	0.0012	iii	19.21	-25.8	0.04	RL, $\log P_\nu \approx 33.8$ , $\log R_K \approx 2.65$
SDSSJ214009.01-064403.9	2.0808	0.0038	iii	19.26	-25.3	0.56	BAL
SDSSJ220119.62-083911.6	2.1840	0.0015	iii	18.68	-26.1	0.35	BAL
SDSSJ222753.07-092951.7	2.1639	0.0010	iii	19.11	-25.6	0.10	
SDSSJ233132.83+010620.9	2.6271	0.0038	iii	19.19	-26.1	0.33	RL, $\log P_\nu \approx 34.1$ , $\log R_K \approx 2.93$
SDSSJ234657.25+145736.0	2.1682	0.0007	iii	19.11	-25.7	0.29	RI, $\log P_\nu \approx 32.7$ , $\log R_K \approx 1.67$

NOTES. Columns are as follows: (1) SDSS coordinate name. (2) Redshift. (3) Redshift uncertainty. (4) Lines used for redshift determination: i: C<sub>IV</sub>λ1335, ii: OI+SiII λ1305, iii: AlIIIλ1860+SiIIIλ1892. (5) Apparent Johnson  $V$  magnitude as reported by Véron-Cetty & Véron (2010). (6) Absolute  $B$  magnitude according to Véron-Cetty & Véron (2010). (7)  $g - r$  color in magnitudes. (8) Quasar classification: BAL QSO: Broad Absorption Line Quasar; Mini-BAL: Mini Broad Absorption Line Quasar, radio-loud (RL) and radio-intermediate (RI).  $P_\nu$  is power per unit frequency at 1.4 GHz in units of  $\text{erg s}^{-1} \text{Hz}^{-1}$ .

been found at high- $z$  and high- $L$  (Negrete et al. 2012, MS14). Another important issue at high- $L$  concerns the HIL blueshifts. While at low- $L$  large blueshifts ( $v_r \lesssim -1000 \text{ km s}^{-1}$ ) are confined to Pop. A (Sulentic et al. 2007; Richards et al. 2011), at high- $L$  they are ubiquitous (Coatman et al. 2016; Bischetti et al. 2017; Bisogni et al. 2017), even if Pop. A sources still show the largest blueshift amplitudes among all quasars (Sulentic et al. 2017, hereafter S17). At any rate, several recent studies confirm that  $H\beta$ , observed in NIR spectra in quasars at  $z \gtrsim 1$  shows fairly symmetric and unshifted profiles suggesting that the broadening is mainly due to virial motions of the line emitting gas (Marziani et al. 2009; Bisogni et al. 2017; Shen et al. 2016; Vietri et al. 2018, S17). We will show in this paper that this is probably true also for high- $L$  xA sources.

Goals of this paper include testing the effectiveness of 4DE1 selection criteria for identifying high  $L/L_{\text{Edd}}$  xA sources at  $z \sim 2.4$ . This will enable us to analyze spectral properties of the identified xA quasars in the rest-frame UV region. A sample of candidate xA sources (hereafter GTC-xA) was observed with the Gran Telescopio de Canarias (GTC) using the OSIRIS spectrograph. We apply the 4DE1 selection criterion defined by MS14 using UV diagnostic ratios:  $\text{AlIII}\lambda 1860/\text{SiIII}\lambda 1892 \geq 0.5$  and  $\text{CIII}\lambda 1909/\text{SiIII}\lambda 1892 \leq 1.0$ . The selected sources are intended to represent an xA population for which  $R_{\text{FeII}}$  is expected to be larger than 1, with no limitation on line FWHM. The sample selection is described in Sect. §2. The observations and the data reduction are presented in Sect. §3. We perform a multi-component fitting and build Monte Carlo (MC) simulations to estimate measurement uncertainties (Sect. §4 and Appendix B). Spectra and line measures are presented for the 1900Å blend, the blend C<sub>IV</sub>λ1549 + HeIIλ1640, and the Si<sub>IV</sub>λ1397 region in Sect. §5. A comparison with control samples at low- $z$  and/or  $L$  is described in the Section §6.1. A composite spectrum for the

GTC-xA sample (Sect. §5.1) allows us to carry out a comparison with low- $z$ /low- $L$  samples (Sect. §6.2). We discuss the low-ionization spectra and identify FeII and FeI features that are prominent in our spectra (Sect. §6.3 and Sect. §6.4). After estimating the main accretion parameters (§6.5), we briefly analyze the relation of xA sources to Weak Line Quasars (WLQs) – a related class of quasars with extreme properties (Sect. §6.6).

xA sources are especially important because they are the quasars radiating at the highest luminosity per unit mass. The extreme radiative properties of xAs make them prime candidates for maximum feedback effects on host galaxies. We briefly analyze the possibility of significant feedback effects in Sect. §6.7. We conclude the paper with some consideration on the possible cosmological exploitation of xA quasars at high- $z$  (Sect. §7).

## 2. Sample description

MS14 extracted 3000 quasar spectra from the SDSS DR6 archive which provided coverage of the 1900Å blend for sources in the redshift range  $2.0 < z < 2.9$ , with  $g < 19.5$ . Intensity measures of AlIIIλ1860, SiIIIλ1892 and CIIIλ1909 were carried out with an automatic SPLAT procedure within the IRAF reduction package. The majority of SDSS spectra are quite noisy making them suitable for identifying samples of candidate sources, but not providing accurate spectroscopic measures (Sulentic & Marziani 2015). A preliminary selection of extreme Eddington candidates was made and sources were then vetted according to the xA selection criteria ( $\text{AlIII}\lambda 1860/\text{SiIII}\lambda 1892 \geq 0.5$  and  $\text{CIII}\lambda 1909/\text{SiIII}\lambda 1892 \leq 1.0$ ). MS14 considered the brightest candidates with moderate S/N ( $\geq 15$ ) spectra, leaving 58 candidate xA quasars whose relative AlIIIλ1860, SiIIIλ1892 and CIIIλ1909 intensities satisfied the selection criteria, but whose S/N was too poor to make an accurate measurement of individ-

ual lines in the 1900Å blend. Hence the need for new GTC spectroscopic observations. This paper presents an analysis of 19 of the sources that constitute our GTC-xA sample.

Table 1 gives source identifications and basic properties including: redshift and uncertainty (Col. 2 & Col. 3), emission line used for the redshift estimate (Col. 4), apparent  $V$  magnitude and absolute  $B$  magnitude  $M_B$  (Col. 5 & 6) as given in Véron-Cetty & Véron (2010), and  $(g - r)$  color index from the SDSS photometry (Col. 7). Column 8 identifies other observed features like its classification as BAL (Broad Absorption Line) or mini-BAL quasar, and the radio properties, if they are detected. Low- $z$  studies (Zamfir et al. 2008) define a radio-loud (RL) quasar with a radio/optical flux ratio  $\log R_K \geq 1.8$  (Kellermann et al. 1989), or better a radio power measure  $\log P_\nu > 31.6$  [erg s<sup>-1</sup> Hz<sup>1</sup>], independent of uncertainties in optical flux measures.  $R_K$  was obtained normalizing the  $k$ -corrected radio flux at 1.4GHz by the  $k$ -corrected  $B$  magnitude. Two radio-detected sources exceed the  $R_K$  limit with the third one close to the limit. Only for SDSSJ233132.83+010620.9 was possible to estimate the radio spectral index from one observation at 8.4 GHz (Cegłowski et al. 2015):  $\alpha_r \approx 0.44$ , that places it in the compact steep-spectrum radio domain. SDSSJ234657.25+145736.0 has only a low-resolution NRAO VLA Sky Survey (NVSS) map available. In this case,  $P_\nu$  and  $R_K$  have been computed assuming  $\alpha_r = 0$ . All of them exceed the  $\log P_\nu$  limit, then we could have three RL quasars in GTC-xA sample. Since low- $z$  RL are almost never found in the xA domain in 4DE1, it is possible either that the three radio detected sources are not xA extreme accretors or that high- $z$  Pop. A quasars are more frequently RL. At this point the three RL sources in the sample must be treated with caution.

### 2.1. The FOS-A, S14-A and FOS-xA “control” samples

The quasars in the GTC-xA sample are thought to be highly accreting, with an average bolometric luminosity of  $\log L \sim 47$  [erg s<sup>-1</sup>]. Absolute magnitude  $M_B \approx -26$ , before extinction correction, corresponds to a comoving space density of  $\sim 10^{-6}$  mag<sup>-1</sup> Mpc<sup>-3</sup> just beyond the turnover at the high luminosity end of the 2dF luminosity function (Boyle et al. 2000).

In order to compare the behavior of the GTC-xA sample, we consider the FOS sample from Sulentic et al. (2007) as a control sample at low- $L$  and low- $z$ . For the sake of the present paper, we restrict the control FOS sample to 28 Pop. A RQ sources covering the Civ and 1900Å blend spectral range and with previous measures for the H $\beta$  profile and  $R_{FeII}$  (Marziani et al. 2003a). 23 objects are classified as Pop. A1-A2 sources (henceforth FOS-A sample) and 5 as xA sources (hereafter FOS-xA sample) including I Zw 1. The FOS sample has a typical bolometric luminosity  $\log L \sim 45.2$  [erg s<sup>-1</sup>] and a redshift  $z \lesssim 0.5$ .

The sample presented in Sulentic et al. (2014a, hereafter S14) covers a similar range in redshift ( $z \sim 2.3$ , corresponding to a lookback time of  $\approx 10$  Gyr), and are in turn a factor  $\sim 10$  less luminous ( $\log L \sim 46$ ) than the GTC-xA sample. S14 is representative of a general population of faint, moderately accreting quasars that are also found at intermediate redshift (Fraix-Burnet et al. 2017). S14 includes both Pop. A and B sources, but no xA sources. Restriction to the 11 Pop. A quasars (hereafter S14-A) offers a high- $z$  counterpart to the FOS-A sample, and therefore suitable for a comparison between xAs and a sample of Pop. A quasars of moderate  $L$  at  $z \sim 2 - 2.5$ , that represents a population expected to be relatively common ( $\Phi \sim 10^{-6}$  mag<sup>-1</sup> Mpc<sup>-3</sup>).

## 3. Observations and Data Reduction

Long slit observations were carried out in service mode using the OSIRIS spectrograph at the 10.4m GTC telescope of the Roque de los Muchachos Observatory. Grisms R1000B and R1000R with 2x2 CCD binning were used for the observations, depending on the redshift of the source. The majority of the observations employed R1000B that covered the wavelength range from 3650–7400 Å with a reciprocal dispersion of 2.1 Å per pixel ( $R \approx 1000$ ). In our highest  $z$  sources, SDSS J105806.16+600826.9, SDSS J125659.79-033813.8, and SDSS J233132.83+010620.9 with  $z \approx 2.94$ , 2.98 and 2.63 respectively, we used the R1000R grism with reciprocal dispersion of 2.6 Å/pixel and spectral coverage 5100 - 10000Å. These two spectral ranges correspond the rest-frame region covering the UV spectral features of interest such as the Siiv $\lambda$ 1397, Civ $\lambda$ 1549 and the 1900Å blend. The spectra were obtained with an 0.6 arcsec slit width oriented at the parallactic angle in order to minimize atmospheric differential refraction. Table 2 contains a summary of the observations including: SDSS identification, observation date, grism employed, total exposure time for the 3 individual exposures on each source, seeing estimated from the FWHM of field stars in the acquisition image, and the estimated S/N in the 1450Å continuum region on the blue side of Civ $\lambda$ 1549.

**Table 2.** Log of Observations

SDSS Identification (1)	Obs. Date (2)	Grism (3)	Exp. Time (4)	Seeing (5)	S/N (6)
SDSSJ000807.27-103942.7	22/07/2015	R1000B	1800	1.13	29
SDSSJ004241.95+002213.9	22/07/2015	R1000B	1440	1.18	39
SDSSJ021606.41+011509.5	14/08/2015	R1000B	1440	0.80	70
SDSSJ024154.42-004757.5	14/08/2015	R1000B	2700	1.17	40
SDSSJ084036.16+235524.7	18/04/2015	R1000B	1800	1.13	31
SDSSJ101822.96+203558.6	12/06/2015	R1000B	1440	1.40	33
SDSSJ103527.40+445435.6	23/03/2015	R1000B	1800	1.33	32
SDSSJ105806.16+600826.9	12/06/2015	R1000R	2700	1.22	22
SDSSJ110022.53+484012.6	18/04/2015	R1000B	1890	0.74	38
SDSSJ125659.79-033813.8	23/05/2015	R1000R	1440	1.29	12
SDSSJ131132.92+052751.2	23/05/2015	R1000B	1800	1.42	31
SDSSJ143525.31+400112.2	11/04/2015	R1000B	2340	0.98	60
SDSSJ144412.37+582636.9	25/06/2015	R1000B	1980	0.89	23
SDSSJ151258.36+352533.2	11/04/2015	R1000B	2340	0.86	37
SDSSJ214009.01-064403.9	05/08/2015	R1000B	1800	1.13	60
SDSSJ220119.62-083911.6	17/06/2015	R1000B	2340	0.83	75
SDSSJ222753.07-092951.7	26/06/2015	R1000B	1800	0.91	62
SDSSJ233132.83+010620.9	05/08/2015	R1000R	1800	1.29	22
SDSSJ234657.25+145736.0	16/07/2015	R1000B	1440	0.96	54

NOTES. Columns are as follows: (1) SDSS identification. (2) Observation Date. (3) Grism. (4) Exposure time in seconds. (5) Seeing in arcseconds. (6) S/N measured in the continuum at 1450 Å.

Data reduction was carried out in a standard way using the IRAF package. Bias subtraction and flat-fielding correction were performed nightly. Wavelength calibration was obtained using Hg+Ar and Ne lamps observed with the same configuration and slit width used for source observations. Wavelength calibration rms was less than 0.1 Å. We checked the wavelength calibration for individual exposures with sky lines before source extraction, background subtraction, and final combination. Scatter of the sky line wavelength peaks was  $\lesssim 20$  km s<sup>-1</sup>. This value provides a realistic estimate of the wavelength scale uncertainty including zero point error. Spectral resolution estimated from FWHM of the skylines is  $\sim 230$  km s<sup>-1</sup> and 250 km s<sup>-1</sup> for grisms R1000B and R1000R, respectively.

Instrumental response and flux calibration were obtained nightly with observations of the spectrophotometric standard stars Ross 640, GD24-9, Feige 110, Hiltner 600, and G158-100. In order to improve flux calibration, we also included two additional flux standard stars, LDS749B and HZ21, as target objects. They were observed with both grisms and two slits: 0.6 arcsec (as used for quasar observations) and 5 arcsecs. A comparison between the different slits gives a change in the absolute flux calibration  $\sim 10\%$ . Spectra were corrected for light losses due to the narrow slit width employed and also to differential light loss as a function of wavelength. Taking into account the ratio between the slit width and the seeing during an observation allowed correction for the scale factor and for the wavelength dependence of the seeing following the method described by Bellazzini (2007). Telluric absorptions, that affect our spectra mainly beyond 7600 Å, were also corrected using the standard stars to obtain a normalized template of the absorption features. The template was shifted if needed, and scaled for each individual source in an iterative and interactive procedure, until the residuals in the telluric correction were negligible. Spectra were finally deredshifted as explained in Section §4.1.

### 3.1. Extinction estimation

It is visually apparent from examining our spectra that some of them (e.g., SDSSJ021606.41+011509.5, Fig. A.3) show a flatter continuum than cannot be modeled with a single power-law over the observed spectral range. This effect, as well as the presence of BALs, has been interpreted in the literature as indicating the presence of dust or internal reddening. In order to assess the importance of internal reddening on the observed fluxes and derived magnitudes in our xA sample, we have estimated the reddening in each source by fitting its UV continuum with quasar templates excluding spectral regions with broad emission lines (e.g., Ly $\alpha$ , SiIV $\lambda$ 1397, CIV $\lambda$ 1549 and the 1900Å blend). We used four QSO templates: 1) a median composite spectrum representative of the xA quasar population and built with extreme accretor sources identified in the SDSS DataBase by MS14 excluding BALs; 2) a template involving the composite FIRST Bright Quasar Survey spectrum (FBQS; Brotherton et al. (2001)); 3) the composite spectrum provided by Harris et al. (2016) with BOSS spectra of quasars in the redshift range  $2.1 < z < 3.5$ , and 4) the SDSS composite quasar spectra (Vanden Berk et al. 2001). We reddened the templates using an SMC extinction law (Gordon & Clayton 1998) which appears to be the most appropriate reddening law for modeling quasar spectra (York et al. 2006; Gallerani et al. 2010). We assumed a  $R_V$  coefficient of 3.07 for the extinction law.

**Table 3.** Extinction measures for the xA sources

SDSS identification	$A_V$ (mag.)
SDSSJ021606.41+011509.5	0.270
SDSSJ103527.40+445435.6	0.178
SDSSJ131132.92+052751.2	0.325
SDSSJ144412.37+582636.9	0.106
SDSSJ214009.01-064403.9	0.250
SDSSJ233132.83+010620.9	0.150

In general, best fits were obtained with the xA composite although there were no appreciable differences and all the fittings


yielded similar results. For the majority of sources (12) no additional extinction was needed and the continuum was well represented by the templates. In 6 cases the reddening has a significant effect on the spectrum, amounting to  $A_V=0.1$  to  $A_V=0.32$ . Table 3 reports reddening estimates parametrized by the  $A_V$  value. In the case of SDSSJ220119.62-083911.6, classified as BAL, the spectrum shows a broad and deep absorption in the blue wings of CIV $\lambda$ 1549, SiIV $\lambda$ 1397 and Ly $\alpha$  with an evident flattening of the continuum at wavelengths shorter than 1600 Å. However, from 1700 Å the spectrum shows a similar slope to the templates and does not show evidence for reddening. Hence no internal reddening was applied to this source and the continuum at 1350 Å needed for estimation of the luminosity was obtained by extrapolating the power law of the fit applied to the red spectral region.

## 4. Data Analysis

### 4.1. Redshift determination

Accurate redshift estimates are very important because redshift defines the quasar rest-frame from which emission line shifts can be measured. Shifts are particularly important for HILs like CIV $\lambda$ 1549 (e.g., Gaskell 1982; Espey et al. 1989; Carswell et al. 1991; Marziani et al. 1996). Redshift estimates are chiefly obtained from narrow emission lines (or the narrow core of H $\beta$ ) for low- $z$  sources (e.g., Eracleous & Halpern 2003; Hu et al. 2008). In the UV region covered by our spectra of  $z \approx 2.3$ , narrow LILs are not present. We must resort to broad LILs and estimate the redshifts using three features: AlIII $\lambda$ 1860, CIV $\lambda$ 1335 and OI $\lambda$ 1304+SiII $\lambda$ 1306. The strongest and hence most often detected LIL involves AlIII $\lambda$ 1860, which emerged in lower  $z$  studies as a kind of UV surrogate H $\beta$  (Bachev et al. 2004). While often detected, it is always part of the 1900Å blend albeit on the blue end of it. Although AlIII $\lambda$ 1860 and SiII $\lambda$ 1892 are blended lines, they are stronger than CIV $\lambda$ 1335 and OI $\lambda$ 1304+SiII $\lambda$ 1306 and in many of the spectra their peaks are clearly seen. We performed a multicomponent fit for each source, considering all the lines in the region of the 1900Å blend (See Section §4.2). The peaks of AlIII $\lambda$ 1860 and SiII $\lambda$ 1892 were unconstrained in intensity in the fitting and the adopted model of the blend was the one with minimum  $\chi^2$ . CIV $\lambda$ 1335 is the only isolated LIL in the observed UV range and is detected in only 5 sources (e. g. SDSSJ021606.41+011509.5, See Fig. A.3). The OI $\lambda$ 1304+SiII $\lambda$ 1306 blend is also well seen in a few sources where CIV $\lambda$ 1335 is detected. In the rest of the sample these lines are too weak to be useful and are often affected by absorption features.

We constructed synthetic Gaussian profiles for CIV $\lambda$ 1335 and OI $\lambda$ 1304+SiII $\lambda$ 1306 lines using the IRAF task MK1DSPEC assuming FWHM  $\sim 4000$  km s $^{-1}$ . We computed a redshift from the peak of the synthetic features. When the AlIII $\lambda$ 1860 redshift was compared with the one of CIV $\lambda$ 1335 and OI $\lambda$ 1304+SiII $\lambda$ 1306, we find three cases where the difference is less than 100 km s $^{-1}$ . In these cases, we kept the redshift value determined by the CIV $\lambda$ 1335 and OI $\lambda$ 1304+SiII $\lambda$ 1306 (label i and ii in Table 1). In the cases where the difference is larger than 100 km s $^{-1}$ , we considered the redshift given by AlIII $\lambda$ 1860 and SiII $\lambda$ 1892 (label iii in Table 1), since the peak of these lines is clearly observed. The uncertainty reported in Column 4 of Table 1 has been computed from the redshift difference between CIV $\lambda$ 1335, OI $\lambda$ 1304+SiII $\lambda$ 1306, and AlIII $\lambda$ 1860+SiII $\lambda$ 1892.



**Fig. 1.** Top panel: rest-frame composite spectrum (See Sect. §5.1). Abscissa corresponds to vacuum rest-frame wavelength in Å, while ordinate is in arbitrary intensity. Dot-dashed vertical lines identify the position at rest-frame of the strongest emission lines. Bottom panels: multicomponent fits after continuum subtraction for the 1400Å blend, Civλ1549 and 1900Å blend spectral regions (Sect. §4.2). In all the panels continuous black line marks the broad component at rest-frame associated to Sivλ1397, Civλ1549, AlIIIλ1860 and SiIII]λ1892 respectively, while the blue one corresponds to the blueshifted component associated to each emission. Dashed pink line marks the fit to the whole spectrum. Dot-dashed vertical lines correspond to the rest-frame wavelength of each emission line. In the Sivλ1397 spectral range, the cyan line marks the contribution of OIλ1304+SiIIλ1306 blend, while the magenta line corresponds to the CIIλ1335 emission line. In the Civλ1549 region, the yellow one corresponds to the OIII]λ1663 + AlIIIλ1670 blend. In the 1900Å blend range, FeII and FeI contributions are traced by dark and pale green lines respectively, magenta line marks the NIII]λ1750 and the yellow one corresponds to the SiIIλ1816. Lower panels correspond to the residuals, abscissa is in radial velocity units km s<sup>-1</sup>.

#### 4.2. Multicomponent fitting

Emission lines can be distinguished by their ionization potential (IP). The UV range covered in our spectra is populated by intermediate (IP  $\sim$  20–40 eV) and high ionization lines (IP > 40 eV). They offer an opportunity to characterize the behavior of different ionic species at the same time. In order to analyze the emission lines in our spectra, we carried out multicomponent fits using the SPECFIT routine from IRAF (Kriss 1994). This routine simultaneously fits the continuum, and emission/absorption line components. The best model is indicated by the minimum  $\chi^2$  over a spectral range where all components are included.

The main continuum source in the UV region is thought to arise from the accretion disk (Malkan & Sargent 1982). In the absence of extinction the continuum can be modeled by a single power-law over the full observed spectral range. However, due to the presence of absorptions (BAL sources) or dust extinction the continuum is flattened out in several sources (Sect. §3.1). Whenever possible, we fit the entire spectra range with a single power-law or a linear continuum. Otherwise we estimate locally the continuum. We divide the observed spectral range in three parts, which are centered on the most important emission lines relevant to our work.

**REGION 1: 1700–2200Å.** This range is dominated by the 1900Å blend which includes AlIIIλ1860, SiIII]λ1892, CIII]λ1909 and FeII lines (See Sect. §5.6 and Appendix B for FeII line identification). On the blue side of the blend SiIIλ1816 and NIII]λ1750 are also detected. AlIII, SiIII] and CIII] are intermediate-ionization lines (IIL) and according to Negrete et al. (2012, hereafter N12) can be well-modeled with Lorentzian profiles. The strengths of the three lines are allowed to vary freely in our SPECFIT model. FWHM AlIII and SiIII] were assumed equal, while FWHM CIII] was unconstrained. SiIIλ1816 and NIII]λ1750 were also modeled with Lorentzian profiles with flux and FWHM allowed to vary freely. All Lorentzian profile peaks were fixed at rest-frame. FeII makes an important contribution in the range 1715–1785 Å. We tried to use templates available in the literature (Brühweiler & Verner 2008; Mejía-Restrepo et al. 2016), but we could not reproduce the observed contribution. If that template is scaled to reproduce these features, the FeII emission around the MgIIλ2800 line would be overestimated by a large factor. On the converse if the template is normalized to FeII in the proximity of MgIIλ2800, the FeII emission in the spectral region 1700 – 2200 Å is negligible. We therefore decided to fit isolated Gaussian profiles for FeII at 1715 and 1785

Å. Their flux and FWHM vary freely.  $\text{FeIII}$  emission makes a larger contribution than  $\text{FeII}$  and appears especially strong on the red side of the 1900Å blend. We modeled the emission of this ion with the Vestergaard & Wilkes (2001) template and included an extra component at 1914 Å following N12 (the motivation for this choice is discussed in Sect. §6.3). Around 2020–2080 Å we were forced to include extra Gaussians in order to obtain a good fit (we found excess emission with respect to the  $\text{FeIII}$  template). The flux and FWHM of the features at 2020–2080 Å were allowed also to vary freely.

**REGION 2:** 1450–1700 Å. The  $\text{CIV}\lambda 1549$  emission line dominates this region and is accompanied by  $\text{HeII}\lambda 1640$ ,  $\text{OIII}\lambda 1663$  and  $\text{AlII}\lambda 1670$ . The broad component (BC) of  $\text{CIV}$  is modeled by a Lorentzian profile fixed at the rest-frame. The flux of the  $\text{CIV}_{\text{BC}}$  is free and FWHM is assumed to be the same as  $\text{AlIII}\lambda 1860$  and  $\text{SiIII}\lambda 1892$ . All the  $\text{CIV}$  profiles in our sample show a blueshift/bluward asymmetry. In order to model it with SPECFIT, we used one or two blueshifted skewed Gaussian profiles. The flux, FWHM, asymmetry and shift were unconstrained.  $\text{HeII}\lambda 1640$  was modeled assuming components similar to those of  $\text{CIV}\lambda 1549$ : Lorentzian and skewed Gaussian profiles for the BC and blueshifted components, respectively. The FWHM, shift and asymmetry were assumed equal to those of  $\text{CIV}\lambda 1549$ , but the flux varies freely.  $\text{OIII}\lambda 1663$  and  $\text{AlII}\lambda 1670$  were also modeled with unshifted Lorentzian profiles with their fluxed and FWHM varying freely.

**REGION 3:** 1300–1450 Å. The dominant emission in this region involves  $\text{SiIV}\lambda 1397 + \text{OIV}\lambda 1402$  (the 1400Å blend), and is accompanied by weaker  $\text{SiII}\lambda 1306$ ,  $\text{OI}\lambda 1304$  and  $\text{CII}\lambda 1335$  lines. The underlying assumption of the blend modeling is that the BC emission is dominated by  $\text{SiIV}\lambda 1397$  due to collisional de-excitation of the inter combination  $\text{OIV}\lambda 1402$  multiplet (Wills & Netzer 1979), while the blue shifted component is due to an inextricable contribution of both  $\text{OIV}\lambda 1402 + \text{SiIV}\lambda 1397$ . The  $\text{SiIV}\lambda 1397 + \text{OIV}\lambda 1402$  feature is a high-ionization blend, and shows a blueshifted, asymmetric profile not unlike  $\text{CIV}\lambda 1549$ . The broad component was modeled with the same emission components of  $\text{CIV}$ , whenever possible (in several cases the blue side of the  $\text{SiIV}\lambda 1397 + \text{OIV}\lambda 1402$  blend was strongly contaminated by absorption features making a good fit impossible), with only the flux varying freely. In the case of strong absorption the blueshifted emission was modeled independently from the one of  $\text{CIV}$ .

For lines that are composed of more than one component ( $\text{CIV}\lambda 1549$ ,  $\text{HeII}\lambda 1640$  and  $\text{SiIV}\lambda 1397 + \text{OIV}\lambda 1402$ ) the total profile parameters were also computed: FWHM, centroid at half maximum ( $c(\frac{1}{2})$ ), and asymmetry index (AI) defined by Zamfir et al. (2010). Unlike the SPECFIT components, these parameters provide a description of the profile that is not dependent on the model decomposition of the profile.

Figure 1 shows the multicomponent fitting made to the composite spectrum obtained by combining the normalized spectra of the GTC-xA sample (See Section §5.1 for a complete description). The bottom panels of the figure present the fits performed on the  $\text{SiIV}\lambda 1397$ ,  $\text{CIV}\lambda 1549$  and  $\text{AlIII}\lambda 1860$  spectral regions of the continuum subtracted spectrum. Residuals of the fits are shown in the lower part of the bottom panels. Spectra and multicomponent fits for the individual 19 quasars analyzed in this paper are shown from Fig. A.1 to A.19 in the Appendix A. Error estimations of blended emission lines were evaluated by building Monte-Carlo (MC) simulations as explained in Appendix B.

## 5. Results

### 5.1. Composite spectrum

We constructed a median composite spectrum from the individual observations in order to emphasize the main emission features (composites efficiently remove narrow absorption lines) and carry out a comparison with other samples (See Sect. §6.2). The composite GTC-xA spectrum corresponds to the median of all the normalized individual spectra including BALs. This simple approach produced a spectrum that reflects the behavior of the xA objects (See Fig. 1).

In xA sources, the 1900Å blend of the composite spectrum shows a high contribution of  $\text{AlIII}\lambda 1860$  and  $\text{SiIII}\lambda 1892$  compared to  $\text{CIII}\lambda 1909$ . Most of the emission on the red side of  $\text{SiIII}\lambda 1892$  can be attributed to  $\text{FeIII}$  excess in addition to the template, which is required to minimize the fit  $\chi^2$ . A fit with  $\text{CIII}\lambda 1909$  only (with no  $\text{FeIII}\lambda 1914$ ) would leave a large residual at 1915–1920 Å. Strong  $\text{FeIII}$  emission is confirmed also by the prominent bump at 2080 Å predominantly ascribed to the  $\text{FeIII}$  multiplet #48.


On the blue side of  $\text{CIV}\lambda 1549$ , a smooth and shallow trough is due to the combined effect of broad absorption lines that are frequent shortwards of  $\text{CIV}\lambda 1549$ . The emission profile of  $\text{CIV}\lambda 1549$  is in any case almost fully blueshifted, as observed in the majority of sources. Also the  $\text{SiIV}\lambda 1397$  median profile shows a blueshift asymmetry even if it is affected by the heavy absorptions frequently observed on the blue side of this line. The net effect is that  $\text{SiIV}\lambda 1397 + \text{OIV}\lambda 1402$  appear more symmetric because their blue wings are truncated by narrow and broad absorptions. The composite also clearly shows the prominent low-ionization features associated with  $\text{CII}\lambda 1335$ , and  $\text{OI}\lambda 1304$  blended with  $\text{SiII}\lambda 1306$ .  $\text{CIV}\lambda 1549 + \text{HeII}\lambda 1640$  shows low equivalent width, close to the boundary of WLQs (See Sect. §6.6). The composite spectrum is especially helpful for the increase in S/N that allows to trace the broad and faint  $\text{HeII}\lambda 1640$  profile, which shows a flat topped appearance. It is interpreted in the multicomponent fits as due to a strong blueshifted component, blended with a faint BC and with  $\text{OIII}\lambda 1663$  and  $\text{AlII}\lambda 1670$ .

### 5.2. Consistency of selection criteria for xA sources

Using lines in the 1900Å blend, MS14 proposed that xA sources with  $R_{\text{FeIII}} > 1$  show flux ratios  $\text{AlIII}\lambda 1860/\text{SiIII}\lambda 1892 \gtrsim 0.5$  and  $\text{CIII}\lambda 1909/\text{SiIII}\lambda 1892 \lesssim 1$ . Our sample was selected considering these criteria applied to SDSS noisy spectra previously excluded by MS14. The high S/N spectra of the GTC sample presented in this work confirm the defining criteria for the identification of the highly accreting sources in the UV range. Figure 2 shows the location of the 19 sources of our sample in the plane defined by  $\text{CIII}\lambda 1909/\text{SiIII}\lambda 1892$  vs  $\text{AlIII}\lambda 1860/\text{SiIII}\lambda 1892$ . In order to compare the behavior of the xA with the rest of the Pop. A, in Figure 2 are also represented control samples (FOS-xA, FOS-A and S14-A) described in Section §2.1.

Comparing the flux ratios shown by the low and high- $z$  Pop. A and xA samples, like MS14 have done, we observe a clear difference between the two kind of populations. The FOS-xA and GTC-xA samples show flux ratios different to the ones of FOS-A and S14-A samples. The GTC-xA sources occupy a very well defined region in the bottom right side of the panel, while the Pop. A sample occupy the left top space. Two of the objects (SDSSJ222753.07-092951.7 and SDSSJ125659.79-033813.8,  $\sim 10\%$  of our sample) do not rigorously satisfy the

selection criteria, although they present similar spectral properties to ones observed in the rest of the sample. The measured line ratios are actually borderline: the  $\text{AlIII}\lambda 1860/\text{SiIII}\lambda 1892$  ratio of the BAL SDSSJ125659.79-033813.8 is  $\approx 0.40 \pm 0.10$ , the  $\text{CIII}\lambda 1909/\text{SiIII}\lambda 1892$  ratio of SDSSJ222753.07-092951.7 is  $\approx 1.36 \pm 0.23$ . Therefore the two quasars will be considered along with all other GTC-xA sources.



**Fig. 2.** Relation between intensity ratios  $\text{AlIII}\lambda 1860/\text{SiIII}\lambda 1892$  and  $\text{CIII}\lambda 1909/\text{SiIII}\lambda 1892$ . Black, blue, green and red diamonds correspond to normal, BAL, mini-BAL and Radio-Loud quasars respectively for the present GTC-xA sample. Gray and cyan dots correspond to the FOS-A and FOS-xA sample respectively. Gray triangles correspond to S14-A sample. Yellow star marks the position of I Zw 1. Gray area represents the parameter space occupied by xA sources.

### 5.3. 1900Å Blend

Table 4 reports emission lines measurements corresponding to the 1900Å blend region:  $\text{AlIII}\lambda 1860$ ,  $\text{SiIII}\lambda 1892$ ,  $\text{CIII}\lambda 1909$ ,  $\text{NIII}\lambda 1750$  and  $\text{SiII}\lambda 1816$ . The first column lists the SDSS name. The equivalent width ( $W$ ), flux ( $F$ ) and FWHM of  $\text{AlIII}\lambda 1860$  are reported in Columns 2, 3, and 4 respectively. Equivalent width and flux are reported for  $\text{SiIII}\lambda 1892$  in Col. 5 and 6, since its FWHM is assumed equal to the one of  $\text{AlIII}\lambda 1860$ .  $W$ ,  $F$  and FWHM of  $\text{CIII}\lambda 1909$  are reported in Col. 7, 8 and 9 respectively. They are followed by the flux of the two faint features outside of the blend:  $\text{NIII}\lambda 1750$  and  $\text{SiII}\lambda 1816$  (Col. 10 and 11). All the parameters have uncertainties at  $1\sigma$  confidence level, which were obtained from the MC simulations. The MC method followed to estimate the uncertainties is explained in Appendix B.

In several instances the  $\text{AlIII}\lambda 1860$  doublet is partly resolved and is assumed to be at the rest-frame along with  $\text{SiIII}\lambda 1892$ . In general, the  $\text{AlIII}\lambda 1860$  flux, FWHM and  $W$  can be estimated with good accuracy as the doublet can be efficiently deblended save for the cases that the lines are very broad (FWHM  $\gtrsim 6000$  km s $^{-1}$ ).  $\text{AlIII}\lambda 1860$  is well fit with a symmetric and unshifted Lorentzian. Typical uncertainties in FWHM are  $\lesssim 10\%$ . Only

in one case (SDSSJ084036.16+235524.7) there is evidence for a blue asymmetry in the 1900Å blend although the  $\text{AlIII}\lambda 1860$  BC can be isolated with good accuracy. The fraction of the blueshifted to the total  $\text{AlIII}$  emission would of  $\approx 0.25$ , small compared to a median value of  $\approx 0.46$  for Civ. The presence of a blue shifted component in  $\text{AlIII}\lambda 1860$  is a rare occurrence. Only in one source of the S17 sample shows evidence of a blueward  $\text{AlIII}\lambda 1860$  excess (HE0359-3959, Martínez-Aldama et al. 2017).

The peak of the 1900Å blend is often not in correspondence of the  $\text{CIII}\lambda 1909$  wavelength. If the blend of  $\text{SiIII}$ ,  $\text{CIII}$  and  $\text{FeII}$  is peaked, the peak is preferentially redshifted beyond 1909 Å. The 1900Å blend profile may show excess emission on its red side (See e.g., SDSSJ021606.41+011509.5, Fig. A.3). In these cases, we suggest that the main contribution is coming from the  $\text{FeII}\lambda 1914$  line, modeled with a Lorentzian in excess to the  $\text{FeII}$  template (Sections §5.6 and §6.4), or by unresolved  $\text{FeII}$  emission.

The weaker lines  $\text{NIII}\lambda 1750$  and  $\text{SiII}\lambda 1816$  are detected in about 2/3 of the sample, e.g., they are both prominent in the case of the quasar SDSSJ021606.41+011509.5. The  $\text{NIII}\lambda 1750$  line shows a wide range in  $W$ . It is not detected in four sources but the detections show  $W$  in the range 0.5 – 5 Å.

### 5.4. Civλ1549 and HeIIλ1640

Measurements associated with the emission lines of the Civλ1549 spectral region are reported in Table 5. The first column lists the SDSS name. Column 2 reports the rest-frame specific continuum flux ( $f_c$ ) at 1350 Å. No error estimate is provided; the inter calibration with different standard star spectra suggests an uncertainty around 10%, which should include all main source of errors in the absolute flux scale. Cols. 3 to 7 tabulate  $W$ , flux, FWHM, centroid at half intensity ( $c(\frac{1}{2})$ ) and asymmetry index (A.I.) of the Civλ1549 total profile (broad plus blueshifted components). Col. 8 and 9 list the equivalent width and  $F$  of the broad component of Civ. Its FWHM is not reported, because it is assumed to be the same as FWHM  $\text{AlIII}\lambda 1860$ , due to the BC have to be emitted by the same zone. The equivalent width,  $F$ , FWHM and  $c(\frac{1}{2})$  of the blueshifted component of Civ are given in columns 10 to 13, respectively.

Table 6 reports the measurements associated with HeIIλ1640 and contaminant lines. Columns 2 to 7 list the  $W$  and flux of HeIIλ1640 total profile, HeIIλ1640 BC and HeIIλ1640 blue shifted component respectively. Col.8 lists the  $\text{OIII}\lambda 1663 + \text{AlIII}\lambda 1670$  flux. The last column reports the  $\text{NIV}\lambda 1486$  flux. The  $\text{NIV}\lambda 1486$  flux is highly uncertain due to absorption lines and to the blending with the blue wing of Civ, then in many of the cases only upper limits are reported.

On average, the HeIIλ1640 total flux is  $\sim 20\%$  that of Civλ1549. Since the typical  $W$  Civλ1549  $\approx 10 - 15$  Å, the typical  $W$  HeIIλ1640 is around  $\approx 3$  Å distributed over a broad profile with  $4000 \lesssim \text{FWHM} \lesssim 11000$  km s $^{-1}$ . The weakness of the HeIIλ1640 emission with respect to Civλ1549, with Civλ1549/HeIIλ1640  $\approx 5 - 6 \gtrsim 3$  is out of the question. It is important however to stress that our approach tends to maximize the contribution of the Civ<sub>BC</sub> to the total Civλ1549 emission. This may lead to an overestimation of the Civλ1549/HeIIλ1640 BC ratio. Nevertheless, in several cases (e.g., SDSS J214009.1-064403.9) we observed a significant Civλ1549 emission at rest-frame, implying that the Civλ1549/HeIIλ1640 estimate  $\approx 5 - 6$  is a safe one. We mention the Civλ1549/HeIIλ1640 ratio because its value is rather high for the low ionization parameter expected

**Table 4.** Measurements in the 1900Å blend region

SDSS Identification	Alm 1860			Sim 1892		Cm 1909			Nm 1750	Si 1816
	W	F	FWHM	W	F	W	F	FWHM	F	F
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
SDSSJ000807.27-103942.7	9.6 ± 1.2	11 ± 0.8	5410 ± 630	12.1 ± 1.9	13.2 ± 1.6	11.9 ± 1.9	12.7 ± 1.6	4810 ± 790	6.5 ± 0.9	3.1 ± 1.0
SDSSJ004241.95+002213.9	9.6 ± 1.3	25.2 ± 2.2	5060 ± 340	11.8 ± 1.5	30.2 ± 2.5	1.9 ± 1.0	4.9 ± 2.6	3060 ± 1620	4.8 ± 1.5	2.2 u
SDSSJ021606.41+011509.5	11.3 ± 1.3	33.2 ± 1.7	3190 ± 220	9.0 ± 1.2	25.7 ± 2.1	3.7 ± 1.0	11.6 ± 2.8	2690 ± 850	7.4 ± 1.8	8.1 ± 1.5 <sup>a</sup>
SDSSJ024154.42-004757.5	5.1 ± 1.1	9.5 ± 1.9	3340 ± 320	4.2 ± 0.6	7.7 ± 0.8	1.2 u	2.1 u	...	...	6.6 ± 0.8
SDSSJ084036.16+235524.7	6.1 ± 1.2	11.5 ± 2	3420 ± 380	5.3 ± 1.2	9.8 ± 2	2.0 ± 0.9	3.5 ± 1.6	2850 ± 420	...	3.3 ± 0.6 <sup>a</sup>
SDSSJ101822.96+203558.6	8.5 ± 1.2	26.7 ± 2.8	4870 ± 340	11.1 ± 1.8	33.4 ± 4.2	1.7 u	4.9 u	...	3 u	6 ± 2.4
SDSSJ103527.40+445435.6	7.9 ± 1.4	19 ± 2.8	7060 ± 480	11.5 ± 1.7	27.3 ± 2.9	...	...	...	...	...
SDSSJ105806.16+600826.9	5.1 ± 0.8	12.8 ± 1.6	4180 ± 650	7.9 ± 1.1	19.2 ± 1.7	0.8 u	2 u	...	4.2 ± 2.1	6.7 ± 1.3
SDSSJ10022.53+484012.6	9.1 ± 0.8	21.8 ± 2.2	4900 ± 390	15.6 ± 2.0	36.3 ± 2.9	2.6 ± 1.2	5.9 ± 2.7	3510 ± 1970	5.3 ± 1.3	2.4 ± 1.7
SDSSJ125659.79-033813.8	13.4 ± 3.0 <sup>a</sup>	24.1 ± 4.9 <sup>a</sup>	5950 ± 600	34.9 ± 6.4	60.9 ± 9.4	9.5 ± 4.3	16.2 ± 7.2	4000 ± 2090	7.1 ± 4.5 <sup>a</sup>	7.8 ± 3.7 <sup>a</sup>
SDSSJ131132.92+052751.2	5.8 ± 0.7	16.4 ± 1.3	3760 ± 250	6.2 ± 0.8	17.3 ± 1.6	...	...	...	...	...
SDSSJ143525.31+400112.2	4 ± 0.5	25.8 ± 2.3	3350 ± 270	3.8 ± 0.6	23.4 ± 2.5	0.4 u	2.3 u	...	5.3 ± 2.8	4.4 ± 1.9
SDSSJ144412.37+582636.9	13 ± 2.7	12.5 ± 2.3	8080 ± 1560	7.9 ± 1.9	7.5 ± 1.6	3.8 u	3.6 u	...	5.8 ± 0.5	1.6 ± 1.0
SDSSJ151258.36+352533.2	4.1 ± 0.6	10.8 ± 1.1	2780 ± 330	4.6 ± 0.7	11.8 ± 1.2	1.9 ± 0.6	4.6 ± 1.3	2110 ± 700	5.8 ± 1.2	5.3 ± 1.2
SDSSJ14009.01-064403.9	7.7 ± 0.9	13.9 ± 0.9	2620 ± 160	6.3 ± 0.9	11.3 ± 1.1	3.5 ± 1.2	7.4 ± 2.1	3500 ± 1150	0.8 u	1.5 ± 0.7
SDSSJ220119.62-083911.6	6.3 ± 1.1	26.7 ± 4	7250 ± 1050	9.3 ± 1.9	38.6 ± 6.9	7.4 ± 1.8	30 ± 6.5	6110 ± 1670	6.7 ± 2.4	...
SDSSJ222753.07-092951.7	6 ± 0.8	11.2 ± 1	4500 ± 420	7.8 ± 1.2	14.1 ± 1.7	10.8 ± 1.6	19.2 ± 2.2	4550 ± 590	2.9 ± 1.2	0.5 u
SDSSJ23132.83+010620.9	5.5 ± 1.5 <sup>a</sup>	38.5 ± 10.2 <sup>a</sup>	6310 ± 1640 <sup>a</sup>	8.8 ± 2.7	60.7 ± 17.4	3.7 ± 1.3	25.2 ± 8.6	5530 ± 4700	10.2 ± 5.5 <sup>a</sup>	6.6 u <sup>a</sup>
SDSSJ234657.25+145736.0	6.9 ± 0.8	21.1 ± 1.3	3660 ± 560	4.1 ± 0.9	12.4 ± 2.5	2.8 ± 1.1	8.3 ± 3.2	3040 ± 1450	2.8 ± 1.4 <sup>a</sup>	5.3 ± 1.4 <sup>a</sup>

NOTES. Columns are as follows: (1) SDSS name. (2), (5) and (7) report the equivalent width in units of Å. (3), (6), (8), (10) and (11) list fluxes in units of  $\text{erg s}^{-1} \text{cm}^{-2}$ . (4) and (9) correspond to the FWHM of Alm and Cm in units of  $\text{km s}^{-1}$ . <sup>a</sup> means that measurement is contaminated by absorption lines. A *u* letter marks an upper limit to the measurement.

in the emitting regions, if metallicity is solar or slightly supersolar (Sect. §6.3, N12).

An important feature in our spectra involves the strong blueshifts and asymmetries associated with the HILs. In ~50% of our sample, the C<sub>IV</sub> blueshifted component contributes ~50% of the total profile. He<sub>II</sub>λ1640 also shows a highly blueshifted component, contributing ~60% of the total flux. The A.I. values also support the presence of outflows: the blueshifted component pumped by radiative forces is more prominent than the virial component located at rest-frame. This is a major difference between High-Ionization Lines (e.g., C<sub>IV</sub>λ1549) on the one side, and low and intermediate-ionization lines on the other (e.g., Almλ1860). The last one are basically symmetric and show no evidence of large shifts with respect to the rest-frame.

An indication of the strength of the outflow is provided by the shift amplitude of the blue shifted component or by the shift amplitude of the total C<sub>IV</sub> profile measured by the centroid at half intensity,  $c(\frac{1}{2})$ . The most negative values indicate stronger blueshifts suggesting stronger outflows. BAL quasars tend to show smaller values of  $c(\frac{1}{2})$  (closer to 0) in the C<sub>IV</sub> profile, due to the presence of the absorption features in the blue side of the profile. The non-BAL quasars show values  $1000 \lesssim |c(\frac{1}{2})| \lesssim 5000 \text{ km s}^{-1}$ . He<sub>II</sub>λ1640 shows even stronger asymmetries than C<sub>IV</sub>λ1549, although they could be related to the faintness of He<sub>II</sub>λ1640 and the deblending uncertain of broad and blue components. The strong outflows could be important for feedback effects on the host galaxy (Section §6.7).

### 5.5. Si<sub>IV</sub>λ1397+O<sub>IV</sub>]λ1402

Table 7 reports the values of the emission lines in the Si<sub>IV</sub>λ1397+O<sub>IV</sub>]λ1402 spectral region (1400Å blend). The blue side of the 1400Å spectral region is seriously affected by absorption lines. The Si<sub>IV</sub>λ1397+O<sub>IV</sub>]λ1402 blend is formed by high-ionization lines. A blueshift asymmetry is expected in the profile, but in ~40% of the sample the blueshifted component can not be detected due to strong absorption lines. In the rest of the sample, the measurements of the  $c(\frac{1}{2})$  and FWHM

are also affected to the ever-present absorptions. C<sub>III</sub>λ1335 and O<sub>I</sub>+Si<sub>II</sub>λ1304 are also affected by the absorption. In the majority of the sample, they cannot be detected or their flux uncertainty is large.

### 5.6. The Fe<sub>II</sub> and Fe<sub>III</sub> emission

Fe<sub>II</sub> emission is usually not very strong in the range 1700–2100 Å, even in the case of the strongest optical Fe<sub>II</sub> emitters. Only a few features are detected at ~ 1715 Å, 1785 Å and 2020 Å. We modeled these individual features with single Gaussians, without using a model template emission. On the other hand, the Fe<sub>III</sub> emission has a most important role in xA sources. It was modeled using the scaled and broadened template of Vestergaard & Wilkes (2001) based on I Zw 1. However, it is not enough to reproduce the observed Fe<sub>III</sub> emission. We add two extra emission component at 1914 and 2080 Å (modeled by two Gaussians), to account for the line of Fe<sub>III</sub> UV #34 line that may be enhanced by Ly $\alpha$  fluorescence, and for the broad hump at 2080 Å, mainly ascribed to the blended features of Fe<sub>III</sub> UV #48. Most relevant Fe<sub>II</sub> and Fe<sub>III</sub> features between 1700 and 2100 Å are individually discussed Appendix C.1.

Table 8 reports the equivalent widths and the fluxes of the most important UV Fe<sub>II</sub> multiplet #191 (Fe<sub>II</sub> 1785 Å) and the ones of the total Fe<sub>II</sub> emission, i.e., the sum of the Fe<sub>II</sub> features described above: at 1715, 1785 and 2020 Å. Table 9 reports equivalent widths and fluxes of the most important individual features ascribed to Fe<sub>III</sub> (1914 and 2080 Å) and the total Fe<sub>III</sub> emission: the sum of the individual features and the template contribution in the full range. It is interesting to note that Fe<sub>III</sub> emission is almost always stronger than Fe<sub>II</sub>, by a factor ~2–3. The strong Fe<sub>III</sub> emission will be discussed in Sect. §6.4 and Appendix C.2.

**Table 5.** Measurements on C<sub>IV</sub>λ1549 region

SDSS Identification	$f_{\lambda}(1350\text{\AA})$	C <sub>IV</sub> 1549 <sub>TOTAL</sub>					C <sub>IV</sub> 1549 <sub>BC</sub>		C <sub>IV</sub> 1549 <sub>BLUE</sub>			
		W	F	FWHM	$c(\frac{1}{2})$	A.I.	W	F	W	F	FWHM	$c(\frac{1}{2})$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
SDSSJ000807.27-103942.7	2.9	37.7 ± 6.6 <sup>a</sup>	71.9 ± 10.4 <sup>a</sup>	6200 ± 420	-1130 ± 210	-0.044 ± 0.103	27.4 ± 6.0 <sup>a</sup>	51.8 ± 10.1 <sup>a</sup>	10.3 ± 5.3 <sup>a</sup>	20.1 ± 10.2 <sup>a</sup>	4770 ± 300	-2740 ± 150
SDSSJ004241.95+002213.9	3.7	19.5 ± 4.3	61.8 ± 12.2	10080 ± 340	-4530 ± 170	-0.438 ± 0.087	5.2 ± 4.4	16.4 ± 13.6	14.3 ± 2.7	45.4 ± 7.4	9550 ± 250	-5170 ± 130
SDSSJ021606.41+011509.5	3.5	19.1 ± 2.6 <sup>b</sup>	62.4 ± 5.5 <sup>b</sup>	4130 ± 230	-850 ± 110	-0.074 ± 0.086	13.4 ± 2.5	43.8 ± 6.8	5.7 ± 1.1 <sup>b</sup>	18.6 ± 3 <sup>b</sup>	3020 ± 160	-1760 ± 80
SDSSJ024154.42-004757.5	3.1	10.0 ± 1.8 <sup>a</sup>	25.4 ± 3.7 <sup>a</sup>	6830 ± 450	-2420 ± 220	-0.375 ± 0.073	3.9 ± 1.3 <sup>a</sup>	9.8 ± 3.2 <sup>a</sup>	6.1 ± 1.9 <sup>a</sup>	15.6 ± 4.7 <sup>a</sup>	6570 ± 380	-3560 ± 1909
SDSSJ084036.16+235524.7	3.1	14.4 ± 2.2	38.1 ± 4.5	6630 ± 370	-1670 ± 180	-0.378 ± 0.070	9.9 ± 2.4	26.6 ± 5.9	4.5 ± 2	11.5 ± 4.8	4590 ± 240	-3850 ± 120
SDSSJ101822.96+203558.6	4.4	12.1 ± 2.7 <sup>a</sup>	47.5 ± 9.5 <sup>a</sup>	9580 ± 390	-3990 ± 200	-0.533 ± 0.069	2.6 ± 1.7 <sup>a</sup>	10.0 ± 6.6 <sup>a</sup>	9.5 ± 2.1 <sup>a</sup>	37.5 ± 7.4 <sup>a</sup>	9770 ± 320	-4520 ± 160
SDSSJ103527.40+445435.6	3.0	12.2 ± 4.2 <sup>a</sup>	33.4 ± 11.1 <sup>a</sup>	9390 ± 450	-3460 ± 230	0.193 ± 0.073	5.9 ± 3.3	16.2 ± 8.8	6.3 ± 2.5 <sup>a</sup>	17.2 ± 6.7 <sup>a</sup>	6350 ± 320	-5050 ± 160
SDSSJ105806.16+600826.9	4.6	11.4 ± 2.2	39.3 ± 6.4	6880 ± 400	-2890 ± 200	-0.405 ± 0.076	1.5 ± 1.4	5.0 ± 4.8	9.9 ± 1.9	38.4 ± 5.4	6840 ± 380	-3160 ± 190
SDSSJ110022.53+484012.6	3.7	24.3 ± 2.8	75.9 ± 4.1	8010 ± 540	-4430 ± 270	-0.203 ± 0.071	7.1 ± 1.7	22.0 ± 4.9	17.2 ± 2	53.9 ± 3.4	7140 ± 410	-5120 ± 210
SDSSJ125659.79-033813.8	2.8	47.4 ± 9.9 <sup>a</sup>	111.2 ± 20.4 <sup>a</sup>	5650 ± 460	-2020 ± 230	-0.186 ± 0.091	25.8 ± 11.8	60.5 ± 27.1 <sup>a</sup>	21.6 ± 8.2 <sup>a</sup>	50.7 ± 18.5	5000 ± 400	-2950 ± 200
SDSSJ131132.92+052751.2	2.0	6.6 ± 0.9 <sup>a,b</sup>	17.6 ± 1.8 <sup>a,b</sup>	5700 ± 320	-1180 ± 160	-0.220 ± 0.088	4.7 ± 0.9 <sup>a</sup>	12.5 ± 2.2 <sup>a</sup>	1.9 ± 0.7 <sup>b</sup>	5.1 ± 1.7 <sup>b</sup>	4320 ± 220	-2710 ± 110
SDSSJ143525.31+400112.2	10.3	9.9 ± 1.3 <sup>a</sup>	85.2 ± 7.2 <sup>a</sup>	8210 ± 680	-3090 ± 340	-0.214 ± 0.074	3.7 ± 1.2	31.6 ± 9.5	6.2 ± 1.0 <sup>a</sup>	53.6 ± 6.8 <sup>a</sup>	6990 ± 430	-4330 ± 220
SDSSJ144412.37+582636.9	1.0	13.9 ± 1.8 <sup>a</sup>	15.0 ± 1.2 <sup>a</sup>	8570 ± 450	-1990 ± 230	0.020 ± 0.085	9.0 ± 2.0 <sup>a</sup>	9.7 ± 2 <sup>a</sup>	4.9 ± 1.7 <sup>a</sup>	5.3 ± 1.7 <sup>a</sup>	6310 ± 320	-3480 ± 160
SDSSJ151258.36+352533.2	4.7	8.8 ± 1.4 <sup>a</sup>	32.6 ± 4.0 <sup>a</sup>	5300 ± 220	-1880 ± 110	-0.022 ± 0.090	3.3 ± 1.1 <sup>a</sup>	12.1 ± 3.7 <sup>a</sup>	5.5 ± 1.2 <sup>a</sup>	20.5 ± 4.2 <sup>a</sup>	3990 ± 200	-2630 ± 100
SDSSJ214009.01-064403.9	1.9	16.1 ± 2.4 <sup>b</sup>	29.7 ± 3.4 <sup>b</sup>	5260 ± 770	-1170 ± 380	-0.424 ± 0.069	10.8 ± 2.6 <sup>b</sup>	20.0 ± 4.3 <sup>b</sup>	5.3 ± 1.8 <sup>b</sup>	9.7 ± 3.2 <sup>b</sup>	5350 ± 270	-3770 ± 140
SDSSJ220119.62-083911.6	5.5	18.4 ± 3.8 <sup>b</sup>	103.4 ± 19.0 <sup>b</sup>	4430 ± 510	-830 ± 250	0.002 ± 0.092	15.2 ± 4.9 <sup>b</sup>	85.7 ± 26.4 <sup>b</sup>	3.2 ± 2.7 <sup>b</sup>	17.7 ± 15 <sup>b</sup>	2920 ± 170	-1160 ± 90
SDSSJ222753.07-092951.7	2.8	31.9 ± 5.5	77.9 ± 11	5260 ± 410	-1190 ± 200	-0.277 ± 0.051	16.3 ± 4.8	39.5 ± 11.2	15.7 ± 6.8	38.4 ± 16.2	5820 ± 420	-2340 ± 210
SDSSJ233132.83+010620.9	8.3	21.4 ± 4.4	174.2 ± 35.8	11230 ± 700	-2430 ± 350	-0.333 ± 0.097	16.1 ± 5	132.0 ± 39.1	5.3 ± 1.6	42.2 ± 12.4	6990 ± 380	-6960 ± 190
SDSSJ234657.25+145736.0	4.4	11.1 ± 1.7 <sup>a</sup>	41.5 ± 4.6 <sup>a</sup>	6390 ± 800	-1420 ± 400	-0.392 ± 0.071	7.2 ± 1.5 <sup>a</sup>	26.9 ± 4.9 <sup>a</sup>	3.9 ± 1.5 <sup>a</sup>	14.6 ± 5.3 <sup>a</sup>	6670 ± 340	-4750 ± 170

NOTES. Columns are as follows: (1) SDSS name. (2) Continuum flux measured at 1350 Å. (3), (8) and (10) report the equivalent width in unit of Å. (4), (9) and (11) list line fluxes in units of erg s<sup>-1</sup> cm<sup>-2</sup>. (5) and (12) list the FWHM of the corresponding line in km s<sup>-1</sup>. (6) and (13) report the centroid at half intensity in units of km s<sup>-1</sup>. (7) lists the asymmetry index (A.I.). <sup>a</sup>Measurement contaminated by narrow absorption lines. <sup>b</sup>Measurement contaminated by broad absorption lines.

**Table 6.** Measurements on He II λ1640 region

SDSS Identification	He II 1640 <sub>TOTAL</sub>		He II 1640 <sub>BC</sub>		He II 1640 <sub>BLUE</sub>		Al III + O III] 1667	N IV] 1486
	W	F	W	F	W	F	F	F
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
SDSSJ000807.27-103942.7	7.9 u	13.2 u	2.5 u	4.0 u	5.4 u	9.2 u	8.5 ± 4.1	2.4 u <sup>a</sup>
SDSSJ004241.95+002213.9	2.7 ± 2.4	8.3 ± 7.2	1.0 u	3.1 u	1.7 ± 1.2	5.2 ± 3.6	2.7 u	7.8 u
SDSSJ021606.41+011509.5	2.1 ± 1.9	6.6 ± 5.8	1.3:	4.1:	0.8:	2.5:	2.2 ± 1.4	...
SDSSJ024154.42-004757.5	3.3 ± 0.9	7.7 ± 2	0.5	1.1:	2.8 ± 0.9	6.6 ± 2.1	4.5 u	5.5 ± 3.5 <sup>a</sup>
SDSSJ084036.16+235524.7	2.3 ± 1.6	5.6 ± 3.7	1.2 ± 1	3.0 ± 2.5	1.1 ± 1	2.6 u 2.2	2.5 ± 1.9	3.7 u
SDSSJ101822.96+203558.6	2.2 u	8.2 u	0.7 u	2.5 u	1.5 ± 0.7	5.7 ± 2.5	5.5 u	8 u
SDSSJ103527.40+445435.6	2.5 u	6.5 ± 2.3	1.6 u	4.2 u	0.9 ± 0.4	2.3 ± 1	5.1 ± 2.6	16.2 u <sup>a</sup>
SDSSJ105806.16+600826.9	1.1 u	3.2 u	0.5:	1.5:	0.6 u	1.7 u	3 u	10 u
SDSSJ110022.53+484012.6	2.8 ± 1.0 <sup>a</sup>	8.2 ± 2.8 <sup>a</sup>	1.0: <sup>a</sup>	2.9: <sup>a</sup>	1.8 ± 0.9	5.3 ± 2.6	1.7 u	...
SDSSJ125659.79-033813.8	7.0 u <sup>a</sup>	15.5 u <sup>a</sup>	2.7:	6.0:	4.3 u <sup>a</sup>	9.5 u <sup>a</sup>	5.5 u	...
SDSSJ131132.92+052751.2	4.0 ± 1.0 <sup>a</sup>	10.4 ± 2.3 <sup>a</sup>	1.2 ± 0.9	3.1 ± 2.3	2.8 ± 1.1 <sup>a</sup>	7.3 ± 2.7 <sup>a</sup>	6.7 ± 2.5 u <sup>a</sup>	...
SDSSJ143525.31+400112.2	2.3 ± 0.9 <sup>a</sup>	18.2 ± 6.5 <sup>a</sup>	1.3 ± 0.8 <sup>a</sup>	10.3 ± 5.9 <sup>a</sup>	1 ± 0.5	7.9 ± 3.7	9.2 ± 4.1 <sup>a</sup>	5.8 u <sup>a</sup>
SDSSJ144412.37+582636.9	1.3 ± 0.7	4.0 ± 2.5	1.1 ± 0.8	2.9 ± 2.2	0.2:	1.1:	4.9 u	5.7 ± 4.2 u <sup>a</sup>
SDSSJ151258.36+352533.2	2.0 u <sup>a</sup>	6.6 u <sup>a</sup>	1 ± 0.7 <sup>a</sup>	3.2 ± 2.3 <sup>a</sup>	1 u <sup>a</sup>	3.4 u <sup>a</sup>	5.3 ± 4.1	...
SDSSJ214009.01-064403.9	3.8 ± 1.4	7.0 ± 2.5	1.9 ± 0.8	3.5 ± 1.4	1.9 ± 1.2	3.5 ± 2.2	7.6 u	...
SDSSJ220119.62-083911.6	...	...	...	...	...	...	...	...
SDSSJ222753.07-092951.7	5.7 ± 2.6	13.1 ± 5.7	1.4 ± 1.3	3.2 ± 2.9	4.3 ± 2.7	9.9 ± 6	2.7 u	6.8 u <sup>a</sup>
SDSSJ233132.83+010620.9	6.3 ± 2.9	46.9 ± 21.4	2.6 ±	21.0 u	3.7 ± 1.8	25.9 ± 12	17.8 u	9.3 u
SDSSJ234657.25+145736.0	1.7 ± 1.1 <sup>a</sup>	5.8 ± 3.7 <sup>a</sup>	1.7 ± 1.1 <sup>a</sup>	5.8 ± 3.7 <sup>a</sup>	...	...	7.5 ± 3.6	6.7 u <sup>a</sup>

NOTES. Columns are as follows: (1) SDSS name. Equivalent width are listed in col. (2), (4) and (6) in units of Å. Line fluxes are reported in cols. (3), (5), (7), (8) and (9) in units of erg s<sup>-1</sup> cm<sup>-2</sup>. <sup>a</sup>Measurement contaminated by narrow absorption lines. Two colons (:) indicate a high uncertainty in the measurement. Letter *u* marks an upper limit to the measurement.

### 5.7. The origin of the C<sub>IV</sub>λ1549 line broadening

The GTC-xA sample shows a very tight correlation between FWHM and  $c(\frac{1}{2})$  of the total C<sub>IV</sub>λ1549 profile (Fig. 3). The best fitting to our new sample is  $c(\frac{1}{2}) \approx (-0.613 \pm 0.081) \cdot \text{FWHM} + (1893 \pm 545) \text{ km s}^{-1}$ . This relation is consistent within uncertainties with the one found by S17 involving Pop. A sources. The range of blueshifts is from  $\approx -1000$  to  $-5000 \text{ km s}^{-1}$  in both cases; however, the distribution of the GTC-xA is weighted in favor of larger shifts and broader FWHM with respect to the one of S17. As noted by S17, the correlation of Fig. 3 justify the decomposition of C<sub>IV</sub>λ1549 profile into the BC and a blueshifted

excess, which becomes the dominant source of broadening in the case of the largest FWHM.

### 5.8. Al III λ1860 as a High Redshift Virial Estimator

Identification of a UV virial estimator for high redshift quasars ( $z \geq 2$ ) has been an important quest for several years (e.g., review by Marziani et al. 2017b). Once Hβ is lost to ground based spectroscopy, Mg II λ2800 can be exploited up to  $z \sim 2.0$  (Trakhtenbrot & Netzer 2012; Shen & Liu 2012; Marziani et al. 2013). C<sub>IV</sub>λ1549 is the obvious candidate for quasars be-

**Table 7.** Measurements on 1400Å blend region

SDSS Identification	Siv + Oriv] 1400 <sub>TOTAL</sub>		Siv + Oriv] 1400 <sub>BC</sub>		Siv + Oriv] 1400 <sub>BLUE</sub>		CII 1332	OI + S II 1304
	W	F	W	F	W	F	F	F
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
SDSSJ000807.27-103942.7	15.9 ± 2.1 <sup>a</sup>	40.0 ± 3.3 <sup>a</sup>	15.9 ± 2.1 <sup>a</sup>	40.0 ± 3.3 <sup>a</sup>	...	...	4.8 ± 3.6 <sup>a</sup>	1.9 ± 1.1 <sup>a</sup>
SDSSJ004241.95+002213.9	11.1 ± 1.8 <sup>a</sup>	38.0 ± 5.0 <sup>a</sup>	7.7 ± 1.9 <sup>a</sup>	26.3 ± 5.9 <sup>a</sup>	3.4 ± 0.9 <sup>a</sup>	11.7 ± 2.8 <sup>a</sup>	...	...
SDSSJ021606.41+011509.5	19.5 ± 2.2 <sup>b</sup>	67.6 ± 3.8 <sup>b</sup>	19.5 ± 2.2 <sup>b</sup>	67.6 ± 3.8 <sup>b</sup>	...	...	11.7 ± 5.1	10 ± 3.8
SDSSJ024154.42-004757.5	9.0 ± 1.5 <sup>a</sup>	25.9 ± 3.4 <sup>a</sup>	6.4 ± 1.5 <sup>a</sup>	18.4 ± 4 <sup>a</sup>	2.6 ± 1.2	7.5 ± 3.4	7.9 ± 2.7 <sup>a</sup>	8.3 ± 2.1 <sup>a</sup>
SDSSJ084036.16+235524.7	7.1 ± 1	21.0 ± 2.1	7.1 ± 1	21.0 ± 2.1	...	...	5.4 ± 2.8 <sup>a</sup>	3.9 ± 1.8
SDSSJ101822.96+203558.6	9.1 ± 1.3 <sup>a</sup>	39.3 ± 3.8 <sup>a</sup>	5.3 ± 1.3 <sup>a</sup>	23.0 ± 5 <sup>a</sup>	3.8 ± 0.7 <sup>a</sup>	16.3 ± 2.7 <sup>a</sup>	5.3 ± 2.1 <sup>a</sup>	11.4 ± 3.5 <sup>a</sup>
SDSSJ103527.40+445435.6	9.1 ± 1.7 <sup>a</sup>	26.4 ± 4.2 <sup>a</sup>	6.8 ± 2.1 <sup>a</sup>	19.8 ± 5.7 <sup>a</sup>	2.3 ± 0.8 <sup>a</sup>	6.6 ± 2.2 <sup>a</sup>	2.6 u	4 u
SDSS105806.16+600826.9	12.1 ± 1.8	49.7 ± 5.4	8 ± 1.7	33.0 ± 6	4.1 ± 1.2	16.7 ± 4.7	14.3 ± 4.7	8 ± 2.1
SDSSJ110022.53+484012.6	12.8 ± 1.7	41.8 ± 3.8	5.7 ± 1.1	22.1 ± 3.6	7.1 ± 1.3	19.7 ± 3	12.8 ± 6.1	28 ± 11.9
SDSSJ125659.79-033813.8	21.0 ± 3.3 <sup>a</sup>	55.2 ± 6.8 <sup>a</sup>	15.8 ± 10.0 <sup>a</sup>	41.7 ± 26.1 <sup>a</sup>	5.2 ± 2.9	13.5 ± 7.3	14.9 ± 10.8 <sup>a</sup>	27.6 ± 11.1 <sup>a</sup>
SDSSJ131132.92+052751.2	5.7 <sup>b</sup>	10.7 <sup>b</sup>	5.7 <sup>b</sup>	10.7 <sup>b</sup>	...	...	...	...
SDSSJ143525.31+400112.2	7.2 ± 1.9 <sup>a</sup>	70.2 ± 17.4 <sup>a</sup>	5.4 ± 2.4 <sup>a</sup>	52.2 ± 22.9 <sup>a</sup>	1.8 ± 1.7 <sup>a</sup>	18.0 ± 16.7 <sup>a</sup>	14.2 ± 10.2	23.4 ± 18.6 <sup>a</sup>
SDSSJ144412.37+582636.9	12.7 ± 3.7 <sup>b</sup>	14.7 ± 4.0 <sup>b</sup>	12.7 ± 3.7 <sup>b</sup>	14.7 ± 4.0 <sup>b</sup>	...	...	...	...
SDSSJ151258.36+352533.2	6.4 ± 1.2 <sup>a</sup>	28.5 ± 4.7 <sup>a</sup>	4.9 ± 1.5 <sup>a</sup>	21.8 ± 6.2 <sup>a</sup>	1.5 ± 0.7 <sup>a</sup>	6.7 ± 3.2 <sup>a</sup>	...	12.4 ± 4.3 <sup>a</sup>
SDSSJ214009.01-064403.9	20.5 ± 2.9 <sup>b</sup>	39.1 ± 3.8 <sup>b</sup>	18 ± 3.2	34.3 ± 5.1	2.5 ± 1.8 <sup>b</sup>	4.8 ± 3.4 <sup>b</sup>	3.7 ± 2.1 <sup>b</sup>	8.7 ± 2.8
SDSSJ220119.62-083911.6	13.4 ± 6.6 <sup>b</sup>	71.7 ± 34.4 <sup>b</sup>	13.4 ± 6.6 <sup>b</sup>	71.7 ± 34.4 <sup>b</sup>	...	...	...	...
SDSSJ222753.07-092951.7	10.3 ± 1.9 <sup>a</sup>	28.2 ± 4.4 <sup>a</sup>	10.3 ± 1.9 <sup>a</sup>	28.2 ± 4.4 <sup>a</sup>	...	...	...	...
SDSSJ233132.83+010620.9	...	...	...	...	...	...	...	...
SDSSJ234657.25+145736.0	10.3 ± 1.6 <sup>a</sup>	41.7 ± 4.8 <sup>a</sup>	9.8 ± 1.4 <sup>a</sup>	39.7 ± 3.9 <sup>a</sup>	0.5 <sup>a</sup>	2.0 <sup>a</sup>	20.5 ± 4.1 <sup>a</sup>	...


NOTES. Columns are as follows: (1) SDSS name. Cols. (3), (5), (7), (8) and (9) list line fluxes in units of  $\text{erg s}^{-1} \text{cm}^{-2}$ . Cols. (2), (4) and (6) list the equivalent width in Å. <sup>a</sup>Measurement contaminated by narrow absorption lines. <sup>b</sup>Measurement contaminated by broad absorption lines. Two colons (: ) indicates a high uncertainty in the measurement. Letter *u* marks an upper limit to the measurement.

**Table 8.** Flux and Equivalent width of FeII contribution

SDSS Identification	FeII $\lambda 1785$		FeII <sub>TOTAL</sub>	
	W	F	W	F
(1)	(2)	(3)	(4)	(5)
SDSSJ000807.27-103942.7	3.5 ± 0.7	4.4 ± 0.7	11.2 ± 2.8	14.0 ± 3.2
SDSSJ004241.95+002213.9	2.1 ± 0.6	5.7 ± 1.4	6.2 ± 2.0	16.7 ± 5.2
SDSSJ021606.41+011509.5	2.5 ± 0.5	7.0 ± 1.1	9.7 ± 2.0	29.3 ± 5.1
SDSSJ024154.42-004757.5	1.5 ± 0.2	3.1 ± 0.4	6.5 ± 1.1	12.4 ± 1.8
SDSSJ084036.16+235524.7	1.1 ± 0.3	2.2 ± 0.6	7.9 ± 1.9	16.0 ± 3.5
SDSSJ101822.96+203558.6	1.7 ± 0.6	5.5 ± 1.8	6.8 ± 1.9	21.0 ± 5.5
SDSSJ103527.40+445435.6	2.7 ± 0.6	6.6 ± 1.2	4.8 ± 1.1	12.1 ± 2.5
SDSS105806.16+600826.9	0.4:	1.0:	2.5 ± 0.8	7.0 ± 2.2
SDSSJ110022.53+484012.6	1.3 ± 0.4	3.2 ± 1.0	3.4 ± 1.2	8.7 ± 3.1
SDSSJ125659.79-033813.8	...	...	9.9 ± 4.3	16.0 ± 6.7
SDSSJ131132.92+052751.2	1.0 ± 0.4	2.9 ± 1.0	2.3 ± 0.7	6.4 ± 1.9
SDSSJ143525.31+400112.2	1.1 ± 0.3	7.4 ± 1.6	3.8 ± 1.0	23.6 ± 5.4
SDSSJ144412.37+582636.9	...	...	5.5 ± 2.2	5.0 ± 1.9
SDSSJ151258.36+352533.2	1.9 ± 0.4	5.3 ± 1.0	3.8 ± 0.8	11.0 ± 1.9
SDSSJ214009.01-064403.9	0.7 ± 0.3	1.2 ± 0.5	6.5 ± 1.5	11.7 ± 2.4
SDSSJ220119.62-083911.6	...	...	1.1 ± 0.5	3.8 ± 1.8
SDSSJ222753.07-092951.7	0.2 ± 0.1	0.4 ± 0.2	5.3 ± 1.3	10.5 ± 2.3
SDSSJ233132.83+010620.9	1.3 ± 0.9	9.6 ± 6.7	5.5 ± 2.9	35.5 ± 18.0
SDSSJ234657.25+145736.0	...	...	1.3 ± 0.5	4.5 ± 1.5

NOTES. Columns are as follows: (1) SDSS name. (2) and (4) report the equivalent width in units of Å. (3) and (5) correspond to the flux in units of  $\text{erg s}^{-1} \text{cm}^{-2}$ .

yond  $z \sim 2.0$ , but the Civ line profiles often show large blue shifts/asymmetries whose amplitude is a function of source location along the 4DE1 sequence and also of luminosity (S17). This implies that  $M_{\text{BH}}$  estimation using FWHM Civ can be uncertain by a large factor (Sulentic et al. 2007; Netzer et al. 2007; Mejía-Restrepo et al. 2018). There have been several recent attempts to account for the bias associated with the non-virial broadening of Civ  $\lambda 1549$  (Park et al. 2013; Matsuoka et al. 2013; Brotherton et al. 2015; Denney et al. 2016; Coatman et al. 2017), however no correction yields safe  $M_{\text{BH}}$  estimates especially if Population A and B sources are not considered separately. We focus here on extreme Pop. A (xA) sources and attempt to use



**Fig. 3.** Relation between the  $c(\frac{1}{2})$  and the FWHM of the total Civ  $\lambda 1549$  emission. Dashed gray line corresponds to the best fit using an orthogonal method. Black, blue, green and red filled circles correspond respectively to normal, BAL, mini-BAL and RL quasars for the GTC-xA sample.

AlIII  $\lambda 1860$  as an H $\beta$  virial surrogate. The first composites of low- $z$  FOS sources suggested that AlIII  $\lambda 1860$  becomes more prominent at the extreme Pop. A end of the 4DE1 sequence (Bachev et al. 2004). The AlIII  $\lambda 1860$  line shows a FWHM similar to the one of H $\beta$ , which suggests that both lines are emitted in similar dynamical condition (Negrete et al. 2013). Analysis of VLT ISAAC + FORS spectra of high- $z$  Pop. A sources (Marziani et al. 2017b) show also that FWHM AlIII  $\lambda 1860$  agrees well with FWHM H $\beta$ . Previously, the emission lines associated with the

**Table 9.** Flux and Equivalent width of FeIII contribution

SDSS Identification	Fe III $\lambda 1914$		Fe III $\lambda 2080$		Fe III TOTAL	
	W	F	W	F	W	F
(1)	(2)	(3)	(4)	(5)	(6)	(7)
SDSSJ000807.27-103942.7	1.9 ± 0.7	1.9 ± 0.7	7.5 ± 0.8	5.9 ± 0.6	9.9 ± 2.0	17.1 ± 3.0
SDSSJ004241.95+002213.9	2.9 ± 0.8	7.1 ± 2.0	6.4 ± 1.1	13.8 ± 2.5	4.5 ± 1.3	31.6 ± 8.8
SDSSJ021606.41+011509.5	5.8 ± 0.9	16.2 ± 2.4	6.6 ± 0.7	17.7 ± 1.9	25.1 ± 3.4	103.9 ± 9.8
SDSSJ024154.42-004757.5	2.3 ± 1.0	3.8 ± 1.7	0.1 u	0.2 u	12.6 ± 3.9	24.6 ± 7.2
SDSSJ084036.16+235524.7	3.4 ± 1.8	6.1 ± 3.1	...	...	23.0 ± 3.2	44.5 ± 4.3
SDSSJ101822.96+203558.6	4.7 ± 0.9	13.7 ± 2.7	6.8 ± 1.3	17.2 ± 3.2	11.8 ± 2.7	63.9 ± 13.1
SDSSJ103527.40+445435.6	5.8 ± 0.7	13.1 ± 1.7	9.0 ± 0.8	18.3 ± 1.6	3.1 ± 0.6	38.2 ± 6.7
SDSSJ105806.16+600826.9	3.5 ± 0.5	8.0 ± 1.1	1.8 ± 0.9	3.8 ± 1.8	12.9 ± 2.7	40.6 ± 7.3
SDSSJ110022.53+484012.6	0.9 ± 0.6	2.0 ± 1.4	3.2 ± 1.0	6.1 ± 1.9	5.6 ± 2.5	20.0 ± 8.6
SDSSJ125659.79-033813.8	2.6 ± 2.3	4.3 ± 3.8	6.5 ± 2.7	9.0 ± 3.8	7.8 ± 6.0	25.7 ± 19.5
SDSSJ131132.92+052751.2	3.1 ± 0.4	8.6 ± 1.0	3.1 ± 0.5	8.1 ± 1.2	4.4 ± 0.9	28.7 ± 5.0
SDSSJ143525.31+400112.2	3.2 ± 0.7	19 ± 4.4	2.1 ± 0.6	11.0 ± 3.2	8.7 ± 2.0	79.4 ± 16.1
SDSSJ144412.37+582636.9	3.6 ± 1.3	3.2 ± 1.1	4.0 ± 2.1	3.2 ± 1.7	4.3 ± 3.3	10.2 ± 7.9
SDSSJ151258.36+352533.2	3.0 ± 0.6	7.3 ± 1.5	2.1 ± 0.6	4.5 ± 1.3	6.7 ± 1.7	27.2 ± 6.3
SDSSJ214009.01-064403.9	3.7 ± 0.8	6.5 ± 1.4	5.3 ± 0.8	9.0 ± 1.4	4.3 ± 1.2	21.9 ± 5.9
SDSSJ220119.62-083911.6	2.0 ± 0.6	7.6 ± 2.5	1.1 ± 0.7	3.6 ± 2.5	8.0 ± 2.6	41.2 ± 13.0
SDSSJ222753.07-092951.7	1.0 ± 0.5	1.8 ± 0.9	7.1 ± 1.1	10.4 ± 1.6	8.0 ± 2.1	25.4 ± 6.1
SDSSJ233132.83+010620.9	1.3 ± 0.3	8.7 ± 2.3	3.3 ± 0.8	20.5 ± 4.9	4.4 ± 2.2	57.8 ± 28.8
SDSSJ234657.25+145736.0	2.1 ± 0.3	6.0 ± 1.0	1.1 ± 0.8	2.7 ± 2.1	8.6 ± 2.4	32.9 ± 8.7


NOTES. Columns are as follows: (1) SDSS name. Cols. (2), (4) and (6) list the equivalent width in units of Å. Cols. (3), (5) and (7) correspond to the flux in units of  $\text{erg s}^{-1} \text{cm}^{-2}$ .

1900Å blend such as AlIII $\lambda$ 1860 or CIII] $\lambda$ 1909 were avoided due to severe blending (Shen & Liu 2012), and the 1900Å blend remains poorly studied to date. Our earlier work motivated us to identify xA source candidates and to obtain high S/N spectra of CIV $\lambda$ 1549 and the 1900Å blend. In the majority of the xA candidates observed so far the AlIII $\lambda$ 1860 profile is well defined and is not strongly affected by absorption features (See Section §5.3).

The left panel of Figure 4 shows a comparison between FWHM CIV $\lambda$ 1549 and FWHM AlIII $\lambda$ 1860 for the xA sample. The ratio between both FWHMs as a function of the FWHM AlIII $\lambda$ 1860 is shown in the bottom part of the panel. Coatman et al. (2017) proposed an empirical correction to the CIV profile to obtain the contribution associated with the virial component. Its factor is based on a measure of the CIV blueshift analogs to the  $c(\frac{1}{2})$  we measure. We then applied a correction to the FWHM CIV $\lambda$ 1549 proportional to  $c(\frac{1}{2})$  CIV $\lambda$ 1549 to correct for the excess broadening due to the blueshifted component. The result is shown in the right panel of Figure 4. All CIV $\lambda$ 1549 measurements have FWHM larger than FWHM AlIII $\lambda$ 1860 before correction. Sources with FWHM AlIII $\lambda$ 1860  $\approx$  3000  $\text{km s}^{-1}$  show a bias factor of  $\approx$  2.5-3 in CIV FWHM, implying almost an order-of-magnitude excess in  $M_{\text{BH}}$  estimates. After correction, even if the bias is almost zeroed, we find no significant improvement over the uncorrected panel. This is especially true when we examine the spectra of the 4-6 sources with largest differences with FWHM AlIII $\lambda$ 1860. CIV, in most of them, is affected by absorptions. Most of the highest confidence measures involve the cluster of sources in the lower left corner of the figure, which can be argued to be the highest confidence xA sources. However the sample is too small to allow us to go further. In a forthcoming paper comparing FWHM H $\beta$  and FWHM AlIII $\lambda$ 1860 measures for our low and higher  $z$  samples (Marziani et al. 2018, in preparation), we find a clear correlation for Pop. A sources. FWHM AlIII $\lambda$ 1860 provides the safest equivalent of FWHM H $\beta$  for virial estimation in Pop. A sources.

### 5.9. Prevalence and properties of absorbed systems

The CIV $\lambda$ 1549 emission line profile of our GTC-xA sources frequently shows absorptions: only in 9 cases the profile is not contaminated by deep broad absorptions and in only 5 cases the profile is not affected by any narrow or broad absorption. In order to ascertain whether our sample suffers from unusually strong line absorption compared to a general quasar population at  $2.0 \lesssim z \lesssim 2.9$ , we considered estimates of the absorption index (AI) and Balnicity index (BI) as defined in Scaringi et al. (2009). The AI (Hall et al. 2002) includes semibroad absorptions continuously detected over a range of at least 1000  $\text{km s}^{-1}$ , from 0 to  $-29000 \text{ km s}^{-1}$ ; the BI (Weymann et al. 1991) measures contiguous absorption features over 2000  $\text{km s}^{-1}$  in the range of  $-25000$  to  $-3000 \text{ km s}^{-1}$ , and is intended to identify quasars with broad through features almost certainly not associated with external absorbers. Table 10 reports the AI and BI measures for our sample. Large CIV $\lambda$ 1549 BI absolute values ( $\lesssim -1000 \text{ km s}^{-1}$ ) are associated with Pop. A and with “strong” BALs with large equivalent widths and high up to terminal velocity, while mini-BALs are also found at low- $z$  among Pop. B and RL objects (Sulentic et al. 2006a). Trump et al. (2006) provide both AI and BI measurements obtained from a SDSS DR3 catalog containing 5719 quasars in redshift range  $z \sim 2 - 2.6$  (Schneider et al. 2005). There are 542 sources with CIV $\lambda$ 1549 BI  $< 0$  suggesting  $\approx 9.5\%$  for “strong” BAL sources, consistent with conventionally quoted prevalence values for BAL QSOs. In the GTC-xA sample, 5 over 19 sources show CIV $\lambda$ 1549 BI  $< 0$ . If we also include one source with CIV BI=0 (SDSSJ144412.37+582636.9), but with strong absorption in the blue side of SiIV $\lambda$ 1397+Orv] $\lambda$ 1402 blend at 1400 Å and BI  $\ll 0$  (See Fig. A.13), we obtain a prevalence of BAL QSOs around 25 – 30%. AI is less than 0 in 10 out of 19 sources of the GTC-xA sample; about 30% of Trump et al. (2006) satisfy the same condition. If one considers that about 1/2 of SDSS quasars are Pop. A quasars (Zamfir et al. 2010), the prevalence of strong BAL sources should be doubled to  $\sim 20\%$ . We can therefore conclude that the GTC-xA sample does not show an unusually large fraction of intrinsically absorbed sys-



**Fig. 4.** Relation between the FWHM  $\text{AlIII}\lambda 1860$  and FWHM  $\text{CIV}\lambda 1549$  before (top) and after (bottom) Coatman et al. (2017) correction. Dashed gray line corresponds to the 1:1 line. The bottom panels show the ratio between the FWHM  $\text{CIV}\lambda 1549$  and  $\text{AlIII}\lambda 1860$ . The colors are the same as the Figure 2 and 3.

**Table 10.** Balnicity and Absorption indices

SDSS Identification (1)	BI (2)	AI (3)
SDSSJ000807.27-103942.7	0	0
SDSSJ004241.95+002213.9	0	0
SDSSJ021606.41+011509.5	-675	-2575
SDSSJ024154.42-004757.5	0	0
SDSSJ084036.16+235524.7	0	-210
SDSSJ101822.96+203558.6	0	0
SDSSJ103527.40+445435.6	0	-160
SDSS105806.16+600826.9	0	0
SDSSJ110022.53+484012.6	0	0
SDSSJ125659.79-033813.8	-720	-970
SDSSJ131132.92+052751.2	-2440	-5430
SDSSJ143525.31+400112.2	0	0
SDSSJ144412.37+582636.9	0	-770
SDSSJ151258.36+352533.2	0	-270
SDSSJ214009.01-064403.9	-2530	-3620
SDSSJ220119.62-083911.6	-5800	-7460
SDSSJ222753.07-092951.7	0	0
SDSSJ233132.83+010620.9	0	0
SDSSJ234657.25+145736.0	0	-240

NOTES. Columns are as follows: (1) SDSS name. (2) Balnicity index in units of  $\text{km s}^{-1}$ . (3) Absorption index in units of  $\text{km s}^{-1}$ .

tems, possibly at a prevalence somewhat higher than the general quasar population.


## 6. Discussion

### 6.1. Equivalent width distributions

Fig. 5 shows a comparison between the equivalent width distributions of  $\text{AlIII}\lambda 1860$ ,  $\text{SiIII}\lambda 1892$ ,  $\text{CIII}\lambda 1909$  and  $\text{CIV}\lambda 1549$  for the GTC-xA sources and the three control samples: the low- $z$  (FOS-A and FOS-xA) and the high- $z$  (S14-A) samples. The  $\text{AlIII}\lambda 1860$  line of the GTC-xA shows an enhancement relative to the other samples (upper left panel). The  $\text{AlIII}$  values distribution for the GTC-xA sources is significantly different more than  $6\sigma$  respect to the ones observed in FOS-A and S14-A, according to a T-test ( $6.7\sigma$  and  $6.1\sigma$  respectively). In fact, a Kolmogorov-Smirnov (KS) test gives a very low probability of being equal population between GTC-xA, and FOS-A and S14-A ( $8 \cdot 10^{-6}$  and  $5 \cdot 10^{-6}$  respectively). Meanwhile, there is no statistically significant difference between the two Pop. A samples (FOS-A and S14-A) or the two Pop. xA (GTC-xA and FOS-xA) samples.

The  $\text{SiIII}\lambda 1892$  equivalent width (upper right panel) covers a wide range of values with no statistical significant difference between the samples. Only at a level of  $2\sigma$ , the GTC-xA shows slightly higher values of  $W_{\text{SiIII}\lambda 1892}$  than FOS-A sample. But neither the T-test nor the KS test yield a significant probability of being different populations.

Instead, in the GTC-xA sample the  $W_{\text{CIII}\lambda 1909}$  distribution (bottom left panel) appears to be markedly weaker than a Pop. A. Both T-test and KS-test give significant differences at  $6.5\sigma$  and  $5.2\sigma$  level for the GTC-xA sample to be a different population respect to FOS-A and S14-A respectively. The probability ac-



**Fig. 5.** Rest-frame equivalent width distributions of the  $\text{Al III } \lambda 1860$ ,  $\text{Si III } \lambda 1892$ ,  $\text{C III } \lambda 1909$  and  $\text{C IV } \lambda 1549$  for the GTC-xA (gray), FOS-xA (blue), FOS-A (not filled) and S14-A (magenta) samples.

According to the KS test is very low ( $3.2 \cdot 10^{-6}$  and  $1 \cdot 10^{-4}$ , respectively). In xA sources  $\text{C III } \lambda 1909$  is weak and blended with the  $\text{Fe II}$  contribution, to the point of not being clearly detectable in several cases. On the other hand, in the Pop. A samples (FOS-A and S14-A) this ion is perfectly appreciable by naked eye and specially prominent in Pop. B sources (e.g., Bachev et al. 2004; Kuraszkiwicz et al. 2004).

Bottom right panel of Fig. 5 shows a comparison between the equivalent width distributions of  $\text{C IV } \lambda 1549$  for xA and Pop. A sources. xA values are the ones with the lowest  $W_{\text{C IV } \lambda 1549}$  among Pop. A sources, with a wide majority of sources distributed below  $\sim 40 \text{\AA}$ . On the converse, the FOS-A and S14-A samples show a much broader distribution, with no evidence of a systematic difference between them, within the limitations of the relatively small sample sizes. The statistical tests give a significant difference between GTC-xA and FOS-A and S14-A at  $5\sigma$  and  $7\sigma$ , and a KS-test a low probability ( $\sim 10^{-5}$ ) to be the same population.


These histograms give a first hint to the fact that xA quasars show a different spectroscopic behavior compared with the one shown by samples of the full Population A. Due to there are not any significant differences in the equivalent width distributions of two xA samples (GTC-xA and FOS-xA) and neither between the two Pop. A samples (FOS-A and S14-A), we can consider a sample with 24 xA sources (19+5) and 34 Pop. A sources (23+11). A comparison of the  $\text{Al III } \lambda 1860$  equivalent width of the 24 xA respect to 34 Pop. A sources shows a significant difference between them at a level of  $7.8\sigma$  with a low probability to be the same distribution ( $7 \cdot 10^{-8}$ ). There is also a statistically

significant difference in the distributions of  $\text{C III } \lambda 1909$  and  $\text{C IV } \lambda 1549$ , but in the opposite direction: values of the xA sources are significantly lower than Pop. A ( $6.6\sigma$  and  $5\sigma$  level respectively with probabilities lower than  $2 \cdot 10^{-7}$ ).

## 6.2. Composite spectra at low and high- $z$

With the objective of highlight main features observed in the xA spectrum and make a comparison with control samples, we built a composite spectrum using the full GTC-xA sample. Multicomponent fits and its analysis are shown in Section §5.1. At the top level of Figure 6 is represented the composite GTC-xA spectrum. It clearly shows an enhancement of  $\text{Al III}$  and  $\text{Si III}$ , in contrast with the weakness of  $\text{C III}$ , which has mean  $W_{\text{C III}} \sim 4 \text{\AA}$ . Also,  $\text{Fe II}$  becomes the largest contribution in the red side of the  $1900 \text{\AA}$  blend, which is also prominent in the  $\text{Fe II}$  feature detected at  $\sim 2080 \text{\AA}$ . Another noticeable feature is the almost fully blueshifted  $\text{C IV } \lambda 1549$  profile, which shows a low equivalent width. A blueshift is also observed in  $\text{Si IV } \lambda 1397 + \text{O IV } \lambda 1402$ , although in this case the profile is more symmetric due to the presence of strong absorptions in its blue side. As we point out in Section §5.4,  $\text{He II } \lambda 1640$  is very weak with a flat topped appearance profile. Low-ionization features,  $\text{C II } \lambda 1335$  and  $\text{O I } \lambda 1304 + \text{Si II } \lambda 1306$ , are also clearly appreciated.

In order to compare the general behavior of xA sources, we also built composite spectra for two comparison samples: FOS-xA and S14-A samples, middle and bottom level of Figure 6, respectively. A multicomponent analysis was also applied to composite control samples. The width line of  $\text{Al III } \lambda 1860$  in FOS-



**Fig. 6.** From top to bottom, rest-frame composite spectra of the GTC-xA, FOS-xA, and S14-A. Intensity units are arbitrary flux. The GTC-xA and FOS-xA have been displaced by a constant value for clarity.

xA ( $\sim 2740 \text{ km s}^{-1}$ ) is narrower than GTC-xA ( $\sim 4500 \text{ km s}^{-1}$ ). The average bolometric luminosity of the FOS-xA sample is  $\sim 2.4 \cdot 10^{45} \text{ erg s}^{-1}$  and  $\sim 10^{47} \text{ erg s}^{-1}$  for the GTC-xA, therefore a FWHM change can be associated with the higher luminosity in the GTC-xA sample.

The S14-A composite spectrum shows a comparable luminosity ( $\log L \approx 46$ ) than the FOS-xA, and a similar redshift that GTC-xA sample. S14-A shows a strong and symmetric C IV  $\lambda 1549$  profile with a median equivalent width of  $47 \text{ \AA}$ , more than two times large than the median values obtained in xA samples:  $14$  and  $23 \text{ \AA}$  for GTC-xA and FOS-xA, respectively. Conversely, the  $1900 \text{ \AA}$  blend of S14-A composite spectrum is dominated by C III] and Si III] profiles. The xA samples show a weaker C III], with median values of equivalent width  $2.6 \text{ \AA}$  for GTC-xA,  $1.8 \text{ \AA}$  for FOS-xA and  $11 \text{ \AA}$  for S14-A. Also they show a stronger Al III  $\lambda 1860$  than Pop. A samples, with the next median values:  $7 \text{ \AA}$  for the GTC-xA,  $6 \text{ \AA}$  for FOS-xA and  $2 \text{ \AA}$  for S14-A. Other notable features of xA sources include a smaller He II  $\lambda 1640$ , as well as a higher (O IV]  $\lambda 1402 +$  Si IV  $\lambda 1397)/\text{C IV} \lambda 1549_{\text{BC}}$  ratio. Features like O I  $\lambda 1304 +$  Si II  $\lambda 1306$  and C II  $\lambda 1335$  are also clearly observed, supporting the inference of a low-ionization degree for xA objects (See Section §6.3). There is a fundamental caveat between GTC-xA and S14-A samples: among the faint sources of S14-A there is no xA sources. A systematic difference in  $L/L_{\text{Edd}}$  of  $\approx 2$  can be accounted, if the Eddington ratio of GTC-xA sources is  $L/L_{\text{Edd}} \approx 1$  and if the S14-A are instead radiating at  $L/L_{\text{Edd}} \approx 0.3$  (Sulentic et al. 2014a), without invoking any effect of luminosity.

### 6.3. A low-ionization spectrum with prominent Fe III

The UV spectral range provides several diagnostic ratios that can constrain physical parameters. For instance the C III]  $\lambda 1909/\text{Si III] } \lambda 1892$  and Al III  $\lambda 1860/\text{Si III] } \lambda 1892$  ratios are sensitive to hydrogen density, C IV  $\lambda 1549/\text{Al III} \lambda 1860$  and C IV  $\lambda 1549/\text{Si III] } \lambda 1892$  to ionization, and He II  $\lambda 1640/\text{C IV} \lambda 1549$


and (O IV]  $\lambda 1402 + \text{Si IV} \lambda 1397)/\text{C IV} \lambda 1549$  to metallicity (N12, Negrete et al. 2013, 2014). The diagram of Fig. 7 has been computed from an array of CLOUDY 13.05 (Ferland et al. 2013) simulations covering the range in ionization parameter  $-4.5 \leq \log U \leq 0$  and density  $7 \leq n_{\text{H}} \leq 14 \text{ cm}^{-3}$ , with steps of  $0.25 \text{ dex}$ . A total of 551 simulations were carried out assuming a standard AGN continuum appropriate for Pop. A, a column density  $N_{\text{c}} = 10^{23} \text{ cm}^{-2}$  and a metallicity of  $100Z_{\odot}$ . The median flux ratios given by Al III  $\lambda 1860/\text{Si III] } \lambda 1892$ , C IV  $\lambda 1549/\text{Al III} \lambda 1860$ , and C IV  $\lambda 1549/\text{Si III] } \lambda 1892$  converge toward  $\log U \approx -3$  and  $\log n_{\text{H}} \approx 13 [\text{cm}^{-3}]$ . The suggestion of low-ionization and high density is in agreement with the previous study of two xA sources by N12. Consistently with the inference of high density, C III]  $\lambda 1909$  is usually weak although not entirely negligible with a  $\log \text{C III] } \lambda 1909/\text{Si III] } \lambda 1892 \approx -0.42$ . Due to the critical electronic density of the C III]  $\lambda 1909$  is  $n_{\text{H}} = 10^{10} \text{ cm}^{-3}$ , this line is expected to be collisionally suppressed with respect to Al III  $\lambda 1860$  and Si III]  $\lambda 1892$ , as it is shown by the GTC-xA composite spectrum. The strong Fe III contribution observed in the GTC-xA sources also supports the existence of a very low-ionization spectrum,  $\log U \leq -2$ . A more detailed analysis of Fe III emission is presented in Appendix C.

The intensity ratios considered above are not strongly sensitive to metallicity except for the He II  $\lambda 1640/\text{C IV} \lambda 1549$  and the (O IV]  $\lambda 1402 + \text{Si IV} \lambda 1397)/\text{C IV} \lambda 1549$  ratios. According to CLOUDY simulations, at solar metallicity we expect C IV  $\lambda 1549/\text{He II} \lambda 1640 \approx 1$  for the low- $U$  and high density solution, which is very far from the value that we derive for the xA sample ( $\sim 0.19$ ). A possible solution is offered by a large increase in metallicity above solar values. The He II  $\lambda 1640/\text{C IV} \lambda 1549$  is the ratio between a recombination and a collisionally excited line: at high- $Z$ , the metals absorb a larger fraction of the ionizing flux, and the He recombination line become weaker. This effect leads to a decrease of the He II  $\lambda 1640/\text{C IV} \lambda 1549$  over the value expected at solar  $Z$  (Hamann & Ferland 1993, 1999). In this case, the  $Z$ -sensitive He II  $\lambda 1640/\text{C IV} \lambda 1549$  would be small ( $\sim 0.2$ ) even at low ionization. An increment in the metallicity to  $Z \approx 100Z_{\odot}$  gives such low He II  $\lambda 1640/\text{C IV} \lambda 1549$  ratios (Fig. 7). The value  $Z \approx 100Z_{\odot}$  may appear unrealistically high. However, N12 found evidence of high- $Z$ , as well as of possible abundance anomalies. It appears reasonable to assume very high values of  $Z$  for xA sources: Pop. A sources at low- $z$  and high- $z$  have N V  $\lambda 1240/\text{C IV} \lambda 1549$  and (O IV]  $\lambda 1402 + \text{Si IV} \lambda 1397)/\text{C IV} \lambda 1549$  implying  $Z \gtrsim 10Z_{\odot}$  (Shin et al. 2013, S14). The median ratio Si IV  $\lambda 1397/\text{C IV} \lambda 1549 \sim 1$  may suggest a metallicity significantly higher than 10 times solar for the GTC-xA sample. The high Si IV  $\lambda 1397 + \text{O IV] } \lambda 1402/\text{C IV} \lambda 1549$  ratio also supports this suggestion. In agreement with the interpretation of Hamann & Ferland (1999), the equivalent width of He II  $\lambda 1640$  is extremely low, making the line almost undetectable above noise.

We should also consider that our approach may overestimate the C IV<sub>BC</sub> intensity, as it attempts to maximize its contribution attempting to account for the red side of the profile, even when the profile is almost fully blue shifted. There is the possibility that the C IV<sub>BC</sub> is significantly weaker. A slightly lower value of  $Z = 50Z_{\odot}$  is again in agreement with the observations assuming an overestimation of the C IV<sub>BC</sub> intensity by factor 2. Therefore, a  $Z$  value of several tens  $Z_{\odot}$  might be considered as a lower limit if we consider a systematic overestimation of the C IV<sub>BC</sub><sup>3</sup>. More

<sup>3</sup> It is unlikely that the overestimate might be much larger than a factor 2, as the C IV<sub>BC</sub> is clearly detectable while He II  $\lambda 1640$  equivalent width is extremely low.

precise inferences require dedicated photoionization computations beyond the scope of this paper.



**Fig. 7.** Isocontours in the plane  $\log$  of ionization parameter ( $U$ ) versus  $\log$  of hydrogen density ( $n_{\text{H}}$ ) from CLOUDY simulations for the line intensity ratios  $\text{AlIII}\lambda 1860/\text{SiIII}\lambda 1892$  (blue),  $\text{HeII}\lambda 1640/\text{CIV}\lambda 1549$  (green),  $\text{CIV}\lambda 1549/\text{AlIII}\lambda 1860$  (yellow),  $\text{CIV}\lambda 1549/\text{SiIII}\lambda 1892$  (red) using  $Z = 100Z_{\odot}$ . Labels report the median values of the intensity ratios for the objects of the GTC-xA sample.

#### 6.4. Evidence of 2080 Å enhancement

Several sources show a significant excess with respect to the template at around  $\sim 2080$  Å, indicating that UV #48 is selectively enhanced with respect to the “normal” emission in xA sources as represented by Vestergaard & Wilkes (2001) template. In eight sources the equivalent width of the feature in excess respect to the template is  $W(2080) \geq 5$  Å. This excess is evident from the spectra, and is even more outstanding because the HILs in the adjacent spectral regions are of low equivalent width. A mock profile built from the transition of multiplet  $\text{FeII}$  UV #48 between the terms  $a^5S - z^5P^o$ , assuming relative intensity for optically thin transitions (Kovačević et al. 2010) accounts for the 2080 Å feature: the mock profile resulting from the lines broadened to  $4000 \text{ km s}^{-1}$  is a feature at  $\approx 2073$  Å with total FWHM  $\approx 5000$  Å, consistent with the observations.


Figure 8 shows the equivalent width of the  $\text{FeIII}\lambda 1914$  and of the  $\text{FeII}$  UV #191 multiplet versus the equivalent width of the 2080 Å feature. Only 10 sources satisfy the detection limit we imposed at  $W \approx 1$  Å. Strong  $\text{FeIII}\lambda 2080$  is usually associated with strong  $\text{FeIII}\lambda 1914$ . According to an orthogonal fit taking into account errors in both axis, these two emission lines follow the relation  $\text{FeIII}\lambda 1914 = (0.46 \pm 0.14) \cdot \text{FeIII}\lambda 2080 + (1.29 \pm 0.55)$ , which verified the null hypothesis of a slope different from 0 at  $3.3\sigma$  significant level. There is a clear relation, although the errors associated with  $\text{FeIII}\lambda 1914$  are high because of the blend with  $\text{CIII}\lambda 1909$  and the contribution of the  $\text{FeII}$  template. On the another hand, the  $\text{FeII}_{\text{UV}}$  #191 is generally rather weak, but the 5 cases with a clear detection ( $W \geq 2$  Å) are associated with strong 2080 Å features. An orthogonal fit gives a relation  $\text{FeII}_{\text{UV}} = (0.27 \pm 0.06) \cdot \text{FeIII}\lambda 2080 + (0.57 \pm 0.30)$  with a slope different from 0 at a  $4\sigma$  significance level. This relation consid-

**Table 11.** Accretion parameters

SDSS Identification (1)	$\log L_{\text{Bol}}$ (2)	$\log M_{\text{BH, AIII}}$ (3)	$L/L_{\text{Edd}}$ (4)
SDSSJ000807.27-103942.7	46.9	9.29	0.29
SDSSJ004241.95+002213.9	47.0	9.24	0.34
SDSSJ021606.41+011509.5	47.5	9.13	1.54
SDSSJ024154.42-004757.5	46.9	8.88	0.77
SDSSJ084036.16+235524.7	46.9	8.88	0.70
SDSSJ101822.96+203558.6	47.1	9.28	0.42
SDSSJ103527.40+445435.6	47.1	9.63	0.21
SDSS105806.16+600826.9	47.2	9.21	0.65
SDSSJ110022.53+484012.6	47.0	9.22	0.37
SDSSJ125659.79-033813.8	47.0	9.41	0.26
SDSSJ131132.92+052751.2	47.3	9.19	0.93
SDSSJ143525.31+400112.2	47.4	9.14	1.31
SDSSJ144412.37+582636.9	46.7	9.49	0.10
SDSSJ151258.36+352533.2	47.1	8.80	1.31
SDSSJ214009.01-064403.9	47.2	8.79	1.61
SDSSJ220119.62-083911.6	47.3	9.72	0.23
SDSSJ222753.07-092951.7	46.9	9.09	0.39
SDSSJ233132.83+010620.9	47.7	9.84	0.50
SDSSJ234657.25+145736.0	47.1	9.02	0.73

NOTES. Columns: (1) SDSS identification. (2) Log of Bolometric luminosity in unit of  $\text{erg s}^{-1}$ . The associated uncertainty is 10% the luminosity value. (3) Black hole mass in unit of  $M_{\odot}$ . (4) Eddington ratio.

ers measurements with  $W > 1.0$  Å of both  $\text{FeII}$  UV #191 and  $\text{FeIII}\lambda 1914$  (Figure 8). A fluorescence mechanism (described in Appendix C.2) may account for the selective enhancement of the 2080 Å feature.



**Fig. 8.** Relation between intensity of the  $\text{FeIII}$  feature at 2080 Å with  $\text{FeII}$  UV 1785 Å (blue) and  $\text{FeIII}\lambda 1914$  (red). Squares, up and down triangles, and circles mark the normal, BAL, mini-BAL and RL quasars, respectively. The gray shaded area indicates an approximate detection limit. Blue and red lines mark the orthogonal fit for each case.

#### 6.5. Estimates of the accretion parameters

##### 6.5.1. Bolometric correction

There has been no accurate determination of the bolometric correction (hereafter B.C.) specific to xA sources. An attempt has been made to compute the bolometric correction from the model SEDs of MS14. MS14 suggested that the difference in the optical/UV/X range between a typical NLSy1 and an extreme

NLSy1 is about 10% (see their section 6.1.2.3). It is unlikely that the uncertainty in the B.C. is much larger than  $\approx 20\%$  (at  $1\sigma$  confidence level). SDSS quasars have with  $B.C._{1350} = 3.5$  and a scatter  $\sim 20\%$  (Richards et al. 2006). SDSS quasars are however an heterogeneous sample composed by quasar of different spectral types. On the contrary, the GTC-xA sample is composed by similar sources. If we consider 27 spectra of sample 3 of MS14 i.e., at redshift in the range 2.0 – 2.6, the rms averaged between  $\approx 1100$  and  $2900 \text{ \AA}$  is  $\approx 10\%$ .

Considering that the bulk of the ionizing energy is coming from the big blue bump in correspondence of the Lyman limit, the SED is not significantly different from the one of Mathews & Ferland (1987). The MS14 SED differs from the reference Mathews & Ferland (1987) SED only for the slope in the X-ray domain, and yields a  $B.C._{1350} \approx 5.4$ . The B.C. value is consistent with the one of Elvis et al. (1994). The value of Elvis et al. (1994) may perhaps provide a more appropriate comparison, as the sources in that paper implied a detection from the IPC instrument on board the Einstein observatory (Giacconi et al. 1979), i.e., in the soft X-ray domain. Bolometric luminosities  $L_{\text{Bol}}$  of the GTC-xA sources are presented in Table 11. They have been computed through the monochromatic luminosities at  $1350 \text{ \AA}$  ( $\lambda L_{\lambda}(1350\text{\AA})$ ) and a  $B.C.=5.4 \pm 0.6$ . A more careful estimate would require a detailed analysis of the mid- and far-IR domain, where part of the accretion luminosity is reradiated, and is deferred to an eventual work.

### 6.5.2. Black hole mass

We estimate black hole mass ( $M_{\text{BH}}$ ) using the UV  $\text{Civ}$  scaling relation of Vestergaard & Peterson (2006)[Eq. 7] applied to  $\text{FWHM}(\text{AlIII})^4$ . The  $\text{AlIII}$  profile is usually not affected by the strong outflows often observed in the  $\text{Civ}\lambda 1549$  profile, thus providing a good approximation to the virial motion of the BLR (e.g., Marziani et al. 2017b, see also Section §5.8). We adopt  $M_{\text{BH,AlIII}}$  as the virial mass estimator. Second column of Table 11 show the  $M_{\text{BH}}$  estimations. Typical formal uncertainties in the estimation of the masses, through error propagation, are about 25% taking into account an uncertainty of 10% in the flux at  $1350\text{\AA}$ , and including the estimated errors for the  $\text{FWHM AlIII}\lambda 1860$ .

The GTC-xA sample covers a black hole mass range of  $8.8 \leq M_{\text{BH}} \leq 9.8 M_{\odot}$ , with a median value of  $\langle M_{\text{BH}} \rangle \sim 9.2 M_{\odot}$ . The GTC-xA sample is covering the high- $M_{\text{BH}}$  counterparts of the FOS-xA sample. Also these values confirm the well-known tendency of larger black hole mass ( $\sim 10^9 M_{\odot}$ ) at high redshift ( $z \approx 2$ ) (Shemmer et al. 2004; Netzer et al. 2007; Trakhtenbrot & Netzer 2012), a redshift-dependent selection effect common to all flux-limited samples (Sulentic et al. 2014a).

### 6.5.3. Eddington ratio

Eddington ratios ( $L/L_{\text{Edd}}$ ) have been calculated based on masses obtained from the  $\text{FWHM AlIII}$ , explained in previous section and following the relation:  $L_{\text{Edd}} \approx 1.5 \times 10^{38} \cdot (M_{\text{BH}}/M_{\odot}) [\text{erg s}^{-1}]$ , as recommended by Netzer & Marziani (2010).  $L/L_{\text{Edd}}$  values for our sample are reported in Table 11. Applying the usual method of propagation of errors and using previous formulae, the typical mean uncertainty in the Eddington ratio is about 30-

35% (only in one case the estimated error is larger than the 50%, in SDSSJ233132.83+010620.9  $L/L_{\text{Edd}} \approx 0.5 \pm 0.3$ ).

The average  $L/L_{\text{Edd}}$  value for the GTC-xA sample is  $\langle L/L_{\text{Edd}} \rangle \approx 0.66$ , comparable to the one obtained for I Zw 1, a prototype highly accreting quasar at low- $z$ . An appropriate sample for a comparison is also the sample #3 of MS14 that involves xA quasars in the redshift range  $2 \leq z \leq 2.6$  with the  $\text{AlIII}\lambda 1860$  used as a virial broadening estimator. The obtained average  $L/L_{\text{Edd}}$  of the GTC-xA sample is consistent with the value obtained for sample #3 of MS14 ( $L/L_{\text{Edd}} \approx 0.7$ ), once the estimates are computed with the same bolometric correction (5.4 in this work versus 6.3 in MS14), and the same  $L/L_{\text{Edd}}$  constant (MS14 used  $1.3 \cdot 10^{38} \text{ erg s}^{-1}$ ). A Welch-test on the average  $L/L_{\text{Edd}}$  confirms that two averages are not significantly different, and therefore the xA nature of the sources in the present sample.


The scatter in  $L/L_{\text{Edd}}$  in the GTC-xA sample after extinction correction is  $\sigma_{L/L_{\text{Edd}}} \sim 0.47$  which is somewhat above the one of sample-#3 of MS14 ( $\sigma_{L/L_{\text{Edd}}} \approx 0.16$ ), although this may be in part due to the small size of the GTC-xA, with only 19 objects. Also the larger scatter in FWHM may contribute to the  $L/L_{\text{Edd}}$  scatter: average  $\text{FWHM AlIII} \sim 4220 \pm 900 \text{ km s}^{-1}$  for MS14 versus an average  $\text{FWHM AlIII} \sim 4720 \pm 1600 \text{ km s}^{-1}$  for the GTC-xA sample. As an example, SDSSJ144412.37+582636.9 has a  $\text{FWHM AlIII} \sim 8080 \text{ km s}^{-1}$  which yields the minimum  $L/L_{\text{Edd}}$  value (0.1) in the sample. Additionally this source is a mini-BAL with clear absorptions in its spectrum (See Fig. A.13) introducing additional uncertainties in the extinction correction. In the absence of an orientation correction, a large scatter in the observed FWHM translates into a large scatter in  $L/L_{\text{Edd}}$ .

A correlation between the  $c(\frac{1}{2})$  shift and  $L/L_{\text{Edd}}$  (Sulentic et al. 2017) implies that highest  $c(\frac{1}{2})$  values are associated with the highest  $L/L_{\text{Edd}}$ . Figure 9 shows the plane  $c(\frac{1}{2})$  vs.  $L/L_{\text{Edd}}$ , where GTC-xA sources occupy a fairly extreme position, where also I Zw 1 is located. Two-third of the 11 sample (unabsorbed sources) have  $c(\frac{1}{2}) \lesssim -2000 \text{ km s}^{-1}$ , and the blueshifts distribution is more skewed toward high  $c(\frac{1}{2})$  values than in the sample of S17. Comparing with the HE sample (Sulentic et al. 2017) and FOS-xA and FOS-A sample (Sulentic et al. 2007), we observe that FOS-xA sources (the pale blue squares around  $L/L_{\text{Edd}} \sim 1$ ) also cover a similar domain of blueshift amplitudes, while Pop. A sources show lower amplitude shifts,  $-900 \text{ km s}^{-1}$  on average. This diagram provides further support to the idea that the Eddington ratio is the physical driver of the outflows, and that xA sources are associated with high Eddington ratio values.


### 6.6. Weak-lined and xA quasars

The Weak-Line Quasars (WLQs) show very low equivalent width in the  $\text{Civ}\lambda 1549$  and other UV emission lines (Diamond-Stanic et al. 2009; Wu et al. 2012; Shemmer et al. 2010). A standard definition of WLQ involves sources with  $W(\text{Civ}) \leq 10 \text{ \AA}$ . WLQs also show strong FeII emission and Civ blueshift. Luo et al. (2015) provides one of the largest samples of WLQ candidates, with detailed measures on shifts and  $W$  of Civ for 21 of them. It is an almost exclusively RQ sample as are 4DE1 Population A, and especially xA quasars. Most WLQ candidates are high- $z$  making estimates of source rest-frame uncertain in many cases. When  $\text{H}\beta$  measures are available (Plotkin et al. 2015; Sulentic et al. 2007, 2017), we confirm that WLQs belong to the Population A domain of 4DE1 with  $R_{\text{FeII}} \gtrsim 1.0$ , independently of  $z$  and  $L$  (Fig. 9 of Marziani et al. 2016). At least 80% of WLQs are xA sources and, if doubtful cases (e.g., affected by absorptions, so most likely not real WLQs) are excluded all WLQs

<sup>4</sup>  $\log M_{\text{BH}}(\text{AlIII}) = \log \left\{ \left[ \frac{\text{FWHM}(\text{AlIII})}{1000 \text{ km s}^{-1}} \right]^2 \left[ \frac{\lambda L_{\lambda}(1350\text{\AA})}{10^{44} \text{ erg s}^{-1}} \right]^{0.53} \right\} + (6.73 \pm 0.01)$ .



**Fig. 9.** Behavior of the centroid at 1/2 of total  $\text{CIV}\lambda 1549$  profile as a function of Eddington ratio. Hexagons represent the GTC-xA sample. The colors of hexagons are the same as Figures 2 and 3. Blue and red pale symbols correspond to the Population A and B, respectively. Squares and stars correspond to the FOS-A and xA samples (Sulentic et al. 2007). Circles are from HE sample Sulentic et al. (2017) sample. Yellow star marks the position of I Zw 1. The black dashed horizontal line indicates  $c(\frac{1}{2})=0 \text{ km s}^{-1}$ . Gray and black dashed vertical lines indicate  $L/L_{\text{Edd}}=0.2$  and 1.0, respectively.



**Fig. 10.** Relation between rest-frame equivalent width of  $\text{CIV}\lambda 1549$  and  $c(\frac{1}{2})$  in  $\text{km s}^{-1}$ . Black squares: GTC-xA. Gray squares: BAL GTC-xA. Blue triangles: Plotkin et al. (2015), magenta: Wu et al. (2012) and cyan: Luo et al. (2015). Red and green circles correspond to the Pop. A samples from S17 and FOS-A sample, respectively. Circled symbols correspond to FOS-xA and S17-xA.

might belong to the xA population. The majority of WLQ show

$\text{FWHM H}\beta \gtrsim 2000 \text{ km s}^{-1}$ . WLQ are simply Pop. A sources with strong  $R_{\text{FeII}}$ , high amplitude  $\text{Crv}$  blueshift and the lowest values for  $W \text{Civ}$ .

According to the standard WLQ definition our GTC-xA sample includes three WLQs, excluding BALs. If we consider the prevalence of WLQs among xAs in different samples, we found: 4 out of 42 in MS14; 1 out of 3 (S17); 3 out of 13 in the present sample (excluding BAL and mini-BAL QSOs), and 2 out of 10 (S07), for a total  $f_{\text{WLQ}} = n_{\text{WLQ}}/n_{\text{xA}} = 10/68 \approx 15\%$ .

As we see in Section §6.1, the peak of the GTC-xA  $W \text{Civ}\lambda 1549$  distribution is slightly above  $10 \text{ \AA}$  (See also bottom right panel of Fig. 5). There is no solution of continuity at  $W \text{Civ}\lambda 1549 \approx 10 \text{ \AA}$ . WLQs are located at the low end of the  $W \text{Civ}\lambda 1549$  distribution of the GTC-xA sources and there is no suggestion that they constitute a special class. Moreover, as in the case xA sources, WLQs are characterized by extreme outflow velocities. Figure 10 shows the  $W \text{Civ}\lambda 1549$  vs.  $\text{Civ}\lambda 1549$  shift domain for the 19 GTC-xA sources of our sample and several WLQs, Pop. xA and Pop. A samples: 19 WLQs from Luo et al. (2015), 6 WLQs from Plotkin et al. (2015), 4 WLQ from Wu et al. (2012, which includes PHL 1811 from Wu et al. 2011), 3 Pop. xA and 11 Pop. A from Sulentic et al. (2017), and 5 xA and 23 Pop. A from S07. The measurements in our samples are the centroid at half maximum, while Luo et al. (2015) consider interactive measurement, presumably of the peak shift. In the context of the present analysis, we expect that the effect may not be relevant because, in the case of the largest blueshift, the total profile is shifted so that peak and centroid at half maximum yield consistent values.

The distributions of xA and WLQ blueshifts are largely overlapping, with an excess of WLQs showing blueshifts in the range  $5000\text{--}10000 \text{ km s}^{-1}$ . A trend between  $W \text{Civ}\lambda 1549$  and  $\text{Civ}\lambda 1549$  shift amplitude is suggested, in the sense that the blueshift amplitude appears to increase among weaker  $\text{Civ}\lambda 1549$  emitters. It is important to consider that this is part of a general trend among quasars, with a much higher statistical significance if Pop. A and B at high luminosity are included (Marziani et al. 2016).

Restricting attention to xA and WLQs means to consider only the extreme end of the equivalent width-shift distribution. Both xAs and WLQs show a coupling between physical and dynamical conditions of the outflowing gas which seems a well-defined, common phenomenon affecting outflow-dominated sources (S17). WLQs can therefore interpreted as extreme xA sources in the context of 4DE1. The previous discussion also validates the assumption that at least the wide majority (if not all) of RQ WLQs are xA quasars. In addition, it suggests that WLQs are an ill-defined class, and that their observational properties should be analyzed within the context of the wider class of xA sources.

At the present time we suggest that observational biases may be playing a serious role. If one considers the data for the most extreme WLQs, extremely low  $W$  and extremely high  $\text{Crv}$  blueshifts, we find that some of the lowest  $W$  values in Fig. 10 involve sources with  $\text{Civ}$  absorptions, that reduce  $W \text{Civ}$ , and UV redshift uncertainties, that affect estimates of  $\text{Civ}$  shifts (Luo et al. 2015). Only a few WLQ candidates have data available for the  $\text{H}\beta$  region (Plotkin et al. 2015). Two of these sources report doubtful detections of an extremely broad and very low  $W \text{Civ}$  (SDSSJ094533.98+100950.1 and SDSSJ141730.92+073320.7). Sources with  $c(\frac{1}{2}) < -5000 \text{ km s}^{-1}$  and/or  $W \text{Civ} < 3\text{--}4 \text{ \AA}$  require high S/N confirmatory spectra. Even if confirmed, we see no evidence for a break in the distribution at  $W \text{Civ} = 10 \text{ \AA}$  suggesting

that WLQ are the most extreme Pop. A quasars (xA), with a lowest values for  $W_{\text{Civ}}$ .

### 6.7. Outflows and feedback

Hereafter we will consider only median luminosity estimates after extinction correction excluding BAL and mini-BAL sources. The median value of the blueshifted component (excluding the BAL cases) is  $\log L_{\text{BLUE}} \approx 44.0 \pm 0.3$  [erg s<sup>-1</sup>]. It is relatively straightforward to compute the mass of ionized gas needed to sustain the C<sub>IV</sub>λ1549 blue shifted component luminosity:  $\dot{M}^{\text{ion}} \sim 95 Z_{10}^{-1} n_9^{-1} M_{\odot}$ , where the density is assumed  $10^9$  cm<sup>-3</sup>, and the relation is normalized to a metal content ten times solar (e.g., Cano-Díaz et al. 2012; Marziani et al. 2016, 2017a). Computation of the mass outflow rate, of its thrust, and of its kinetic power requires several assumptions (discussed in the recent review of Marziani et al. 2017a). In the present context, we assume a BLR radius estimated from the radius luminosity relation derived by Kaspi et al. (2007). At the average continuum luminosity after extinction correction  $\lambda L_{\lambda}(1350) \approx 2.4 \cdot 10^{46}$  erg s<sup>-1</sup>, the  $r_{\text{BLR}} \approx 2.6 \cdot 10^{17}$  cm  $\approx 0.086$  pc. The mass outflow rate can then be written as:  $\dot{M}^{\text{ion}} \sim 150 L_{45} v_{0,5000} r_{\text{BLR},0.1\text{pc}}^{-1} Z_{10}^{-1} n_9^{-1} M_{\odot}$  [yr<sup>-1</sup>], where the outflow radial velocity is in units of 5000 km s<sup>-1</sup>. For the median centroid displacement of the blue shifted component  $\approx -3560$  km s<sup>-1</sup>, we obtain a  $\dot{M}^{\text{ion}} \approx 12.5 M_{\odot}$  yr<sup>-1</sup>. The final thrust can be expressed as:  $\dot{M}^{\text{ion}} k v_0 \sim 5 \cdot 10^{36} L_{45} k v_{0,5000}^2 r_{\text{BLR},0.1\text{pc}}^{-1} Z_{10}^{-1} n_9^{-1}$  [g cm s<sup>-2</sup>], where we have now introduced a factor  $k$  that takes into account that the gas within the BLR is still being accelerated by radiation forces. In the scenario of Netzer & Marziani (2010), for Eddington ratio  $\rightarrow 1$ , as it is likely the case of xA sources,  $k$  can be as high as 10. We prudentially assume  $k = 5$ , and that the full blue shifted component is accelerated to  $k v_0$ . For the average values of the xA sources we obtain  $\dot{M}^{\text{ion}} k v_0 \sim 1.5 \cdot 10^{36}$  g cm s<sup>-2</sup>. The kinetic power of the outflow is  $\dot{\epsilon} \sim 1.2 \cdot 10^{45} L_{45} k^2 v_{0,5000}^3 r_{\text{BLR},0.1\text{pc}}^{-1} Z_{10}^{-1} n_9^{-1}$  erg s<sup>-1</sup>, and becomes  $\dot{\epsilon} \sim 1.3 \cdot 10^{45}$  erg s<sup>-1</sup> for the average parameter values of the xA sample.

The large outflow radial velocity (the average  $c(\frac{1}{2})$  of the blue shifted component is close to  $\sim -4000$  km s<sup>-1</sup>), as well as the high fraction of the blue shifted component to the total C<sub>IV</sub>λ1549 luminosity ( $\sim 0.45$ , Sect. §5.4) ensure the maximization of the quasar mechanical feedback even if  $W_{\text{Civ}\lambda 1549}$  is in general relatively low (c.f., King & Muldrew 2016). The ratio  $\dot{\epsilon}/L \sim 0.01$  comes close to the threshold value for exerting significant mechanical feedback on the entire host galaxies especially if the quasar is radiating at high Eddington ratio and at high luminosity (e.g., King & Pounds 2015, and references therein). High luminosity outflows may be even more powerful if one considers that the C<sub>IV</sub>λ1549 emission traces only a phase of multiphase outflowing medium (Tombesi et al. 2010; Harrison et al. 2014; Ciccone et al. 2014; Carniani et al. 2015; Feruglio et al. 2015). On the other hand, if  $Z$  is as high as  $100 Z_{\odot}$ , then all estimates should be reduced by a factor of 10. This emphasizes the need of accurate metallicity estimates not only to study the enrichment of the ISM by nuclear outflows, but also to ascertain their dynamical and evolutionary relevance on the host galaxy.

## 7. Conclusion

We present a sample of 19 quasar satisfying the 4DE1 selection criteria for finding high redshift highly accretors xA sources. We provide a description of the rest-frame UV spectra of our candidate sources.

- Our GTC-xA sample shows the same characteristics of xA quasars at low- $z$ , of which a prototype is I Zw 1. Figure 2 shows that GTC-xA sample occupies a very restricted Al<sub>III</sub>λ1860/Si<sub>III</sub>λ1892 vs. C<sub>III</sub>λ1909/Si<sub>III</sub>λ1892 domain space, compared to more general quasar populations. GTC-xA sources show different spectroscopic properties than other Pop. A (Bins A1, A2) quasars. We show that our sample is likely radiating close to the Eddington limit ( $L/L_{\text{Edd}} \geq 0.2$ ) with an average consistent with the one computed on the MS14 xA sample.
- Al<sub>III</sub>λ1860 and Si<sub>III</sub>λ1892 lines are well-modeled with a symmetric profile that is narrower than that of C<sub>IV</sub>λ1549 total profile. These results support our previous suggestions that Al<sub>III</sub>λ1860 is likely a reliable virial estimator in analogue to Hβ for lower  $z$  quasars. On the converse, the FWHM C<sub>IV</sub> remains unreliable as a virial estimator showing no significant correlation with FWHM Al<sub>III</sub>λ1860.
- C<sub>IV</sub>λ1549 profiles in GTC-xA sources show large blueshifts,  $|c(\frac{1}{2})| = 1000\text{--}5000$  km s<sup>-1</sup>. Figure 3 shows evidence for a correlation between FWHM and  $c(\frac{1}{2})$  for C<sub>IV</sub>λ1549. The uniformity of the source distribution along the full blueshift range can be used to argue for a quasi isotropic wind in xA quasars. Another possibility sees a slight source excess around  $c(\frac{1}{2}) \sim -2000$  km s<sup>-1</sup>. Only a larger sample can permit a more robust speculation.
- The emission line spectrum of the GTC-xA sources is characterized by low equivalent width of the most prominent emission lines. WLQs appear as extreme xA sources and not as an independent class. All WLQ might be xA sources, if we consider observations with the Hβ spectral range covered.
- The appearance of the xA spectrum at high- $L$  is of very low ionization. Fe<sub>I</sub> and Fe<sub>II</sub> contributions appear to be more prominent in the spectra, although this is in part due to the low equivalent width of all the emission features. The Fe<sub>II</sub> strength is only in apparent contradiction with very low values of the ionization parameter: Fe<sup>+2</sup> becomes the dominant ionization stage of iron at the expense of Fe<sup>+3</sup>, therefore increasing Fe<sup>+2</sup> with respect to Fe<sup>+1</sup>.

Numerous results merit discussion and further study. Among them, we list two that are relevant to our global understanding of the black hole-host galaxy connection.

- The C<sub>IV</sub> shift and the luminosity suggest high values of mass outflow rate, thrust, and kinetic power of the outflow. This is expected for sources of high- $L$  and maximized radiation forces per unit mass. However, more refined computations are needed to assess the relevance of the outflow dynamical parameters for feedback effects on the host galaxy.
- The intensity ratio He<sub>I</sub>λ1640/C<sub>IV</sub>λ1549 requires high  $Z$  ( $\geq 50 Z_{\odot}$ ) to be consistent with a low-ionization, high density ( $\log U \sim -3$ ,  $\log n_{\text{H}} \sim 13$  cm<sup>-3</sup>) that we found from our data and that is consistent with the analysis of (N12). This result claims for accurate estimates of the line emitting region chemical composition that are still missing today.

This preliminary paper provides a description of several key properties of xA at high- $L$ . One of the most interesting result is that Al<sub>III</sub>λ1860 is providing a tracer of the dynamics of a virialized LIL-emitting region. If this is true at least approximately Al<sub>III</sub>λ1860 may offers a virial broadening estimator. The Al<sub>III</sub>λ1860 is a resonant doublet which implies resonant absorption and therefore efficient acceleration by continuum scattering. Then a non virial contribution due to outflows may remain unresolved in the Al<sub>III</sub>λ1860 profile. Simultaneous observations of

the  $\text{Alm}\lambda 1860$  and of the  $H\beta$  spectral ranges are needed to make an exhaustive assessment of  $\text{Alm}\lambda 1860$  as a virial broadening estimator. The conceptual validity of luminosity estimates based on the  $H\beta$  line width measurements has been confirmed by the Hubble diagrams shown by MS14 and Negrete et al. (2017). The large scatter displayed by quasars with respect to supernovae demands that random and systematic sources of error should be understood and, if possible, corrected or reduced before the  $x\text{A}$  sources can be exploited as useful distance indicators for cosmology.

A reliable virial broadening estimator would make possible an application of the virial luminosity estimates proposed by MS14 to a large number of quasars in the redshift range  $1 \lesssim z \lesssim 3.5$ . This redshift range is especially well suited for obtaining constraints on the cosmic density of matter  $\Omega_M$ , since the effect of the cosmological constant in the dynamics of the Universe is felt only at  $z \lesssim 1$ .


**Acknowledgements.** MLMA acknowledges a CONACyT postdoctoral fellowship. AdO, MLMA and JWS acknowledge financial support from the Spanish Ministry for Economy and Competitiveness through grants AYA2013-42227-P and AYA2016-76682-C3-1-P. MLMA, PM and MDO acknowledge funding from the INAF PRIN-SKA 2017 program 1.05.01.88.04. JP acknowledged financial support from the Spanish Ministry for Economy and Competitiveness through grants AYA2013-40609-P and AYA2016-76682-C3-3-P. MLMA is thankful for the kind hospitality at the Padova Astronomical Observatory, PM for the hospitality at IAA-CSIC. DD and AN acknowledge support from CONACyT through grant CB221398. DD and AN thank also for support from grant IN108716 PAPIIT, UNAM. We would like to thank Drs. J. Masegosa, I. Marquez, M. Povic and S. Cazzoli for all the fruitful discussions on the GANG meetings. This work is based on observations made with the Gran Telescopio Canarias (GTC), installed in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias, in the island of La Palma. We thank all the GTC Staff for their support with the observations. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

## References

- Andrae, R. 2010, ArXiv e-prints, [arXiv:1009.2755]
- Bachev, R., Marziani, P., Sulentic, J. W., et al. 2004, ApJ, 617, 171
- Baldwin, J. A., Ferland, G. J., Korista, K. T., et al. 1996, ApJ, 461, 664
- Baldwin, J. A., Ferland, G. J., Korista, K. T., Hamann, F., & LaCluyzé, A. 2004, ApJ, 615, 610
- Barlow, R. 1989, Statistics. A guide to the use of statistical methods in the physical sciences
- Bellazzini, M. 2007, Light losses due to variation of seeing with wavelength. Technical Notes GAIA
- Bensch, K., del Olmo, A., Sulentic, J., Perea, J., & Marziani, P. 2015, Journal of Astrophysics and Astronomy, 36, 467
- Bischetti, M., Piconcelli, E., Vietri, G., et al. 2017, A&A, 598, A122
- Bisogni, S., di Serego Alighieri, S., Goldoni, P., et al. 2017, A&A, 603, A1
- Boroson, T. A. 2002, ApJ, 565, 78
- Boroson, T. A. & Green, R. F. 1992, ApJS, 80, 109
- Boyle, B. J., Shanks, T., Croom, S. M., et al. 2000, MNRAS, 317, 1014
- Brotherton, M. S., Runnoe, J. C., Shang, Z., & DiPompeo, M. A. 2015, MNRAS, 451, 1290
- Brotherton, M. S., Tran, H. D., Becker, R. H., et al. 2001, ApJ, 546, 775
- Brühweiler, F. & Verner, E. 2008, ApJ, 675, 83
- Cano-Díaz, M., Maiolino, R., Marconi, A., et al. 2012, A&A, 537, L8
- Carniani, S., Marconi, A., Maiolino, R., et al. 2015, A&A, 580, A102
- Carswell, R. F., Mountain, C. M., Robertson, D. J., et al. 1991, ApJL, 381, L5
- Cegłowski, M., Kunert-Bajraszewska, M., & Roskowiński, C. 2015, MNRAS, 450, 1123
- Cicone, C., Maiolino, R., Sturm, E., et al. 2014, A&A, 562, A21
- Clowes, R. G., Habertzettl, L., Raghunathan, S., et al. 2016, MNRAS, 460, 1428
- Coatman, L., Hewett, P. C., Banerji, M., & Richards, G. T. 2016, MNRAS, 461, 647
- Coatman, L., Hewett, P. C., Banerji, M., et al. 2017, MNRAS, 465, 2120
- Cracco, V., Ciroi, S., Berton, M., et al. 2016, MNRAS, 462, 1256
- Czerny, B., Beaton, R., Beijer, M., et al. 2018, SSR, 214, 32
- Denney, K. D., Horne, K., Brandt, W. N., et al. 2016, ApJ, 833, 33
- Diamond-Stanic, A. M., Fan, X., Brandt, W. N., et al. 2009, ApJ, 699, 782
- D’Onofrio, M. & Burigana, C. 2009, Questions of Modern Cosmology: Galileo’s Legacy
- Du, P., Lu, K.-X., Hu, C., et al. 2016a, ApJ, 820, 27
- Du, P., Wang, J.-M., Hu, C., et al. 2016b, ApJL, 818, L14
- Elvis, M., Wilkes, B. J., McDowell, J. C., et al. 1994, APJS, 95, 1
- Eracleous, M. & Halpern, J. P. 2003, ApJ, 599, 886
- Espey, B. R., Carswell, R. F., Bailey, J. A., Smith, M. G., & Ward, M. J. 1989, ApJ, 342, 666
- Ferland, G. J., Porter, R. L., van Hoof, P. A. M., et al. 2013, RevMexA&Ap, 49, 137
- Feruglio, C., Fiore, F., Carniani, S., et al. 2015, A&A, 583, A99
- Fraix-Burnet, D., Marziani, P., D’Onofrio, M., & Dultzin, D. 2017, Frontiers in Astronomy and Space Sciences, 4, 1
- Gallerani, S., Maiolino, R., Juarez, Y., et al. 2010, A&A, 523, A85
- Gaskell, C. M. 1982, ApJ, 263, 79
- Gaskell, C. M., Brandt, W. N., Dietrich, M., Dultzin-Hacyan, D., & Eracleous, M., eds. 1999, Astronomical Society of the Pacific Conference Series, Vol. 175, Structure and Kinematics of Quasar Broad Line Regions
- Giacconi, R., Branduardi, G., Briel, U., et al. 1979, ApJ, 230, 540
- Gordon, K. D. & Clayton, G. C. 1998, ApJ, 500, 816
- Graham, M. J., Clowes, R. G., & Campusano, L. E. 1996, MNRAS, 279, 1349
- Hall, P. B., Anderson, S. F., Strauss, M. A., et al. 2002, ApJS, 141, 267
- Hamann, F. & Ferland, G. 1993, ApJ, 418, 11
- Hamann, F. & Ferland, G. 1999, ARA&A, 37, 487
- Harris, D. W., Jensen, T. W., Suzuki, N., et al. 2016, AJ, 151, 155
- Harrison, C. M., Alexander, D. M., Mullaney, J. R., & Swinbank, A. M. 2014, MNRAS, 441, 3306
- Hu, C., Wang, J.-M., Ho, L. C., et al. 2008, ApJL, 683, L115
- Johansson, S., Brage, T., Leckrone, D. S., Nave, G., & Wahlgren, G. M. 1995, ApJ, 446, 361
- Johansson, S. & Hansen, J. E. 1988, in Astrophysics and Space Science Library, Vol. 138, IAU Colloq. 94: Physics of Formation of FE II Lines Outside LTE, ed. R. Viotti, A. Vittone, & M. Friedjung, 235–241
- Kaspi, S., Brandt, W. N., Maoz, D., et al. 2007, ApJ, 659, 997
- Kellermann, K. I., Sramek, R., Schmidt, M., Shaffer, D. B., & Green, R. 1989, AJ, 98, 1195
- King, A. & Muldrew, S. I. 2016, MNRAS, 455, 1211
- King, A. & Pounds, K. 2015, ARA&A, 53, 115
- Kovačević, J., Popović, L. Č., & Dimitrijević, M. S. 2010, ApJS, 189, 15
- Krisz, G. 1994, Astronomical Data Analysis Software and Systems III, A.S.P. Conference Series, 61, 437
- Kuraszkiewicz, J. K., Green, P. J., Crenshaw, D. M., et al. 2004, ApJS, 150, 165
- Kurucz, R. & Bell, B. 1995, Atomic Line Data (R.L. Kurucz and B. Bell) Kurucz CD-ROM No. 23. Cambridge, Mass.: Smithsonian Astrophysical Observatory, 1995., 23
- La Franca, F., Bianchi, S., Ponti, G., Branchini, E., & Matt, G. 2014, ApJL, 787, L12
- Laor, A., Jannuzi, B. T., Green, R. F., & Boroson, T. A. 1997, ApJ, 489, 656
- Luo, B., Brandt, W. N., Hall, P. B., et al. 2015, ApJ, 805, 122
- Malkan, M. A. & Sargent, W. L. W. 1982, ApJ, 254, 22
- Marinello, A. O. M., Rodríguez-Ardila, A., García-Rissmann, A., Sigut, T. A. A., & Pradhan, A. K. 2016, ApJ, 820, 116
- Martínez-Aldama, M. L., Del Olmo, A., Marziani, P., et al. 2017, Frontiers in Astronomy and Space Sciences, 4, 29
- Martínez-Aldama, M. L., Dultzin, D., Marziani, P., et al. 2015, ApJS, 217, 3
- Marziani, P., Dultzin, D., Sulentic, J. W., et al. 2018, ArXiv e-prints, arXiv:180205575M
- Marziani, P., Martínez Carballo, M. A., Sulentic, J. W., et al. 2016, Ap&SS, 361, 29
- Marziani, P., Negrete, C. A., Dultzin, D., et al. 2017a, Frontiers in Astronomy and Space Sciences, 4, 16
- Marziani, P., Olmo, A., Martínez-Aldama, M., et al. 2017b, Atoms, 5, 33
- Marziani, P. & Sulentic, J. W. 2014, MNRAS, 442, 1211
- Marziani, P., Sulentic, J. W., Dultzin-Hacyan, D., Calvani, M., & Moles, M. 1996, ApJS, 104, 37
- Marziani, P., Sulentic, J. W., Negrete, C. A., et al. 2010, MNRAS, 409, 1033
- Marziani, P., Sulentic, J. W., Plauchu-Frayn, I., & del Olmo, A. 2013, ApJ, 764
- Marziani, P., Sulentic, J. W., Stirpe, G. M., Zamfir, S., & Calvani, M. 2009, A&Ap, 495, 83
- Marziani, P., Sulentic, J. W., Zamanov, R., et al. 2003a, ApJS, 145, 199
- Marziani, P., Sulentic, J. W., Zwitter, T., Dultzin-Hacyan, D., & Calvani, M. 2001, ApJ, 558, 553
- Marziani, P., Zamanov, R. K., Sulentic, J. W., & Calvani, M. 2003b, MNRAS, 345, 1133
- Mathews, W. G. & Ferland, G. J. 1987, ApJ, 323, 456
- Matsuoka, K., Silverman, J. D., Schramm, M., et al. 2013, ApJ, 771, 64
- Mejía-Restrepo, J. E., Trakhtenbrot, B., Lira, P., & Netzer, H. 2018, MNRAS

- Mejía-Restrepo, J. E., Trakhtenbrot, B., Lira, P., Netzer, H., & Capellupo, D. M. 2016, *MNRAS*, 460
- Mineshige, S., Kawaguchi, T., Takeuchi, M., & Hayashida, K. 2000, *PASJ*, 52, 499
- Negrete, A., Dultzin, D., Marziani, P., & Sulentic, J. 2012, *ApJ*, 757, 62
- Negrete, C. A., Dultzin, D., Marziani, P., & et al. 2018, Submitted
- Negrete, C. A., Dultzin, D., Marziani, P., & Sulentic, J. W. 2013, *ApJ*, 771, 31
- Negrete, C. A., Dultzin, D., Marziani, P., & Sulentic, J. W. 2014, *ApJ*, 794, 95
- Negrete, C. A., Dultzin, D., Marziani, P., et al. 2017, *Frontiers in Astronomy and Space Sciences*, 4, 59
- Netzer, H., Lira, P., Trakhtenbrot, B., Shemmer, O., & Cury, I. 2007, *ApJ*, 671, 1256
- Netzer, H. & Marziani, P. 2010, *ApJ*, 724, 318
- Park, D., Woo, J.-H., Denney, K. D., & Shin, J. 2013, *ApJ*, 770, 87
- Penston, M. V. 1987, *MNRAS*, 229, 1P
- Plotkin, R. M., Shemmer, O., Trakhtenbrot, B., et al. 2015, *ApJ*, 805, 123
- Richards, G. T., Kruczek, N. E., Gallagher, S. C., et al. 2011, *AJ*, 141, 167
- Richards, G. T., Lacy, M., Storrie-Lombardi, L. J., et al. 2006, *ApJS*, 166, 470
- Scaringi, S., Cottis, C. E., Knigge, C., & Goad, M. R. 2009, *MNRAS*, 399, 2231
- Schneider, D. P., Hall, P. B., Richards, G. T., et al. 2005, *AJ*, 130, 367
- Shemmer, O., Netzer, H., Maiolino, R., et al. 2004, *ApJ*, 614, 547
- Shemmer, O., Trakhtenbrot, B., Anderson, S. F., et al. 2010, *ApJL*, 722, L152
- Shen, Y., Brandt, W. N., Richards, G. T., et al. 2016, *ApJ*, 831, 7
- Shen, Y. & Liu, X. 2012, *ApJ*, 753, 125
- Shin, J., Woo, J.-H., Nagao, T., & Kim, S. C. 2013, *ApJ*, 763, 58
- Sigut, T. A. A. & Pradhan, A. K. 1998, *ApJL*, 499, L139
- Sigut, T. A. A., Pradhan, A. K., & Nahar, S. N. 2004, *ApJ*, 611, 81
- Sniegowska, M., Czerny, B., You, B., et al. 2017, ArXiv e-prints
- Sulentic, J. & Marziani, P. 2015, *Frontiers in Astronomy and Space Sciences*, 2, A6
- Sulentic, J. W., Bachev, R., Marziani, P., Negrete, C. A., & Dultzin, D. 2007, *ApJ*, 666, 757
- Sulentic, J. W., del Olmo, A., Marziani, P., et al. 2017, *A&A*, 608, A122
- Sulentic, J. W., Dultzin-Hacyan, D., Marziani, P., et al. 2006a, *Revista Mexicana de Astronomia y Astrofisica*, 42, 23
- Sulentic, J. W., Martínez-Carballo, M. A., Marziani, P., et al. 2015, *MNRAS*, 450, 1916
- Sulentic, J. W., Marziani, P., del Olmo, A., et al. 2014a, *A&A*, 570, A96
- Sulentic, J. W., Marziani, P., Dultzin, D., D'Onofrio, M., & del Olmo, A. 2014b, *Journal of Physics Conference Series*, 565, 012018
- Sulentic, J. W., Marziani, P., & Dultzin-Hacyan, D. 2000a, *ARA&A*, 38, 521
- Sulentic, J. W., Marziani, P., Zamanov, R., et al. 2002, *ApJL*, 566, L71
- Sulentic, J. W., Repetto, P., Stirpe, G. M., et al. 2006b, *A&Ap*, 456, 929
- Sulentic, J. W., Zamfir, S., Marziani, P., et al. 2003, *ApJL*, 597, L17
- Sulentic, J. W., Zamfir, S., Marziani, P., & Dultzin, D. 2008, in *Revista Mexicana de Astronomia y Astrofisica Conference Series*, Vol. 32, 51–58
- Sulentic, J. W., Zwitter, T., Marziani, P., & Dultzin-Hacyan, D. 2000b, *ApJL*, 536, L5
- Sun, J. & Shen, Y. 2015, *ApJL*, 804, L15
- Tombesi, F., Sambruna, R. M., Reeves, J. N., et al. 2010, *ApJ*, 719, 700
- Trakhtenbrot, B. & Netzer, H. 2012, *MNRAS*, 427, 3081
- Trump, J. R., Hall, P. B., Reichard, T. A., et al. 2006, *ApJS*, 165, 1
- Tully, R. B. & Fisher, J. R. 1977, *A&A*, 54, 661
- Vanden Berk, D. E., Richards, G. T., Bauer, A., et al. 2001, *AJ*, 122, 549
- Verner, E., Bruhweiler, F., Verner, D., et al. 2004, *ApJ*, 611, 780
- Véron-Cetty, M.-P. & Véron, P. 2010, *A&A*, 518, A10
- Vestergaard, M. & Peterson, B. M. 2006, *ApJ*, 641, 689
- Vestergaard, M. & Wilkes, B. J. 2001, *ApJS*, 134, 1
- Vietri, G., Piconcelli, E., Bischetti, M., et al. 2018, ArXiv e-prints
- Wang, J.-M., Du, P., Hu, C., et al. 2014, *ApJ*, 793, 108
- Wang, J.-M., Du, P., Valls-Gabaud, D., Hu, C., & Netzer, H. 2013, *Physical Review Letters*, 110, 081301
- Weymann, R. J., Morris, S. L., Foltz, C. B., & Hewett, P. C. 1991, *ApJ*, 373, 23
- Wills, D. & Netzer, H. 1979, *ApJ*, 233, 1
- Wu, J., Brandt, W. N., Anderson, S. F., et al. 2012, *ApJ*, 747, 10
- Wu, J., Brandt, W. N., Hall, P. B., et al. 2011, *ApJ*, 736, 28
- York, D. G., Khare, P., Vanden Berk, D., et al. 2006, *MNRAS*, 367, 945
- Zamfir, S., Sulentic, J. W., & Marziani, P. 2008, *MNRAS*, 387, 856
- Zamfir, S., Sulentic, J. W., Marziani, P., & Dultzin, D. 2010, *MNRAS*, 403, 1759

**Appendix A:**  
**Rest-frame spectra and multicomponent fits**



**Fig. A.1.** Top panel: calibrated rest-frame spectrum of SDSS000807.27-103942.7 before continuum subtraction. Abscissa is rest-frame wavelength in  $\text{\AA}$ , while ordinate is specific flux in units  $10^{-15} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ Hz}^{-1}$ . Global or local continuum are specified by a continuous line. Dot-dashed vertical lines identify the position at rest-frame of the strongest emission lines. Bottom: multicomponent fits after continuum subtraction for the SiIV $\lambda$ 1397, CIV $\lambda$ 1549 and 1900 $\text{\AA}$  blend spectral ranges. In all the panels the continuous black line marks the broad component at rest-frame associated to SiIV $\lambda$ 1397, CIV $\lambda$ 1549, AlIII $\lambda$ 1860 and SiIII $\lambda$ 1892 respectively, while the blue one corresponds to the blueshifted component associated to each emission. Dashed pink line marks the fit to the whole spectrum. Absorption lines are indicated by a brown line. Dot-dashed vertical lines correspond to the rest-frame of each emission line. In the SiIV $\lambda$ 1397 spectral range, the cyan line marks the contribution of OI + S II 1304 blend, while the magenta line corresponds to the CII $\lambda$ 1335 emission line. In the CIV $\lambda$ 1549 region, NIV $\lambda$ 1486 is represented by a cyan line, while the yellow one corresponds to the OIII $\lambda$ 1663 + AlIII $\lambda$ 1670 blend. In the 1900 $\text{\AA}$  blend range, FeIII and FeII contributions are traced by dark and pale green lines respectively, magenta lines marks the NIII $\lambda$ 1750 and the yellow one corresponds to the SiII $\lambda$ 1816. Lower panels correspond to the residuals, abscissa is in radial velocity units  $\text{km s}^{-1}$ .




Fig. A.2. (cont.) Same of the previous panel, for SDSSJ004241.95+002213.9.




Fig. A.3. (cont.) Same of the previous panel, for SDSSJ021606.41+011509.5.




Fig. A.4. (cont.) Same of the previous panel, for SDSSJ024154.42-004757.5.




Fig. A.5. (cont.) Same of the previous panel, for SDSSJ084036.16+235524.7.




Fig. A.6. (cont.) Same of the previous panel, for SDSSJ101822.96+203558.6.




Fig. A.7. (cont.) Same of the previous panel, for SDSSJ103527.40+445435.6.




Fig. A.8. (cont.) Same of the previous panel, for SDSSJ105806.16+600826.9.




Fig. A.9. (cont.) Same of the previous panel, for SDSSJ110022.53+484012.6.




Fig. A.10. (cont.) Same of the previous panel, for SDSSJ125659.79-033813.8.




Fig. A.11. (cont.) Same of the previous panel, for SDSSJ131132.92+052751.2.




Fig. A.12. (cont.) Same of the previous panel, for SDSSJ143525.31+400112.2.




Fig. A.13. (cont.) Same of the previous panel, for SDSSJ144412.37+582636.9.




Fig. A.14. (cont.) Same of the previous panel, for SDSSJ151258.36+352533.2.




Fig. A.15. (cont.) Same of the previous panel, for SDSSJ214009.01-064403.9.




Fig. A.16. (cont.) Same of the previous panel, for SDSSJ220119.62-083911.6.




Fig. A.17. (cont.) Same of the previous panel, for SDSSJ222753.07-092951.7.




Fig. A.18. (cont.) Same of the previous panel, for SDSSJ233132.83+010620.9.




Fig. A.19. (cont.) Same of the previous panel, for SDSSJ234657.25+145736.0.


## Appendix B: Error estimates

Emission lines of the xA spectra are blended in almost all cases. For example, in the 1900Å blend the emission of CIII]λ1909 and FeII λ1914 cannot be deblended; or in the case of CIV the separation between broad and blue components is not obvious. We built Monte Carlo (MC) simulations to determine the error estimates, in order to consider the effects of blending. The MC method allows us to consider all emission components and include a variation of the flux and FWHM simultaneously. The method is based on minimizing the  $\chi^2$  (Barlow 1989). A variation in parameters of the line properties causes changes in the  $\chi^2$ . Then, the parameters at  $1\sigma$  confidence level are those that satisfy the constraint:  $\chi_{1\sigma}^2 \leq \chi_{\min}^2 + 1$  (Andrae 2010).

The MC method is applied to the same spectral regions identified for the multicomponent fits with SPECFIT (Section §4.2). All line components considered in modeling the observed quasar emission spectrum were taken into account. For example, in the 1900Å blend region variations in flux and FWHM of AlIII, SiIII], CIII], SiIIλ1816, NIII]λ1750, FeII and FeIII are considered for each simulation. Parameters vary around the values given by the SPECFIT model without any constraint and the simulations are therefore totally independent of the SPECFIT model.

We ran 5000 simulations for each quasar in order to obtain good sampling. Left panels of Figure B.1 show the distributions of  $\chi^2$  for the flux of AlIII (top) and for the total CIV profile (bottom), for the quasar SDSSJ222753.07-092951.7. Usually  $\sim 1000 - 3000$  simulations satisfy the criterion  $\chi_{1\sigma}^2 \leq \chi_{\min}^2 + 1$ . The total width of the distribution at  $\chi_{\min}^2 + 1$  is the  $1\sigma$  error confidence level we consider for each parameter. The central panels of Figure B.1 show the 1900Å blend (top) and CIV (bottom) spectra, and the quasar emission associated with the  $1\sigma$  confidence level derived from the MC simulations (gray shadow regions).

Strongest emission lines like AlIII or the total emission of CIV show Gaussian distributions for the flux and FWHM MC replications (See right panels of Figure B.1). However, for weak lines like CIII]λ1909, NIII]λ1750 or HeIIλ1640<sub>BC</sub>, the distribution has an asymmetric or flat behavior. This is because faint lines are strongly affected by the presence of the stronger ones (for example SiIIλ1816 affected by AlIII). In these cases, we consider the flux found by SPECFIT as an upper limit. We report the FWHM values followed by a double colon to signify a high uncertain value (Tables 4, 5, 6 and 7).



**Fig. B.1.** Left panels: distribution of  $\chi^2$  as a function of the flux of AlIII (top) and total CIV profile (bottom). Black continuous lines mark the normal distribution associated with simulations. Gray horizontal lines indicate the value at  $\chi^2_{\min}$  and at  $\chi^2_{\min} + 1$ . The gray vertical line indicates the mean of the normal distribution. Middle panels: spectra of the quasar SDSSJ222753.07-092951.7 in the 1900Å blend (top) and total CIV $\lambda$ 1549 (bottom) spectral ranges. Gray shadow regions correspond to the models at the  $\chi^2_{\min} + 1$  confidence level obtained from the Monte Carlo simulations. The blue line corresponds to the SPECFIT result. Right panels show the distributions of the fluxes of AlIII (top) and total CIV (bottom) for all the models under the  $1\sigma$  confidence level plotted in the left panels. Vertical lines has the same meaning as in left panels.

## Appendix C: FeII and FeIII

### C.1. Identification of FeII and FeIII transitions

In next lines are described the most important FeII and FeIII transitions in the spectral range 1700 – 2100 Å. Table C.1 reports the approximate central wavelength of the FeII and FeIII features, the multiplet identification, and the sources where they are prominent.

#### FeII

[ $\lambda 1715 \text{ \AA}$ ] – This feature has not been identified previously as FeII emission. Graham et al. (1996) suggest AlII associated with the  $3p^3P^o - 3d^3D$  transitions. However, the wavelength consistency is poor, the AlII transition is fairly high level, and its intensity is predicted negligible even for very low ionization parameter ( $\sim 10^{-3}$ ). We therefore ascribe the feature to FeII UV multiplet #38 (associated with the transition between terms  $a^4F - 4D^o$ ), and specifically to the transition  $4D_{3/2}^o \rightarrow 4F_{7/2}^o$  that has the higher oscillator strength among all FeII lines listed by Kurucz & Bell (1995,  $\log gf \approx -0.395$ ).

[ $\lambda 1785 \text{ \AA}$ ] – The UV multiplet #191 is strong in several quasars, most notably I Zw 1 (Marziani et al. 1996; Laor et al. 1997; Vestergaard & Wilkes 2001). The emission is believed to be enhanced by a Ly $\alpha$  fluorescence mechanism (Baldwin et al. 2004) also operating in symbiotic stars and in stars with chromospheric activity (Johansson & Hansen 1988; Johansson et al. 1995).

[ $\lambda 2020 \text{ \AA}$ ] – This feature is never strong ( $\lesssim 2 \text{ \AA}$ ), but is apparently detected in a few objects (Tab. C.1). The main feature could be associated with the line at  $2020.739 \text{ \AA}$  and other two lines of FeII multiplet UV #83.

#### FeIII

[ $\lambda 1914 \text{ \AA}$ ] – The line of the FeIII UV multiplet #34 is included in the template. We add a component to take into account the possibility of extraordinary enhancement due to Ly $\alpha$  fluorescence, as outlined in Marziani et al. (2010).

[ $\lambda 2005 \text{ \AA}$  &  $\lambda 2045 \text{ \AA}$ ] – These features have been identified as due to FeIII UV #55 (Graham et al. 1996; Vestergaard & Wilkes 2001).

[ $\lambda 2080 \text{ \AA}$ ] – This feature is perhaps the most prominent one among the FeIII features. Its identification FeIII UV #48 is supported by the consistency in wavelength of the three multiplet components. The peculiarity of the GTC-xA sources is that the feature is not reproduced well by the template: an additional component is needed. The analysis of the  $2080 \text{ \AA}$  feature is reported in Section §6.4, and its origin is discussed in Appendix C.2 below.

[ $\lambda 2093 \text{ \AA}$ ] – We identify this feature as due to FeIII UV #77 (Vestergaard & Wilkes 2001). A previous identification as FeI seems less likely, considering that FeI features are predicted to be exceedingly weak (Sigut et al. 2004) even at the lowest ionization degrees.

[ $\lambda 2115 \text{ \AA}$ ] – Vestergaard & Wilkes (2001) associates this feature with FeIII UV #58.

### C.2. On the origin of FeIII emission

The strength of FeIII emission is surprising, given that the overall appearance of the spectrum is suggestive of extremely low

ionization. The FeIII features have also been observed in Pop. A sources or associated with extreme Pop. A (e.g., Baldwin et al. 1996; Graham et al. 1996; Richards et al. 2011). The FeIII lines are mainly emitted in a region at the boundary between the fully/partially ionized zone. At ionization parameter  $\log U \gtrsim -2$ , the region of FeIII line formation is usually a region where the opacity in the Lyman continuum grows to values  $\tau_{912} \gtrsim 1$  (i.e., a small fraction of the geometric depth of the emitting gas of slab). Deeper into the slab from its illuminated face, the dominant ionization stage is Fe $^{+3}$ . A significant fraction of FeIII is present at the illuminated face of the cloud within the HII zone, but only if the ionization parameter is very low ( $\log U \lesssim -3$ , Fig. 3 of Sigut et al. 2004). Fe $^{+2}$  is  $\lesssim 2\%$  of Fe $^{+1}$  in the partially-ionized zone (PIZ): Fe $^{+1}$  still remains the dominant ionization stage of iron in the PIZ, as required by the strong optical FeII emission required for these sources. In the spectral range between 1800 and 2100 Å, however, the dominant contribution may be due to FeII, as predicted by the models of Sigut et al. (2004), which assumes  $\log U \sim -3$ .

The role of Ly $\alpha$  fluorescence in explaining the UV and IR FeII emission has been known since the late 1980s from photoionization simulations (e.g., Sigut & Pradhan 1998; Verner et al. 2004; Sigut et al. 2004) as well as from observational evidence (e.g., Penston 1987; Marinello et al. 2016; Clowes et al. 2016, and references therein). As mentioned, a specific feature that is believed to be enhanced by Ly $\alpha$  fluorescence is the FeII<sub>UV</sub> #191 multiplet at  $1785 \text{ \AA}$ . Also the FeII line at  $\lambda 1914$ , ascribed to the transition of the FeII<sub>UV</sub> #34 multiplet  $z^7P_3^o \rightarrow a^7S_3$ , is expected to be enhanced by Ly $\alpha$  fluorescence. Although this is not a resonant line (the lower level is  $\sim 3 \text{ eV}$  above ground), the line appears often stronger than in the Vestergaard & Wilkes (2001) template, presumably because the upper level is populated by Ly $\alpha$  fluorescence (Marziani et al. 2010). Figure 8 suggest a close association between the 2080 feature and the FeII and FeIII features enhanced by Ly $\alpha$  fluorescence.


Generally speaking, we expect that Ly $\alpha$  fluorescence may be also relevant in the explanation of the overall FeIII spectrum, considering the large number of radiative transitions possible for the electronic configuration of the Fe $^{++}$ . The FeIII high ionic fraction in the HII zone and at the transition zone between HII and PIZ are expected to be conditions that make the fluorescent absorption of Lyman continuum photons by FeIII ions especially efficient. In the specific case of the 2080 feature, we suggest that a mechanism of enhancement is due to fluorescence with the Lyman continuum ionization edge. The FeIII UV #48 multiplet is radiatively linked to the multiplets #118 and #119 between the terms  $z^5P^o e^5D$  and  $e^5S$ , respectively. The energy difference between the lower level of the lower  $a^5S$  term of multiplet #48 and the upper terms of #118 and #119, is 13.65 and 13.67 eV respectively, slightly above the ionization potential of hydrogen (13.5984 eV). Figure C.1 schematically shows the energy levels associated with the spectroscopic terms. Even if the energy is not strictly coincident, the Lyman continuum emission (which is expected to decrease with  $\nu^{-3}$ ) may act as a broadened pseudo-line.

If this mechanism is indeed operating, we expect FeIII emission from multiplets UV #118 and #119 in the range 1595 – 1610 Å. A faint hump present on the composite spectrum in the predicted range could be due to some FeIII emission as well as to the blueshifted component of HeII $\lambda 1640$  blending with the red side of Civ $\lambda 1549$ . Considering the weakness of the features, and the severe blending with HeII $\lambda 1640$  and Civ $\lambda 1549$ , the predic-

**Table C.1.** Identification of FeII and FeIII features

Feature [Å]	Identification	Sources where is detected
$\lambda 1715$	FeII UV #38	J000807, J021606, J084036, J214009, J222753
$\lambda 1785$	FeII UV #191	J125659, J101822, J144412
$\lambda 2005$	FeII UV #55	J233132
$\lambda 2020$	FeII UV #83	J021606, J233132, J214009
$\lambda 2045$	FeII UV #60	J222753, J110022, J151258
$\lambda 2080$	FeII UV #48	J000807, J004241, J021606, J101822, J103527, J125659, J214009, J222753
$\lambda 2093$	FeII UV #77	J105806
$\lambda 2115$	FeII UV #58	J220119, J222753, J110022

tion of FeIII emission around 1600 Å should be verified on high S/N spectra of sources with narrower lines.



**Fig. C.1.** Partial Grotrian diagram showing the energy levels and the spectroscopic terms associated with the emission of FeII multiplet #48, #118 (top left) and #119 (top right). Energy level separation is not drawn to scale for clarity.