



Publication Year	2023
Acceptance in OA	2024-12-18T09:34:40Z
Title	The chemical DNA of the Magellanic Clouds: II. High-resolution spectroscopy of the SMC globular clusters NGC 121, NGC 339, and NGC 419
Authors	Mucciarelli, A., Minelli, A., Lardo, C., MASSARI, DAVIDE, BELLAZZINI, Michele, ROMANO, Donatella, ORIGLIA, Livia, Ferraro, F. R.
Publisher's version (DOI)	10.1051/0004-6361/202347120
Handle	http://hdl.handle.net/20.500.12386/35531
Journal	ASTRONOMY & ASTROPHYSICS
Volume	677

The chemical DNA of the Magellanic Clouds

II. High-resolution spectroscopy of the SMC globular clusters NGC 121, NGC 339 and NGC 419 [★]

A. Mucciarelli^{1,2}, A. Minelli^{1,2}, C. Lardo¹, D. Massari², M. Bellazzini², D. Romano², L. Origlia², and F. R. Ferraro^{1,2}

¹ Dipartimento di Fisica e Astronomia “Augusto Righi”, Alma Mater Studiorum, Università di Bologna, Via Gobetti 93/2, I-40129 Bologna, Italy

² INAF - Osservatorio di Astrofisica e Scienza dello Spazio di Bologna, Via Gobetti 93/3, I-40129 Bologna, Italy

July 10, 2023

ABSTRACT

The Small Magellanic Cloud (SMC) is the host of a rich system of globular clusters (GCs) that span a wide age range. The chemical composition of the SMC clusters is still poorly understood, despite their significance to chemical evolution studies. Here, we provide the first detailed chemical study of evolved giants in three distinct clusters, NGC 121 (10.5 Gyr), NGC 339 (6 Gyr), and NGC 419 (1.4 Gyr). Results are based on high-resolution spectra obtained with FLAMES at the Very Large Telescope.

The chemical fingerprints of these clusters closely resemble those of SMC field stars, supporting the SMC’s specific history of chemical enrichment relative to the Milky Way. The approximately solar-scaled $[\alpha/\text{Fe}]$ observed in all three clusters, independent of their $[\text{Fe}/\text{H}]$, demonstrate the SMC’s low star formation efficiency. Compared to their Milky Way counterparts, elements primarily produced by massive stars are severely underrepresented. Particularly, the young cluster NGC 419’s extremely low $[\text{Zn}/\text{Fe}]$ shows that hypernovae have contributed relatively little during the past two Gyr. The three GCs have high $[\text{Eu}/\text{Fe}]$ values regardless of their age. This suggests that the production of the r-process elements in the SMC was extremely efficient up to 1.5 Gyr ago, with an enrichment timescale comparable to that from Type Ia supernovae.

When the properties of the oldest SMC object NGC 121 are compared to those of in-situ Milky Way clusters and accreted clusters linked to the Gaia-Enceladus merger event, it is shown that the SMC had already attained the same metallicity as Gaia-Enceladus but with lower $[\text{Fe}/\text{H}]$ ratios at the age of NGC 121. This suggests that the chemical enrichment histories of the early SMC and Gaia-Enceladus differed, and that the SMC probably had a lower mass in its early ages than Gaia-Enceladus.

Key words. Galaxies: Magellanic Clouds; Techniques: spectroscopic; Stars: abundances

1. Introduction

The Milky Way’s (MW) most massive satellites, the Large and Small Magellanic Clouds (LMC and SMC, respectively), offer the chance to study a pair of interacting galaxies before they merge and are cannibalised by the Galaxy. The currently prevailing view for their formation and evolution is that the two galaxies evolved independently and only recently formed a binary pair (Diaz & Bekki 2011; D’Onghia & Fox 2016). This binary system is probably at its first peri-Galactic passage (Shuter 1992; Byrd et al. 1994; Besla et al. 2007, 2012). The tidal interactions occurring between these galaxies are responsible for the disturbed structure of the SMC, as well as of the LMC disc warp and of the complex star formation (SF) history of both galaxies, characterised by different, synchronous SF bursts (Harris & Zaritsky 2009; Rubele et al. 2012; Nidever et al. 2020; Massana et al. 2022).

In the last decade, there has been a renewed interest in the turbulent evolution of these irregular galaxies and their star populations, as evidenced by several dedicated photometric surveys, as VMC (VISTA survey of the Magellanic Clouds system, Cioni et al. 2011), STEP (the SMC in Time: Evolution of a Prototype

interacting late-type dwarf galaxy, Ripepi et al. 2014), SMASH (Survey of the MAGellanic Stellar History, Nidever et al. 2017), VISCACHA (VISible Soar photometry of star Clusters in tApid and Coxi HuguA, Maia et al. 2019). Additionally, they will be privileged targets for forthcoming multi-object spectrographs, like 4MOST (see Cioni et al. 2019) and MOONS (see Gonzalez et al. 2020).

The study of the chemical composition of SMC stars has received limited attention, despite its importance to understand the chemical enrichment history of this galaxy. Chemical analyses of high-resolution spectra of SMC red giant branch (RGB) stars have been presented only recently. The APOGEE-2 survey measured abundances for ~1000 SMC RGB stars, in particular Fe and α -elements (Nidever et al. 2020), Al, Ni, Ce (Hasselquist et al. 2021) and Mn (Fernandes et al. 2023). This dataset shows a quite flat behaviour of $[\alpha/\text{Fe}]$ ratios in the range of $[\text{Fe}/\text{H}]$ between -1.2 and -0.2 dex, and a knee (the metallicity corresponding to the decrease in $[\alpha/\text{Fe}]$ abundance ratios) likely located at $[\text{Fe}/\text{H}] < -2.2$ dex. All the measured abundance ratios reveal a clear difference with respect to the MW stars, pointing to a slower SF efficiency in the SMC with respect to the MW. Reggiani et al. (2021) derived the chemical composition of four SMC metal-poor stars ($[\text{Fe}/\text{H}] < -2$ dex) and for two of them measured also the abundance of the r-process element Eu. They found that the SMC is more enriched in Eu than the MW and

[★] Based on observations collected at the ESO-VLT under the programs 072.D-0507, 073.D-0211, 083.D-0208, 085.D-0375, 086.D-0665, 093.B-0583, 095.D-0290 and 188.B-3002.

that elements like Ba and La are produced from this nucleosynthesis channel at these metallicities. These two stars are the only ones in the SMC where Eu has been measured. Mucciarelli et al. (2023, hereafter Paper I) measured Na, α , iron-peak and s-process elements for 206 SMC stars, belonging to three distinct fields around as many globular clusters (GCs). They found that different regions in the SMC are characterised by different metallicity and radial velocity distributions. Some systematic differences in some abundance ratios are identified in different fields. This suggests that the chemical enrichment history in the SMC has not been uniform, with the presence of possible chemically and kinematic distinct substructures. Also, all the abundance ratios of species produced by massive stars are significantly lower than those measured in the MW, indicating a contribution of these stars to the chemical enrichment of the SMC lower than in the MW, according to the low star formation rate expected for this galaxy.

Globular clusters are an excellent tool in the reconstruction of the origin and evolution of the stellar populations in a galaxy as they enable the simultaneous derivation of ages and metallicities and therefore the determination of a reliable age-metallicity relation (AMR). In particular, at large distances (as for external resolved galaxies), the ages of GCs are more accurate and precise than those of field stars that are largely affected by uncertainties arising from reddening and distance. Therefore, the abundances derived in GCs provide time-resolved chemical information and are complementary to those of the field stars. The SMC is the only galaxy in the Local Group that has formed and preserved GCs more or less continuously over the past 11 Gyr. As opposed to that, the LMC hosts 15 old GCs, coeval to the MW ones (Brocato et al. 1996; Olsen et al. 1998; Wagner-Kaiser et al. 2017), and a populous family of clusters younger than ~ 3 Gyr, with a lack of GCs over a broad age range – the so-called Age Gap; the few clusters (ESO 121-SC03, KMHK 1592 and KMHK 1762) falling in this age interval have been likely accreted by the SMC (Mackey et al. 2006; Piatti 2022; Gatto et al. 2022). Several SMC GCs fill the age range corresponding to the LMC Age Gap. The metallicities of SMC GCs have been widely investigated, using photometry (Glatt et al. 2008a,b; Narloch et al. 2021) and low-resolution spectroscopy (Da Costa & Hatzidimitriou 1998; Parisi et al. 2009, 2015; Dias et al. 2021; Parisi et al. 2022; Dias et al. 2022), to explore the global chemical evolution of the galaxy and its AMR. The missing pieces of evidence in our current comprehension of the SMC GCs (and of the SMC stellar populations in general) are their abundance patterns. The only chemical analyses of SMC GCs based on high-resolution spectroscopy available so far are for the very young (~ 30 Myr) cluster NGC 330 (Hill 1999) and for the old cluster NGC 121 (Dalessandro et al. 2016, limited to Fe, Na, O, Mg and Al). Consequently, the investigation of their chemical composition is still entirely unexplored.

This paper is part of a series aimed at deriving the finer chemical details of field and stellar clusters in the Magellanic Clouds, providing new clues to the evolution of these galaxies. In Paper I, we analysed the properties of 206 stars located in three fields around three GCs. In this work, we examine the chemical composition of these three SMC GCs (namely, NGC 121, NGC 339, and NGC 419) covering a wide age range. The aim is of studying the SMC's chemical enrichment history at various ages and to compare the chemical patterns of these clusters with those of field stars derived in Paper I. In this work we discuss 18 species in order to study different sites/channels of nucleosynthesis. We do not discuss the light elements (O, Na, Mg and Al) involved in the CNO-cycle and observed to vary from star to star in GCs;

this phenomenon is usually referred to as multiple populations (see Gratton et al. 2012; Bastian & Lardo 2018, and references therein) and will be discussed in a companion paper. The paper is structured as follows: in Section 2 we summarise the observations and describe the spectroscopic dataset. Sections 3 and 4 present the atmospheric parameters determination and the chemical analysis, respectively. We discuss our results in Sections 5, 6, 7, 8. Finally we summarise our findings and draw our conclusions in Section 9.

2. Spectroscopic dataset

NGC 121 is the oldest SMC GC, with an age of 10.5 ± 0.5 Gyr (Glatt et al. 2008a) and a metallicity of -1.28 ± 0.03 dex (Dalessandro et al. 2016). According to the classification proposed by Parisi et al. (2022), this cluster is associated with the SMC West Halo, a part of the SMC moving outward with respect to the main body of the galaxy (Dias et al. 2016) and probably part of the more extended Counter Bridge (Diaz & Bekki 2012). NGC 339 is a cluster with an age of 6 ± 0.5 Gyr (Glatt et al. 2008b) and $[\text{Fe}/\text{H}] = -1.12 \pm 0.10$ dex (Da Costa & Hatzidimitriou 1998) that has been found to be dynamically unevolved based on its blue stragglers spatial distribution (Dresbach et al. 2022). It is an example of the SMC intermediate-age clusters lacking in the LMC. This cluster is associated with the Southern Bridge, a second branch of the stellar Magellanic Bridge (Belokurov et al. 2017). No high-resolution spectroscopic analysis of this cluster is available in the literature.

NGC 419 is the youngest among the three observed GCs, with an age of 1.4 ± 0.2 Gyr and $[\text{Fe}/\text{H}] = -0.67 \pm 0.12$ dex (Glatt et al. 2008b). Parisi et al. (2022) associated NGC 419 with the main body of the SMC, and stars from the Magellanic bridge have been found in its field of view (Massari et al. 2021). Also for this cluster, no high-resolution spectroscopic data exist.

The spectra analysed in this work were collected with the multi-object optical spectrograph FLAMES (Pasquini et al. 2002) mounted at the Very Large Telescope of the European Southern Observatory under the program 086.D-0665 (PI: Mucciarelli). The observations with the UVES-FLAMES spectrograph have been performed adopting the Red Arm 580 UVES setup (spectral resolution of $R = 47000$ and spectral coverage $\sim 4800 - 6800$ Å). The observations made with the GIRAFFE spectrograph were obtained with the MEDUSA setups HR11 ($5597 - 5840$ Å and $R = 24200$) and HR13 ($6120 - 6406$ Å and $R = 22500$). The exposure times and the number of individual exposures for each setup and cluster are reported in Table 1. In particular, NGC 121 has been observed with 12 exposures of 2700 sec and one of 2200 sec, NGC 339 with 14 of 2700 sec and NGC 419 with 10 of 2700 sec. For each cluster the same targets have been observed during all the exposures.

After the reduction, performed with the dedicated ESO pipelines¹ (including bias subtraction, flat-fielding, wavelength calibration, spectral extraction, and order merging), the individual spectra of each exposure have been cleaned from the sky contribution by subtracting the spectra of some close sky regions observed at the same time as the science targets. Subsequently, single exposures of the same target have been combined in an individual spectrum for each star.

Target stars were selected from the near-infrared photometric catalogues obtained with SofI at the New Technology Telescope (Mucciarelli et al. 2009) in the brightest portion of the RGB ($K_s \sim 13-14$). Cluster membership was determined using the derived

¹ <http://www.eso.org/sci/software/pipelines/>




Fig. 1. $(K, J-K_s)$ colour-magnitude diagrams of the three SMC GCs are plotted as small grey dots. Large coloured circles in each panel represent the spectroscopic targets analysed in this work.

radial velocities (RVs) and metallicities information, considering the targets located within the cluster tidal radius (according to Glatt et al. 2009). In particular, stars located within the cluster tidal radius with discrepant values of RVs and/or $[\text{Fe}/\text{H}]$ have been considered as SMC field stars and not cluster member stars and they have been discussed in Paper I (see Table 2 of Paper I). The positions in the $(K, J-K_s)$ colour-magnitude diagrams of the cluster member stars analysed in this work are shown in Fig. 1.

The final sample of cluster stars includes five members of NGC 121 (all observed with UVES), seven stars of NGC 339 (four observed with UVES and three with GIRAFFE) and eight stars of NGC 419 (five observed with UVES and three with GIRAFFE).

Similar to what we did in our previous works (Minelli et al. 2021; Mucciarelli et al. 2021, Paper I), we defined a control sample of MW GCs analysed with the same approach as the SMC targets. The MW control sample includes spectra of giant stars of 16 GCs retrieved from the ESO archive and obtained with the same UVES-FLAMES configuration (Red Arm 580) used for the majority of target stars of this work. The clear advantage of defining a MW control sample is to erase the systematics of the analysis when comparing the measured abundances in SMC clusters to those of their MW counterparts. These systematics arise mainly from the zero-point of the adopted T_{eff} scale, the adopted solar reference values, the used atomic data (in particular the $\log g_f$ values), the model atmospheres, and the code used for the analysis, so that a simple re-scaling to the same solar reference values does not suffice to put different datasets on the same scale.

3. Atmospheric parameters

For consistency with Paper I, where the effective temperature (T_{eff}) have been derived adopting the colour- T_{eff} transformations by Mucciarelli, Bellazzini & Massari (2021), here we

used $(J - K_s)_0 - T_{\text{eff}}$ relation by González Hernández, & Bonifacio (2009) because these two sets of transformations are based on the same calibration stars. We used the near-infrared photometry by Mucciarelli et al. (2009). Since the colour- T_{eff} relation has a mild dependence on stellar metallicity, we adopted the $[\text{Fe}/\text{H}]$ values available in the literature as a first step and then we recomputed T_{eff} with metallicity values determined from the spectroscopic analysis (anyway leading to variation of less than 20 K with respect to the first run). Colour excess $E(B-V)$ values are from the reddening maps by Schlafly & Finkbeiner (2011), e.g., $E(B-V)=0.028, 0.042,$ and 0.089 mags for NGC 121, NGC 339, and NGC 419 respectively. We compared these values with those of the reddening maps by Skowron et al. (2021) finding an excellent agreement: the colour excess values towards NGC 121 and NGC 339 differ by less than 0.01 mag, and a difference of 0.03 mag is found for NGC 419. The latter difference in $E(B-V)$ leads to variations in T_{eff} of about 30 K and in $\log g$ of 0.01 dex, with a negligible impact on the derived abundances.

Surface gravity ($\log g$) values are then derived by projecting in T_{eff} on the appropriate isochrone (in terms of age and metallicity) for each GC, computed with the Dartmouth Stellar Evolution Database (Dotter et al. 2008). In order to choose the most appropriate isochrone, we first used the age and metallicity determined by Glatt et al. (2008b), refining the metallicity value using those determined in the subsequent analysis. The final values chosen for the isochrones are: 10.5 Gyr and $[\text{Fe}/\text{H}] = -1.2$ dex for NGC 121, 6 Gyr and $[\text{Fe}/\text{H}] = -1.2$ dex for NGC 339 and 1.4 Gyr and $[\text{Fe}/\text{H}] = -0.6$ dex for NGC 419. We used solar-scaled isochrones according to our results for the α -elements abundances (see Section 7).

Finally, the microturbulent velocities v_t for UVES spectra have been determined spectroscopically with the code GALA (Mucciarelli et al. 2013), minimising the slope between the abundances from Fe I lines and the logarithm of the EWs normalized to the wavelength. For the GIRAFFE spectra, the spectroscopically v_t risk to be affected by uncertainties due to the low number of available Fe lines. Therefore we compute them from the $\log g - v_t$ relation provided by Mucciarelli & Bonifacio (2020). We checked for the UVES targets that the two approaches provide consistent results, with an average difference between v_t from the relation and those obtained from the optimisation process of $+0.14 \text{ km s}^{-1}$ ($\sigma=0.13 \text{ km s}^{-1}$). Final atmospheric parameters for the target stars are listed in Table 2.

4. Chemical analysis

4.1. Determination of the chemical abundances

In order to select for the chemical analysis only unblended and linear/poorly saturated lines, we compared the observed spectra with synthetic ones computed with the appropriate atmospheric parameters and metallicity by using the code SYNTH (Kurucz 2005). Model atmospheres have been calculated with the code ATLAS9 (Kurucz 1993, 2005). As a first guess we assumed for the model atmospheres of all the target clusters an α -enhanced chemical mixture. The results of the first run of chemical analysis indicated that these stars have solar-scaled $[\alpha/\text{Fe}]$ abundance ratios and we repeated the analysis adopting solar-scaled model atmospheres.

Atomic and molecular data (such as excitation potential χ , $\log g_f$, damping constants and hyperfine/isotopic splitting) used for synthetic spectra are from the last release of the Kurucz/Castelli linelists², with some exceptions for more recent or more accu-

² <http://wwwuser.oats.inaf.it/castelli/linelists.html>

Table 1. Coordinates of the FLAMES pointing, number of exposures, and exposure times for the GIRAFFE and UVES setups.

Field	RA	Dec	HR11	HR13	UVES
	(J2000)	(J2000)			
NGC 121	00:26:49.0	-71:32:09.9	7x2700sec 1x2200sec	5x2700sec	12x2700 sec 1x2200 sec
NGC 339	00:57:48.9	-74:28:00.1	9x2700sec	5x2700sec	14x2700 sec
NGC 419	01:08:17.7	-72:53:02.7	6x2700sec	4x2700sec	10x2700 sec

Table 2. Adopted atmospheric parameters for SMC GCs targets. The ID numbers are referred to SofI photometric catalogs. The adopted spectrograph is reported in the last column.

ID	RA (Degrees)	Dec (Degrees)	T_{eff} (K)	$\log g$	v_t (km/s)	RV (km/s)	spectrum
NGC 121							
9	6.6842639	-71.5367593	3990	0.53	1.60	+143.71±0.05	UVES
14	6.7033858	-71.5295107	4070	0.66	1.80	+142.12±0.05	UVES
18	6.7233897	-71.5450034	4110	0.73	1.70	+145.35±0.05	UVES
31	6.6845726	-71.5315588	4250	0.97	1.60	+144.67±0.07	UVES
35	6.6970187	-71.5477942	4240	0.95	1.40	+147.80±0.07	UVES
NGC 339							
219	14.4805308	-74.4371549	4140	0.72	1.80	+115.30±0.14	GIRAFFE
466	14.4346714	-74.4407691	4000	0.50	1.90	+112.60±0.12	GIRAFFE
535	14.4182781	-74.4629265	4000	0.50	1.70	+110.83±0.03	UVES
835	14.5722246	-74.4602041	4000	0.50	1.90	+115.70±0.11	GIRAFFE
893	14.4515721	-74.4772827	4050	0.58	1.60	+114.63±0.03	UVES
958	14.4013939	-74.4746562	4210	0.83	1.50	+112.90±0.04	UVES
1076	14.4236180	-74.4731685	4290	0.97	1.50	+111.66±0.05	UVES
NGC 419							
345	17.1136244	-72.9015548	4320	1.41	1.56	+189.90±0.12	GIRAFFE
616	17.0741376	-72.8648011	4050	0.98	1.72	+189.30±0.13	GIRAFFE
727	17.0584817	-72.8773208	4270	1.33	1.60	+183.92±0.05	UVES
732	17.0569039	-72.8916523	4110	1.07	1.50	+186.03±0.04	UVES
852	17.0417327	-72.8819103	4145	1.13	1.70	+187.37±0.06	UVES
885	17.0363101	-72.8775638	4190	1.20	1.64	+191.32±0.13	GIRAFFE
1384	17.0854887	-72.8729310	4150	1.13	1.70	+187.69±0.03	UVES
1633	17.0871666	-72.8803339	4240	1.28	1.60	+190.41±0.06	UVES

rate data for some transitions of Fe, Si, Ca, Ti, Ni, Ba and Eu (see Mucciarelli et al. 2017b, for additional references). Hyperfine and/or isotopic splitting are taken into account for Sc, V, Mn, Co, Cu, Ba, La, Eu. The synthetic spectra have been convolved with a Gaussian profile in order to reproduce the observed broadening of GIRAFFE and UVES spectra. The final linelists include transitions of elements belonging to the main groups, such as α , iron-peak and neutron-capture, in particular, 18 elements for UVES spectra and 12 elements for GIRAFFE spectra.

For species with unblended lines (Fe, Ca, Ti, Si, Cr, Ni, Zr, Y and Nd), we measured their EWs with DAOSPEC (Stetson, & Pancino 2008) through the wrapper 4DAO (Mucciarelli 2013), which also provides RVs of sample stars. Chemical abundances are derived from the measured EWs by using the code GALA (Mucciarelli et al. 2013). The line fitting has been visually inspected line by line, with the purpose to identify possible poorly fitted lines or erroneous determination of the local continuum.

A different approach is adopted for the species with lines characterised by hyperfine/isotopic structure (Sc, V, Mn, Co, Cu, Ba, La, Eu) or transitions located in noisy/complex spectral regions (Zn). Their abundances have been derived with our own code SALVADOR that performs a χ^2 -minimisation between the observed line and a grid of synthetic models computed with the

SYNTHE code, for which we only vary the abundance of the investigated element.

Solar values are taken from Grevesse & Sauval (1998) for all the elements. All the abundance ratios for individual stars are listed in Tab. 3 and 4, together with the corresponding uncertainties as described in Section 4.2. We adopted a maximum likelihood algorithm (see e.g. Pryor & Meylan 1993; Walker et al. 2006; Mucciarelli et al. 2012) to estimate for each cluster the average abundance ratios and their intrinsic scatter together with associated uncertainty, obtained by taking into account the uncertainties of abundance ratios in individual stars. Table 5 lists the average abundance ratios for all the targets, together with the intrinsic (σ_{int}) and observed (σ_{obs}) spread.

4.2. Error estimates

Abundance uncertainties have been computed combining in quadrature the errors arising from the measure of lines strength and the uncertainties arising from atmospheric parameters (see Paper I, for a detailed description). Internal errors due to the measurements have been estimated as the standard error of the mean of the abundances of individual stars. For abundances from only one line, we propagated the error in the EW as provided

Table 3. Chemical abundances for the SMC GCs stars, first part.

ID	[Fe/H]	[Fe II/H]	[Si/Fe]	[Ca/Fe]	[Sc II/Fe]	[Ti/Fe]	[V/Fe]	[Cr/Fe]	[Mn/Fe]	[Co/Fe]
NGC 121										
9	-1.22 ± 0.04	-1.34 ± 0.12	+0.13 ± 0.08	+0.02 ± 0.06	-0.22 ± 0.06	+0.05 ± 0.09	-0.22 ± 0.07	-0.08 ± 0.14	-0.50 ± 0.09	-0.16 ± 0.05
14	-1.21 ± 0.06	-1.35 ± 0.11	+0.16 ± 0.09	+0.05 ± 0.06	-0.13 ± 0.06	+0.02 ± 0.10	-0.18 ± 0.07	-0.07 ± 0.15	-0.55 ± 0.07	-0.10 ± 0.07
18	-1.13 ± 0.07	-1.19 ± 0.08	+0.08 ± 0.10	+0.06 ± 0.05	-0.20 ± 0.09	+0.03 ± 0.07	-0.30 ± 0.12	-0.24 ± 0.10	-0.63 ± 0.12	-0.18 ± 0.08
31	-1.10 ± 0.06	-1.09 ± 0.10	+0.05 ± 0.12	+0.08 ± 0.05	-0.21 ± 0.07	+0.07 ± 0.08	-0.12 ± 0.06	-0.17 ± 0.13	-0.52 ± 0.15	-0.12 ± 0.06
35	-1.18 ± 0.06	-1.28 ± 0.07	-0.05 ± 0.11	+0.01 ± 0.04	-0.23 ± 0.06	+0.04 ± 0.06	-0.26 ± 0.09	-0.19 ± 0.14	-0.64 ± 0.09	-0.11 ± 0.08
NGC 339										
219	-1.26 ± 0.11	-	+0.10 ± 0.12	+0.18 ± 0.08	-	+0.11 ± 0.12	-0.29 ± 0.16	-0.22 ± 0.12	-	-0.05 ± 0.09
466	-1.24 ± 0.08	-	+0.26 ± 0.14	+0.30 ± 0.10	-	+0.10 ± 0.15	-0.17 ± 0.18	-0.19 ± 0.13	-	-0.10 ± 0.09
535	-1.24 ± 0.02	-1.27 ± 0.06	+0.17 ± 0.04	+0.05 ± 0.06	-0.28 ± 0.07	-0.05 ± 0.04	-0.40 ± 0.06	-0.18 ± 0.05	-0.53 ± 0.03	-0.14 ± 0.03
835	-1.24 ± 0.08	-	+0.23 ± 0.12	+0.22 ± 0.12	-	+0.05 ± 0.14	-0.35 ± 0.18	-0.02 ± 0.11	-	-0.07 ± 0.08
893	-1.25 ± 0.03	-1.36 ± 0.05	+0.10 ± 0.05	+0.11 ± 0.05	-0.32 ± 0.05	+0.02 ± 0.05	-0.32 ± 0.07	-0.11 ± 0.05	-0.53 ± 0.03	-0.27 ± 0.03
958	-1.20 ± 0.05	-1.32 ± 0.05	+0.18 ± 0.06	+0.11 ± 0.06	-0.10 ± 0.09	+0.04 ± 0.05	-0.22 ± 0.06	-0.05 ± 0.06	-0.48 ± 0.09	-0.04 ± 0.05
1076	-1.27 ± 0.04	-1.38 ± 0.05	+0.18 ± 0.05	+0.16 ± 0.04	-0.35 ± 0.07	+0.04 ± 0.03	-0.27 ± 0.07	-0.13 ± 0.05	-0.55 ± 0.04	-0.06 ± 0.04
NGC 419										
345	-0.54 ± 0.11	-	-0.08 ± 0.17	-0.01 ± 0.10	-	+0.01 ± 0.13	-0.15 ± 0.17	-0.18 ± 0.15	-	-0.19 ± 0.09
616	-0.62 ± 0.09	-	+0.04 ± 0.13	+0.01 ± 0.14	-	+0.01 ± 0.15	-0.32 ± 0.17	-0.11 ± 0.13	-	-0.05 ± 0.09
727	-0.60 ± 0.05	-0.70 ± 0.07	+0.02 ± 0.08	+0.04 ± 0.08	-0.19 ± 0.05	-0.08 ± 0.09	-0.42 ± 0.11	-0.10 ± 0.08	-0.50 ± 0.08	-0.30 ± 0.05
732	-0.53 ± 0.05	-0.67 ± 0.11	-0.04 ± 0.07	-0.07 ± 0.10	-0.32 ± 0.06	+0.00 ± 0.10	-0.36 ± 0.12	-0.06 ± 0.08	-0.51 ± 0.10	-0.23 ± 0.05
852	-0.61 ± 0.05	-0.57 ± 0.09	+0.11 ± 0.08	-0.09 ± 0.11	-0.12 ± 0.07	-0.22 ± 0.09	-0.52 ± 0.09	-0.15 ± 0.08	-0.54 ± 0.08	-0.30 ± 0.05
885	-0.55 ± 0.11	-	+0.07 ± 0.15	+0.17 ± 0.14	-	+0.00 ± 0.14	-0.31 ± 0.17	-0.08 ± 0.20	-	-0.10 ± 0.10
1384	-0.56 ± 0.04	-0.67 ± 0.08	+0.01 ± 0.06	-0.11 ± 0.08	-0.22 ± 0.05	-0.18 ± 0.08	-0.36 ± 0.09	-0.12 ± 0.07	-0.48 ± 0.05	-0.25 ± 0.04
1633	-0.65 ± 0.05	-0.72 ± 0.07	+0.02 ± 0.08	-0.05 ± 0.07	-0.26 ± 0.06	-0.12 ± 0.08	-0.35 ± 0.09	-0.11 ± 0.08	-0.64 ± 0.05	-0.21 ± 0.05

Table 4. Chemical abundances for the SMC GCs stars, second part.

ID	[Ni/Fe]	[Cu/Fe]	[Zn/Fe]	[Y/Fe]	[Zr/Fe]	[Ba II/Fe]	[La II/Fe]	[Nd II/Fe]	[Eu II/Fe]
NGC 121									
9	-0.17 ± 0.03	-1.00 ± 0.22	-	-0.42 ± 0.15	-0.01 ± 0.14	-0.16 ± 0.09	+0.04 ± 0.08	+0.50 ± 0.09	+0.45 ± 0.11
14	-0.17 ± 0.04	-1.13 ± 0.23	-	-0.12 ± 0.17	-0.09 ± 0.16	-0.10 ± 0.12	+0.03 ± 0.08	+0.50 ± 0.09	+0.49 ± 0.12
18	-0.11 ± 0.03	-0.81 ± 0.23	-	-	-0.17 ± 0.14	-0.01 ± 0.09	-0.05 ± 0.10	+0.66 ± 0.08	+0.56 ± 0.12
31	-0.20 ± 0.04	-	-	-	0.15 ± 0.18	-0.03 ± 0.11	-0.08 ± 0.08	+0.57 ± 0.10	+0.50 ± 0.12
35	-0.10 ± 0.04	-	-	-	-0.05 ± 0.15	-0.18 ± 0.10	+0.00 ± 0.09	+0.69 ± 0.09	+0.37 ± 0.11
NGC 339									
2199	-0.17 ± 0.05	-0.67 ± 0.12	-	-	-0.01 ± 0.20	+0.33 ± 0.09	+0.07 ± 0.11	-	-
4669	-0.22 ± 0.04	-0.59 ± 0.12	-	-	+0.01 ± 0.21	+0.16 ± 0.10	+0.36 ± 0.11	-	-
5353	-0.21 ± 0.02	-0.65 ± 0.07	-0.24 ± 0.09	-0.24 ± 0.05	+0.00 ± 0.08	+0.24 ± 0.06	+0.19 ± 0.05	+0.28 ± 0.04	+0.67 ± 0.07
8358	-0.14 ± 0.05	-0.63 ± 0.11	-	-	-0.09 ± 0.22	+0.07 ± 0.11	+0.32 ± 0.11	-	-
8933	-0.22 ± 0.02	-0.64 ± 0.08	-0.33 ± 0.09	-0.22 ± 0.05	+0.00 ± 0.09	+0.09 ± 0.06	+0.17 ± 0.05	+0.31 ± 0.04	+0.63 ± 0.07
9585	-0.17 ± 0.02	-0.57 ± 0.11	-0.21 ± 0.14	-0.09 ± 0.11	+0.02 ± 0.11	+0.18 ± 0.09	+0.20 ± 0.07	+0.45 ± 0.05	+0.71 ± 0.11
10764	-0.15 ± 0.02	-0.80 ± 0.08	-0.35 ± 0.10	-0.27 ± 0.06	+0.06 ± 0.09	+0.15 ± 0.07	+0.18 ± 0.07	+0.33 ± 0.05	+0.72 ± 0.10
NGC 419									
345	-0.07 ± 0.08	-0.67 ± 0.15	-	-	-0.01 ± 0.20	+0.27 ± 0.08	+0.39 ± 0.10	-	-
616	-0.15 ± 0.07	-0.65 ± 0.10	-	-	+0.03 ± 0.20	+0.28 ± 0.07	+0.25 ± 0.09	-	-
727	-0.22 ± 0.03	-0.88 ± 0.15	-0.66 ± 0.15	-0.11 ± 0.08	+0.09 ± 0.16	+0.28 ± 0.08	+0.29 ± 0.07	+0.42 ± 0.06	+0.54 ± 0.10
732	-0.28 ± 0.03	-	-0.70 ± 0.15	-0.33 ± 0.07	+0.20 ± 0.16	+0.34 ± 0.06	+0.35 ± 0.07	+0.31 ± 0.06	+0.46 ± 0.07
852	-0.15 ± 0.04	-0.99 ± 0.32	-0.69 ± 0.19	-0.16 ± 0.08	-0.10 ± 0.15	+0.29 ± 0.13	+0.26 ± 0.09	+0.35 ± 0.06	+0.53 ± 0.12
885	-0.25 ± 0.08	-0.87 ± 0.15	-	-	-0.02 ± 0.20	+0.20 ± 0.09	+0.23 ± 0.11	-	-
1384	-0.26 ± 0.03	-0.83 ± 0.15	-0.66 ± 0.15	-0.30 ± 0.06	-0.02 ± 0.13	+0.34 ± 0.06	+0.24 ± 0.06	+0.27 ± 0.05	+0.57 ± 0.07
1633	-0.22 ± 0.03	-	-0.21 ± 0.08	-0.21 ± 0.08	+0.06 ± 0.16	+0.21 ± 0.10	+0.30 ± 0.07	+0.27 ± 0.08	+0.44 ± 0.10

Table 5. Average abundance ratios for the SMC GCs, together with the error of the mean, the typical uncertainty of individual stars and the number of stars.

	NGC 121				NGC 339				NGC 419			
	$\langle \rangle$	σ_{int}	σ_{obs}	N_{\star}	$\langle \rangle$	σ_{int}	σ_{obs}	N_{\star}	$\langle \rangle$	σ_{int}	σ_{obs}	N_{\star}
[Fe/H]	-1.18 ± 0.02	0.00 ± 0.06	0.05	5	-1.24 ± 0.01	0.00 ± 0.02	0.02	7	-0.58 ± 0.02	0.00 ± 0.03	0.04	8
[Fe II/H]	-1.24 ± 0.04	0.00 ± 0.09	0.11	5	-1.34 ± 0.03	0.00 ± 0.04	0.05	4	-0.68 ± 0.04	0.00 ± 0.05	0.06	5
[Si/Fe]	+0.09 ± 0.04	0.00 ± 0.06	0.08	5	+0.16 ± 0.02	0.00 ± 0.03	0.06	7	+0.02 ± 0.03	0.00 ± 0.04	0.06	8
[Ca/Fe]	+0.04 ± 0.02	0.00 ± 0.03	0.03	5	+0.14 ± 0.02	0.00 ± 0.04	0.08	7	-0.03 ± 0.03	0.00 ± 0.05	0.09	8
[Sc II/Fe]	-0.20 ± 0.03	0.00 ± 0.04	0.04	5	-0.29 ± 0.03	0.00 ± 0.10	0.11	7	-0.22 ± 0.03	0.00 ± 0.10	0.07	8
[Ti/Fe]	+0.04 ± 0.03	0.00 ± 0.03	0.02	5	-0.01 ± 0.02	0.00 ± 0.10	0.05	4	-0.10 ± 0.04	0.00 ± 0.06	0.09	5
[V/Fe]	-0.20 ± 0.03	0.00 ± 0.06	0.07	5	-0.30 ± 0.03	0.03 ± 0.05	0.08	7	-0.38 ± 0.04	0.00 ± 0.07	0.10	8
[Cr/Fe]	-0.17 ± 0.06	0.00 ± 0.07	0.07	5	-0.13 ± 0.02	0.00 ± 0.04	0.07	7	-0.11 ± 0.03	0.00 ± 0.03	0.04	8
[Mn/Fe]	-0.57 ± 0.04	0.00 ± 0.05	0.06	5	-0.53 ± 0.02	0.00 ± 0.02	0.03	4	-0.54 ± 0.03	0.04 ± 0.04	0.06	5
[Co/Fe]	-0.13 ± 0.03	0.00 ± 0.03	0.03	5	-0.10 ± 0.02	0.00 ± 0.08	0.04	7	-0.23 ± 0.02	0.00 ± 0.06	0.09	8
[Ni/Fe]	-0.15 ± 0.02	0.01 ± 0.04	0.04	5	-0.18 ± 0.01	0.02 ± 0.01	0.03	7	-0.22 ± 0.02	0.03 ± 0.02	0.07	8
[Cu/Fe]	-0.98 ± 0.13	0.00 ± 0.16	0.16	3	-0.66 ± 0.03	0.00 ± 0.06	0.07	7	-0.76 ± 0.06	0.00 ± 0.11	0.13	6
[Zn/Fe]	—	—	—	—	-0.29 ± 0.05	0.00 ± 0.06	0.07	4	-0.68 ± 0.08	0.00 ± 0.08	0.02	4
[Y/Fe]	-0.29 ± 0.11	0.00 ± 0.30	0.21	2	+0.23 ± 0.03	0.00 ± 0.03	0.08	4	-0.23 ± 0.04	0.04 ± 0.05	0.09	5
[Zr/Fe]	-0.04 ± 0.07	0.00 ± 0.09	0.12	5	+0.01 ± 0.04	0.00 ± 0.04	0.05	7	+0.03 ± 0.06	0.00 ± 0.07	0.09	8
[Ba II/Fe]	-0.10 ± 0.04	0.00 ± 0.07	0.08	5	+0.17 ± 0.03	0.02 ± 0.08	0.09	7	+0.30 ± 0.03	0.00 ± 0.04	0.05	8
[La II/Fe]	-0.01 ± 0.04	0.00 ± 0.05	0.05	5	+0.19 ± 0.03	0.00 ± 0.03	0.10	7	+0.29 ± 0.03	0.00 ± 0.04	0.06	8
[Nd II/Fe]	+0.58 ± 0.04	0.00 ± 0.10	0.09	5	+0.33 ± 0.03	0.04 ± 0.04	0.07	4	+0.32 ± 0.03	0.00 ± 0.12	0.06	5
[Eu II/Fe]	+0.47 ± 0.05	0.00 ± 0.06	0.07	5	+0.67 ± 0.04	0.00 ± 0.05	0.04	4	+0.51 ± 0.04	0.00 ± 0.05	0.06	5

by DAOSPEC or, for abundances from spectral synthesis, we rely on Montecarlo simulations (see Minelli et al. 2021, for details). The uncertainties arising from the atmospheric parameters have been computed deriving the abundance variation due to the change of only one parameter at a time, keeping the others fixed, except for T_{eff} which affects the values derived for $\log g$ and v_t that had to be changed accordingly.

The $\log g$ value depends on T_{eff} and on the isochrone adopted. Uncertainties related to the choice of a specific isochrone are derived from the age and metallicity values adopted for each cluster. In particular, we compute the variation of $\log g$ determined by a change of 1 Gyr in age and 0.1 dex in metallicity, which are the typical errors associated with these parameters. Specifically, variations in metallicity mainly affect the typical error associated with $\log g$ measurements, which is of the order of 0.07 dex.

Finally, typical errors associated with v_t measurements are again different for GIRAFFE and UVES spectra, depending on the method used to derive their values. For UVES spectra, v_t is derived spectroscopically and the typical uncertainties are of the order of 0.1 km s⁻¹. For GIRAFFE targets v_t values have been derived from the $\log g$ - v_t relation by Mucciarelli & Bonifacio (2020) and the adopted uncertainty is of 0.15 km s⁻¹, taking into account the errors in $\log g$ and in the adopted calibration.

5. [Fe/H] abundances and RVs

Here we discuss iron abundances and the systemic RVs for the three target clusters, comparing them with estimates from literature. For each cluster we provide the average [Fe/H] and RV together with the standard error of the average and the observed standard deviation (σ_{obs}).

For NGC 121 we derive an iron abundance of [Fe/H] = -1.18 ± 0.02 (σ_{obs} = 0.05) dex. Da Costa & Hatzidimitriou (1998) derived for this cluster [Fe/H] = -1.19 ± 0.12 dex using Ca II triplet. The same high-resolution dataset discussed here has been previously analysed by Dalessandro et al. (2016). They find an average [Fe/H] of -1.28 ± 0.03 dex (σ = 0.06 dex). The small difference between the two values is due to the differences in T_{eff} . Indeed, the average difference between the T_{eff} values derived in this study and those by Dalessandro et al. (2016) is +122 K, reflecting the difference in the zero-point of the adopted T_{eff}

scales – Alonso, Arribas & Martinez-Roger (1999) in Dalessandro et al. (2016) and González Hernández, & Bonifacio (2009) in this study. The mean RV of this cluster is +144.7 ± 0.9 km s⁻¹ (σ_{obs} = 1.9 km s⁻¹), in good agreement with previous estimates by low-resolution spectra. RVs measured from integrated spectra of NGC 121 provide +139 ± 20 km s⁻¹ (Zinn & West 1984), +138 ± 15 km s⁻¹ (Hesser et al. 1986) and +147 ± 2 km s⁻¹ (Dubath et al. 1997), while measures based on Ca II triplet lines of individual stars provide +138 ± 4 km s⁻¹ (Da Costa & Hatzidimitriou 1998).

The intermediate-age GC NGC 339 has a metallicity similar to that of NGC 121, with [Fe/H] = -1.24 ± 0.01 (σ_{obs} = 0.02) dex. This value is consistent with the ones obtained from the Ca II triplet, [Fe/H] = -1.19 ± 0.10 dex (Da Costa & Hatzidimitriou 1998) and photometry, [Fe/H] = -1.10 ± 0.12 dex (Narloch et al. 2021). The mean RV of NGC 339 is +113.4 ± 0.7 km s⁻¹ (σ_{obs} = 1.9 km s⁻¹) in excellent agreement with the value by Song et al. (2021) of +112.9 ± 0.4 km s⁻¹ obtained from high-resolution spectra collected with M2FS on the Magellan/Clay Telescope. On the other hand, the average RV provided by Parisi et al. (2022) from low-resolution spectra is ~10 km/s lower than ours. They attribute the similar difference with respect to the Song et al. (2021) estimate to mis-centering of their stars in the slit.

Finally, the intermediate-age GC NGC 419 has a metallicity significantly higher than the other two clusters ([Fe/H] = -0.58 ± 0.02 dex, σ_{obs} = 0.04 dex). For this cluster Parisi et al. (2022) provided [Fe/H] = -0.62 ± 0.02 dex from Ca II triplet. Its mean RV is +188.2 ± 0.9 km s⁻¹ (σ_{obs} = 2.5 km s⁻¹). Similar to NGC 339, we have a good agreement with the measure by Song et al. (2021) providing 189.9 ± 0.3 km s⁻¹, while the value by Parisi et al. (2022) is lower by ~17 km s⁻¹.

6. Age-metallicity relation

The left panel of Fig. 2 shows the behaviour of the measured [Fe/H] for the three target GCs as a function of the age. Age estimates are from Glatt et al. (2008a,b). Thanks to NGC 121, the metallicity reached by the SMC ~2–3 Gyr after the earliest SF bursts may be precisely determined. Its metallicity ([Fe/H] = -1.18 dex) is lower than that reached by the LMC (Pagel & Tautvaisiene 1998; Harris & Zaritsky 2009) and MW (Haywood et




Fig. 2. Age-metallicity relation of the SMC clusters. Left panel: average metallicity as a function of the age (Glatt et al. 2009) for the three SMC GCs analysed in this work (same colour code of Fig. 1). SMC GCs from previous low-resolution works (Parisi et al. 2009, 2015; Dias et al. 2021; Parisi et al. 2022) are plotted as open circles. Theoretical AMRs calculated by Pagel & Tautvaišienė (1998) are also reported (the solid curve is the bursting model and the dashed line the continuous model). Right panel: metallicity distribution of the SMC field stars (Paper I).

al. 2013; Snaith et al. 2015) at the same epoch, as expected for a galaxy characterised by a lower SF efficiency. In Section 8 we compare in details the properties of NGC 121 with the empirical AMR defined by the MW clusters.

The $[\text{Fe}/\text{H}]$ distribution of the SMC field stars analysed in Paper I is displayed in the right panel of Fig. 2. The comparison between this $[\text{Fe}/\text{H}]$ distribution and the metallicity reached by the SMC at the different ages of the three target GCs allows us to constrain the age of the most metal-poor SMC field stars. In fact, it is likely that SMC field stars with $[\text{Fe}/\text{H}] < -1.4/-1.5$ dex (considerably lower than that of NGC 121) were created in the early epochs of the galaxy, showing its early enrichment (Nidever et al. 2020; Reggiani et al. 2021, Paper I).

Despite the large difference in their ages (~ 4 -5 Gyr), NGC 121 and NGC 339 have similar $[\text{Fe}/\text{H}]$, hinting to a very low SF efficiency at those epochs that leads to an almost constant $[\text{Fe}/\text{H}]$ over a large age range, or, perhaps, the effects of outflows. However, it is worth noting that the population of SMC GCs coeval to NGC 339 exhibits a significant spread in $[\text{Fe}/\text{H}]$ as estimated using the CaII triplet strength (Parisi et al. 2009, 2015; Dias et al. 2021; Parisi et al. 2022). This $[\text{Fe}/\text{H}]$ dispersion can be only partially explained by the metallicity gradient inside the SMC (see e.g. Carrera et al. 2008; Parisi et al. 2016; Grady et al. 2021, Paper I), and it may be indicative that different regions of the SMC could be characterised by different chemical enrichment histories and AMR (as already suggested by Dias et al. 2016, and Paper I). The metallicity of NGC 339 only is not sufficient to disentangle the origin of the metallicity spread at those ages, but it could suggest a low SF efficiency in the South-

ern Bridge (the region where NGC 339 is located) with respect to other regions of the SMC.

Finally, NGC 419 has a significant higher $[\text{Fe}/\text{H}]$ abundance likely due to the sequence of intense bursts of SF occurring in the last 3 Gyr (Harris & Zaritsky 2009; Rubele et al. 2012; Massana et al. 2022), confirming that the chemical enrichment of the SMC continues at a quicker pace in the last Gyr, increasing $[\text{Fe}/\text{H}]$ up to ~ -0.6 dex.

In Fig. 2 we compare our results with the two theoretical AMR trends computed by Pagel & Tautvaišienė (1998), the first one assuming a closed-box model with a continuous SF rate and the second one with a long period without SF (or characterised by a very low-level activity) between 12 and 4 Gyr ago and a SF burst 4 Gyr ago. Other AMR have been proposed in the literature, both theoretical (Tsujiimoto & Bekki 2009) and based on observed SF histories (Harris & Zaritsky 2004; Cignoni et al. 2013) and that are consistent with or intermediate between the two models by Pagel & Tautvaišienė (1998). The two models by Pagel & Tautvaišienė (1998) are similar with each other at the ages of NGC 121 and NGC 419 while they clearly disagree at intermediate ages (corresponding to the age of NGC 339 and where a large $[\text{Fe}/\text{H}]$ scatter among GCs has been observed). This highlights the importance to derive with high-resolution spectroscopy the chemical composition of other clusters coeval to NGC 339.

7. Abundance ratios

We discuss the abundance ratios of the three target GCs for α (Si, Ca, Ti), iron-peak (Sc, V, Cr, Mn, Co, Ni, Cu, Zn), s-process (Y, Zr, Ba, La, Nd) and r-process (Eu) elements. For all the abundances ratios discussed in this section, the observed scatters are compatible with null intrinsic scatter (see Table 5). Figs. 3 - 5 show the individual abundance ratios as a function of $[\text{Fe}/\text{H}]$ for the SMC GCs, the MW GCs of the control sample and the SMC field stars from Paper I. It is worth noting that we consider the stars of Paper I regardless of their position within the galaxy. However, differences in the metallicity distribution of different regions, as well as some small systematic differences in the abundance patterns of some elements have been detected, enforcing the idea that the SMC has been characterised by a spatially non-uniform chemical enrichment history. In addition, also literature abundances from high-resolution spectra for MW field stars are shown as reference. Mean abundance ratios for the MW control sample are listed in Appendix A.

7.1. α -elements

Fig. 3 shows the abundance ratios of the explosive α -elements Si, Ca, Ti and their average abundance. These elements are mainly produced in massive stars and ejected in the interstellar medium via CC-SNe, with a small but not negligible contribution by SNe Ia, especially for Ca and Ti (see e.g. Kobayashi et al. 2020).

The abundance ratios for the SMC GCs nicely agree with those of the SMC field stars. All the clusters display $[\alpha/\text{Fe}]$ lower than those measured in MW (field and GCs) stars of similar metallicity. In particular, NGC 121 has nearly solar-scaled $[\alpha/\text{Fe}]$ abundance ratios, pointing out that at that age (~ 10.5 Gyr) the SMC has been already enriched by SN Ia (see also Section 8).

NGC 339 shows only marginally higher (by ~ 0.1 dex) $[\text{Si}/\text{Fe}]$ and $[\text{Ca}/\text{Fe}]$ values and similar $[\text{Ti}/\text{Fe}]$. We can consider that these two clusters have similar $[\alpha/\text{Fe}]$. This is evidence for a poor enrichment both in Fe and α -elements over the large range

of time covered by these two clusters, thus pointing to a low SF rate when the SMC evolved in isolation. Lower $[\alpha/\text{Fe}]$ values are found in NGC 419, demonstrating that also at that age the chemical enrichment is dominated by SN Ia. Overall, the run of $[\alpha/\text{Fe}]$ with $[\text{Fe}/\text{H}]$ for these three GCs agrees with the results from the APOGEE survey (Nidever et al. 2020; Hasselquist et al. 2021; Fernandes et al. 2023).

7.2. Iron-peak elements

Iron-peak elements are mainly produced in CC-SNe (both standard Type II and HNe) with an important contribution by SN Ia for some of them. In particular, Sc, Cu and Zn are almost completely produced by massive stars, V and Co are mainly produced by massive stars but with a small contribution by SN Ia while most of Cr, Mn and Ni are from SN Ia (Romano et al. 2010; Kobayashi et al. 2020; Palla et al. 2020). We stress that abundances of these elements in the SMC are still poorly investigated: apart from the Sc, V, Ni and Cu abundances we derived in Paper I, only abundances of Ni for SMC field stars obtained by the APOGEE survey has been discussed (Hasselquist et al. 2021).

Almost all the iron-peak elements measured in these GCs exhibit abundance ratios lower than those observed in MW stars. In particular, these differences are more pronounced in those elements produced mainly in massive stars, like Sc, V, and Zn. NGC 419 generally shows the most pronounced discrepancies, with differences reaching -0.8 and -0.7 dex for $[\text{Cu}/\text{Fe}]$ and $[\text{Zn}/\text{Fe}]$, respectively. In the following we discuss in details the abundances of three key iron-peak elements that are crucial to constrain the chemical enrichment history of the SMC, namely Mn (produced almost totally in SN Ia), Cu (produced via s-process in massive stars) and Zn (almost totally built by HNe).

In our Galaxy $[\text{Mn}/\text{Fe}]$ is sub-solar until $[\text{Fe}/\text{H}] \sim -1$ dex because at these metallicities it is produced only by CC-SNe and in a small amount. At higher metallicities, $[\text{Mn}/\text{Fe}]$ significantly increases due to the dominant contribution by SNe Ia. The Mn yields increase with the metallicity of the SN Ia progenitor and are also sensitive to the explosion mechanism of the SNe-Ia (Badenes et al. 2008). This behaviour is well reproduced by our MW control sample (see also Sobeck et al. 2006). Concerning the SMC, NGC 121 and NGC 339 have $[\text{Mn}/\text{Fe}]$ values compatible with those of the MW control sample, indicating a similar low production of Mn at early times. At higher metallicities, NGC 419 has a $[\text{Mn}/\text{Fe}]$ comparable with the other two GCs and in stark disagreement with the MW stars of similar $[\text{Fe}/\text{H}]$. This suggests that at these ages the enrichment of Mn has been dominated by metal-poor SN Ia (or at least more-metal poor than those dominating the Mn production in our Galaxy).

Copper is mainly produced in massive stars via the weak s-process. In our Galaxy $[\text{Cu}/\text{Fe}]$ increases with the metallicity reflecting the metallicity dependence of the Cu yields (Romano & Matteucci 2007). As discussed in Paper I, the SMC field stars exhibit $[\text{Cu}/\text{Fe}]$ lower than the MW stars, despite the large star-to-star scatter limits our conclusions about the average trend of $[\text{Cu}/\text{Fe}]$ with $[\text{Fe}/\text{H}]$. Concerning the SMC clusters, NGC 121 and NGC 419 have $[\text{Cu}/\text{Fe}]$ significantly lower than the MW stars and GCs, while NGC 339 has a value barely consistent with the MW control sample but still compatible also with the SMC field stars. In general, the low $[\text{Cu}/\text{Fe}]$ values in these clusters, in particular in NGC 419, support a scenario where the contribution by massive stars is significantly reduced in the SMC.

Zinc is produced mainly in HNe, explosions with energy significantly larger than those of standard CC-SNe and related

to stars more massive than $\sim 30\text{-}35 M_{\odot}$ (Kobayashi et al. 2006; Nomoto et al. 2013). Therefore, Zn offers crucial hints regarding the contribution of HNe and high-mass stars. We are able to measure Zn only in NGC 339 and NGC 419, both having $[\text{Zn}/\text{Fe}]$ values lower than the MW stars. In particular, NGC 419 has $[\text{Zn}/\text{Fe}] \approx -0.7$ dex, while the MW stars at similar metallicities have nearly solar abundance ratios. Also this element supports a chemical enrichment in the SMC where the contribution by massive stars (and in particular by HNe) is significantly reduced with respect to the MW. Low $[\text{Zn}/\text{Fe}]$ abundance ratios have been measured in other dwarf galaxies characterised by a low SF efficiency and a small contribution by massive stars, like Sagittarius (Sbordone et al. 2007; Minelli et al. 2021), Sculptor (Skúladóttir et al. 2017) and the LMC (Mucciarelli et al. 2021), confirming that systems less massive than the MW have a lower contribution by HNe.

7.3. Slow neutron-capture elements

The derived abundance ratios for Y, Zr, Ba, La and Nd are reported in Fig. 5 as a function of the metallicity. These elements are mainly produced through s-processes in Asymptotic Giant Branch (AGB) stars. In particular the light s-process elements Y and Zr are produced by AGB stars over a large mass range, while the heavy s-process elements Ba, La and Nd are mainly produced by AGB stars with masses lower than $4 M_{\odot}$. The ratio of heavy to light s-process elements is sensitive to the metallicity (Gallino et al. 1998; Busso et al. 1999): increasing the metallicity of the AGB stars, the production of light s-process elements is favoured against the heavy ones.

The SMC GCs have similar $[\text{Y}/\text{Fe}]$ and $[\text{Zr}/\text{Fe}]$, consistent with the MW stars, suggesting a similar enrichment by high-metallicity and/or intermediate-mass AGB stars. Interesting differences are observed for Ba and La (Nd is known to be equally produced by s- and r-processes). In fact, NGC 121 has $[\text{Ba}/\text{Fe}]$ and $[\text{La}/\text{Fe}]$ slightly lower than the MW GCs, while NGC 339 has $[\text{Ba}/\text{Fe}]$ and $[\text{La}/\text{Fe}]$ ratios higher by ~ 0.3 dex than MW clusters at similar $[\text{Fe}/\text{H}]$. NGC 419 has enhanced values for both the abundance ratios.

We use the $[(\text{Ba}+\text{La})/(\text{Y}+\text{Zr})]$ abundance ratio to evaluate the relative contribution of heavy to light s-process elements. This ratio increases with decreasing the cluster age, from $\sim +0.1$ dex in NGC 121 up to $+0.4$ dex in NGC 419. This indicates that at younger ages the production of these elements is dominated by metal-poor AGB stars that produce especially heavy s-process elements, as expected.

7.4. Rapid neutron-capture elements

The r-processes occur in the presence of a very high neutron flux ($> 10^{20}$ neutron/cm³). Several sites of production have been proposed, as neutron star mergers (Lattimer & Schramm 1974), collapsars (Siegel et al. 2019) and proto-magnetars (Nishimura et al. 2015). The detection of r-process elements in the kilonovae event associated to the gravitational wave GW170817 (Abbott et al. 2017) confirmed that neutron star mergers are an important r-process production site, even if the time delay distribution for the merging process remains unknown. However, chemical evolution models including neutron star mergers as the only r-process source are not able to reproduce the Galactic $[\text{Eu}/\text{Fe}]$ distribution (see e.g. Molero et al. 2021).

We derived abundances for the pure r-process element Eu. The only measures of Eu abundances in the SMC available so far




Fig. 3. α -elements abundance ratios (Si, Ca, Ti and their mean value, from top left panel to bottom right panel, respectively) as a function of $[\text{Fe}/\text{H}]$ for SMC GCs (coloured squares: NGC 121 in red, NGC 339 in green and NGC 419 in blue), SMC field stars (light blue open dots, Mucciarelli et al. 2021), MW GCs from the control sample (grey squares) and MW field stars (grey dots). Arrows indicate upper limits. MW field stars are from Edvardsson et al. (1993); Fulbright (2000); Stephens & Boesgaard (2002); Gratton et al. (2003); Reddy et al. (2003, 2006); Barklem et al. (2005); Bensby et al. (2005); Adibekyan et al. (2012); Roederer & Lawler (2012); Mishenina et al. (2013); Reggiani et al. (2017). The abundance errors for the SMC GCs are reported as error bars, where the thin bar is the typical total uncertainty for individual stars and the thick bar is the error of the mean value of the GC.

are provided by Reggiani et al. (2021), who measured $[\text{Eu}/\text{Fe}]$ in two metal-poor ($[\text{Fe}/\text{H}] < -2.0$ dex) SMC field stars. These two stars are characterised by high value of $[\text{Eu}/\text{Fe}]$ ($\sim +0.8/+1.0$ dex), similar to those measured in LMC and MW stars of similar metallicity. The three SMC GCs discussed here display enhanced $[\text{Eu}/\text{Fe}]$ values compatible with those of the MW GCs control sample. This finding suggests that the production of r-process is very efficient in the SMC over a large range of ages, in particular it continues also at an age of ~ 1.5 Gyr (the age of NGC 419) where the contribution of SNe Ia is significant, as demonstrated by its low $[\alpha/\text{Fe}]$ ratios. The consistent $[\text{Eu}/\text{Fe}]$ at various $[\text{Fe}/\text{H}]$ ratios seems to indicate that the timescales for the synthesis of Eu and Fe (the production of the latter being dominated by SNe Ia), are fairly similar. The measured $[\text{Eu}/\text{Fe}]$ ratios thus provide a crucial constraint to the timescale of Eu production and seem to suggest a dominant contribution from neutron star mergers. This issue will be further analysed by us by means of detailed chemical evolution models (D. Romano et al., in prep.).

7.5. Relative contribution of r- and s-processes

The simultaneous measure of s-process elements produced by AGB stars and the pure r-process element Eu allows us to study

the contribution of the r-process at different ages. In fact, because elements like Ba and La are mainly produced by AGB stars, this contribution starts to dominate the enrichment with a delay after the onset of the SF. Therefore, at low metallicity, the production of Ba and La is dominated by the r-processes. The ratio $[(\text{Ba}+\text{La})/\text{Eu}]$ provides the relative contribution of these two processes. NGC 121 and NGC 339 have similar $[(\text{Ba}+\text{La})/\text{Eu}]$ ratios (~ -0.5 dex), close to the theoretical values predicted in the case of pure r-process (see e.g. Bisterzo et al. 2014). This finding suggests that the production of Ba and La in these clusters is still dominated by the r-processes and the contribution by AGB stars is negligible. However, in NGC 419, this ratio climbs to -0.2 dex, indicating a rising contribution from AGB stars. A similar increase of the ratio between s-process and Eu with the metallicity has been observed in other dwarf galaxies (see e.g. Van der Swaelmen et al. 2013; McWilliam et al. 2013; Hill et al. 2019; Minelli et al. 2021).

8. Comparison between NGC 121, Milky Way in-situ and Gaia-Enceladus clusters

The old cluster NGC 121 allows us to compare the chemical properties of the SMC in its early stages with those of MW GCs of similar ages. The SMC reached an iron content of $[\text{Fe}/\text{H}] \sim$




Fig. 4. Iron-peak elements abundance ratios as a function of $[\text{Fe}/\text{H}]$, for SMC and MW GCs and field stars. Same symbols and data as in figure 3. MW field stars data are from Fulbright (2000); Stephens & Boesgaard (2002); Gratton et al. (2003); Reddy et al. (2003, 2006); Bensby et al. (2005); Adibekyan et al. (2012); Roederer & Lawler (2012); Reggiani et al. (2017).

1.2 dex at an age of ~ 10.5 Gyr. We compare the properties (age, $[\text{Fe}/\text{H}]$ and $[\alpha/\text{Fe}]$) of NGC 121 with those of the MW in-situ GCs and those associated to the main merger event of the MW, named Gaia-Enceladus (G-E, Helmi et al. 2018). It has been suggested that the stellar mass of the progenitor of G-E should be comparable with the present-day stellar mass of the SMC, $\sim 5 - 6 \times 10^8 M_{\odot}$ (van der Marel et al. 2009; Helmi et al. 2018).

We consider the trends of age versus metallicity for the MW in-situ and G-E clusters adopting the dynamical selection by Massari et al. (2019). The upper panel of Fig. 6 shows the behaviour of age from VandenBerg et al. (2013) as a function of $[\text{Fe}/\text{H}]$ for the MW in-situ and G-E clusters, in comparison with NGC 121. For $[\text{Fe}/\text{H}] > -1.5$ dex, the MW in-situ and G-E clusters define two well-distinct sequences in the age- $[\text{Fe}/\text{H}]$ plane. It is remarkable that the MW in-situ GCs at a comparable age of that of NGC 121 are significantly more metal-rich, reaching $[\text{Fe}/\text{H}] \sim -0.5$ dex. On the other hand, NGC 121 well matches with the AMR described by G-E GCs. Yet, despite the consistency of NGC 121 with the G-E AMR might suggest that its former progenitor (the SMC) followed an evolution similar to that of G-E, a significant difference arises when considering their α elements, in the form of $[(\text{Si}+\text{Ca})/\text{Fe}]$ ratios (we consider only these two α -elements because they are available for a large number of GCs, while Ti abundances are not available for some of them). The lower panel of Fig. 6 shows the behaviour of $[(\text{Si}+\text{Ca})/\text{Fe}]$ as a function of $[\text{Fe}/\text{H}]$ for MW in-situ, G-E clusters and for NGC 121. The clusters associated to G-E, despite starting to deviate from the sequence of the MW in-situ GCs at lower $[\text{Fe}/\text{H}]$, still show values of $[(\text{Si}+\text{Ca})/\text{Fe}] \sim 0.2$ dex at the metallicity of NGC 121. This cluster, instead, shows a nearly solar value, indicating a formation environment already dominated by SNIa enrichment, at variance with the G-E clusters for which SN Ia have just started to pollute their CC-SNe dominated star-forming gas. This comparison provides robust evidence for distinct chemical evolution histories between the SMC and GE, with the SMC having experienced a slower and less ef-

ficient star formation compared to G-E. This is consistent with the fact that the SMC mass at ~ 10.5 Gyr was likely significantly lower than that of G-E. A similar result has been suggested also by Hasselquist et al. (2021) and Fernandes et al. (2023) based on the chemical patterns of SMC and G-E field stars (hence without taking the stellar ages into account, which can be done via GCs). It should be emphasised that the interaction of G-E with the MW during the cannibalisation process might have triggered a SF burst and, hence, higher SF rates than experienced by the SMC during its quiet evolution in isolation. A sudden SF burst is expected to restore the $[\alpha/\text{Fe}]$ ratios to values more typical of those of CC-SN ejecta.

9. Summary

This study provides a time-resolved reconstruction of the chemical enrichment history of the SMC galaxy by presenting for the first time the detailed chemical composition based on high-resolution spectra of three clusters in a wide range interval. The following is a summary of the main findings:

- We can place constraints on the metallicity that the SMC attained at certain ages using the derived $[\text{Fe}/\text{H}]$ of the three GCs. The relationship between $[\text{Fe}/\text{H}]$ and cluster age is in excellent accord with the theoretical AMR proposed by Pagel & Tautvaisiene (1998), which incorporated a sizeable halt in star formation between the first burst and the subsequent one at 4 Gyr. However, given the wide metallicity variance seen at any age (see e.g. Parisi et al. 2022), our sample of clusters should not be seen as being as indicative of the entire SMC. This spread may have been influenced by the SMC’s metallicity gradient and the existence of varied chemical enrichment histories in different SMC sites (as originally mentioned in Paper I), which reflect turbulent interactions between the two Clouds.




Fig. 5. Neutron-capture elements abundance ratios (Y, Zr, Ba, La, Nd and Eu, from top left panel to bottom right panel, respectively) as a function of $[Fe/H]$, for SMC and MW GCs and field stars. Same symbols and data as in figure 3. MW field stars data are from Edvardsson et al. (1993); Burris et al. (2000); Fulbright (2000); Stephens & Boesgaard (2002); Reddy et al. (2003, 2006); Barklem et al. (2005); Bensby et al. (2005); Roederer & Lawler (2012); Mishenina et al. (2013); Yan et al. (2015); Battistini & Bensby (2016); Reggiani et al. (2017); Forsberg et al. (2019).




Fig. 6. Behaviour of age and $[(\text{Si}+\text{Ca})/\text{Fe}]$ as a function of $[\text{Fe}/\text{H}]$ (upper and lower panel, respectively) for the MW in-situ GCs (grey diamonds), from Gaia-Enceladus (green diamonds) and for NGC 121 (red square). For MW GCs ages are from Vandenberg et al. (2013). The chemical abundances are from the MW control sample of this study and from Marino et al. (2011); Carretta et al. (2013); Muñoz et al. (2013); Carretta et al. (2014); Koch & McWilliam (2014); Marino et al. (2015); San Roman et al. (2015); Johnson et al. (2016); Rojas-Arriagada et al. (2016); Villanova et al. (2016); Massari et al. (2017); Mura-Guzmán et al. (2018); Masseron et al. (2019); Koch-Hansen et al. (2021); Marino et al. (2021). Dynamical association of the GCs are from Massari et al. (2019). In the lower panel, $[(\text{Si}+\text{Ca})/\text{Fe}]$ from the APOGEE results from 17th Data Release of the Sloan Digital Sky Survey (Abdurro'uf et al. 2022) are shown for the sake of comparison, considering MW giant stars within 2 kpc (grey points) and Gaia-Enceladus stars (light green circles) selected according to Hasselquist et al. (2021).

- We derived the abundance ratios for elements belonging to the main groups (α , iron-peak and neutron-capture) and we compared them with the abundances of SMC field stars (Paper I). The two samples have similar (homogenous) abundances in all the elements, indicating that GCs and field stars experienced a similar chemical enrichment. The comparison between abundances of SMC and MW GCs highlights their different chemical enrichment histories. In particular, the enrichment of the SMC (at least in the age range explored in this study) is dominated by SN Ia and displays a lower contribution by massive stars.
- Despite having an age difference of about 4-5 Gyr, the ancient cluster NGC 121 and the intermediate-age cluster NGC 339 exhibit identical $[\text{Fe}/\text{H}]$ values and similar abundances for the majority of the measured elements. Differ-

ences of about 0.2-0.3 dex are observed only for heavy s-process elements. These results suggest a similar chemical enrichment by CC-SNe and SN Ia but an increase of the contribution of the metal-poor AGB stars with the age.

- The young (1.4 Gyr) cluster NGC 419 has chemical patterns that are markedly different from those of the other GCs, including higher $[\text{Fe}/\text{H}]$, lower $[\alpha/\text{Fe}]$ and $[\text{Zn}/\text{Fe}]$ ratios, and higher heavy s-process elements. These abundances show that the metal-poor AGB stars and SN Ia made a significant contribution to the evolution of the SMC at that age, although HNe made a less significant contribution, as shown by the extremely low $[\text{Zn}/\text{Fe}]$ ratio. It is interesting to note that $[\text{Eu}/\text{Fe}]$ in this cluster is enhanced, indicating that the production of r-process in the SMC is extremely efficient throughout a wide range of ages, including at an age of around 1.5 Gyr where the contribution of SNe Ia is considerable as evidenced by its low $[\alpha/\text{Fe}]$ ratios.
- We contrasted the characteristics of the old SMC cluster NGC 121 with those of MW clusters formed in-situ and clusters connected to the G-E merger event. The early SMC and G-E have both attained the same metallicity at the same age as NGC 121, but their $[\alpha/\text{Fe}]$ ratios are different, with the latter displaying enhanced values in contrast to NGC 121, which has solar-scaled $[\alpha/\text{Fe}]$ ratios. This shows that the SMC and G-E have distinct histories of chemical enrichment, most likely a quiet and a bursty one, respectively, because of the interaction of G-E with the MW as opposed to the evolution in isolation of the SMC. Furthermore, the SMC had at early ages had a mass lower than G-E.

Acknowledgements. We thanks the anonymous referee for the useful and constructive suggestions. This research is funded by the project "Light-on-Dark", granted by the Italian MIUR through contract PRIN-2017K7REXT. C. Lardo acknowledges funding from Ministero dell'Università e della Ricerca through the Programme *Rita Levi Montalcini* (grant PGR18YRML1).

References

- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017, *Phys. Rev. Lett.*, 119, 161101. doi:10.1103/PhysRevLett.119.161101
- Abdurro'uf, Accetta, K., Aerts, C., et al. 2022, *ApJS*, 259, 35. doi:10.3847/1538-4365/ac4414
- Adibekyan, V. Z., Sousa, S. G., Santos, N. C., et al. 2012, *A&A*, 545, A32. doi:10.1051/0004-6361/201219401
- Alonso, A., Arribas, S., & Martínez-Roger, C., 1999, *A&As*, 140, 261
- Badenes, C., Bravo, E., & Hughes, J. P. 2008, *ApJ*, 680, L33. doi:10.1086/589832
- Barklem, P. S., Christlieb, N., Beers, T. C., et al. 2005, *A&A*, 439, 129. doi:10.1051/0004-6361:20052967
- Bastian, N. & Lardo, C. 2018, *ARA&A*, 56, 83. doi:10.1146/annurev-astro-081817-051839
- Battistini, C. & Bensby, T. 2016, *A&A*, 586, A49. doi:10.1051/0004-6361/201527385
- Belokurov, V., Erkal, D., Deason, A. J., et al. 2017, *MNRAS*, 466, 4711. doi:10.1093/mnras/stw3357
- Bensby, T., Feltzing, S., Lundström, I., et al. 2005, *A&A*, 433, 185. doi:10.1051/0004-6361:20040332
- Besla, G., Kallivayalil, N., Hernquist, L., et al. 2007, *ApJ*, 668, 949
- Besla, G., Kallivayalil, N., Hernquist, L., et al. 2012, *MNRAS*, 421, 2109. doi:10.1111/j.1365-2966.2012.20466.x
- Bihain, G., Israelian, G., Rebolo, R., et al. 2004, *A&A*, 423, 777. doi:10.1051/0004-6361:20035913
- Bisterzo, S., Travaglio, C., Gallino, R., et al. 2014, *ApJ*, 787, 10. doi:10.1088/0004-637X/787/1/10

- Brocato, E., Castellani, V., Ferraro, F. R., et al. 1996, *MNRAS*, 282, 614. doi:10.1093/mnras/282.2.614
- Burris, D. L., Pilachowski, C. A., Armandroff, T. E., et al. 2000, *ApJ*, 544, 302. doi:10.1086/317172
- Busso, M., Gallino, R., & Wasserburg, G. J. 1999, *ARA&A*, 37, 239. doi:10.1146/annurev.astro.37.1.239
- Byrd, G., Valtonen, M., McCall, M., et al. 1994, *AJ*, 107, 2055
- Carrera, R., Gallart, C., Aparicio, A., et al. 2008, *AJ*, 136, 1039. doi:10.1088/0004-6256/136/3/1039
- Carretta, E., Bragaglia, A., Gratton, R. G., et al. 2013, *A&A*, 557, A138. doi:10.1051/0004-6361/201321905
- Carretta, E., Bragaglia, A., Gratton, R. G., et al. 2014, *A&A*, 564, A60. doi:10.1051/0004-6361/201323321
- Cignoni, M., Cole, A. A., Tosi, M., Gallagher, J. S., Sabbi, E., Anderson, J., Grebel, E. K. & Nota, A., 2013, *ApJ*, 775, 83
- Cioni, M.-R. L., Clementini, G., Girardi, L., et al. 2011, *A&A*, 527, A116. doi:10.1051/0004-6361/201016137
- Cioni, M.-R. L., Storm, J., Bell, C. P. M., et al. 2019, *The Messenger*, 175, 54. doi:10.18727/0722-6691/5128
- Da Costa, G. S. & Hatzidimitriou, D. 1998, *AJ*, 115, 1934. doi:10.1086/300340
- Dalessandro, E., Lapenna, E., Mucciarelli, A., et al. 2016, *ApJ*, 829, 77. doi:10.3847/0004-637X/829/2/77
- Dias, B., Kerber, L., Barbuy, B., et al. 2016, *A&A*, 591, A11. doi:10.1051/0004-6361/201527558
- Dias, B., Angelo, M. S., Oliveira, R. A. P., et al. 2021, *A&A*, 647, L9. doi:10.1051/0004-6361/202040015
- Dias, B., Parisi, M. C., Angelo, M., et al. 2022, *MNRAS*, 512, 4334. doi:10.1093/mnras/stac259
- Diaz, J. & Bekki, K. 2011, *MNRAS*, 413, 2015. doi:10.1111/j.1365-2966.2011.18289.x
- Diaz, J. D. & Bekki, K. 2012, *ApJ*, 750, 36. doi:10.1088/0004-637X/750/1/36
- Dresbach, F., Massari, D., Lanzoni, B., et al. 2022, *ApJ*, 928, 47. doi:10.3847/1538-4357/ac5406
- D'Onghia, E. & Fox, A. J. 2016, *ARA&A*, 54, 363. doi:10.1146/annurev-astro-081915-023251
- Dotter, A., Chaboyer, B., Jevremović, D., et al. 2008, *ApJS*, 178, 89. doi:10.1086/589654
- Dubath, P., Meylan, G., & Mayor, M. 1997, *A&A*, 324, 505
- Edvardsson, B., Andersen, J., Gustafsson, B., et al. 1993, *A&A*, 500, 391
- Fernandes, L., Mason, A. C., Horta, D., et al. 2023, *MNRAS*, 519, 3611. doi:10.1093/mnras/stac3543
- Forsberg, R., Jönsson, H., Ryde, N., et al. 2019, *A&A*, 631, A113. doi:10.1051/0004-6361/201936343
- Fulbright, J. P. 2000, *AJ*, 120, 1841. doi:10.1086/301548
- Gallino, R., Arlandini, C., Busso, M., et al. 1998, *ApJ*, 497, 388. doi:10.1086/305437
- Gatto, M., Ripepi, V., Bellazzini, M., et al. 2022, *A&A*, 664, L12. doi:10.1051/0004-6361/202243993
- Glatt, K., Gallagher, J. S., Grebel, E. K., et al. 2008, *AJ*, 135, 1106. doi:10.1088/0004-6256/135/4/1106
- Glatt, K., Grebel, E. K., Sabbi, E., et al. 2008, *AJ*, 136, 1703. doi:10.1088/0004-6256/136/4/1703
- Glatt, K., Grebel, E. K., Gallagher, J. S., et al. 2009, *AJ*, 138, 1403
- Gonzalez, O. A., Mucciarelli, A., Origlia, L., et al. 2020, *The Messenger*, 180, 18. doi:10.18727/0722-6691/5196
- González Hernández, J. I., & Bonifacio, P. 2009, *A&A*, 497, 497
- Grady, J., Belokurov, V., & Evans, N. W. 2021, *ApJ*, 909, 150. doi:10.3847/1538-4357/abd4e4
- Gratton, R. G., Carretta, E., Claudi, R., et al. 2003, *A&A*, 404, 187. doi:10.1051/0004-6361/20030439
- Gratton, R. G., Carretta, E., & Bragaglia, A. 2012, *A&A Rev.*, 20, 50. doi:10.1007/s00159-012-0050-3
- Grevesse, N., & Sauval, A. J. 1998, *Space Sci. Rev.*, 85, 161
- Johnson, C. I., Caldwell, N., Rich, R. M., et al. 2016, *AJ*, 152, 21. doi:10.3847/0004-6256/152/1/21
- Harris, J. & Zaritsky, D. 2004, *AJ*, 127, 1531. doi:10.1086/381953
- Harris, J., & Zaritsky, D. 2009, *AJ*, 138, 1243
- Hasselquist, S., Hayes, C. R., Lian, J., et al. 2021, *ApJ*, 923, 172. doi:10.3847/1538-4357/ac25f9
- Haywood, M., Di Matteo, P., Lehnert, M. D., et al. 2013, *A&A*, 560, A109. doi:10.1051/0004-6361/201321397
- Helmi, A., Babusiaux, C., Koppelman, H. H., et al. 2018, *Nature*, 563, 85. doi:10.1038/s41586-018-0625-x
- Hesser, J. E., Shawl, S. J., & Meyer, J. E. 1986, *PASP*, 98, 403. doi:10.1086/131771
- Hill, V. 1999, *A&A*, 345, 430
- Hill, V., Skúladóttir, Á., Tolstoy, E., et al. 2019, *A&A*, 626, A15. doi:10.1051/0004-6361/201833950
- Kobayashi, C., Umeda, H., Nomoto, K., et al. 2006, *ApJ*, 653, 1145. doi:10.1086/508914
- Kobayashi, C., Karakas, A. I., & Lugaro, M. 2020, *ApJ*, 900, 179. doi:10.3847/1538-4357/abae65
- Koch, A. & McWilliam, A. 2014, *A&A*, 565, A23. doi:10.1051/0004-6361/201323119
- Koch-Hansen, A. J., Hansen, C. J., & McWilliam, A. 2021, *A&A*, 653, A2. doi:10.1051/0004-6361/202141130
- Kurucz, R. L. 1993, *VizieR Online Data Catalog*, VI/39
- Kurucz, R. L. 2005, *Memorie della Societa Astronomica Italiana Supplementi*, 8, 14
- Lattimer, J. M. & Schramm, D. N. 1974, *ApJ*, 192, L145. doi:10.1086/181612
- Mackey, A. D., Payne, M. J., & Gilmore, G. F. 2006, *MNRAS*, 369, 921. doi:10.1111/j.1365-2966.2006.10342.x
- Maia, F. F. S., Dias, B., Santos, J. F. C., et al. 2019, *MNRAS*, 484, 5702. doi:10.1093/mnras/stz369
- Marino, A. F., Sneden, C., Kraft, R. P., et al. 2011, *A&A*, 532, A8. doi:10.1051/0004-6361/201116546
- Marino, A. F., Milone, A. P., Karakas, A. I., et al. 2015, *MNRAS*, 450, 815. doi:10.1093/mnras/stv420
- Marino, A. F., Milone, A. P., Renzini, A., et al. 2021, *ApJ*, 923, 22. doi:10.3847/1538-4357/ac282c
- Massana, P., Ruiz-Lara, T., Noël, N. E. D., et al. 2022, *MNRAS*, 513, L40. doi:10.1093/mnras/slac030
- Massari, D., Mucciarelli, A., Dalessandro, E., et al. 2017, *MNRAS*, 468, 1249. doi:10.1093/mnras/stx549
- Massari, D., Koppelman, H. H., & Helmi, A. 2019, *A&A*, 630, L4. doi:10.1051/0004-6361/201936135
- Massari, D., Raso, S., Libralato, M., et al. 2021, *MNRAS*, 500, 2012. doi:10.1093/mnras/staa3497
- Masseron, T., García-Hernández, D. A., Mészáros, S., et al. 2019, *A&A*, 622, A191. doi:10.1051/0004-6361/201834550
- McWilliam, A., Wallerstein, G., & Mottini, M. 2013, *ApJ*, 778, 149. doi:10.1088/0004-637X/778/2/149
- Milone, A. P., Piotto, G., Renzini, A., et al. 2017, *MNRAS*, 464, 3636
- Minelli, A., Mucciarelli, A., Romano, D., et al. 2021, *ApJ*, 910, 114. doi:10.3847/1538-4357/abe3f9
- Mishenina, T. V., Pignatari, M., Korotin, S. A., et al. 2013, *A&A*, 552, A128. doi:10.1051/0004-6361/201220687
- Molero, M., Simonetti, P., Matteucci, F., et al. 2021, *MNRAS*, 500, 1071. doi:10.1093/mnras/staa3340
- Mucciarelli, A., Origlia, L., Maraston, C., & Ferraro, F. R., 2009, *ApJ*, 690, 288
- Mucciarelli, A., Bellazzini, M., Ibata, R., et al. 2012, *MNRAS*, 426, 2889. doi:10.1111/j.1365-2966.2012.21847.x
- Mucciarelli, A. 2013, *arXiv e-prints*, arXiv:1311.140
- Mucciarelli, A., Pancino, E., Lovisi, L., et al. 2013, *ApJ*, 766, 78
- Mucciarelli, A., Monaco, L., Bonifacio, P., et al. 2017, *A&A*, 603, L7
- Mucciarelli, A. & Bonifacio, P. 2020, *A&A*, 640, A87. doi:10.1051/0004-6361/202037703
- Mucciarelli, A., Bellazzini, M., & Massari, D. 2021, *A&A*, 653, A90. doi:10.1051/0004-6361/202140979
- Mucciarelli, A., Massari, D., Minelli, A., et al. 2021, *Nature Astronomy*, 5, 1247. doi:10.1038/s41550-021-01493-y
- Mucciarelli, A., Minelli, A., Bellazzini, M., et al. 2023, *A&A*, 671, A124. doi:10.1051/0004-6361/202245133
- Muñoz, C., Geisler, D., & Villanova, S. 2013, *MNRAS*, 433, 2006. doi:10.1093/mnras/stt868
- Mura-Guzmán, A., Villanova, S., Muñoz, C., et al. 2018, *MNRAS*, 474, 4541. doi:10.1093/mnras/stx2918
- Narloch, W., Pietrzyński, G., Gieren, W., et al. 2021, *A&A*, 647, A135. doi:10.1051/0004-6361/202039623

- Nidever, D. L., Olsen, K., Walker, A. R., et al. 2017, *AJ*, 154, 199. doi:10.3847/1538-3881/aa8d1c
- Nidever, D. L., Hesselquist, S., Hayes, C. R., et al. 2020, *ApJ*, 895, 88. doi:10.3847/1538-4357/ab7305
- Nishimura, N., Takiwaki, T., & Thielemann, F.-K. 2015, *ApJ*, 810, 109. doi:10.1088/0004-637X/810/2/109
- Nomoto, K., Kobayashi, C., & Tominaga, N. 2013, *ARA&A*, 51, 457. doi:10.1146/annurev-astro-082812-140956
- Olsen, K. A. G., Hodge, P. W., Mateo, M., et al. 1998, *MNRAS*, 300, 665. doi:10.1046/j.1365-8711.1998.01860.x
- Pagel, B. E. J. & Tautvaisiene, G. 1998, *MNRAS*, 299, 535. doi:10.1046/j.1365-8711.1998.01792.x
- Palla, M., Matteucci, F., Spitoni, E., et al. 2020, *MNRAS*, 498, 1710. doi:10.1093/mnras/staa2437
- Parisi, M. C., Grocholski, A. J., Geisler, D., et al. 2009, *AJ*, 138, 517. doi:10.1088/0004-6256/138/2/517
- Parisi, M. C., Geisler, D., Clariá, J. J., et al. 2015, *AJ*, 149, 154
- Parisi, M. C., Geisler, D., Carraro, G., et al. 2016, *AJ*, 152, 58
- Parisi, M. C., Gramajo, L. V., Geisler, D., et al. 2022, *A&A*, 662, A75. doi:10.1051/0004-6361/202142597
- Pasquini, L., Avila, G., Blecha, A., et al. 2002, *The Messenger*, 110, 1
- Piatti, A. E. 2022, *MNRAS*, 511, L72. doi:10.1093/mnras/511/1/L72
- Pryor, C. & Meylan, G. 1993, *Structure and Dynamics of Globular Clusters*, 50, 357
- Reddy, B. E., Tomkin, J., Lambert, D. L., et al. 2003, *MNRAS*, 340, 304. doi:10.1046/j.1365-8711.2003.06305.x
- Reddy, B. E., Lambert, D. L., & Allende Prieto, C. 2006, *MNRAS*, 367, 1329. doi:10.1111/j.1365-2966.2006.10148.x
- Reggiani, H., Meléndez, J., Kobayashi, C., et al. 2017, *A&A*, 608, A46. doi:10.1051/0004-6361/201730750
- Reggiani, H., Schlaufman, K. C., Casey, A. R., et al. 2021, *AJ*, 162, 229. doi:10.3847/1538-3881/ac1f9a
- Ripepi, V., Cignoni, M., Tosi, M., et al. 2014, *MNRAS*, 442, 1897. doi:10.1093/mnras/stu918
- Roederer, I. U. & Lawler, J. E. 2012, *ApJ*, 750, 76. doi:10.1088/0004-637X/750/1/76
- Rojas-Arriagada, A., Zoccali, M., Vásquez, S., et al. 2016, *A&A*, 587, A95. doi:10.1051/0004-6361/201527351
- Romano, D. & Matteucci, F. 2007, *MNRAS*, 378, L59. doi:10.1111/j.1745-3933.2007.00320.x
- Romano, D., Karakas, A. I., Tosi, M., et al. 2010, *A&A*, 522, A32. doi:10.1051/0004-6361/201014483
- Rubele, S., Kerber, L., & Girardi, L. 2010, *MNRAS*, 403, 1156. doi:10.1111/j.1365-2966.2010.16201.x
- Rubele, S., Kerber, L., Girardi, L., et al. 2012, *A&A*, 537, A106. doi:10.1051/0004-6361/201117863
- San Roman, I., Muñoz, C., Geisler, D., et al. 2015, *A&A*, 579, A6. doi:10.1051/0004-6361/201525722
- Sbordone, L., Bonifacio, P., Buonanno, R., et al. 2007, *A&A*, 465, 815. doi:10.1051/0004-6361:20066385
- Schlafly, E. F. & Finkbeiner, D. P. 2011, *ApJ*, 737, 103
- Shuter, W. L. H. 1992, *ApJ*, 386, 101
- Siegel, D. M., Barnes, J., & Metzger, B. D. 2019, *Nature*, 569, 241. doi:10.1038/s41586-019-1136-0
- Skowron, D. M., Skowron, J., Udalski, A., et al. 2021, *ApJS*, 252, 23. doi:10.3847/1538-4365/abcb81
- Skúladóttir, Á., Tolstoy, E., Salvadori, S., et al. 2017, *A&A*, 606, A71. doi:10.1051/0004-6361/201731158
- Snaith, O., Haywood, M., Di Matteo, P., et al. 2015, *A&A*, 578, A87. doi:10.1051/0004-6361/201424281
- Sobeck, J. S., Ivans, I. I., Simmerer, J. A., et al. 2006, *AJ*, 131, 2949. doi:10.1086/503106
- Song, Y.-Y., Mateo, M., Bailey, J. I., et al. 2021, *MNRAS*, 504, 4160. doi:10.1093/mnras/stab1065
- Stephens, A. & Boesgaard, A. M. 2002, *AJ*, 123, 1647. doi:10.1086/338898
- Stetson, P. B., & Pancino, E. 2008, *PASP*, 120, 1332
- Tsujimoto, T. & Bekki, K. 2009, *ApJ*, 700, L69. doi:10.1088/0004-637X/700/2/L69
- VandenBerg, D. A., Brogaard, K., Leaman, R., et al. 2013, *ApJ*, 775, 134. doi:10.1088/0004-637X/775/2/134
- van der Marel, R. P., Kallivayalil, N., & Besla, G. 2009, *The Magellanic System: Stars, Gas, and Galaxies*, 256, 81. doi:10.1017/S1743921308028299
- Van der Swaelmen, M., Hill, V., Primas, F., et al. 2013, *A&A*, 560, A44. doi:10.1051/0004-6361/201321109
- Villanova, S., Monaco, L., Moni Bidin, C., et al. 2016, *MNRAS*, 460, 2351. doi:10.1093/mnras/stw1146
- Wagner-Kaiser, R., Mackey, D., Sarajedini, A., et al. 2017, *MNRAS*, 471, 3347. doi:10.1093/mnras/stx1702
- Walker, M. G., Mateo, M., Olszewski, E. W., et al. 2006, *AJ*, 131, 2114. doi:10.1086/500193
- Yan, H. L., Shi, J. R., & Zhao, G. 2015, *ApJ*, 802, 36. doi:10.1088/0004-637X/802/1/36
- Zinn, R. & West, M. J. 1984, *ApJS*, 55, 45. doi:10.1086/190947

Appendix A: Chemical abundance ratios for the MW control sample

The MW control sample includes spectra of giant stars of 16 GCs retrieved from the ESO archive. All the spectra have been secured with UVES-FLAMES adopting the Red Arm 580 configuration. We refer to Mucciarelli et al. (2021) for further details on the MW GC sample (i.e. the number of observed stars and the corresponding observative programs) and its analysis.

In tables A.1, A.2, A.3 and A.4 we list for each MW target cluster the average abundance ratios together with the corresponding standard error and the observed dispersion of the mean, σ_{obs} .

Appendix B: Comparison with APOGEE abundances

We compare the average abundance ratios for the three SMC target clusters discussed in this paper with the abundances derived from the APOGEE survey for SMC field stars using H-band high-resolution spectra. We used the results from 17th Data Release of the Sloan Digital Sky Survey (Abdurro'uf et al. 2022), adopting the selection provided by Hasselquist et al. (2021) for the SMC stars. The elements in common between our analysis and APOGEE are the explosive α -elements Si, Ca and Ti, and the iron-peak elements V, Mn and Ni. Fig. B.1 shows the behaviour of the abundance ratios of these six elements as a function of [Fe/H] for the three SMC clusters, the SMC field stars measured in Paper I and the SMC field stars measured by APOGEE.

[Si/Fe], [Ca/Fe] and [Ni/Fe] from APOGEE and FLAMES spectra agree well each other. [Ti/Fe] measured by APOGEE is 0.2-0.3 dex lower than the values obtained with FLAMES and disagree with the average values of the other $[\alpha/Fe]$ measured by APOGEE. We note as the abundances of the three explosive α -elements measured in Paper I are in better agreement each other. This different behaviour is reported also in the APOGEE documentation³ where the Ti abundances in giant stars are classified as "deviant" because the measured trends are discrepant with respect to literature expectations. [V/Fe] exhibits a significant star-to-star scatter in the APOGEE data, larger than that of Paper I. According to the APOGEE documentation, V abundances in giant stars are classified as "less reliable". Finally, [Mn/Fe] is measured by APOGEE but not in Paper I. There is an offset of ~ 0.3 dex between the APOGEE [Mn/Fe] values and those measured in our SMC clusters. Mn abundances by APOGEE are classified as "most reliable" and we interpret this offset as due to a systematic between the optical and near-infrared Mn transitions.

³ <https://www.sdss4.org/dr17/irspec/abundances/>

Table A.1. Milky Way globular clusters: average abundance ratios of iron and α elements (Si, Ca, Ti I and Ti II) together with the corresponding standard error and dispersion of the mean.

Cluster	[Fe/H]	σ_{obs}	[Si/Fe]	σ_{obs}	[Ca/Fe]	σ_{obs}	[Ti/Fe]	σ_{obs}	[Ti II/Fe]	σ_{obs}
NGC 104	-0.75±0.01	0.03	+0.28±0.01	0.03	+0.21±0.02	0.07	+0.33±0.01	0.03	+0.18±0.01	0.04
NGC 288	-1.24±0.01	0.04	+0.33±0.01	0.03	+0.27±0.01	0.03	+0.24±0.01	0.03	+0.26±0.01	0.02
NGC 1851	-1.13±0.01	0.04	+0.25±0.01	0.05	+0.18±0.01	0.05	+0.09±0.01	0.04	+0.25±0.01	0.04
NGC 2808	-1.06±0.02	0.07	+0.26±0.01	0.04	+0.21±0.01	0.02	+0.14±0.01	0.03	+0.18±0.01	0.02
NGC 4590	-2.28±0.01	0.05	+0.35±0.04	0.06	+0.23±0.01	0.04	+0.16±0.01	0.04	+0.23±0.01	0.04
NGC 5634	-1.80±0.02	0.05	+0.29±0.01	0.04	+0.22±0.01	0.03	+0.21±0.01	0.04	+0.35±0.03	0.07
NGC 5824	-1.92±0.02	0.04	+0.36±0.03	0.08	+0.24±0.01	0.02	+0.18±0.02	0.05	+0.19±0.02	0.05
NGC 5904	-1.22±0.01	0.03	+0.29±0.01	0.03	+0.21±0.01	0.03	+0.15±0.01	0.03	+0.25±0.01	0.05
NGC 6093	-1.76±0.01	0.03	+0.35±0.01	0.04	+0.28±0.01	0.03	+0.15±0.01	0.04	+0.34±0.01	0.04
NGC 6397	-2.01±0.01	0.03	+0.37±0.02	0.08	+0.26±0.01	0.03	+0.19±0.01	0.02	+0.22±0.01	0.04
NGC 6752	-1.48±0.01	0.03	+0.29±0.01	0.03	+0.28±0.01	0.02	+0.24±0.01	0.03	+0.22±0.01	0.05
NGC 6809	-1.73±0.01	0.03	+0.26±0.01	0.04	+0.25±0.01	0.03	+0.15±0.01	0.02	+0.28±0.01	0.03
NGC 7078	-2.42±0.02	0.07	+0.47±0.03	0.08	+0.28±0.01	0.02	+0.23±0.02	0.06	+0.29±0.02	0.05
NGC 7099	-2.31±0.01	0.05	+0.45±0.01	0.02	+0.28±0.01	0.03	+0.20±0.01	0.05	+0.26±0.01	0.04

Table A.2. Milky Way globular clusters: average abundance ratios of iron-peak elements (Sc, V, Cr and Mn) together with the corresponding standard error and dispersion of the mean.

Cluster	[Sc II/Fe]	σ_{obs}	[V/Fe]	σ_{obs}	[Cr/Fe]	σ_{obs}	[Mn/Fe]	σ_{obs}
NGC 104	+0.21±0.02	0.06	+0.32±0.02	0.07	-0.03±0.02	0.07	-0.28±0.01	0.04
NGC 288	+0.13±0.01	0.03	-0.01±0.02	0.06	+0.00±0.01	0.03	-0.47±0.01	0.03
NGC 1851	+0.08±0.01	0.06	-0.13±0.02	0.08	-0.14±0.01	0.06	-0.51±0.02	0.04
NGC 1904	+0.06±0.01	0.03	-0.15±0.01	0.05	-0.12±0.01	0.03	-0.56±0.01	0.05
NGC 2808	+0.12±0.01	0.06	-0.04±0.04	0.10	-0.09±0.01	0.05	-0.47±0.01	0.03
NGC 4590	+0.07±0.01	0.04	+0.09±0.02	0.03	-0.21±0.01	0.07	-0.45±0.01	0.04
NGC 5634	+0.01±0.03	0.07	-0.12±0.03	0.06	-0.07±0.01	0.04	-0.48±0.03	0.07
NGC 5824	+0.02±0.03	0.06	-0.07±0.03	0.06	-0.11±0.01	0.01	-0.53±0.03	0.07
NGC 5904	+0.13±0.01	0.03	-0.07±0.01	0.04	-0.08±0.01	0.05	-0.50±0.01	0.03
NGC 6093	+0.10±0.01	0.04	-0.12±0.01	0.04	-0.14±0.01	0.04	-0.54±0.01	0.02
NGC 6397	+0.03±0.01	0.04	+0.03±0.02	0.07	-0.18±0.01	0.03	-0.46±0.03	0.10
NGC 6752	+0.03±0.01	0.03	-0.02±0.01	0.04	-0.05±0.01	0.03	-0.46±0.01	0.03
NGC 6809	+0.04±0.01	0.04	-0.09±0.01	0.03	-0.10±0.01	0.04	-0.57±0.01	0.05
NGC 7078	+0.09±0.02	0.07	-0.06±0.06	0.10	-0.19±0.01	0.05	-0.41±0.03	0.07
NGC 7099	+0.07±0.01	0.04	-0.03±0.02	0.05	-0.22±0.01	0.03	-0.51±0.02	0.06

Table A.3. Milky Way globular clusters: average abundance ratios of iron-peak elements (Co, Ni, Cu and Zn) together with the corresponding standard error and dispersion of the mean.

Cluster	[Co/Fe]	σ_{obs}	[Ni/Fe]	σ_{obs}	[Cu/Fe]	σ_{obs}	[Zn/Fe]	σ_{obs}
NGC 104	+0.24±0.01	0.04	-0.02±0.01	0.02	—	—	-0.03±0.03	0.09
NGC 288	+0.02±0.01	0.03	-0.01±0.01	0.03	-0.24±0.02	0.05	-0.18±0.04	0.14
NGC 1851	-0.05±0.01	0.04	-0.10±0.01	0.02	—	—	+0.05±0.03	0.14
NGC 1904	-0.09±0.02	0.07	-0.11±0.01	0.02	-0.71±0.01	0.04	-0.04±0.02	0.14
NGC 2808	-0.04±0.01	0.04	-0.09±0.01	0.03	-0.37±0.02	0.12	+0.04±0.05	0.17
NGC 4590	+0.12±0.02	0.03	-0.01±0.01	0.05	-0.68±0.02	0.04	+0.07±0.03	0.10
NGC 5634	+0.03±0.01	0.04	+0.03±0.01	0.04	-0.52±0.04	0.11	-0.03±0.05	0.15
NGC 5824	+0.04±0.03	0.08	-0.03±0.01	0.02	-0.60±0.04	0.11	-0.07±0.03	0.07
NGC 5904	-0.05±0.01	0.03	-0.09±0.01	0.02	-0.47±0.02	0.06	-0.02±0.02	0.09
NGC 6093	-0.06±0.01	0.03	-0.07±0.01	0.02	-0.58±0.01	0.03	-0.08±0.02	0.07
NGC 6397	+0.16±0.02	0.04	-0.02±0.01	0.03	-0.73±0.04	0.09	+0.00±0.02	0.06
NGC 6752	+0.04±0.01	0.03	-0.08±0.01	0.02	-0.47±0.01	0.06	-0.02±0.03	0.12
NGC 6809	+0.03±0.02	0.06	+0.00±0.01	0.02	-0.66±0.01	0.05	-0.06±0.01	0.05
NGC 7078	+0.10±0.01	0.02	-0.02±0.02	0.06	-0.66±0.03	0.07	+0.09±0.03	0.12
NGC 7099	+0.23±0.03	0.08	-0.02±0.01	0.03	-0.73±0.03	0.10	+0.08±0.02	0.08

Table A.4. Milky Way globular clusters: average abundance ratios of neutron-capture elements (Y, Zr, Ba, La, Nd and Eu) together with the corresponding standard error and dispersion of the mean.

Cluster	[Y II/Fe]	σ_{obs}	[Zr/Fe]	σ_{obs}	[Ba II/Fe]	σ_{obs}	[La II/Fe]	σ_{obs}	[Nd II/Fe]	σ_{obs}	[Eu II/Fe]	σ_{obs}
NGC 104	-0.33 ± 0.04	0.11	$+0.23\pm 0.05$	0.15	$+0.15\pm 0.01$	0.03	$+0.15\pm 0.02$	0.07	$+0.08\pm 0.01$	0.05	$+0.44\pm 0.01$	0.04
NGC 288	-0.02 ± 0.03	0.10	$+0.23\pm 0.03$	0.14	$+0.38\pm 0.02$	0.07	$+0.31\pm 0.01$	0.02	$+0.19\pm 0.01$	0.03	$+0.53\pm 0.02$	0.06
NGC 1851	-0.15 ± 0.03	0.13	-0.12 ± 0.03	0.13	$+0.57\pm 0.03$	0.12	$+0.41\pm 0.03$	0.17	$+0.33\pm 0.02$	0.12	$+0.73\pm 0.01$	0.05
NGC 1904	-0.31 ± 0.02	0.05	$+0.15\pm 0.04$	0.13	$+0.22\pm 0.02$	0.05	$+0.24\pm 0.03$	0.09	$+0.09\pm 0.02$	0.07	$+0.55\pm 0.04$	0.10
NGC 2808	-0.25 ± 0.03	0.11	$+0.15\pm 0.06$	0.15	$+0.19\pm 0.03$	0.10	$+0.24\pm 0.01$	0.07	$+0.20\pm 0.02$	0.07	$+0.64\pm 0.02$	0.08
NGC 4590	-0.52 ± 0.02	0.06	—	—	-0.14 ± 0.02	0.09	—	—	$+0.25\pm 0.02$	0.05	$+0.42\pm 0.01$	0.04
NGC 5634	$+0.21\pm 0.06$	0.17	$+0.13\pm 0.09$	0.21	$+0.22\pm 0.07$	0.19	$+0.18\pm 0.03$	0.07	$+0.15\pm 0.03$	0.08	$+0.52\pm 0.03$	0.07
NGC 5824	-0.36 ± 0.01	0.03	—	—	-0.04 ± 0.05	0.10	$+0.06\pm 0.03$	0.07	$+0.04\pm 0.04$	0.12	$+0.28\pm 0.03$	0.07
NGC 5904	-0.10 ± 0.06	0.25	$+0.20\pm 0.05$	0.21	$+0.17\pm 0.02$	0.07	$+0.16\pm 0.02$	0.07	$+0.12\pm 0.01$	0.04	$+0.58\pm 0.02$	0.07
NGC 6093	-0.17 ± 0.03	0.11	$+0.24\pm 0.04$	0.11	$+0.29\pm 0.05$	0.14	$+0.29\pm 0.05$	0.12	$+0.29\pm 0.04$	0.12	$+0.64\pm 0.02$	0.05
NGC 6397	-0.34 ± 0.03	0.10	—	—	$+0.09\pm 0.02$	0.07	$+0.24\pm 0.03$	0.08	$+0.10\pm 0.03$	0.10	$+0.52\pm 0.03$	0.07
NGC 6752	-0.17 ± 0.02	0.06	$+0.10\pm 0.04$	0.16	$+0.24\pm 0.03$	0.10	$+0.21\pm 0.01$	0.04	$+0.11\pm 0.01$	0.03	$+0.45\pm 0.01$	0.04
NGC 6809	$+0.18\pm 0.01$	0.03	—	—	$+0.25\pm 0.03$	0.09	$+0.26\pm 0.01$	0.04	$+0.22\pm 0.01$	0.04	$+0.59\pm 0.01$	0.05
NGC 7078	-0.19 ± 0.02	0.09	—	—	$+0.24\pm 0.04$	0.16	$+0.53\pm 0.06$	0.11	$+0.35\pm 0.02$	0.06	$+0.93\pm 0.03$	0.09
NGC 7099	-0.33 ± 0.02	0.06	—	—	-0.06 ± 0.02	0.10	$+0.05\pm 0.02$	0.05	$+0.45\pm 0.02$	0.05	$+0.46\pm 0.02$	0.05




Fig. B.1. Behaviour of [Si/Fe], [Ca/Fe], [Ti/Fe], [V/Fe], [Mn/Fe] and [Ni/Fe] as a function of [Fe/H] for the three SMC target clusters (squares, same colours in Fig. 1.), SMC field stars of Paper I (cyan circles) and SMC field stars measured by APOGEE (orange circles).