



Publication Year	2015
Acceptance in OA	2020-03-16T18:08:34Z
Title	Comparing masses in literature (CoMaLit) - I. Bias and scatter in weak lensing and X-ray mass estimates of clusters
Authors	Sereno, Mauro, ETTORI, STEFANO
Publisher's version (DOI)	10.1093/mnras/stv810
Handle	http://hdl.handle.net/20.500.12386/23296
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
Volume	450

Comparing masses in literature (CoMaLit) – I. Bias and scatter in weak lensing and X-ray mass estimates of clusters

Mauro Sereno^{1,2★} and Stefano Ettori^{2,3}

¹*Dipartimento di Fisica e Astronomia, Alma Mater Studiorum – Università di Bologna, viale Berti Pichat 6/2, I-40127 Bologna, Italia*

²*INAF, Osservatorio Astronomico di Bologna, via Ranzani 1, I-40127 Bologna, Italia*

³*INFN, Sezione di Bologna, viale Berti Pichat 6/2, I-40127 Bologna, Italia*

Accepted 2015 April 4. Received 2015 March 14; in original form 2014 July 24

ABSTRACT

The first building block to use galaxy clusters in astrophysics and cosmology is the accurate determination of their mass. Two of the most well-regarded direct mass estimators are based on weak lensing (WL) determinations or X-ray analyses assuming hydrostatic equilibrium (HE). By comparing these two mass measurements in samples of rich clusters, we determined the intrinsic scatters, $\sigma_{\text{WL}} \sim 15$ per cent for WL masses and $\sigma_{\text{HE}} \sim 25$ per cent for HE masses. The certain assessment of the bias is hampered by differences as large as ~ 40 per cent in either WL or HE mass estimates reported by different groups. If the intrinsic scatter in the mass estimate is not considered, the slope of any scaling relation ‘observable–mass’ is biased towards shallower values, whereas the intrinsic scatter of the scaling is overestimated.

Key words: gravitational lensing: weak – methods: statistical – galaxies: clusters: general – galaxies: clusters: intracluster medium.

1 INTRODUCTION

Usage of clusters of galaxies in cosmology and astrophysics relies on precise determination of their masses (Voit 2005; Limousin et al. 2013). In the context of ongoing and future large surveys (Laureijs et al. 2011), cluster properties which can be easily measured, e.g. optical richness, X-ray luminosity, Sunyaev-Zel’dovich (SZ) flux, are used as mass proxies. This requires an accurate calibration of the observable through comparison with direct mass estimates (Andreon & Bergé 2012; Ettori 2013).

The assessment of scaling relations is the foundation for investigating the physics of the baryons and of the dark matter at the cluster scale (Pratt et al. 2009; Arnaud et al. 2010; Giodini et al. 2013). Cosmological parameters can be constrained with cluster abundances and the observed growth of massive galaxy clusters (Vikhlinin et al. 2009; Mantz et al. 2010; Planck Collaboration XX 2014) or with gas fractions (Ettori et al. 2009).

Two of the most well-regarded mass estimates are the weak lensing (WL) mass and the X-ray mass exploiting hydrostatic equilibrium (HE). WL observations of the shear distortion of background galaxies trace the gravitational field of the matter distribution of the lens (Hoekstra et al. 2012; von der Linden et al. 2014; Umetsu et al. 2014). The physics behind gravitational lensing is very well understood and WL provides unbiased estimates of the total mass along the line of sight. The problem is to single out the contribution of

the lens and to de-project the information to get the intrinsic mass, which can then be confronted with theoretical predictions.

Under the assumption that HE holds between the intracluster medium (ICM) and the gravitational potential, the cluster mass can be recovered from observations of the spatially resolved spectroscopic data and the X-ray surface brightness (LaRoque et al. 2006; Donahue et al. 2014). However, deviations from equilibrium or non-thermal contributions to the pressure are difficult to quantify and can bias the mass estimate.

Methods based on spectroscopic measurements of galaxies velocities, such as the caustic technique (Rines & Diaferio 2006) or approaches exploiting the Jeans equation (Lemze et al. 2009; Biviano et al. 2013), can be effective too but they are hindered by the expensive observational requirements and are mostly limited to low-redshift haloes.

In principle, either WL or HE can provide accurate and unbiased mass measurements, but the approximations that have to be used (e.g. spherical symmetry, smooth density distributions, thermal equilibrium) may bias and scatter the results. These effects must be accurately quantified to calibrate other mass proxies. Due to scatter, WL and HE masses are proxies to the true mass themselves.

Numerical studies argued that lensing masses obtained from the fit of the cluster tangential shear profiles with Navarro-Frenk-White (Navarro, Frenk & White 1996, hereafter NFW) functionals are biased low by ~ 5 –10 per cent with a scatter of ~ 10 –25 per cent (Meneghetti et al. 2010; Becker & Kravtsov 2011; Rasia et al. 2012). The main sources of uncertainty in deprojected WL mass measurements are due to the presence of substructures and

★ E-mail: mauro.sereno@unibo.it

triaxiality. Lensing properties depend on the orientation of the cluster with respect to the line of sight (Oguri et al. 2005; Sereno 2007; Sereno & Umetsu 2011; Limousin et al. 2013). For systems whose major axis points towards the observer, 3D masses derived under the standard assumption of spherical symmetry are typically overestimated. The opposite occurs for clusters elongated in the plane of the sky, which are in the majority if the selected sample is randomly oriented.

The presence of substructures in the cluster surroundings may dilute the tangential shear signal (Meneghetti et al. 2010; Giocoli et al. 2012, 2014). Severe mass underestimations may come from either massive sub-clumps (Meneghetti et al. 2010) or uncorrelated large-scale matter projections along the line of sight (Becker & Kravtsov 2011).

The scatter is less significant in optimally selected clusters either having regular morphology or living in substructure-poor environments (Rasia et al. 2012).

The origins of bias and scatter of X-ray masses are well understood too, even though they are difficult to quantify (Rasia et al. 2012). They are strictly connected to non-thermal sources of pressure in the gas, to temperature inhomogeneity, and, to a lesser degree and mainly in the external regions, to the presence of clumps. Even if the cluster is in HE, the assumption that all the pressure is thermal biases the HE mass low. Large-scale, unvirialized bulk motions and subsonic turbulence contribute kinetic pressure (Battaglia et al. 2012).

Furthermore, structures in the temperature distribution bias low the temperature estimate. In fact, the X-ray detectors of *Chandra* and *XMM-Newton* (*X-ray Multi-Mirror Mission*) have a higher efficiency in the soft band and, thus, weight more colder gas (Mazzotta et al. 2004).

Numerical simulations showed that X-ray masses based on HE are biased low by a large amount of ~ 25 – 35 per cent (Piffaretti & Valdarnini 2008; Rasia et al. 2012, 2014). The bias grows from the inner to the outer regions of the clusters, where the presence of non-thermal sources of pressure in the ICM and temperature inhomogeneity play a larger role (Rasia et al. 2012).

Since the intrinsic scatters in either WL or HE masses have different origins, they are mostly uncorrelated. Scatter in WL masses is mainly due to triaxiality and substructures in the dark matter halo. On the other hand, the gas distribution approximately follows the gravitational potential and it is rounder than the dark matter one. Dark matter substructures are not necessarily associated with gas clumps. The sources which cause scatter in the HE masses are more related to gas physics and temperature distributions than to the total matter distribution and have a small impact on WL estimates.

On the observational side, the certain assessment of cluster masses is further complicated by instrumental and methodological sources of errors which may cause systematic uncertainties in data analysis (Rozo et al. 2014a).

The main sources of systematics in WL masses are due to selection and redshift estimate of background galaxies, which can be obtained through accurate photometric redshifts and colour–colour selection methods (Medezinski et al. 2010), and to the calibration of the shear signal. A small calibration correction of the order of just a few per cent translates into a typical error of ~ 10 per cent in the estimate of the virial mass (Umetsu et al. 2014).

Instrumental uncertainty has long been recognized as one of the main source of systematics plaguing HE masses. *XMM* cluster temperatures are systematically smaller by 10–20 per cent than *Chandra* estimates (Nevalainen, David & Guainazzi 2010; Donahue et al. 2014). On the other hand, *Chandra* and *XMM* measurements of the

gas distribution are highly consistent with one another (Donahue et al. 2014; Rozo et al. 2014a).

The picture on the inconsistencies between *Chandra* and *XMM* results is still debated. Donahue et al. (2014) found that *Chandra* and *XMM* temperatures of the very massive CLASH (Cluster Lensing And Supernova survey with Hubble; Postman et al. 2012) clusters agree in the core, where photon fluxes are considerable, whereas the regions where the temperature differences are larger are typically ~ 1 arcmin from the much brighter cluster core. Temperature differences persist even in outer regions with large signal-to-background ratio. These temperature discrepancies caused analogue off-sets in the HE mass.

Martino et al. (2014) compared the mass profiles of 21 LoCuSS (Local Cluster Substructure Survey) clusters that were observed with both satellites, extracting surface brightness and temperature profiles from identical regions of the respective data sets and including analytic models that predict the spatial variation of the *Chandra* and *XMM-Newton* backgrounds to $\lesssim 2$ and $\lesssim 5$ per cent precision, respectively. Notwithstanding global *XMM* spectroscopic temperatures lower by ~ 10 per cent, they obtained consistent results for the gas and total hydrostatic cluster masses. Martino et al. (2014) explained this counterintuitive result noticing that temperature discrepancies were significant only above a value of 6 keV. In the outer regions, most of the estimated temperatures were lower than this threshold and *Chandra* and *XMM* temperatures were in good agreement. Furthermore, they argued that larger errors bars are associated with highest temperature, due to the larger difficulty to distinguish the hottest spectra having a flatter shape from the background. The relative statistical weight in a fitting procedure is then lower.

This is the first in a series of papers focused on COmparing MAsses in LITERature (CoMaLit). Here, we look for systematic differences in WL and HE masses obtained from independent analyses and we assess the overall level of bias and intrinsic scatter. According to numerical simulations, the scatter in X-ray masses is supposedly smaller than in WL masses but a definite assessment of the values of bias and scatter of HE masses is still lacking, due to uncertainties in the treatment of the gas physics and to variability caused by the hydrodynamical scheme adopted in numerical simulations (Rasia et al. 2014). In this paper, we provide the first measurements of intrinsic scatters of WL and HE masses of real clusters.

In the second paper of the series (Sereno, Ettori & Moscardini 2015, hereafter CoMaLit-II), the Bayesian method developed to calibrate scaling relations between masses and observables, which fully accounts for intrinsic scatters in both the mass estimate and the scaling relation, was applied to the scaling relation between SZ flux and mass in *Planck* selected clusters of galaxies (Planck Collaboration XXIX 2014). The third paper of the series (Sereno 2015, hereafter CoMaLit-III) presents the Literature Catalogs of weak Lensing Clusters of galaxies (LC²), which are standardized compilations of clusters with measured WL masses. The fourth paper of the series (Sereno & Ettori 2015, hereafter CoMaLit-IV) extends the Bayesian methodology to account for time-evolution of the scaling relation. Products associated with the CoMaLit series, as well as future updates, will be hosted at <http://pico.bo.astro.it/~sereno/CoMaLit/>.

This paper is structured as follows. In Section 2, we discuss how the scatter in mass proxies can be estimated and how it impacts the calibration of scaling relations. Samples of clusters used in the analysis are introduced in Section 3. Comparison among either WL or HE masses from different groups is investigated in Section 4. Section 5 is devoted to the measurements of scatter and bias affecting the mass proxies. Discussion of results is contained in Section 6.

Final considerations can be found in Section 7. Three appendices contain some simplified and ready-to-use formulae to de-bias the scaling relations. A toy-model to illustrate the bias induced by the intrinsic scatter in the mass estimate is discussed in Appendix A. The correction for the widely used Bivariate Correlated Errors and Intrinsic Scatter method (BCES; Akritas & Bershady 1996) is proposed in Appendix B. Appendix C details how WL and HE mass estimates depends on the cosmological parameters.

Throughout the series of papers, we assume a fiducial flat Λ cold dark matter cosmology with density parameter $\Omega_M = 0.3$, and Hubble constant $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$; M_Δ denotes the mass within the radius r_Δ , which encloses a mean overdensity of Δ times the critical density at the cluster redshift, $\rho_{cr} = 3H(z)^2/(8\pi G)$; $H(z)$ is the redshift-dependent Hubble parameter. When H_0 is not specified, h is the Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

The presence of the superscript ‘WL’, ‘HE’ and ‘Tr’, means that M_{500} and r_{500} were determined using the mass estimate from the WL analysis, the X-ray measurements, or the knowledge of the true mass (which is available only for simulated clusters), respectively. ‘log’ is the logarithm to base 10 and ‘ln’ is the natural logarithm.

2 BIASES AND SCATTER INDUCED BIASES

The biases and the scatters of two mass proxies can be estimated by comparing the proxies in a cluster sample. The lensing and the hydrostatic mass approximate the true mass as

$$\ln M^{\text{WL}} \pm \delta_{\text{WL}} = \alpha_{\text{WL}} + \beta_{\text{WL}} \ln M^{\text{Tr}} \pm \sigma_{\text{WL}}, \quad (1)$$

$$\ln M^{\text{HE}} \pm \delta_{\text{HE}} = \alpha_{\text{HE}} + \beta_{\text{HE}} \ln M^{\text{Tr}} \pm \sigma_{\text{HE}}, \quad (2)$$

where the α 's quantify the bias and the β 's embody any deviation from linearity. The intrinsic scatters σ_{WL} and σ_{HE} are due to different physical processes and are assumed to be uncorrelated, see Section 1. The actual WL (HE) mass is known save for a measurement error δ_{WL} (δ_{HE}).

The notation in the left-hand sides of equations (1) and (2) is a shortcut to remind that the measurement processes are affected by errors and uncertainties. The observed masses differ from the ‘true’ values of the WL and HE masses which we would measure

in the ideal case of infinitely accurate and precise observations and systematics free-analyses. The notation in the right-hand sides of equations (1) and (2) reminds us that even in the case of ideal measurements without systematic/statistical errors ($\delta = 0$), the ‘true’ WL or HE masses differ from the ‘true’ mass of the cluster due to intrinsic scatter. In this sense, WL and HE masses are mass proxies. The relations in either the left- or the right-hand side of equations (1) and (2) can be modelled with normal distributions.

Bias and scatter in logarithmic variables differ from analogue quantities in linear variables. We adopt (natural) logarithmic variables for coherence with the standard derivation of scaling relations.

2.1 Eddington-like bias

Intrinsic scatter in the mass proxy induces systematic effects alike to the Eddington bias (Eddington 1913; Jeffreys 1938; Eddington 1940), which was first discussed in relation to observational uncertainties. Due to scatter and even in absence of measurement errors, the average value of an observed proxy differs from the true intrinsic mean for objects of the same class, see Fig. 1. When a subsample is selected according to the measured values of the proxy, X_{Proxy} , the distribution of the differences between the proxy and the true values, X_{True} , may be biased.

For quantities drawn from a limited range, border and selection effects have to be considered. Near a threshold, the symmetry between objects that are scattered into a range of observed values from above and objects that are scattered into from below is broken. This can be accounted for by assuming that the true masses are drawn from a normal rather than a uniform distribution.

Let us assume that the proxy X_{True} is the WL mass. Due to selection effects, the observed sample may be poor in clusters below a given threshold. At the tail at low values, more objects with larger X_{True} are scattered into the subsample from the right-hand side, than from the less populated left-hand side where the X_{True} 's are smaller, see Fig. 1. In a sample with steep bounds in true mass, clusters with very low values of M^{WL} are then of two main kinds. They are either intrinsically less massive clusters, i.e. with low values of M^{Tr} and nearly unbiased values of M^{WL} , or clusters with higher values of M^{Tr} that are scattered to lower values of observed M^{WL} . The mean M^{Tr} is then larger than the measured M^{WL} .




Figure 1. Probability distributions of a quantity X_{True} and of its scattered proxy X_{Proxy} . Left-hand panel: the probability distribution $p(X_{\text{True}})$ is a Gaussian function. $p(X_{\text{Proxy}})$ is Gaussian too. It is smoothed due to the intrinsic (normal) scatter. At the tail at low values, for samples selected according to X_{Proxy} (interval delimited by the blue vertical lines), more objects with larger X_{True} are scattered into the subsample from the right-hand side than from the left-hand side. Right-hand panel: conditional probability of X_{True} given X_{Proxy} , $p(X_{\text{True}}|X_{\text{Proxy}})$ (black line), and of a second proxy $X_{\text{Proxy}}^{(2)}$ given X_{Proxy} , $p(X_{\text{Proxy}}^{(2)}|X_{\text{Proxy}})$ (red line). The scatters in X_{Proxy} and $X_{\text{Proxy}}^{(2)}$ are not correlated. The mean X_{True} given X_{Proxy} is larger than X_{Proxy} , $\langle X_{\text{True}} \rangle(X_{\text{Proxy}}) > X_{\text{Proxy}}$. The second proxy $X_{\text{Proxy}}^{(2)}$ is unbiased with respect to X_{True} .

Let us consider a second proxy such as the HE mass, whose intrinsic scatter is uncorrelated to the WL mass. The second proxy is not biased by selections based on the first proxy. If the clusters are selected according to their M^{WL} , M^{HE} is still an unbiased scattered proxy of the true mass, since the scatters in measurements of WL and X-ray masses are uncorrelated. As a consequence, the mass ratio $M^{\text{HE}}/M^{\text{WL}}$ is biased high for clusters with small WL masses.

The opposite happens at large masses, where the more massive the clusters the rarer. The mass ratio $M^{\text{HE}}/M^{\text{WL}}$ is then biased low for clusters with large WL masses.

2.2 Biased slope

The intrinsic scatter in the mass estimate can make the slope of any scaling relation calibrated with either WL or HE masses shallower if the true masses in the selected sample are not uniformly distributed. This is a ripple effect of the Eddington-like bias. Due to measurement errors, the observed variance of the mass proxies is larger than the variance of the true masses. Slope estimators have to correct for this by de-biasing the sample variance (Akritas & Bershadsky 1996). Here, we are emphasizing the similar effect of the intrinsic scatter. Analogue treatments, which are often focused on observational errors rather than intrinsic scatter, have already been discussed (Andreon & Bergé 2012).

The distribution of the observed mass proxy is smoothed and it has a larger dispersion than the true masses. Due to the finite range, very large (small) measured WL or HE masses likely correspond to smaller (larger) true masses (in arbitrary units), whose observed WL or HE mass were scattered to the tails. If this is not accounted for, the derived slope of the scaling relation is biased towards flatter values.

Let us consider an unbiased (but scattered) mass proxy X , i.e. the (logarithm of the) WL or the HE mass, of the (logarithm of the) true mass Z ,

$$X \pm \delta_X = Z \pm \sigma_{X|Z}, \quad (3)$$

and a second observable quantity Y we want to calibrate,

$$Y \pm \delta_Y = \alpha_{Y|Z} + \beta_{Y|Z}Z \pm \sigma_{Y|Z}. \quad (4)$$

What we usually do is to compare the observable Y to the mass proxy,

$$Y \pm \delta_Y = \alpha_{Y|X} + \beta_{Y|X}X \pm \sigma_{Y|X}. \quad (5)$$

Due to the intrinsic scatter in the mass proxy, $\alpha_{Y|X}$ and $\beta_{Y|X}$ are biased estimates of $\alpha_{Y|Z}$ and $\beta_{Y|Z}$.

The full scheme outlined before can be formalized through a latent variable model (Feigelson & Babu 2012). In this approach, x and y are the measured manifest surrogate variables for the unobserved latent true variables X and Y , respectively. We refer to Feigelson & Babu (2012, chapter 7) for a review of this and other regression methods for astronomy. Other applications of Bayesian techniques to astronomical contexts can be found in Kelly (2007) and Andreon & Hurn (2012), and references therein.

Let us detail our model, which comprises a third latent variable Z together with X and Y . The observed values of the mass proxies X and of the observable Y are distributed according to Gaussian distributions centred on the true corresponding values,

$$x_i \sim \mathcal{N}(X_i, \delta_{x,i}), \quad (6)$$

$$y_i \sim \mathcal{N}(Y_i, \delta_{y,i}). \quad (7)$$

The true values of the mass proxies and of the observables are scattered with respect the true masses. If the scatters are uncorrelated, the distributions are

$$X_i \sim \mathcal{N}(Z_i, \sigma_{X|Z}), \quad (8)$$

$$Y_i \sim \mathcal{N}(\alpha_{Y|Z} + \beta_{Y|Z}Z_i, \sigma_{Y|Z}). \quad (9)$$

The distribution of true masses is approximated as a normal function of mean μ_Z and standard deviation σ_Z ,

$$Z_i \sim \mathcal{N}(\mu_Z, \sigma_Z). \quad (10)$$

The Gaussian distribution provides a good approximation for signal-selected samples. In fact, at low masses the number of clusters is limited by the selection threshold. At high masses, there are a few clusters because of the steepness of the mass function. The resulting mass distribution is then approximately lognormal for realistic cases, see CoMaLit-IV. Furthermore, the Gaussian function is flexible enough to accommodate for a large range of fairly unimodal distributions (Kelly 2007) or even the uniform distribution, which is approximated in case of very large variance.

We chose non-informative priors. For the variances, i.e. the squared scatters, we considered inverse Gamma distribution (Andreon & Hurn 2010),

$$1/\sigma_{X|Z}^2, 1/\sigma_{Y|Z}^2, 1/\sigma_Z^2 \sim \Gamma(\epsilon, \epsilon), \quad (11)$$

where ϵ is a small number. In our calculation we took $\epsilon = 10^{-3}$ (Andreon & Hurn 2010). We adopted uniform priors for the intercept $\alpha_{Y|Z}$ and the mean μ_Z ,

$$\alpha_{Y|Z}, \mu_Z \sim \mathcal{U}(-1/\epsilon, 1/\epsilon). \quad (12)$$

For the slope $\beta_{Y|Z}$, we assumed a Student's t_1 distribution with one degree of freedom, which is equivalent to a uniform prior on the direction angle $\arctan \beta_{Y|Z}$ (Andreon & Hurn 2010),

$$\beta_{Y|Z} \sim t_1. \quad (13)$$

The unknowns of the regression are the intercept $\alpha_{Y|Z}$, the slope $\beta_{Y|Z}$, and the scatter $\sigma_{Y|Z}$ of the scaling $Y-Z$, the scatter $\sigma_{X|Z}$ of the scaling $X-Z$, the values of the independent variable Z_i and the parameters describing their distribution μ_Z and σ_Z , and the values of the covariate variables X_i and Y_i .

We implemented the above Bayesian methodology with JAGS.¹ The scripts used for our analysis can be found at <http://pico.bo.astro.it/~sereno/CoMaLit/JAGS/>.

The bias in the estimate of the slope due to the intrinsic scatter in the mass estimate can be studied through a simple simulation. Let us consider a sample of 100 true (logarithmic) masses drawn from a Gaussian distribution with mean $\mu_Z = 1.0$ and $\sigma_Z = 0.35$. The mass proxies are measured with an observational uncertainty $\delta_x = 0.05$. The intrinsic scatter is $\sigma_{X|Z} = 0.15$. The proxy Y is linearly related to Z with $\alpha_{Y|Z} = -0.2$ and $\beta_{Y|Z} = 1.5$. The scatter around the relation is $\sigma_{Y|Z} = 0.2$. The observational uncertainty is fixed to $\delta_y = 0.05$. We model scatters and errors with normal distributions.

As a first step, we verified that the regression retrieves unbiased parameters when we compare observable Y and true mass Z , see equation (4). In this simple case, we do not need equations (6) and (8). We found $\alpha_{Y|Z} = -0.22 \pm 0.07$, $\beta_{Y|Z} = 1.49 \pm 0.07$ and

¹ The package JAGS (Just Another Gibbs sampler) by M. Plummer performs analysis of Bayesian hierarchical models using Markov Chain Monte Carlo simulation. It is publicly available at <http://mcmc-jags.sourceforge.net/>.

Table 1. Characteristics of the X-ray and WL samples used in the analysis. N_{cl} is the number of clusters in the sample.

Acronym	N_{cl}	WL instrument	WL reference	X-ray instrument	X-ray reference	notes
RA12	60	–	Rasia et al. (2012)	–	Rasia et al. (2012)	Simulations
CCCP-WL	55	CFHT	Hoekstra et al. (2012)	–	–	
CCCP-HE	50	–	–	<i>Chandra</i> , <i>XMM</i>	Mahdavi et al. (2013)	–
WTG	51	Subaru, CFHT	Applegate et al. (2014)	–	–	–
CLASH-WL	20	Subaru	Umetsu et al. (2014)	–	–	–
CLASH-CXO	25	–	–	<i>Chandra</i>	Donahue et al. (2014)	–
CLASH-XMM	18	–	–	<i>XMM</i>	Donahue et al. (2014)	–
E10	44	–	–	<i>XMM</i>	Ettori et al. (2010)	–
L13	35	–	–	<i>Chandra</i>	Landry et al. (2013)	–
B12	25	–	–	<i>Chandra</i>	Bonamente et al. (2012)	Additional SZ data from SZA

$\sigma_{Y|Z} = 0.17 \pm 0.02$. Regression results are statistically consistent with the input parameters.

We then considered the evolution of Y with the mass proxy X , see equation (5). In this case, we are not interested in the true values Z_i and we can substitute equations (9) and (10) with

$$Y_i \sim \mathcal{N}(\alpha_{Y|X} + \beta_{Y|X} X_i, \sigma_{Y|X}), \quad (14)$$

and

$$X_i \sim \mathcal{N}(\mu_X, \sigma_X), \quad (15)$$

respectively. As before, we do not need equations (6) and (8). We found $\alpha_{Y|X} = 0.00 \pm 0.08$, $\beta_{Y|X} = 1.25 \pm 0.07$ and $\sigma_{Y|X} = 0.23 \pm 0.02$. The relation is flatter than the intrinsic one Y – Z and the estimated scatter is larger. On turn, the flatter relation causes a higher intercept.

To avoid biases, we have to consider that X is a scattered proxy of the true mass. Equations (3) and (4) have to be fitted simultaneously and the full scheme in equations (6)–(14) has to be adopted with the additional parameter $\sigma_{X|Z}$. We found $\alpha_{Y|Z} = -0.18 \pm 0.16$, $\beta_{Y|Z} = 1.43 \pm 0.15$, $\sigma_{Y|Z} = 0.15 \pm 0.07$ and $\sigma_{X|Z} = 0.11 \pm 0.05$. Intrinsic parameters are well recovered even though statistical uncertainties are larger.

More details on this statistical model are provided in Appendix A, which also provides some ready-to-use approximate corrections. Correcting $\alpha_{Y|X}$ and $\beta_{Y|X}$ as suggested in equation (A11), we found $\alpha_{Y|Z} \sim -0.22$ and $\beta_{Y|Z} \sim 1.48$, in agreement with the input parameters.

The correction for the biased slope estimated through the widely used BCES method is suggested in Appendix B.

3 CLUSTER SAMPLES

We looked in the literature for public catalogues compiled in the last few years with either WL or HE masses. The main properties of the samples, which we are going to introduce in the following, are summarized in Table 1.

When quoted mass values were provided with asymmetric errors, we estimated the mean value and the standard deviation as suggested in D’Agostini (2004). All the considered masses refer to the fiducial cosmological model. Conversions were performed as described in Appendix C. A full account of references of WL analyses and standardization methods can be found in CoMaLit-III.

3.1 Numerical simulations

Rasia et al. (2012, hereafter RA12) compared the WL and X-ray properties of a sample of 20 numerically simulated massive clusters at redshift $z = 0.25$. The haloes were the most massive

($M_{\text{vir}} > 5 \times 10^{14} M_{\odot} h^{-1}$) from a set of radiative simulations in a cosmological volume of $1(\text{Gpc } h^{-1})^3$, evolved in the framework of a *WMAP-7* normalized cosmology (Fabjan et al. 2011).

Each cluster was later re-simulated at higher resolution and with more complex gas physics. The simulations included: metal-dependent radiative cooling and cooling/heating from a spatially uniform and evolving UV background; a star formation model where a hot ionized phase coexists in pressure equilibrium with a cold phase, which is the reservoir for star formation; a description of metal enrichment from different stellar populations; the effect of supernovae feedback through galactic winds.

The clusters were finally processed to generate optical and X-ray mock observations along three orthogonal projections. The final sample consists of 60 cluster realizations. WL and HE masses are estimated within r_{500}^{Tr} , the overdensity radius corresponding to the true mass.

3.2 Canadian Cluster Comparison Project

The Canadian Cluster Comparison Project (CCCP; Mahdavi et al. 2013) assembled a sample of 50 rich clusters of galaxies in the redshift range $0.15 < z < 0.55$. All of the clusters were observable from the Canada-France-Hawaii Telescope (CFHT), which restricts the sample to systems at $-15 \text{ deg} < \text{declination} < 65 \text{ deg}$. Most of them were selected to have an ASCA (Advanced Satellite for Cosmology and Astrophysics) temperature $k_B T_X > 3 \text{ keV}$. X-ray properties were measured either with *Chandra* or *XMM-Newton*. WL studies for five additional clusters without X-ray analyses can be found in Hoekstra et al. (2012).

Lensing masses were determined with aperture statistics (Hoekstra et al. 2012). This approach relies on shear measurements at large radii and reduces the effects of contamination by cluster members. The 3D masses were computed from the model-independent 2D aperture masses with a de-projection method based on an NFW density profile.

Mahdavi et al. (2013) performed the X-ray analysis of the sample using both *Chandra* and *XMM* observations. They found that due to temperature discrepancies, the *XMM* cluster masses were systematically ~ 15 per cent smaller than *Chandra* masses. In order to combine the data, Mahdavi et al. (2013) down-weighted the high-energy effective area of *Chandra*. X-ray quantities were estimated either within r_{500}^{WL} , the radius evaluated from the WL mass measurement, or r_{500}^{HE} , as evaluated from the mass estimate assuming HE.²

² Values of M_{500}^{WL} and M_{500}^{HE} for the CCCP sample are publicly available at <http://sfstar.sfsu.edu/cccp>.

Mahdavi et al. (2013) identified a subsample of 20 cool-core (CC) systems with core entropy at 20 kpc smaller than 70 keVcm² and eight systems with low offsets between the brightest cluster galaxy (BCG) and the X-ray surface brightness peak, $D_{\text{BCG}} < 10\text{kpc}$.

3.3 Cluster Lensing And Supernova survey with Hubble

The CLASH programme (Postman et al. 2012) has been mapping the matter distribution of 25 rich clusters drawn largely from the Abell and MAssive Cluster Survey (MACS; Ebeling et al. 2010) cluster catalogues. Umetsu et al. (2014) performed a joint shear-and-magnification WL analysis of a subsample of 16 X-ray regular and four high-magnification galaxy clusters in the redshift range $0.19 \lesssim z \lesssim 0.69$. A complementary analysis exploiting strong lensing data was presented in Merten et al. (2014).

To make the comparison with the other data samples easier, we will use the mass estimates in Umetsu et al. (2014), whose methodology exploits only the WL regime, whereas results in Merten et al. (2014) strongly rely on information from the inner regions. Mass estimates in Umetsu et al. (2014) were based on joint WL shear plus magnification measurements based on ground-based wide-field Subaru data. On the other hand, the analysis in Merten et al. (2014) combined the Subaru shear profile with WL constraints from the *Hubble Space Telescope* (HST) in the intermediate regime and strong lensing constraints from HST.

All of the CLASH clusters have been observed with the *Chandra* satellite (Postman et al. 2012). A subsample of 18 clusters was targeted by *XMM* too. The X-ray analysis was presented in Donahue et al. (2014), which computed HE masses and gas fractions.

Based on *Chandra* data, Donahue et al. (2014) identified 10 clusters (nine of them with WL mass) with a strong CC, i.e. with an excess core entropy smaller than 30 keVcm².

3.4 Weighing the Giants

The Weighing the Giants (WTG; von der Linden et al. 2014) programme targeted 51 X-ray luminous clusters from the MACS and the Brightest Cluster Survey (BCS; Ebeling et al. 2000). The clusters span a large range in redshift ($0.15 \lesssim z \lesssim 0.7$) and dynamical state. Seven clusters are classified as relaxed (von der Linden et al. 2014).

The values of the scale radius and the concentration are provided in Applegate et al. (2014, table 4). We derived M_{500}^{WL} and r_{500}^{WL} using the NFW density profile adopted in the WTG analysis.

3.5 X-ray samples

Ettori et al. (2010, E10) studied a sample of 44 X-ray luminous galaxy clusters observed with *XMM-Newton* in the redshift range $0.1 \lesssim z \lesssim 0.3$. They applied two different techniques (the backward ‘method 1’, which we take as the reference method, and the forward ‘method 2’) to recover the gas and the dark mass properties, described with an NFW profile. Clusters were classified according to their core properties. E10 identified a subsample of 16 low-entropy-core systems, which represent the prototype of relaxed clusters with a well-defined CC at low entropy.

Landry et al. (2013, L13) presented *Chandra* X-ray measurements of the hydrostatic mass and of the gas mass fraction out to r_{500} for the complete sample of the 35 most luminous clusters from the BCS and its extension at redshift $0.15 \lesssim z \lesssim 0.30$. The clusters span a large range of dynamical states. The data were analysed using two independent pipelines and two different models for the

gas density and temperature, the ‘Polytropic’ (which we take as our reference case) or the ‘Vikhlinin’ model.

Bonamente et al. (2012, B12) derived the hydrostatic masses and the pressure profile of a sample of 25 massive relaxed galaxy clusters with a simultaneous analysis of SZ data from the Sunyaev-Zel’dovich array (SZA) and archival *Chandra* observations.

4 MASS COMPARISON

Even though in principle WL and X-ray masses could be unambiguously determined from a given set of observations, calibration issues and hidden systematics make these measurements very difficult.

In this section, we compare either WL or HE masses from different catalogues. It is nowadays customary to quote masses within a given overdensity and to derive scaling relations in terms of them. These masses can be related to the virial mass and most cluster properties are expected to be self-similar if rescaled by their value at r_{Δ} . To limit extrapolation of published results, we then considered the masses within r_{500} , rather than extrapolating the results up to a fixed length.

On the other hand, the relationship between M_{Δ} and r_{Δ} exacerbates problems connected to aperture differences, which complicate the comparison between different samples. Since the total mass within a fixed radius scales nearly linearly with the radius, differences in mass within a given overdensity are inflated by $\sim 100/3$ per cent with respect to differences within a fixed physical radius.

Differences among properties measured within a fixed length are not inflated but they refer to physically different regions in differently sized clusters. A promising alternative is to express the results in terms of the circular velocity $v_{\text{circ}}^2 = GM(r)^2/r$. In fact, the circular velocity is almost independent of cosmology and it is nearly unaffected by aperture problems (Donahue et al. 2014). Within a given overdensity radius, the velocity scales with mass as $v_{\Delta} \propto M_{\Delta}^{2/3}$. Quoted results for central estimate and scatter of $\Delta \ln M_{500}$, as well as fractional changes, can be translated in analogue results for $\Delta \ln v_{500}$ by simply multiplying by the factor 2/3.

To compare different samples, we considered the (natural) logarithm of mass ratios (Rozo et al. 2014a). The central estimate and the scatter were computed as bi-weight estimators of the distribution. Uncertainties were estimated with bootstrap resampling with replacement. The main advantage in using logarithms is that their difference is (anti)symmetric. This solves the problem affecting those estimators of ratio which are not symmetric with respect to the exchange of numerator and denominator.

Quoted errors in compiled catalogues may account for different sources of statistical and systematic uncertainties. Furthermore, it can be argued that the published uncertainties are unable to account for the actual variance seen in sample pairs (Rozo et al. 2014a). We then conservatively performed unweighted analyses.

4.1 WL masses

In principle, WL masses could be determined to an accuracy of $\lesssim 8$ per cent (Umetsu et al. 2014; von der Linden et al. 2014), but differences between masses reported by distinct groups are off by ~ 20 –50 per cent (Applegate et al. 2014; Umetsu et al. 2014). On the other hand, comparisons show that mass measurements correlate quite tightly (Applegate et al. 2014).

The CCCP and the WTG samples share 17 clusters, see Table 2. M_{500} from CCCP are smaller by ~ 30 per cent with a scatter of ~ 20 per cent. This difference is way larger than the claimed mass

Table 2. Comparison of WL masses from independent analyses. We quote the mean ln differences in mass for sample pairs. Entries are in the format: $(N_{\text{cl}}), \mu (\pm \delta\mu) \pm \sigma (\pm \delta\sigma)$, where N_{cl} is the number of clusters in common between the samples, μ is the central estimate of the difference in natural logarithm $\ln(M_{500}^{\text{row}}/M_{500}^{\text{col}})$, with associated uncertainty $\delta\mu$; σ is the dispersion with associated uncertainty $\delta\sigma$. M_{500}^{row} (M_{500}^{col}) refers to the sample indicated in the corresponding row (column). Quoted values are the bi-weight estimators.

	CLASH-WL	WTG
	(6)	(17)
CCCP-WL	$-0.45 (\pm 0.12)$ $\pm 0.25 (\pm 0.10)$	$-0.31 (\pm 0.05)$ $\pm 0.21 (\pm 0.09)$
	–	(17)
CLASH-WL	–	$0.01 (\pm 0.10)$
	–	$\pm 0.37 (\pm 0.06)$




Figure 2. Ratio of WL masses estimated by CCCP and WTG as a function of redshift.

calibration uncertainty and highlights the difficulties connected to unbiased calibrations in WL measurements. We found no trend with redshift, see Fig. 2.

The CCCP masses are notably underestimated with respect to the CLASH clusters too. On the other hand, the agreement between the WTG and the CLASH results is substantial, even though the scatter in the mass ratios is quite large. Usually the two mass estimates of a given cluster from WTG and CLASH coincide within $\lesssim 30$ per cent of the combined error.

The scatters in the mass ratio are of the order of 20–40 per cent and are consistent with the quoted statistical uncertainties on the WL mass estimates, which are of the order of 20 per cent or larger. In fact, if statistical errors are properly estimated, the combined scatter in the mass ratios should be approximately given by the quadratic sum of the typical errors of the two considered samples.

4.2 X-ray masses

X-ray properties of galaxy clusters reported by competing groups may reach discrepancies of 50 per cent (Rozo et al. 2014a). Here, we consider the off-set and the scatter in the estimate of HE masses. Discrepancies may stem from either differences in the considered data sets (to the larger extent if taken with different instruments),

Table 3. Comparison of HE masses from independent methods but from the same data sets. E10-M1 and E10-M2 denote the two methods used in Ettori et al. (2010). L13-Vick and L13-Poly denote the two methods used in Landry et al. (2013). N_{cl} (col. 3) is the number of clusters in common between the samples listed in cols. 1 and 2. $M_{500}^{(1)}/M_{500}^{(2)}$ (col. 4) is the central estimate of the ratio between masses in the two samples; σ is the dispersion. Quoted values are the bi-weight estimators.

Sample 1	Sample 2	N_{cl}	$M_{500}^{(1)}/M_{500}^{(2)}$	σ
E10-M2	E10-M1	44	0.98 ± 0.05	0.25 ± 0.03
L13-Vick	L13-Poly	35	0.99 ± 0.04	0.23 ± 0.03

or from not consistent data reduction pipelines, or from different techniques to recover the mass.

This last issue can be quantified by comparing mass estimates obtained from the same data sets but with different methodologies. This is the case of the analyses in either E10 or L13, for which we could compare the scatter in the mass estimate due to the different modelling, see Table 3.

The typical statistical error in an HE mass estimate is of the order of ~ 15 per cent. The observed scatter in the mass ratios is then consistent with the propagation of this error. This comparison suggests that mass estimates are not biased due to different techniques, whose associated variance is negligible with respect to the statistical uncertainty.

Larger variations are mainly related to different data sets, see Table 4. Discrepancies of order of $\gtrsim 30$ per cent may be in place. This is the case for results based on *Chandra* (CLASH-CXO, B12, L13) versus *XMM* analyses (E10, CLASH-*XMM*), whose temperature estimates may disagree at large radii (Donahue et al. 2014).

Each method/analysis may systematically either under- or over-estimate the cluster mass. X-ray masses in the CLASH sample based on *Chandra* (*XMM*) data are systematically larger (smaller) than other estimates. On the other hand, masses from B12 and L13 are lower than other samples.

A significant role can be played by additional data sets exploited in the analysis. The inclusion of SZ data, which are more sensitive to the outer regions, might lower the mass values in B12.

The large differences in estimated masses and the large scatters suggest that quoted formal statistical uncertainties in HE masses, usually of the order of ~ 10 –15 per cent, might be underestimated.

5 REGRESSION RESULTS

We measured biases and intrinsic scatters of WL and HE masses through the statistical model detailed in Section 2. To simplify the analysis, we assumed that the lensing and the hydrostatic masses scale linearly with the true mass, $\beta_{\text{WL}} = 1$ and $\beta_{\text{HE}} = 1$. In relation to the notation in Section 2.2, the logarithm of the WL mass ($\ln M^{\text{WL}}$), of the HE mass ($\ln M^{\text{HE}}$), and of the true mass ($\ln M^{\text{Tr}}$) can be identified with X , Y and Z , respectively.

The true masses are known only in simulations. For observed samples, we can estimate only the relative bias between WL and HE masses and we fixed $\alpha_{\text{WL}} = 0$. The effective bias $M^{\text{HE}}/M^{\text{Tr}}$ ($M^{\text{WL}}/M^{\text{Tr}}$) can be defined as $\exp[\alpha_{\text{HE}}]$ ($\exp[\alpha_{\text{WL}}]$). The relative bias $M^{\text{HE}}/M^{\text{WL}}$ can be defined as $\exp[\alpha_{\text{HE}} - \alpha_{\text{WL}}]$. Bias and scatter are largely uncorrelated. We tested that the estimates of scatter and relative bias do not change whether we consider $\alpha_{\text{HE}} = 0$ rather than $\alpha_{\text{WL}} = 0$.

The intrinsic distribution of the independent variable, $\ln M^{\text{Tr}}$, was approximated with a Gaussian function, see Section 2.2. We tested

Table 4. Comparison of HE masses from independent analyses. For the CCCP-HE sample, we considered masses within r_{500}^{HE} . We quote the mean ln differences in mass for sample pairs. Entries are in the format: (N_{cl}) , $\mu(\pm\delta\mu)\pm\sigma(\pm\delta\sigma)$, where N_{cl} is the number of clusters in common between the samples, μ is the central estimate of the difference in natural logarithm $\ln(M_{500}^{\text{row}}/M_{500}^{\text{col}})$, with associated uncertainty $\delta\mu$; σ is the dispersion with associated uncertainty $\delta\sigma$. M_{500}^{row} (M_{500}^{col}) refers to the sample indicated in the corresponding row (column). Quoted values are the bi-weight estimators.

	CCCP-HE	CLASH-XMM	CLASH-CXO	L13	B12
	(11)	(3)	(3)	(11)	(6)
E10	0.22 (± 0.08) ± 0.28 (± 0.11)	~ 0.17 $\pm(\sim)0.06$	~ -0.15 $\pm(\sim)0.28$	0.35 (± 0.14) ± 0.30 (± 0.09)	0.25 (± 0.10) ± 0.19 (± 0.08)
CCCP-HE	–	(5) 0.03 (± 0.21) ± 0.29 (± 0.16)	(6) –0.38 (± 0.14) ± 0.34 (± 0.21)	(18) 0.12 (± 0.07) ± 0.33 (± 0.14)	(5) 0.24 (± 0.22) ± 0.35 (± 0.17)
CLASH-XMM	–	–	(18) –0.38 (± 0.09) ± 0.35 (± 0.10)	(2) ~ -0.05 $\pm(\sim)0.18$	(10) 0.14 (± 0.15) ± 0.46 (± 0.30)
CLASH-CXO	–	–	–	(4) ~ 0.31 $\pm(\sim)0.01$	(12) 0.45 (± 0.14) ± 0.37 (± 0.13)
L13	–	–	–	–	(4) ~ 0.03 $\pm(\sim)0.06$

Table 5. Intrinsic scatter and bias of mass proxies M_{500}^{WL} and M_{500}^{HE} for the RA12 clusters.

M^{Proxy}	$M_{500}^{\text{Proxy}}/M_{500}^{\text{Tr}}$	σ
M_{500}^{WL}	0.95 ± 0.03	0.13 ± 0.03
M_{500}^{HE}	0.71 ± 0.01	0.11 ± 0.02

that results based on more complex distributions, such as mixtures of Gaussian functions (CoMaLit-II), were indistinguishable from the simplest case.

5.1 Simulated sample

As a first step, we analysed the simulated sample from RA12. In the realm of simulations, we know the true masses of the clusters. We can exploit this information to compute the bias and the intrinsic scatter of each mass proxy separately by direct comparison with the true mass. WL and HE masses can be compared to true masses autonomously. Regression results are in agreement with the original analysis in RA12 and are summarized in Table 5. Note that differently from RA12, we estimated the intrinsic rather than the total scatter and we focused on logarithmic variables.

The level of bias for each proxy is approximately constant with respect to the true mass, see Fig. 3.

The intrinsic scatter plays an important role when we analyse the bias as a function of the mass proxy, see Fig. 4. The decreasing trend of the ratio $M_{500}^{\text{HE}}/M_{500}^{\text{WL}}$ as a function of M_{500}^{WL} is an effect of the scatter of M_{500}^{WL} around the true mass, see Section 2.1.

Due to the combined action of selection effect and intrinsic scatters, at small (large) values of WL masses, M_{500}^{WL} is biased low (high) with respect to the true mass whereas M_{500}^{HE} is unbiased. As a consequence, the ratio $M_{500}^{\text{HE}}/M_{500}^{\text{WL}}$ decreases with M_{500}^{WL} . At intermediate values, clusters can be scattered into a given range in M_{500}^{WL} from either above or below, and the ratio $M_{500}^{\text{HE}}/M_{500}^{\text{WL}}$ is not biased. The larger the scatter, the steeper the trend in the mass ratio as a function




Figure 3. Masses in the RA12 sample. Top panel: WL mass versus HE mass. Clusters are grouped in four bins in true mass (black points). The red line is the bisector $M_{500}^{\text{HE}} = M_{500}^{\text{WL}}$. Lower panel: bias of the proxy as a function of the true mass. Clusters are grouped in four bins in true mass. Black (blue) points correspond to the bias of the WL (HE) mass. The solid error bars denote the 1σ uncertainties for the central estimate. The dashed error bars denote the dispersion. All masses are computed within r_{500}^{Tr} .




Figure 4. Comparison of WL and HE masses for the RA12 sample. Masses are measured within r_{500}^{Tr} , the overdensity radius related to the true mass. In the left- (right-)hand panels, clusters are grouped in four bins according to their measured WL (HE) mass. Green points mark the predictions based on the analytical model discussed in the paper. Top-left panel: M_{500}^{HE} as a function of M_{500}^{WL} . Black (green) points mark clusters (analytical predictions) binned in M_{500}^{WL} . Errors bars for the binned points are computed as the dispersion around the central value. Bottom-left panel: logarithm of $M_{500}^{\text{HE}}/M_{500}^{\text{WL}}$ as a function of M_{500}^{WL} . The solid error bars denote the 1σ uncertainties for the central estimate. The dashed error bars denote the dispersion. Top-right panel: M_{500}^{WL} as a function of M_{500}^{HE} . Black points marks clusters binned in M_{500}^{HE} . Bottom-right panel: logarithm of $M_{500}^{\text{HE}}/M_{500}^{\text{WL}}$ as a function of M_{500}^{HE} .

of the mass proxy. The scatter also flattens the $M_{500}^{\text{HE}}-M_{500}^{\text{WL}}$ relation towards larger values of M_{500}^{WL} . For similar reasons and being M_{500}^{HE} at the denominator, the ratio $M_{500}^{\text{WL}}/M_{500}^{\text{HE}}$ increases as a function of M_{500}^{HE} .

The statistical model summarized in equations (1) and (2) and detailed in Section 2.2 accounts for these effects and can be validated by comparing its predictions to the RA12 sample. We generated a sample of simulated clusters whose true masses are drawn from a Gaussian distribution. The corresponding measured WL and HE masses were simulated assuming intrinsic scatters and observational uncertainties as measured in the RA12 sample. This model successfully reproduces the trends in the observed mass ratio, see Fig. 4.

As a second step, we tested the regression algorithm with the mock observations of the simulated RA12 sample, see Table 6. Differently from the first step, we used the manifest estimated values of the mass proxies but we did not exploit the information on the latent true mass. We could then not calibrate the bias in either M_{500}^{WL} or M_{500}^{HE} in an absolute way, but we had to normalize one bias relatively to the other one. We assumed $\alpha_{\text{WL}} = 0$, i.e. we measured $M_{500}^{\text{HE}}/M_{500}^{\text{Tr}}$ in units of $M_{500}^{\text{WL}}/M_{500}^{\text{Tr}}$. The level of relative bias and the intrinsic scatters are recovered within the statistical uncertainties.

5.2 Observed samples

We considered a number of samples of clusters with observed WL and HE masses: (i) the CCCP sample; (ii) the CLASH sample with X-ray estimates based on either *Chandra* or *XMM* data; (iii) the WTG clusters with HE masses from either B12 (WTG-B12) or L13 (WTG-L13). For the CCCP sample, we could consider either masses within the same radius, i.e. r_{500}^{WL} , or alternatively WL masses within r_{500}^{WL} and HE masses within r_{500}^{HE} .

Results for the real clusters are summarized in Table 6 and in Figs 5–8. Thereafter in the text, when necessary we rescale the values of bias and scatters reported in Table 6 to the corresponding values within the same length by simply multiplying by a factor 2/3.

The parameter determination is not degenerate, see Fig. 5. In fact, when comparing HE to WL masses, the relative intrinsic scatters acts in orthogonal directions.

HE masses are biased low with respect to WL masses by $\lesssim 15$ per cent if we rely on the CCCP sample or by $\lesssim 25$ –35 per cent if we consider the WTG estimates. The results for the CLASH sample depend on the X-ray analysis. The bias is ~ 10 per cent for *Chandra* data and ~ 30 per cent for *XMM* data.

The difference in the level of the bias among the various samples reflects the different absolute mass calibrations in the WL and the X-ray samples, see Section 4. The bias ascertained with either the

Table 6. Biases and intrinsic scatters of the WL and HE masses. Col. 1: sample; col. 2: number of clusters in the sample, N_{cl} ; cols. 3, 4: radius within which the WL lensing and the HE mass were computed, respectively; col. 5: effective ratio between the true mass and the WL mass; the WL mass is assumed to be an unbiased proxy; col. 6: intrinsic scatter of $\ln M_{500}^{WL}/M_{500}^{Tr}$; col. 7: effective ratio $M_{500}^{HE}/M_{500}^{WL}$; col. 8: intrinsic scatter (as in col. 6 but for HE masses). Quoted values are bi-weight estimators of the posterior probability distribution.

Sample	N_{cl}	r_{500}^{WL}	r_{500}^{HE}	$M_{500}^{WL}/M_{500}^{Tr}$	σ_{WL}	$M_{500}^{HE}/M_{500}^{WL}$	σ_{HE}
RA12	60	r_{500}^{Tr}	r_{500}^{Tr}	[1]	0.14 ± 0.04	0.75 ± 0.03	0.13 ± 0.04
CCCP	50	r_{500}^{WL}	r_{500}^{WL}	[1]	0.14 ± 0.06	0.85 ± 0.05	0.24 ± 0.07
CCCP-CC	16	r_{500}^{WL}	r_{500}^{WL}	[1]	0.18 ± 0.10	0.93 ± 0.11	0.24 ± 0.12
CCCP-Low Offset	20	r_{500}^{WL}	r_{500}^{WL}	[1]	0.18 ± 0.10	0.82 ± 0.09	0.30 ± 0.11
CCCP	50	r_{500}^{WL}	r_{500}^{HE}	[1]	0.20 ± 0.09	0.81 ± 0.07	0.45 ± 0.07
CLASH-CXO	20	r_{500}^{WL}	r_{500}^{HE}	[1]	0.17 ± 0.09	0.78 ± 0.09	0.34 ± 0.12
CLASH-CXO-CC	9	r_{500}^{WL}	r_{500}^{HE}	[1]	0.22 ± 0.14	0.77 ± 0.14	0.31 ± 0.17
CLASH-CXO-Low Offset	8	r_{500}^{WL}	r_{500}^{HE}	[1]	0.31 ± 0.17	0.70 ± 0.15	0.34 ± 0.17
CLASH-XMM	16	r_{500}^{WL}	r_{500}^{HE}	[1]	0.17 ± 0.10	0.56 ± 0.08	0.45 ± 0.14
WTG-L13	14	r_{500}^{WL}	r_{500}^{HE}	[1]	0.32 ± 0.14	0.64 ± 0.09	0.16 ± 0.08
WTG-B12	14	r_{500}^{WL}	r_{500}^{HE}	[1]	0.19 ± 0.12	0.47 ± 0.07	0.34 ± 0.15




Figure 5. Probability distributions of the scatter σ_{WL} of WL masses, of the scatter σ_{HE} of HE masses, and of the logarithm of the effective bias, $\alpha_{HE} = \ln M_{500}^{HE}/M_{500}^{WL}$, for the CCCP sample. HE masses were measured within r_{500}^{WL} , the overdensity radius related to the WL mass. The thick (thin) lines in the two-dimensional plots include the $1(2)\sigma$ confidence region in two dimensions, here defined as the region within which the value of the probability is larger than $\exp(-2.3/2)$ ($\exp(-6.17/2)$) of the maximum.




Figure 6. Same as Fig. 4 but for the clusters in the CCCP sample. Masses are measured within r_{500}^{WL} , the overdensity radius related to the WL mass. Blue points plot the results for clusters in the RA12 sample grouped in the same mass bins as the CCCP clusters. Masses for the RA12 sample are computed within r_{500}^{Tr} .

WTG or the CLASH-XMM sample is in agreement with results from numerical simulations whereas results based on the CCCP and CLASH-CXO slightly underestimate it.

Apart from the overall normalization, results from different data sets are qualitatively and quantitatively consistent. The intrinsic scatter on WL masses is of order of ~ 10 – 15 per cent, in very good agreement with numerical simulations. On the other hand, the estimated scatter on HE masses is ~ 25 per cent, a factor of 2 larger than theoretical predictions. The large value of σ_{HE} is evident in the plots of $M_{500}^{\text{WL}}/M_{500}^{\text{HE}}$ versus the HE mass. The observed ratio increases much more steeply than the simulated ratio in RA12.

For the CCCP and the CLASH samples, we could restrict the analysis to either CC clusters or systems with low offsets between the BCG and the X-ray surface brightness peak. The HE mass of CC clusters in the CCCP sample is less biased. Due to the limited number of clusters, we could not confirm this result for the CLASH sample. There is no evident trend for the low off-sets clusters.

We verified a posteriori how well the statistical model reproduce the observed trends of the mass ratio, see Figs 6–8. We generated a number of ‘true’ masses from the normal distributions derived in the regression analyses and the corresponding ‘true’ latent WL and HE masses, scattered according to the measured σ_{WL} and σ_{HE} . The observed manifest WL and HE masses were finally generated considering the measured statistical uncertainties. The masses were then binned as the real clusters. The

observed trends in bias and scatter are well recovered. This test further validates a posteriori the assumptions of the Bayesian modelling, in particular that scatters are approximately lognormal as well as that the mass distribution can be described as a Gaussian.

The observed trends were also compared to the numerical simulations, see Figs 6–8. The extent of this comparison is limited. First, the simulated clusters are not representative of the observed samples, even though both observed and simulated clusters should sample the tail at large values of the halo mass function. The discrepancy is further mitigated since we compared clusters in the same mass bins. Nevertheless, the dynamical and morphological properties of the simulated clusters may differ from the observed ones. Secondly, WL and HE masses of the simulated clusters were measured within the true r_{500} differently from the real clusters, whose overdensity radii were estimated based on the WL/HE mass. This may limit the scatter of the simulated clusters compared to the real ones.

Due to these differences only qualitative considerations can be made. Nevertheless some trends seem to further validate the picture detailed before. In particular, the observed mass bias $M^{\text{HE}}/M^{\text{WL}}$ scales with the WL mass as in simulations, which further supports that the scatter of measured WL masses agree with theoretical predictions. On the other hand, the observed mass bias as a function of the HE mass is significantly steeper than that in the simulated sample, which points to a larger than theoretically predicted scatter in HE masses.




Figure 7. Same as Fig. 6 but for the clusters in the CLASH sample with *Chandra*-based X-ray analyses. WL (HE) masses are measured within r_{500}^{WL} (r_{500}^{HE}).

6 DISCUSSION

The main sources of bias and scatter in WL mass measurements are due to the presence of substructures and triaxiality (RA12; Giocoli et al. 2014). These effects are dominated by the dark matter component and are more easily reproduced in numerical simulations than the more complex processes involving gas physics. Reassuringly, the level of scatter we ascertained from observations is in very good agreement with the theoretical prediction of $\sigma_{\text{WL}} \gtrsim 10$ per cent.

The observed relative bias between HE and WL masses is consistent within the statistical errors with predictions from simulations (RA12). Based on a suite of different numerical simulations (Battaglia et al. 2012; Kay et al. 2012), Planck Collaboration XX (2014) estimated $b = 1 - M_{\text{HE}}^{500}/M_{\text{Tr}}^{500} = 0.2^{+0.1}_{-0.2}$. However, the inherent uncertainty in the HE/WL calibrations prevents firmer conclusions. For some samples, the bias is as large as ~ 50 per cent.

The measured intrinsic scatter in HE masses is larger than the theoretical prediction. The disagreement may hinge on several plausible causes. The formal statistical uncertainty in X-ray mass estimates is usually of the order of ~ 10 – 15 per cent. However, the observed discrepancies among mass estimates from independent analyses are as large as 45 per cent (~ 30 per cent within the same physical radius). The underestimation of the formal error on the HE masses could determine an overestimation of the intrinsic scatter.

6.1 Gas physics

Scatter in simulations may be underestimated due to their current limits (Rasia et al. 2014). Estimates of bias and scatter from

numerical simulations are still uncertain, showing dependences on the physical treatment of the gas, and, possibly, on the hydrodynamical scheme adopted. Each simulation suite has circumstantial prescriptions for gas physics. Different treatments of radiative cooling and cooling/heating from the UV background play an important role. Thermal conduction in hot clusters may be effective in removing cold blobs and in making the thermal structure of the ICM more homogeneous. This leads to an increase of the spectroscopic temperature and therefore of the hydrostatic mass. Feedback from active galactic nuclei and supernovae can significantly reduce the temperature inhomogeneity.

The impact of each ingredient is significant and each process may be more or less effective in different clusters. Theoretical predictions based on specific descriptions may then significantly underestimate the intrinsic scatter in the HE mass.

6.2 Simulation scheme

Some disagreement among theoretical predictions is also caused by the adopted simulation scheme. Smoothed particle hydrodynamics (SPH) simulations produce larger temperature variations connected to the persistence of both substructures and their stripped cold gas than adaptive-mesh-refinement (AMR) codes (Sijacki, Springel & Haehnelt 2011; Rasia et al. 2014), which lead to a more efficient mixing of gas entropy. Low-entropy gas residing in high-density clumps is more efficiently mixed to the high-entropy ICM than in SPH simulations. The simulated temperature distribution is then more homogenous and the relative bias introduced in the estimate of X-ray temperature is smaller (Vazza et al. 2011). Around r_{500} , the




Figure 8. Same as Fig. 6 but for the clusters in the CLASH sample with *XMM*-based X-ray analyses. WL (HE) masses are measured within r_{500}^{WL} (r_{500}^{HE}).

temperature inhomogeneities of the SPH simulations can generate twice the typical HE mass bias of the AMR sample (Rasia et al. 2014).

These variations between simulation schemes make predictions less certain. A better understanding of the physical processes responsible for the complex thermal structure in ICM requires improved resolution and high-sensitivity observations, first of all for higher temperature systems and larger cluster-centric radii (Rasia et al. 2014).

6.3 Mass dependence

A further source of disagreement might be ascribed to any dependence of the bias on cluster mass. Neglecting such dependence can inflate the estimate of the scatter. The massive objects are expected to be the most disturbed ones, and they should have a complex temperature structure (RA12). This would imply a bias larger for the more massive clusters. We tested this hypothesis by repeating the analysis of Section 5 without fixing the slope β_{HE} to unity.

Due to the addition of a new free parameter to be determined with the regression, we could obtain well-constrained results only for the two richer samples. For the CCCP sample (all masses within r_{500}^{WL}), we obtained $\sigma_{\text{WL}} = 0.17 \pm 0.07$, $\beta_{\text{HE}} = 1.19 \pm 0.24$ and $\sigma_{\text{HE}} = 0.20 \pm 0.09$. For the CLASH-CXO sample, we obtained $\sigma_{\text{WL}} = 0.21 \pm 0.10$, $\beta_{\text{HE}} = 1.29 \pm 0.63$ and $\sigma_{\text{HE}} = 0.30 \pm 0.16$.

For both samples, with respect to the results obtained fixing $\beta_{\text{HE}} = 1$, the measured σ_{WL} is slightly larger whereas σ_{HE} is smaller. However, σ_{HE} is still larger than both σ_{WL} and the scatter predicted by numerical simulations.

The estimated β_{HE} is slightly larger than unity, but still consistent within the statistical uncertainty. This scenario would then imply a still larger than expected scatter in HE masses at the expense of a not so plausible bias decreasing with mass ($\beta_{\text{HE}} > 1$). This alternative scenario is then more complex but does not solve the main incongruences it was supposed to address. Since the estimated β_{HE} is consistent with unity within the errors, we then disfavour this scenario.

7 CONCLUSIONS

In this paper, the first in a series which aims to critically revise the status quo in measuring cluster masses and calibrating scaling relations, we studied the biases and the intrinsic scatters of WL and hydrostatic masses. Either WL or HE masses determined from different groups may differ by ~ 40 per cent, which hinders the absolute calibration of any scaling relation and the assessment of the relative bias between WL and HE masses.

We found that the intrinsic scatter of WL masses is of the order of ~ 10 – 15 per cent, in line with theoretical predictions. The intrinsic scatter of HE masses turned out to be larger, ~ 20 – 30 per cent, at odds with results from numerical simulations. The discrepancies may hinge on underestimated statistical uncertainties in HE masses. A better understanding of the physical processes responsible for the complex thermal structure in the ICM and improved simulation schemes are also required to improve the theoretical predictions.

Most of the sources of scatter in the estimates of WL and HE masses are of well-known origin. The assumption of spherical symmetry causes an over- or underestimate of the WL mass whether

the cluster is elongated in the plane of the sky or towards the observer, respectively. Departures from HE or the difficult assessment of non-thermal contribution to the pressure limit the accuracy of HE masses.

Oversimplified modelling inflates the intrinsic scatter. The joint analysis of multiwavelength observations, from the X-ray to the optical band to the SZ effect in the radio, can provide unbiased estimates of the cluster mass (De Filippis et al. 2005; Sereno et al. 2006, 2013; Morandi et al. 2012; Sereno, Ettori & Baldi 2012; Limousin et al. 2013). In fact, the combined information from the different data sets enables us to recover the triaxial structure and the orientation of the cluster and to quantify the non-thermal contributions to the pressure.

An alternative approach is focusing on well-behaved clusters, where bias and scatter are intrinsically small. Biases are lower in morphologically regular and isolated clusters (RA12). However, there are a few of them and they are rare to find. Even apparently spherical clusters with a regular morphology might significantly deviate from hydrostatic thermal equilibrium (Sereno et al. 2013). Furthermore, a projected circular shape is well suited to either spherical systems or strongly prolate haloes elongated along the line of sight towards the observer.

Scatter and bias in WL and X-ray estimates play a fundamental role in the calibration of mass proxies. Ongoing programmes are making significant efforts to understand the sources of systematics and to solve the related calibration issues (Donahue et al. 2014; Rozo et al. 2014b). We quantified the size of the intrinsic scatter in WL and HE masses and discussed the effect of scatter in the determination of scaling relations. The scatter makes relation systematically flatter and more scattered. Proper statistical treatments could and should account for this.

ACKNOWLEDGEMENTS

The authors thank Lauro Moscardini and Elena Rasia for useful discussions. MS acknowledges financial contributions from contracts ASI/INAF n.I/023/12/0 ‘Attività relative alla fase B2/C per la missione Euclid’, PRIN MIUR 2010-2011 ‘The dark Universe and the cosmic evolution of baryons: from current surveys to Euclid’, and PRIN INAF 2012 ‘The Universe in the box: multiscale simulations of cosmic structure’. SE acknowledges the financial contribution from contracts ASI-INAF I/009/10/0 and PRIN-INAF 2012 ‘A unique data set to address the most compelling open questions about X-Ray Galaxy Clusters’.

REFERENCES

Akritas M. G., Bershadsky M. A., 1996, *ApJ*, 470, 706
 Andreon S., Bergé J., 2012, *A&A*, 547, A117
 Andreon S., Hurn M. A., 2010, *MNRAS*, 404, 1922
 Andreon S., Hurn M. A., 2012, preprint ([arXiv:1210.6232](https://arxiv.org/abs/1210.6232))
 Applegate D. E. et al., 2014, *MNRAS*, 439, 48
 Arnaud M., Pratt G. W., Piffaretti R., Böhringer H., Croston J. H., Pointecouteau E., 2010, *A&A*, 517, A92
 Battaglia N., Bond J. R., Pfrommer C., Sievers J. L., 2012, *ApJ*, 758, 74
 Becker M. R., Kravtsov A. V., 2011, *ApJ*, 740, 25
 Biviano A. et al., 2013, *A&A*, 558, A1
 Bonamente M. et al., 2012, *New J. Phys.*, 14, 025010 (B12)
 D’Agostini G., 2004, preprint ([astro-ph/0403086](https://arxiv.org/abs/astro-ph/0403086))
 De Filippis E., Sereno M., Bautz M. W., Longo G., 2005, *ApJ*, 625, 108
 Donahue M. et al., 2014, *ApJ*, 794, 136
 Ebeling H., Edge A. C., Allen S. W., Crawford C. S., Fabian A. C., Huchra J. P., 2000, *MNRAS*, 318, 333

Ebeling H., Edge A. C., Mantz A., Barrett E., Henry J. P., Ma C. J., van Speybroeck L., 2010, *MNRAS*, 407, 83
 Eddington A. S., 1913, *MNRAS*, 73, 359
 Eddington Sir A. S., 1940, *MNRAS*, 100, 354
 Ettori S., 2013, *MNRAS*, 435, 1265
 Ettori S., Morandi A., Tozzi P., Balestra I., Borgani S., Rosati P., Lovisari L., Terenzi F., 2009, *A&A*, 501, 61
 Ettori S., Gastaldello F., Leccardi A., Molendi S., Rossetti M., Buote D., Meneghetti M., 2010, *A&A*, 524, A68 (E10)
 Fabjan D., Borgani S., Rasia E., Bonafede A., Dolag K., Murante G., Tornatore L., 2011, *MNRAS*, 416, 801
 Feigelson E. D., Babu G. J., 2012, *Modern Statistical Methods for Astronomy*. Cambridge Univ. Press, Cambridge
 Giocoli C., Meneghetti M., Ettori S., Moscardini L., 2012, *MNRAS*, 426, 1558
 Giocoli C., Meneghetti M., Metcalf R. B., Ettori S., Moscardini L., 2014, *MNRAS*, 440, 1899
 Giodini S., Lovisari L., Pointecouteau E., Ettori S., Reiprich T. H., Hoekstra H., 2013, *Space Sci. Rev.*, 177, 247
 Hoekstra H., Mahdavi A., Babul A., Bildfell C., 2012, *MNRAS*, 427, 1298
 Jeffreys H., 1938, *MNRAS*, 98, 190
 Kay S. T., Peel M. W., Short C. J., Thomas P. A., Young O. E., Battye R. A., Little A. R., Pearce F. R., 2012, *MNRAS*, 422, 1999
 Kelly B. C., 2007, *ApJ*, 665, 1489
 Landry D., Bonamente M., Giles P., Maughan B., Joy M., Murray S., 2013, *MNRAS*, 433, 2790 (L13)
 LaRoque S. J., Bonamente M., Carlstrom J. E., Joy M. K., Nagai D., Reese E. D., Dawson K. S., 2006, *ApJ*, 652, 917
 Laureijs R. et al., 2011, preprint ([arXiv:1110.3193](https://arxiv.org/abs/1110.3193))
 Lemze D., Broadhurst T., Rephaeli Y., Barkana R., Umetsu K., 2009, *ApJ*, 701, 1336
 Limousin M., Morandi A., Sereno M., Meneghetti M., Ettori S., Bartelmann M., Verdugo T., 2013, *Space Sci. Rev.*, 177, 155
 Mahdavi A., Hoekstra H., Babul A., Bildfell C., Jeltema T., Henry J. P., 2013, *ApJ*, 767, 116
 Mantz A., Allen S. W., Rapetti D., Ebeling H., 2010, *MNRAS*, 406, 1759
 Martino R., Mazzotta P., Bourdin H., Smith G. P., Bartalucci I., Marrone D. P., Finoguenov A., Okabe N., 2014, *MNRAS*, 443, 2342
 Mazzotta P., Rasia E., Moscardini L., Tormen G., 2004, *MNRAS*, 354, 10
 Mezdziński E., Broadhurst T., Umetsu K., Oguri M., Rephaeli Y., Benítez N., 2010, *MNRAS*, 405, 257
 Meneghetti M., Rasia E., Merten J., Bellagamba F., Ettori S., Mazzotta P., Dolag K., Marri S., 2010, *A&A*, 514, A93
 Merten J. et al., 2014, preprint ([arXiv:1404.1376](https://arxiv.org/abs/1404.1376))
 Morandi A. et al., 2012, *MNRAS*, 425, 2069
 Navarro J. F., Frenk C. S., White S. D. M., 1996, *ApJ*, 462, 563 (NFW)
 Nevalainen J., David L., Guainazzi M., 2010, *A&A*, 523, A22
 Oguri M., Takada M., Umetsu K., Broadhurst T., 2005, *ApJ*, 632, 841
 Piffaretti R., Valdarnini R., 2008, *A&A*, 491, 71
 Planck Collaboration XX, 2014, *A&A*, 571, A20
 Planck Collaboration XXIX, 2014, *A&A*, 571, A29
 Postman M. et al., 2012, *ApJS*, 199, 25
 Pratt G. W., Croston J. H., Arnaud M., Böhringer H., 2009, *A&A*, 498, 361
 Rasia E. et al., 2012, *New J. Phys.*, 14, 055018 (RA12)
 Rasia E. et al., 2014, *ApJ*, 791, 96
 Rines K., Diaferio A., 2006, *AJ*, 132, 1275
 Rozo E., Rykoff E. S., Bartlett J. G., Evrard A., 2014a, *MNRAS*, 438, 49
 Rozo E., Bartlett J. G., Evrard A. E., Rykoff E. S., 2014b, *MNRAS*, 438, 78
 Sereno M., 2007, *MNRAS*, 380, 1207
 Sereno M., 2015, *MNRAS*, 450, 3665 (CoMaLit-III)
 Sereno M., Ettori S., 2015, *MNRAS*, 450, 3675 (CoMaLit-IV)
 Sereno M., Umetsu K., 2011, *MNRAS*, 416, 3187
 Sereno M., De Filippis E., Longo G., Bautz M. W., 2006, *ApJ*, 645, 170
 Sereno M., Ettori S., Baldi A., 2012, *MNRAS*, 419, 2646
 Sereno M., Ettori S., Umetsu K., Baldi A., 2013, *MNRAS*, 428, 2241
 Sereno M., Ettori S., Moscardini L., 2015, *MNRAS*, 450, 3649 (CoMaLit-II)
 Sijacki D., Springel V., Haehnelt M. G., 2011, *MNRAS*, 414, 3656
 Umetsu K. et al., 2014, *ApJ*, 795, 163

Vazza F., Dolag K., Ryu D., Brunetti G., Gheller C., Kang H., Pfrommer C., 2011, MNRAS, 418, 960
 Vikhlinin A. et al., 2009, ApJ, 692, 1060
 Voit G. M., 2005, Rev. Mod. Phys., 77, 207
 von der Linden A. et al., 2014, MNRAS, 439, 2

APPENDIX A: BIAS

The intrinsic scatter biases the average of the intrinsic variables with respect to the observable proxy. Here, we focus on intrinsic scatter and we assume that observational uncertainties are negligible. Let us consider a proxy X of the independent variable Z ,

$$X = \alpha_{X|Z} + \beta_{X|Z}Z \pm \sigma_{X|Z}, \quad (\text{A1})$$

where $\alpha_{X|Z}$ is the bias, $\beta_{X|Z}$ accounts for rescaling, and $\sigma_{X|Z}$ is the intrinsic scatter of the linear relation. We assume that the scatter is Gaussian,

$$p(X|Z) \propto \exp\left[-\frac{1}{2}\left(\frac{X - (\alpha_{X|Z} + \beta_{X|Z}Z)}{\sigma_{X|Z}}\right)^2\right]. \quad (\text{A2})$$

The intrinsic scatter can bias average values if the true variables are not uniformly distributed. The average intrinsic value of an ensemble of objects with the same measured X can differ from X . The average is given by

$$\langle Z \rangle(X) \propto \int Z p(X, Z) dZ. \quad (\text{A3})$$

The analytical treatment of the bias is particularly simple when the distribution of the intrinsic variable is Gaussian,

$$p(Z) \propto \exp\left[-\frac{1}{2}\left(\frac{Z - \mu_Z}{\sigma_Z}\right)^2\right]. \quad (\text{A4})$$

A simple application of the Bayes' theorem shows that $p(X)$ is normally distributed too, with the same mean μ_Z and standard deviation

$$\sigma_X = \sqrt{\sigma_Z^2 + \sigma_{X|Z}^2}. \quad (\text{A5})$$

In this case, the integral in equation (A3) can be solved in terms of simple functions,

$$\langle Z \rangle(X) = b_\sigma X + \Delta b_\sigma, \quad (\text{A6})$$

where b_σ is the multiplicative bias due to the intrinsic scatter,

$$b_\sigma = \frac{1}{\beta_{X|Z}} \frac{1}{1 + \sigma_{X|Z}^2 / (\beta_{X|Z} \sigma_Z)^2}, \quad (\text{A7})$$

and Δb_σ is an additive bias,

$$\Delta b_\sigma = \left(\mu_Z \frac{\sigma_{X|Z}^2}{\beta_{X|Z}^2 \sigma_Z^2} - \frac{\alpha_{X|Z}}{\beta_{X|Z}} \right) \frac{1}{1 + \sigma_{X|Z}^2 / (\beta_{X|Z} \sigma_Z)^2}. \quad (\text{A8})$$

The contribution to the bias due to intrinsic scatter is negligible either if the intrinsic scatter is very small ($\sigma_{X|Z} \rightarrow 0$) or if the intrinsic variable is uniformly distributed ($\sigma_Z \rightarrow \infty$).

If we have to calibrate a linear relation,

$$Y = \alpha_{Y|Z} + \beta_{Y|Z}Z, \quad (\text{A9})$$

but we have only measurements of the proxy X instead of Z , we cannot just study the relation

$$Y = \alpha_{Y|X} + \beta_{Y|X}X, \quad (\text{A10})$$

and take $\beta_{Y|X}$ as an unbiased estimator of $\beta_{Y|Z}$. In the Gaussian case,

$$\alpha_{Y|X} = \alpha_{Y|Z} + \Delta b_\sigma \beta_{Y|Z}, \quad (\text{A11})$$

$$\beta_{Y|X} = b_\sigma \beta_{Y|Z}. \quad (\text{A12})$$

The intrinsic scatter apparently flattens the slope of the relation. The most convenient statistical approach requires a proper treatment of the selection effects and of the intrinsic scatter in the linear regression. De-biasing the data as suggested in equation (A6) would correct each measured scattered proxy by the expected mean bias instead of the actual one, which is random for objects with the same measured X . Nevertheless this mean correction may provide a useful tool to quickly evaluate the effect of the intrinsic scatter.

If the intrinsic scatter is lognormal, as it is usually the case, in the above discussion X can be read, for example, as $\ln M^{\text{WL}}$ or $\ln M^{\text{HE}}$, whereas Z stands for $\ln M^{\text{Tr}}$.

APPENDIX B: BCES

The BCES method is a well-known regression technique with good performance for data sets with heteroscedastic and correlated errors on both axes as well as intrinsic scatter in the linear relation (Akritas & Bershady 1996). The slope of the conditional linear relation in equation (A10) can be estimated as

$$\beta(Y|X) = \frac{\sum_i (y_i - \langle y \rangle)(x_i - \langle x \rangle) - \sum_i \delta_{xy,i}}{\sum_i (x_i - \langle x \rangle)^2 - \sum_i \delta_{x,i}^2}, \quad (\text{B1})$$

where x_i (y_i) is the observed values of X_i (Y_i), with associated observational uncertainty $\delta_{x,i}$ ($\delta_{y,i}$), and $\delta_{xy,i}$ is the covariance between errors; $\langle x \rangle$ and $\langle y \rangle$ are the mean values.

If the variable X is scattered too, see equation (A1), the slope in equation (B1) is a biased estimator of $\beta_{Y|Z}$. For $\alpha_{X|Z} = 0$ and $\beta_{X|Z} = 1$, the unbiased slope is

$$\beta(Y|Z) = \frac{\sum_i (y_i - \langle y \rangle)(x_i - \langle x \rangle) - \sum_i \delta_{xy,i}}{\sum_i (x_i - \langle x \rangle)^2 - \sum_i (\delta_{x,i}^2 + \sigma_{X|Z}^2)}. \quad (\text{B2})$$

The modified BCES estimator should be used to evaluate the slope of the conditional linear relation if the covariate X is scattered, i.e. X is the scattered response variable of the covariate Z .

APPENDIX C: MASSES AND COSMOLOGY

Masse estimates depend on the cosmological model. A conversion from other cosmological parameters may be required to convert to a reference model. The mass within a given cosmological overdensity Δ is defined as

$$M_\Delta = \frac{4\pi}{3} \Delta \rho_{\text{cr}} (D_d \theta_\Delta)^3, \quad (\text{C1})$$

where θ_Δ is the angular radius enclosing the overdensity and D_d is the angular diameter distance to the cluster.

Lensing 3D masses within a radius $r = D_d \theta$, where θ is the aperture radius, scale as

$$M^{\text{WL}} \propto \Sigma_{\text{cr}}(D_d \theta_E) D_d \theta f(\theta), \quad (\text{C2})$$

where $\Sigma_{\text{cr}} \equiv (c^2 D_s) / (4\pi G D_d D_{\text{ds}})$ is the critical surface density for lensing, θ_E is the angular Einstein radius and D_s and D_{ds} are the source and the lens-source angular diameter distances, respectively. The function $f(\theta) \sim \theta^{\delta\gamma}$ quantifies the deviation of the mass

profile from the isothermal case. At r_{500} , mass profiles are nearly isothermal, i.e. $\delta\gamma \sim 0$.

Solving for equations (C1) and (C2), we obtain

$$M_{\Delta}^{\text{WL}} \propto D_{\text{d}}^{-\frac{3\delta\gamma}{2-\delta\gamma}} \left(\frac{D_{\text{ds}}}{D_{\text{s}}} \right)^{-\frac{3}{2-\delta\gamma}} H(z)^{-\frac{1+\delta\gamma}{1-\delta\gamma/2}} \quad (\text{C3})$$

$$= \left(\frac{D_{\text{ds}}}{D_{\text{s}}} \right)^{-3/2} H(z)^{-1} \text{ for } \delta\gamma = 0. \quad (\text{C4})$$

Hydrostatic masses within θ scales as

$$M^{\text{HE}} \propto D_{\text{d}} \theta^{1+\delta\gamma}; \quad (\text{C5})$$

the HE mass within a given cosmological overdensity is then

$$M_{\Delta}^{\text{HE}} \propto D_{\text{d}}^{-\frac{3\delta\gamma}{2-\delta\gamma}} H(z)^{-\frac{1+\delta\gamma}{1-\delta\gamma/2}} \quad (\text{C6})$$

$$= H(z)^{-1} \text{ for } \delta\gamma = 0. \quad (\text{C7})$$

For $\delta\gamma = 0$, $M_{\Delta}^{\text{HE}}/r_{\Delta}$ is independent of the adopted cosmology. When it was required, we used the above relations with $\delta\gamma = 0$ to make the proper conversion from different cosmological parameters. We refer to [CoMaLit-III](#) for further details. An analogue treatment for X-ray observables can be found in Mantz et al. (2010).

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.