














Publication Year	2022
Acceptance in OA	2022-03-17T13:37:30Z
Title	Hiding in plain sight: observing planet-starspot crossings with the James Webb Space Telescope
Authors	BRUNO, GIOVANNI, Nikole K. Lewis, Jeff A. Valenti, PAGANO, Isabella, Tom J. Wilson, Everett Schlawin, Joshua Lothringer, LANZA, Antonino Francesco, Jonathan Fraine, SCANDARIATO, GAETANO, MICELA, Giuseppina, CRACCHIOLO, Gianluca
Publisher's version (DOI)	10.1093/mnras/stab3199
Handle	http://hdl.handle.net/20.500.12386/31666
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
Volume	509

Hiding in plain sight: observing planet-starspot crossings with the *James Webb Space Telescope*

Giovanni Bruno ^{1,★}, Nikole K. Lewis ², Jeff A. Valenti ³, Isabella Pagano ¹, Tom J. Wilson ⁴,
Everett Schlawin ⁵, Joshua Lothringer ^{6,7}, Antonino F. Lanza ¹, Jonathan Fraine ^{8,9},
Gaetano Scandariato ¹, Giuseppina Micela ¹¹ and Gianluca Cracchiolo^{10,11}

¹INAF – Catania Astrophysical Observatory, Via Santa Sofia, 78, I-95123 Catania, Italy

²Department of Astronomy and Carl Sagan Institute, Cornell University, 122 Sciences Drive, Ithaca, NY 14853, USA

³Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

⁴University of Exeter, Physics Building, Stocker Road, Exeter EX4 4QL, UK

⁵University of Arizona, 933 Cherry Ave, Tucson, AZ 85721, USA

⁶Department of Physics and Astronomy, Johns Hopkins University, Baltimore, MD 21210, USA

⁷Department of Physics, Utah Valley University, 800 W. University Parkway, MS 179, Orem, UT 84058, USA

⁸Space Science Institute Center for Data Science, 4765 Walnut St., Suite B, Boulder, CO 80301, USA

⁹Space Science Institute Center for Exoplanet and Planetary Science, 4765 Walnut St, Suite B, Boulder, CO 80301, USA

¹⁰Dipartimento di Fisica e Chimica Emilio Segré, Università, di Palermo, Via Archirafi 36, 90123 Palermo, Italy

¹¹INAF – Osservatorio Astronomico di Palermo, P.zza Parlamento 1, I-90134 Palermo, Italy

Accepted 2021 October 27. Received 2021 October 25; in original form 2020 October 1

ABSTRACT

Transiting exoplanets orbiting active stars frequently occult starspots and faculae on the visible stellar disc. Such occultations are often rejected from spectrophotometric transits, as it is assumed they do not contain relevant information for the study of exoplanet atmospheres. However, they can provide useful constraints to retrieve the temperature of active features and their effect on transmission spectra. We analyse the capabilities of the *James Webb Space Telescope* in the determination of the spectra of occulted starspots, despite its lack of optical wavelength instruments on board. Focusing on K and M spectral types, we simulate starspots with different temperatures and in different locations of the stellar disc, and find that starspot temperatures can be determined to within a few hundred kelvins using NIRSpec/Prism and the proposed NIRCам/F150W2+F322W2’s broad wavelength capabilities. Our results are particularly promising in the case of K and M dwarfs of $\text{mag}_K \lesssim 12.5$ with large temperature contrasts.

Key words: techniques: photometric – techniques: spectroscopic – planets and satellites: atmospheres – starspots.

1 INTRODUCTION

The contamination from the stellar signal in transit spectrophotometry is one of the main challenges to precisely characterize exoplanet atmospheres. Stellar active features, such as dark starspots and bright faculae, both hamper the correct measure of transit parameters and introduce spurious spectral features that overlap with the exoplanetary spectral lines. A large body of observational evidence is now available about general starspot properties as a function of stellar type (e.g. Berdyugina 2005; Strassmeier 2009; Balona & Abedigamba 2016; Nielsen et al. 2019; Savanov 2019; Herbst et al. 2021), and important progress in the theoretical starspot modelling has been made (e.g. Yadav et al. 2015; Panja, Cameron & Solanki 2020). Predicting the appearance and properties of stellar active features remains however elusive and, in the interpretation of an exoplanet observation which is potentially contaminated by stellar activity, the statistical knowledge of starspots and faculae

only provides with partial guidance. For example, different starspot sizes and temperatures have been shown to have a dramatically different impact on transmission spectra (e.g. Zellem et al. 2017; Rackham, Apai & Giampapa 2018, 2019, and references therein), and for individual stars we might not be able to recognize the most likely scenario without additional observational constraints (such as e.g. photometric monitoring, Huitson et al. 2013).

Moreover, the impact of stellar active features in spectrophotometry is at least twofold, according to whether they lie on the part of the stellar disc intersected by the transit chord or not. The so-called ‘non-occulted’ features affect several transit parameters and cannot be easily identified with single-transit information alone (e.g. Alonso et al. 2008; Pont et al. 2008, 2013; Czesla et al. 2009; Léger et al. 2009; Silva-Valio & Lanza 2011; Sing et al. 2011; Ballerini et al. 2012; Barros et al. 2013; Csizmadia et al. 2013; McCullough et al. 2014; Oshagh et al. 2014; Barstow et al. 2015; Rackham et al. 2017, 2018, 2019; Zellem et al. 2017; Alam et al. 2018; Wakeford et al. 2019). For this reason, long-duration photometric observations of the star need to be secured around the transits, in order to monitor its activity level. One downfall of this method is, however, that

* E-mail: giovanni.bruno@inaf.it

the coverage fraction of the stellar disc in active features and their temperature are strongly correlated parameters: usually, the starspot temperature is assumed in order to extract a wavelength-dependent correction factor for the transit depth, or vice versa (e.g. Alam et al. 2018). Simultaneous multiwavelength observations were shown to be promising approaches to break this degeneracy (e.g. Rosich et al. 2020; Cracchiolo, Micela & Peres 2021).

‘Occulted’ active features have an opposite effect on the transit depth compared to non-occulted features, and can often be easily identified in the transit profile through a typical flux bump (or dip, depending on whether they are starspots or faculae). They can therefore be modelled (e.g. Huber et al. 2010; Sing et al. 2011; Pont et al. 2013; Béky, Kipping & Holman 2014; Fraine et al. 2014; Montalto et al. 2014; Tregloan-Reed et al. 2015; Bruno et al. 2016; Espinoza et al. 2017; Mancini et al. 2017; Louden et al. 2017; Scandariato et al. 2017) and provide useful constraints to characterize both the planet and the host star. With a good enough signal-to-noise ratio (SNR), their temperature, size, and coordinates can be determined, even if a certain degree of degeneracy is unavoidable (Béky et al. 2014; Mancini et al. 2017; Morris et al. 2017; Scandariato et al. 2017). In combination with out-of-transit photometric observations, occulted starspots can provide some indications on the temperature and size of the active features on the stellar disc, together with some priors on their filling factor. This attractive possibility can be applied when the SNR of the occultation is large, while the impact of occulted starspots and faculae becomes more subtle when they are so small and tightly grouped that they do not stand out against the transit profile. In this case, they can potentially bias the determination of the transit bottom altogether in a similar, even if opposite, way to non-occulted features (e.g. Czesla et al. 2009; Ballerini et al. 2012). The impact of such an unresolved background of active features, similarly to the one of stellar granulation, could be particularly relevant for terrestrial planets orbiting Sun-like stars (Sarkar et al. 2018).

The effect of starspots and faculae on transmission spectroscopy is stronger in the visible, as that is where their brightness contrast with the stellar photosphere is the largest. For this reason photometry, as well as spectroscopy in the visible, were successfully used both from the ground and from space to constrain some of the properties of the occulted active features accidentally detected during transit observations (e.g. Silva-Valio et al. 2010; Sing et al. 2011; Béky et al. 2014; Mancini et al. 2017).

As the starspot brightness contrast is lower in the near-infrared (NIR), the occultation signal often falls below the noise level, or however at a level where it does not enable the robust determination of starspot parameters (e.g. Bruno et al. 2018). For FGK stars, it is generally assumed that both the impact of occulted and non-occulted active features is negligible in the NIR (e.g. Zellem et al. 2017; Rackham et al. 2019), while the contamination spectrum was shown to be much more problematic for M dwarfs (Rackham et al. 2018; Iyer & Line 2020). At the same time, active features produce small but measurable variations in the limb darkening coefficients, which can be measured in this spectral region (Morello et al. 2017).

Exoplanet searches are increasingly focusing on small, cool stars, as they are favourable targets for the detection and characterization of low-mass planets in the habitable zone; we expect many more interesting targets than the ones already known to be revealed by the analysis of the *Transiting Exoplanet Survey Satellite* (TESS) data (Ricker et al. 2014). For this reason, as well as the imminent launch of the *James Webb Space Telescope* (JWST), it is crucial to identify realistic strategies to determine the properties of active features and their level of spectral contamination for exoplanets orbiting low-mass stars. Moreover, high-precision observations of

starspot and facula crossings could also provide important constraints for planets orbiting warmer stars (e.g. Beaulieu et al. 2008; Désert et al. 2011; Bruno et al. 2020), which will become the focus of future searches for long-period, terrestrial planets (such as *PLATO*, Rauer et al. 2014). Such information would come for free from transmission spectroscopy observations, and would be valuable both for the purpose of exoplanet characterization and to better understand stellar activity itself.

In this work, we study the possibilities opened up by *JWST* to constrain the temperature of occulted active features on K and M stars in several geometric configurations and activity levels. We focus on dark starspots, assuming that the same line of reasoning we use here can be adopted for bright faculae. Section 2 describes our working framework, and in Section 3 we present our simulations; the results of our analysis are detailed in Section 4. Section 5 is dedicated to the conclusions that can be drawn from this work, as well as to future perspectives.

2 CONTRAST RATIO OF ACTIVE FEATURES

The brightness contrast between an active feature and the stellar photosphere, $A_{\lambda, \mu}$, is a function of wavelength λ and $\mu \equiv \cos \theta$, where θ is the angle between the normal to the stellar surface and the line of sight. It can be expressed as (e.g. Silva 2003)

$$A_{\lambda, \mu} \equiv 1 - \frac{I_{\bullet}(\lambda, \mu)}{I_{\star}(\lambda, \mu)}, \quad (1)$$

where $I_{\bullet}(\lambda, \mu)$ and $I_{\star}(\lambda, \mu)$ are the active feature and stellar specific intensity, respectively. In this formulation, $A_{\lambda, \mu} = 1$ denotes maximum contrast – a completely dark spot – and $A_{\lambda, \mu} = 0$ no contrast at all – feature and stellar photosphere are indistinguishable. A negative contrast indicates a warmer structure, such as a facula, which is brighter than the stellar photosphere.

The intensity of active features can be represented by specific intensity model spectra with a different effective temperature than the one describing the host star. In addition, starspots are thought to be characterized by a 0.5–1 dex lower $\log g$ than the stellar one, because of the decrease in gas pressure caused by increased magnetic pressure in the darker photospheric upper layers (e.g. Solanki 2003).

As both I_{\bullet} and I_{\star} depend on μ , the effect of limb darkening (LD) needs to be included in the calculation of the contrast (e.g. Dorren 1987). Given this and the complications in describing the limb-brightening behaviour of faculae (e.g. Spruit 1976; Keller et al. 2004; Solov’ev & Kirichek 2019), in our analysis we focused on dark starspots. The limb-angle dependent specific intensity stellar models were computed with the PHOENIX stellar atmosphere models (Hauschildt, Allard & Baron 1999). We calculated a baseline model for both the M-dwarf ($T_{\star} = 3500$ K, $\log g = 5.0$) and K-dwarf ($T_{\star} = 5000$ K, $\log g = 4.5$), and then computed models for the starspots every 100 K between 2300 K and 3400 K for the M-dwarf and between 3600 K and 4900 K for the K-dwarf. The M-dwarf starspot models had a $\log g = 4.5$ and the K-dwarf starspot models had $\log g = 4.0$. All models assumed solar metallicity. For each model, once the temperature structure was converged, we calculated the specific intensity through the atmosphere at 51 different μ values between 0 and 1 at every Ångstrom between 6000 and 5.35 Å.

Fig. 1 presents a set of starspot contrast spectra for a $T_{\star} = 3500$ K and a $T_{\star} = 5000$ K star with increasingly cooler spots, calculated from PHOENIX stellar models. The spectra were downgraded to a resolution $R = 100$, considering the NIRSpc/Prism’s resolution in the spectral range 0.6 – 5.3 μm . Below $\simeq 4$ μm , several absorption features, mainly due to water vapour, increase in strength as




Figure 1. Starspot versus stellar photosphere brightness contrast ratios at the centre of the stellar disc ($\mu = 1$), as a function of wavelength, for different stars and starspot effective temperatures (equation 1). The contrast ratios were obtained with PHOENIX specific intensity models and downgraded to resolution $R = 100$. Different starspot effective temperatures are represented with different colours, as indicated in the top right corner of each panel. *Left:* Case of a 3500 K, $\log g = 5.0$, $[\text{Fe}/\text{H}] = 0.0$ star and starspots with varying T_{\bullet} , $\log g = 4.5$ and the same metallicity. *Right:* Case of a 5000 K, $\log g = 4.5$, $[\text{Fe}/\text{H}] = 0.0$ star, and starspots with varying T_{\bullet} , $\log g = 4.0$ and the same metallicity. Water vapour bands are visible for the 3500 K star, while CO and OH bands can be noticed for the K star.

the temperature contrast between the starspot and the rest of the stellar photosphere increases. Molecular water bands can indeed be observed in stars with spots cooler than ~ 3000 K (e.g. Wallace et al. 1995; Jones et al. 2002), so that they can be used as an indicator of the temperature of active features. For the K star, CO absorption at 2.3 and 4.5 μm , and OH absorption at 2.5 μm , also increase with increasing temperature contrast. While the TiO spectral lines at 567, 705.5, and 886 nm are also known as robust temperature indicators (e.g. Wing, Spinrad & Kuhl 1967; Vogt 1979, 1981; Ramsey & Nations 1980; Neff, O’Neal & Saar 1995; Mirtorabi, Wasatonic & Guinan 2003; O’Neal et al. 2004; Bidaran, Mirtorabi & Azizi 2016), a spectral resolution larger than $\sim 10\,000$ would be needed to achieve significant constraints. Moreover, such lines were only observed in very active stars, where starspots occupy a significant fraction of the visible stellar disc, but which are usually excluded from exoplanet searches.

Starspots that are occulted during transit can be better characterized than those which are not. For example, all latitude values for non-occulted starspots are consistent with observations, other than those spanned by the transit chord; moreover, out-of-transit monitoring with a comparable duration to the stellar rotation period are needed to constrain starspot longitudes. When a starspot is occulted by a planet, its latitude and μ position on the stellar disc can instead be determined from transit information alone.

During the occultation of a starspot, the relative flux received from the system increases, creating a ‘bump’ in the transit profile. The peak of this bump, Δf , strongly depends on T_{\bullet} . Sing et al. (2011) related Δf in the wavelength bin centred at λ and at position μ to the bump at a reference wavelength λ_0 at the same μ (their equation 3):

$$\frac{\Delta f_{\lambda}}{\Delta f_{\lambda_0}} = \frac{1 - I_{T_{\bullet}}(\lambda, \mu)/I_{T_{\star}}(\lambda, \mu)}{1 - I_{T_{\bullet}}(\lambda_0, \mu)/I_{T_{\star}}(\lambda_0, \mu)}. \quad (2)$$

In their formulation, the choice of reference wavelength becomes important, and information is lost in the regions of the contrast spectrum that are around λ_0 .

To avoid any assumption on the reference wavelength, we decided to work with absolute quantities. To derive the flux change that is observed during a starspot occultation, we started by computing the flux observed during a transit if the starspot were at the same μ

position but not occulted, e.g. if it were at an opposite stellar latitude with respect to the centre of the transited stellar belt. In this case, we modelled the nominal transit configuration (e.g. Mandel & Agol 2002), and made the μ -dependent stellar specific intensity I_{\star} explicit, in place of the integrated flux:

$$F_1 = F_{\star+\bullet} - \frac{dA_p \mu_{\bullet}}{R_{\star}^2} I_{\star}(\lambda, \mu_{\bullet}) = F_{\star+\bullet} - \pi \left(\frac{R_p}{R_{\star}} \right)^2 I_{\star}(\lambda, \mu_{\bullet}), \quad (3)$$

where $F_{\star+\bullet}$ denotes the flux emitted by the star when the starspot is visible on its disc before occultation, and $dA_p \mu_{\bullet}/R_{\star}^2$ represents the solid angle intercepted by the planetary disc on the stellar disc. We used the subscript μ_{\bullet} to indicate that the limb angle is evaluated at the centre of the starspot, and we dubbed the planetary surface dA_p to indicate that it is small compared to the stellar surface. This allowed us to assume that the portion of the stellar photosphere covered by the planet can be described by a single μ_{\bullet} value (Marino et al. 1999; Ballerini et al. 2012).

When the starspot is occulted, the stellar specific intensity needs to be replaced by the starspot specific intensity,

$$F_2 = F_{\star+\bullet} - \pi \left(\frac{R_p}{R_{\star}} \right)^2 [\beta I_{\bullet}(\lambda, \mu_{\bullet}) + (1 - \beta) I_{\star}(\lambda, \mu_{\bullet})], \quad (4)$$

where β represents the fraction of the planetary disc that occults the starspot, and $1 - \beta$ the part that covers the unspotted stellar disc. This formalism allows the description of planets whose projection on the stellar disc is larger than the occulted starspot, as in this case $1 - \beta > 0$.

The difference between these two quantities is

$$F_2 - F_1 = \pi \beta \left(\frac{R_p}{R_{\star}} \right)^2 [I_{\star}(\lambda, \mu_{\bullet}) - I_{\bullet}(\lambda, \mu_{\bullet})]. \quad (5)$$

To compute the occultation flux bump in normalised quantities, we then divided $F_2 - F_1$ by the spotted stellar flux $F_{\star+\bullet}$. To do this, we noticed that in an observational context one works with the measured planet-to-stellar surface ratio, which is affected by the presence of starspots. True and observed transit depth can be related

by a wavelength-dependent factor that multiplies the stellar flux,

$$\left(\frac{R_p}{R_\star}\right)_{\text{true}}^2 \equiv \left(\frac{\Delta F}{F_\star}\right) = \alpha(\lambda) \left(\frac{\Delta F}{F_{\star+\bullet}}\right) \equiv \alpha(\lambda) \left(\frac{R_p}{R_\star}\right)_{\text{obs}}^2, \quad (6)$$

where we used the subscripts ‘obs’ for the measured transit depth, F_\star is the unspotted stellar flux, ΔF indicates the stellar flux drop during transit, and $\alpha < 1$ for a dark starspot (e.g. Czesla et al. 2009). Given this definition,

$$\alpha(\lambda) = F_{\star+\bullet}/F_\star. \quad (7)$$

By plugging equation (7) in equation (5), and dividing by $F_{\star+\bullet}$, the two $F_{\star+\bullet}$ terms conveniently cancel out,

$$\Delta f(\lambda) \equiv \frac{F_2 - F_1}{F_{\star+\bullet}} = \pi\beta \left(\frac{R_p}{R_\star}\right)_{\text{obs}}^2 \frac{I_\star(\lambda, \mu_\bullet) - I_\bullet(\lambda, \mu_\bullet)}{F_\star}. \quad (8)$$

This derivation is therefore independent of the presence of additional non-occulted starspots on the stellar disc, as they can be absorbed by the $F_{\star+\bullet}$ parameter, and then removed by the $\alpha(\lambda)$ factor. The last term to be made explicit is the unspotted stellar flux, which was calculated as (e.g. Gray 1976)

$$F_\star = 2\pi \int_0^1 I_\star(\lambda, \mu) \mu d\mu. \quad (9)$$

We again remark that this derivation depends on the assumption that the starspot can be represented by a single $I_\bullet(\lambda, \mu_\bullet)$. In reality, large starspots or starspot groups might be better represented by a combination of the μ values they span. In this regard, the assumptions of our formalism are the weakest when $\mu = 1$: in this case the non-occulted flux F_1 cannot be calculated with equation (3). However, this is not a problem if the resolution of the μ -grid allows both the projected planet and the starspot to lie in the stellar annulus at $\mu = 1$ without overlapping. This holds true in the hypothesis of small planets and starspots compared to the stellar disc.

The quantity $\Delta f(\lambda)$ can be fitted to the relative measured flux bumps in the transit profile, in order to derive T_\bullet (assuming T_\star is known to a sufficient level of precision) and β . The T_\bullet and β parameters will be therefore prone to the degeneracy between the starspot temperature and size. Moreover, the fraction of the spot’s surface which is actually occulted by the planet cannot be constrained from the occultation event alone, but only a lower limit on its size can be placed by the duration of the occultation event, as we will discuss in Sections 3.3 and 4.4.

For our simulations, we chose two *JWST* modes that allow for exoplanet transmission spectroscopy observations over the broadest range of NIR wavelengths. The NIRSpec/Prism mode, in the range 0.6 – 5.3 μm , will offer constraints on starspot and facula contrast spectra for dim targets; also, it will often provide the most of information for planet atmospheres (Batalha & Line 2017). This mode would reach saturation on targets brighter than $\text{mag}_K \simeq 10.5$: hence, we also performed simulations where the NIRCams Dispersed Hartmann Sensor (DHS) would enable simultaneous 1.0 – 2.0 μm and 2.4 – 4.0 μm observations on bright targets by combining the F150W2 and the F322W2 filters (Schlawin et al. 2016), if this mode becomes operational.

3 SIMULATIONS

The goal of our simulations was to predict the constraints that *JWST* NIRSpec/Prism and NIRCams/F150W2 + F322W2 will be able to place on the effective temperature of occulted starspots for stars with different magnitudes, and spanning a wide range of activity levels. For each observing mode, we proceeded as follows:

(i) We considered two systems, a K and an M star, each hosting a planet with a different radius. We simulated wavelength-dependent uncertainties on the transmission spectra for a number of different stellar magnitudes, and assumed the transmission spectra to be featureless (to represent a thin or cloudy planetary atmosphere).

(ii) Given each transmission spectrum, we modelled the spectrophotometric transits for every wavelength bin. We added a starspot occultation to each transit, using the same spot configuration for every point in the transmission spectrum. We repeated the simulation for a range of starspot temperatures, representing different stellar activity levels, as well as different positions on the stellar disc.

(iii) For each one of these scenarios (identified by a stellar type, stellar magnitude, starspot temperature and location), we fitted all associated spectrophotometric transits in order to obtain $\Delta f(\lambda)$ (equation (8)). Such measures were then used to sample the contrast spectra of the occulted starspot, and were fitted to stellar specific intensity spectra to determine the distance from the input starspot temperatures T_\bullet .

3.1 Transit modelling

For NIRSpec/Prism, we used the PANDEXO software (Batalha et al. 2017) to simulate the transmission spectra uncertainties, using BT-Settl stellar spectral energy distributions. PANDEXO includes the contribution from shot, background, and read noise, with a dependence on the ratio between the duration of out- and in-transit observations, that we set equal to 1. We used the optimization function in the software that selects the maximum possible number of groups¹ before saturation (fixed to 80 per cent of the full well), and adopted the NRSRAPID readout pattern. The simulations were run for the SUB512 subarray, and a noise floor of 20 parts per million was adopted. For NIRCams, we used a custom SNR estimator (Greene et al. 2016; Schlawin et al. 2016) and PHOENIX spectra. We assumed the maximum number of groups that can be achieved to be 0.5 magnitudes below the saturation limit for the F322W2 grism time series and the RAPID read mode. For the simultaneous short wavelength DHS observations, we adopted the same exposure parameters as the F322W2 grism because the software requires simultaneous readouts. The size of the subarray determines the number of DHS spectra that may be captured as well as the minimum magnitude that will saturate the detector. The balance between capturing DHS spectra while not saturating subarrays results in 1 DHS for $K \leq 6.0$, 2 DHS for $K > 6$ and ≤ 8.3 , and 10 DHS for $K > 8.3$. The resulting uncertainties as a function of stellar magnitude and wavelength are shown in Fig. 2.

We tested different choices for the spectral resolution, and reduced it until the SNR on the measured ‘flux bump’ (increasing with decreasing spectral resolution) allowed us to achieve a few hundred kelvins precision on the retrieved starspot temperatures in the brightest star cases (as described later in Section 4). In this way, we chose $R \simeq 10$ for our spectra, both for NIRSpec and NIRCams. We modelled the transmission spectrum uncertainties of a 0.75 R_J planet in front of a 5000 K, 0.75 R_\odot star and one of a 0.25 R_J planet in front of a 3500 K, 0.47 R_\odot star. The details of the two scenarios are summarized in Table 1.

For each point of the transmission spectrum, the simulated transit depth uncertainty was used to derive the relative flux uncertainties

¹This is the number of non-destructive frames that are averaged onboard before being recorded; see <https://jwst-docs.stsci.edu/understanding-exposure-times> for more detail.




Figure 2. Transmission spectra uncertainties computed for the NIRSpec/Prism (*top*) and NIRCcam/F150W2 + F322W2 (*bottom*) configurations. Increasingly larger uncertainties correspond, for NIRSpec, to mag $K = 10.5, 11.5, 12.5, 13.5,$ and 14.5 ; for NIRCcam, they correspond to $K = 4.5, 6.0, 7.5,$ and 9.0 . M and K star scenarios are shown on the left and right column, respectively.

Table 1. Simulated scenarios. From left to right, the columns indicate modelled instrument and observing mode, stellar K mag, stellar effective temperature, stellar surface gravity, starspot simulated effective temperature range and step, planet radius, stellar radius, starspot limb angle θ_* , starspot latitude on the stellar disc ξ , and planet orbital inclination i .

Instrument/mode	K mag range	T_* [K]	$\log g$ [cgs]	T_\bullet range (step) [K]	R_p [R_J]	R_* [R_\odot]	θ_* [$^\circ$]	ξ [$^\circ$]	i [$^\circ$]
NIRSpec/Prism	10.5-14.5	5000	4.5	3800-4700 (300)	0.75	0.75	0, 21, 40	0, 21	90
NIRSpec/Prism	10.5-14.5	3500	5.0	2600-3200 (300)	0.25	0.47	0, 40	0	90
NIRCcam/F150W2 + F322W2	4.5-9.0	5000	4.5	3800-4700 (300)	0.75	0.75	0, 21, 40	0, 21	90, 88.1
NIRCcam/F150W2 + F322W2	4.5-9.0	3500	5.0	2600-3200 (300)	0.25	0.47	0, 40	0	90, 88.8

in the corresponding transit. They were calculated as (e.g. Sarkar, Madhusudhan & Papageorgiou 2020)

$$\sigma_s(\lambda) \simeq \frac{\sqrt{N_{\text{int}}}}{2} \sigma_D(\lambda), \quad (10)$$

where N_{int} is the number of exposures in the transit and $\sigma_D(\lambda)$ is the uncertainty on the transit depth at the same wavelength, as shown in Fig. 2. We set an exposure time of 60 s for the simulated light curves, which given the transit parameters (described later in this Section) resulted in $N_{\text{int}} = 432$ and 188 for the K and the M star simulation, respectively. We assumed our mock observations to be already corrected for systematic and astrophysical noise, and did not include the contribution from stellar granulation.

White noise in the wavelength-dependent transits results in the placement of each data point on a transit model to which a Gaussian

scatter calculated with equation (10) is added. By iterating our simulations, we noticed that different noise realizations have an impact on the final measure of T_\bullet . The implications of being limited, in an observational context, to a specific noise instance can be appreciated by running a large number ($\gtrsim 10$) of simulations, each one with a different noise realization, and by merging the posterior distributions on the fitted parameters. Following Feng et al. (2018), we decided to estimate this effect by avoiding the inclusion of Gaussian scatter in the transit data points, and by later evaluating the impact of this choice on a few significant scenarios (Section 4.5).

We then used KSINT (Montalto et al. 2014) to simulate the transits affected by the occultation of an active feature, including the transit depth contamination due to the out-of-transit spot-induced stellar flux variation. This software takes as input the stellar density, which

we calculated from the stellar radius and respective $\log g$ for each scenario (as presented in Table 1). Given the focus on simulations of a single transit event, the planetary orbital period and stellar rotation period did not significantly impact our analysis: we then fixed the orbital period to 2 d and the stellar rotation period to 11 d (this latter choice represents an acceptable value for moderately active stars of the type we explored, e.g. McQuillan, Mazeh & Aigrain 2014). For all targets, we set no spin-orbit misalignment.

There exists a multitude of possible configurations that would produce a measurable starspot crossing, but here we have selected reasonable physically motivated scenarios that can highlight the potential of *JWST* to provide further insights into starspot properties from planetary transits. To start with, we used a 3° -wide spot for our simulations, a value that sits between the one of large spot groups on the Sun (Mandal et al. 2017) and the largest spots observed for M dwarfs (Berdyugina 2011), and which is also consistent with starspots observed on active K stars (e.g. Morris et al. 2017). Secondly, we chose several positions for the starspot: one close to the centre of the stellar disc ($\theta = 0^\circ$, $\mu = 1$) and one closer to the limb ($\theta = 40^\circ$, $\mu = 0.77$) at 0° stellar latitude, associated to an $i = 90^\circ$ planet orbital inclination. We also simulated spots at 21° stellar latitude, corresponding to $\theta = 21^\circ$ ($\mu = 0.93$), occulted by a planet at $i \simeq 88^\circ$. In this case, we introduced a slight displacement between the starspot and the projection of the planet centre on the stellar disc. In the hypothesis of circular starspots, the planet and starspot centres are aligned if

$$i = \arccos\left(\frac{R_\star}{a} \sin \xi\right), \quad (11)$$

where a is the orbital semimajor axis and ξ is the starspot latitude.

We used no starspot other than the occulted one, and the other transit parameters as in Table 1. The orbits were assumed to be circular.

KSINT uses a quadratic LD law: we calculated the corresponding coefficients by fitting PHOENIX specific intensity spectra. For the NIRCcam modes, we downloaded the throughput curves from the SVO Filter Profile Service² (Rieke, Kelly & Horner 2005; Rodrigo, Solano & Bayo 2012; Rodrigo & Solano 2020); those for NIRSspec (CLEAR filter) are available with the PANDEIA reference data (Pontoppidan et al. 2016).

For each wavelength bin, we computed the starspot contrast as a function of wavelength as the ratio of two PHOENIX specific intensity models at different temperatures (T_\bullet versus T_\star) and using a starspot $\log g$ as low as the stellar $\log g - 0.5$ dex.

3.2 Transit fits

We treated the synthetic data the same standard way that transits are typically analysed. We chose a transit modelling approach that allows for efficient iterative-based analyses that is insensitive to the degeneracies between starspot size, location on stellar disc, and inclination of planetary orbit. In place of using KSINT for the transit fit we adopted the BATMAN code, which implements the standard Mandel & Agol (2002)’s transit model (Kreidberg 2015). We then expanded Fraine et al. (2014)’s approach by adding the starspot occultation as a flattened Gaussian function profile to the transit. As

Table 2. Boundaries on the fitted parameters. $\mathcal{U}[a, b]$ and $\mathcal{J}[a, b]$ represent a Uniform and a Jeffreys prior between a and b (inclusive), respectively. $\mathcal{G}(a, b)$ denotes a Gaussian prior with mean a and standard deviation b . Two different occultation mid-times were adopted according to the limb-angle θ of each scenario.

Parameter	Prior
R_p/R_\star	$\mathcal{J}[0.01, 0.2]$
Orbit inclination i [deg]	$\mathcal{U}[80, 90]$
Transit mid-time t_{tr} [days]	$\mathcal{U}[0.08, 0.12]$
LD linear coefficient u_1	$\mathcal{G}(\text{theoretical value}, 0.05)$
LD quadratic coefficient u_2	$\mathcal{G}(\text{theoretical value}, 0.05)$
Gaussian peak Δf	$\mathcal{U}[10^{-6}, 0.1]$
Gaussian flattening factor n	$\mathcal{U}[2, 8]$
Gaussian width w [days]	$\mathcal{U}[7 \times 10^{-4}, 0.02]$
Occultation mid-time t_\bullet [days] ($\theta = 0^\circ$)	$\mathcal{U}[0.09, 0.11]$
Occultation mid-time t_\bullet [days] ($\theta = 21^\circ$ or 40°)	$\mathcal{U}[0.11, 0.13]$
Out-of-transit quadratic term r_0	$\mathcal{U}[-1, 1]$
Out-of-transit linear term r_1	$\mathcal{U}[-1, 1]$
Out-of-transit level r_2	$\mathcal{U}[0, 10]$

a function of time t , this is equal to

$$\Delta f \exp\left[\left(-\frac{|t - t_\bullet|}{w}\right)^n\right], \quad (12)$$

where Δf is the peak of the flux bump during occultation (the main parameter we were interested in), t_\bullet is the occultation mid-time, w is the width of the bump, and n determines the flattening of the bump for starspots that are smaller than the projection of the planet on the stellar disc.

We fitted for the planet-to-star radius ratio $R_p(\lambda)/R_\star$, LD coefficients, starspot bump peak Δf , and for a quadratic trend with time, $r_0 t^2 + r_1 t + r_2$, to account for the out-of-transit stellar flux modulation due to the presence of the starspot. As the orbital inclination i , the transit mid-time t_{tr} , and the spot crossing mid-time t_\bullet are common among all transits of the same transmission spectrum, we fitted these parameters on a band-integrated version (commonly referred to as ‘white light curve’) of the transits for each scenario. We also assumed the bump width w and flatness n of the occultation bump to be independent of wavelength, and fitted for them only in the co-added transit. For the fit of the spectrophotometric transits in each scenario, we then fixed the value of these parameters to the median value obtained on the corresponding band-integrated transits.

Given the complexity of the transit and starspot parameter space, we sampled the parameter posterior distributions with nested sampling, which is more efficient than Markov chain Monte Carlo algorithms in dealing with multimodal posterior distributions (Skilling 2006; Higson et al. 2019). We adopted the ‘standard nested sampling’ implementation in the DYNESTY software (Speagle 2020), using Uniform priors for all transit parameters except R_p/R_\star , for which we used a Jeffreys prior, and the LD coefficients, for which we used Gaussian priors centred on the theoretical values from stellar models. We imposed boundaries on the fitted parameters as indicated in Table 2; in particular, we used two different priors on t_\bullet for the $\theta = 0^\circ$ and the $\theta = 40$ and 21° scenario, to ease convergence. We started the sampling of the band-integrated transits with 200 live points, and reduced this number to 150 for the spectrophotometric transits.

The choice of fitting for the LD coefficients was due to the full transit coverage of future *JWST* observations. Indeed, inaccurate and incomplete opacities in stellar atmosphere models, as well as small variations in the stellar atmospheric parameters due to brightness inhomogeneities on the stellar photosphere, make the LD

²Described at <https://jwst-docs.stsci.edu/near-infrared-camera/nircam-instrumentation/nircam-filters>.




Figure 3. Simulated transit of a $0.25 R_J$ planet orbiting a 10.5-mag_K , $0.47 R_\odot$, 3500 K star, occulting a 2900 K starspot at limb angle $\theta = 40^\circ$, observed with NIRSPEC/Prism in the wavelength bin centred at $\simeq 1.4\ \mu\text{m}$. This plot presents one of the noise realizations to better show the extent of the flux uncertainties. In orange on the upper panel, 300 samples from the posterior distributions obtained with the transit and Gaussian profile modelling are shown. The lower panel shows the residuals of the best-fit model, presenting a white-noise scatter, in parts per million.

coefficients derived from stellar models unreliable for active stars (e.g. Csizmadia et al. 2013; Maxted 2018; Heller 2019). Despite being small variations, such transit profile differences can have a measurable effect on the observed transit depths, and should be taken into account whenever possible. This said, we found that imposing a prior centred on the theoretical values of the LD coefficients provides better results than leaving wide Uniform priors. In our fits, we used the quadratic law in order to improve convergence with respect to the use of the four-coefficients law (Morello et al. 2017).

Fig. 3 presents an example of fit on a simulated transit using the model just described; we there present one of the white noise realizations (Section 4.5) to show an example of simulated transit uncertainties. The simultaneous transit and starspot fit was repeated for all wavelength bins and simulated stellar magnitudes, and the flux bump Δf due to the starspot recorded.

3.3 Starspot contrast spectra fits

For each scenario, specific intensity spectra were used to fit the occultation bump peaks using equation (8). We obtained F_\star using equation (9), relied on the measured $(R_p(\lambda)/R_\star)^2$ and used the specific intensity models $I_\star(\mu_\star)$ and $I_\bullet(\mu_\star)$ at the same μ_\star chosen in input, assuming that μ_\star can be accurately determined from the observed t_\star .

For the stars, we selected a spectrum with the same parameters used in input. For the starspots, we interpolated the stellar models in the range $2300\text{--}3400\text{ K}$ for the M star ($T_\star = 3500\text{ K}$), and in the $3600\text{--}4900\text{ K}$ range for the K star ($T_\star = 5000\text{ K}$), using spectra with the same $[\text{Fe}/\text{H}]$ as the input stellar spectrum and the stellar $\log g - 0.5$ (see Section 2). To do this, we performed a bivariate spline approximation with SCIPY’s RECTBIVARIATESPLINE implementation (Virtanen et al. 2020).

It is well known that the starspot effective temperature and filling factor (here probed by β) are strongly correlated parameters, as a cool and small starspot can cause the same in-transit bump peak as a large and warm spot. In particular, in some of the low-contrast cases we observed bimodal posterior distributions for these two parameters




Figure 4. Example of in-transit starspot crossing flux bumps measured with the model in equation (8) and their uncertainties, for the $\text{mag } K = 7.5$, K star case, in the $T_\star = 3800\text{ K}$, $\mu\text{-angle } \theta = 40^\circ$ scenario without in-transit Gaussian scatter. The models in light blue represent 100 fitting models sampled from the T_\star and β posterior distributions. The result of the nested sampling for fitted parameters, and their 68 percent credible intervals, are indicated in the label. The model spectra were convolved with the NIRCcam filters and degraded to the same resolutions as the observations before performing the fit.

(see Section 4.4). However, observations provide us with a constraint on the lower limit of β , and sometimes allow the choice of one of the modes. To find this value β_{\min} , we first derived the half width at half-maximum (HWHM_\bullet , in time units) of the Gaussian function that we used to describe the occultation bumps in the white light transits. Given equation (12), this was obtained as

$$\text{HWHM}_\bullet = w(\ln 2)^{1/n}. \quad (13)$$

Then, assuming a circular starspot and zero-eccentricity orbit, we determined the minimum starspot radius $R_{\bullet, \min}$ consistent with the observed occultation, by inverting

$$R_p + R_{\bullet, \min} \simeq \frac{3}{2} \text{HWHM}_\bullet v_p = \frac{3}{2} \text{HWHM}_\bullet \frac{2\pi a}{P}, \quad (14)$$

where v_p is the tangential velocity of the planet in its orbit, P is the planet’s orbital period, and we approximated the last moment when the profiles of planet and starspot are in contact with $t_\star + 3 \text{HWHM}_\bullet/2$. By dividing both terms in equation (14) by the stellar radius and the measured planet-to-star radius ratio and taking their square, after some algebra we obtained an expression which is only based on measured quantities:

$$\beta_{\min} \equiv \left(\frac{R_{\bullet, \min}}{R_p} \right)^2 = \left[\frac{3\pi a/R_\star}{P(R_p/R_\star)} \text{HWHM}_\bullet - 1 \right]^2, \quad (15)$$

where a is the orbital semimajor axis. This value is underestimated, as the measured transit depth is affected by the presence of starspots, but we consider it to be a sufficient approximation.

We sampled the T_\star and β posterior distributions with nested sampling, using Uniform priors for both parameters and starting the sampling with 100 live points. The starspot T_\star was let vary across the full interpolated range (different for the M and the K star case), and β was let float between β_{\min} and 1 (inclusive). Fig. 4 presents 100 fitting models from the posterior distributions for one of the K star scenarios without Gaussian scatter, simulated for NIRCcam and downgraded to the resolution of the observations.




Figure 5. Difference between the retrieved starspot temperature T_* (meas) and the true input T_* (true) (y-axis), for the scenarios simulated with NIRSpec/Prism using starspots at 0° latitude; 95 per cent percentiles are reported. The stellar K magnitudes corresponding to the different cases and the true T_* values are indicated with different markers and colours, respectively. The minimum allowed T_* in our retrievals, $T_* - 100$ K, is indicated by a horizontal line with the same colour as the data points for a given scenario (the value for the $T_* = 3800$ K falls off the plot edges). In the title of each plot, the simulated instrument, T_* , and the limb angle θ are reported. Markers with arrows pointing down – such as the lowest-SNR point in the top left-hand panel – represent cases where the occultation flatness parameter n is poorly determined ($\sigma_n \geq 2$), so that the error bars on T_* cannot be considered reliable (see Section 4.3).

4 RESULTS

We present our results as a function of the SNR of the starspot spectrum derived from the transit fits. To do so, we defined a metric that contains information on the starspot contrast decrease from the optical to the NIR, on the contrast ‘baseline’ in the white-light transit, and on the uncertainty in the $\Delta f(\lambda)$ values measured from the spectrophotometric transits. If the starspot contrast spectrum results from N spectrophotometric transit light curves, corresponding to increasing wavelength in bins from 1 to N , the white-light transit is marked with (N) , and Δf_i is the bump peak measured in wavelength bin i , we computed

$$\text{SNR} = \frac{\sqrt{(\Delta f_2 - \Delta f_{N-1})^2 + \Delta f_{(N)}^2}}{\Delta(\Delta f_{N-1})}, \quad (16)$$

where the term on the denominator is the uncertainty on the bump peak in the $(N - 1)$ th wavelength bin. We avoided the use of the shortest and the largest-wavelength transit, as these are the noisiest in the simulated transmission spectra (Fig. 2).

As the SNR of the bump signal increases for larger activity features, this representation adapts to scenarios with varying starspot sizes, even if we here discuss only the simulations we carried out with 3° -wide spots; in particular, the effect of a large spot on the occultation SNR is similar to the one of a bright star.

4.1 Starspots at 0° latitude, planet with $i = 90^\circ$

The overall result of this analysis is that, despite their reduced contrast in the NIR compared to the visible, the effective temperature of starspots observed with JWST can be constrained with an accuracy of a few hundred kelvins in the brightest star and largest contrast cases, and particularly with NIRCам. Figs 5 and 6 present the retrieved values for the temperature contrasts and their 95 per cent credible intervals, as a function of the occultation (equation (16)). The stellar photospheric values are also represented, and indicate when the derived T_* cannot be significantly distinguished from T_* .

At 0° latitude, the limb-angle position of the starspot ($\theta = 0$ or 40°) slightly affects the scatter of the solution with respect to the correct results. For NIRCам/F150W2 + F322W2, simulations produced accurate results for the coolest starspot cases, even for dim ($K = 9$) magnitudes; with NIRSpec/Prism, a few hundred kelvin precision was achieved for the largest temperature contrasts ($T_* - T_* = 900$ and 1200 K for the M and the K star, respectively) for $K \leq 12.5$ magnitudes. For both the M and the K star, the constraints on the temperature contrast become weaker as the true T_* approaches the photospheric temperature, even if the occultation bump is detected in the transits. For the lowest SNR cases, our simulations were only able to place a lower limit to the spot temperature; in the mid-SNR cases, the starspot temperature-size degeneracy caused the T_* posterior distribution to include both very close values to T_* , as well as several hundred kelvin lower values (see Section 4.4). We notice that $T_* - T_* = 1200$ K contrasts are realistic for K dwarfs, while




Figure 6. Same as Fig. 5, but for NIRCcam/F150W2 + F322W2.

larger contrasts than 600 K are unlikely on M stars (Berdyugina 2005; Herbst et al. 2021).

4.2 Starspots at latitude $>0^\circ$, planet with $i \neq 90^\circ$

As described in Section 3.1, we investigated the impact of a different geometrical configuration, including a slight displacement between the starspot and the planet projected centres. For the NIRCcam simulations, which include the largest SNR cases, we simulated a starspot at 21° stellar latitude (requiring also $\theta = 21^\circ$), and added a $\lesssim 1^\circ$ angle to the planet orbital inclination i with respect to the value which results in perfect alignment. Fig. 7 shows that, as expected, the lower SNR of the occultation results in larger scatter and error bars for the retrieved T_s . We found best results for the largest contrast and the brightest ($K = 4.5$ and 6.0) cases, particularly for the K star.

4.3 Low-SNR solutions

We observed that solutions with $\text{SNR} \lesssim 3$ are driven by the choice of priors and cannot be considered reliable. As an example, the solution for the $T_s = 3200$ K, $\theta = 0^\circ$ spot on the mag $K = 14.5$ M star observed with NIRSpec/Prism (Fig. 5, top left-hand panel) has smaller error bars compared to the simulations with the same stellar and spot parameters but brighter stars. This happens because the bump flatness parameter n (equation (12)) is not constrained by the white-light transit fit. The median value of its posterior distribution, used to derive β_{\min} through equations (13) and (15), is then found close to values around the middle value of the prior given to n (set between 2 and 8, as shown in Table 2; Fig. A1 illustrates the posterior distributions for this transit fit). As a result, β_{\min} is found to be close to 1 (i.e. the spot must be almost or completely occulted by the planet) and good fits to the spot contrast spectrum are only those produced by a ‘warm’ activity feature, as shown in the left-hand panel of Fig. 8.

Hence, even if the occultation signal is strong enough to exclude a T_s as high as T_s^* (otherwise, only a lower limit could be attributed to T_s^*), the solution is ‘forced’ around warm spots.

To confirm this, such a solution cannot be reproduced if the constraint on β_{\min} is lifted, i.e. if a $(0, 1]$ prior is set for β . We therefore consider this solution not reliable, and recommend the inspection of the occultation bump fit in white-light transits to correctly interpret the derived T_s values. Cases with a poor fit for the occultation flatness parameter n (defined as where its error bar σ_n is ≥ 2) are marked using arrows pointing down in Figs 5–7, to indicate that cooler spot solutions cannot be excluded.

4.4 Starspot temperature-size degeneracy

For warmer starspots and dimmer stars, the SNR of the occultation feature decreases, and the fitting procedure struggles in distinguishing between the starspot temperature and the stellar photosphere’s. As the lowest allowed T_s in our model was $T_s - 100$ K, we required this latter value to be excluded at the 95 per cent confidence level for the temperature contrast to be detected. The low-SNR cases without such detection were clearly affected by the starspot contrast and size degeneracy, contrarily to the large SNR cases. In some of these cases, the T_s posterior distribution spans a large part of the prior space and does not lead to significant constraints on this parameter. We observed that, in some specific scenarios, the T_s posterior distribution is bimodal, with the correct value lying in one of the modes. One example is the case of the $T_s = 4100$ K starspot on the K star: here, the correct mode cannot be chosen by using the method presented in Section 3.3, as the β values of both modes are enclosed in the prior. Fig. 8, right-hand panel, shows this situation for the mag $K = 7.5$ star observed with NIRCcam/F150W2 + F322W2; in a case like this, the 50 per cent percentile shown in Figs 5–7 is little representative of the shape of the posterior. In Figures B1–B3, we provide the posterior