



<b>Publication Year</b>	2016
<b>Acceptance in OA</b>	2020-05-22T12:49:02Z
<b>Title</b>	Spectral characterization of volcanic rocks in the VIS-NIR for martian exploration
<b>Authors</b>	DE ANGELIS, Simone, CARLI, CRISTIAN, Manzari, Paola, DE SANCTIS, MARIA CRISTINA, CAPACCIONI, FABRIZIO
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/25092">http://hdl.handle.net/20.500.12386/25092</a>

**CONTROL ID:** 2568318

**TITLE:** Spectral characterization of volcanic rocks in the VIS-NIR for martian exploration

**ABSTRACT BODY:**

**Abstract (2,250 Maximum Characters):** Igneous effusive rocks cover much of the surface of Mars [1,2,3]. Initially only two types of lithologies were thought to constitute the Martian crust, i.e. a basaltic one and a more andesitic one [1,2], while more evolved lithologies were ruled out.

Nevertheless a more complex situation is appearing in the last years. Recently several observations have highlighted the presence of evolved, acidic rocks. High-silica dacite units were identified in Syrtis Major caldera by thermal IR data [4]. Outcrops in Noachis Terra were interpreted as constituted of felsic (i.e. feldspar-rich) rocks essentially by the observation of a 1.3- $\mu\text{m}$  spectral feature in CRISM data, attributed to  $\text{Fe}^{2+}$  in feldspars [5]. However different interpretations exist, invoking plagioclase-enriched basalts [6] rather than felsic products.

The increasing of high-resolution and in-situ rover-based observations datasets and the changing of the initial paradigm justify a new systematic spectral study of igneous effusive rocks. In this work we focus on the spectral characterization of volcanic effusive rocks in the 0.35-2.5- $\mu\text{m}$  range. We are carrying out measurements and spectral analyses on a wide ensemble of effusive samples, from mafic to sialic, with variable alkali contents, following the classification in the Total-Alkali-Silica diagram, and discussing the influence on spectral characteristics of different mineral assemblages and/or texture ([7], [8]).

[1] Bandfield J.L., et al., *Science*, 287, 1626, 2000; [2] Christensen P.R., et al., *J. Geophys. Res.*, 105, N.E4, 9609-9621, 2000; [3] Ehlmann B.L. & Edwards C.S., *Annu. Rev. Earth Planet. Sci.*, 42, 291-315, 2014; [4] Christensen P.R., et al., *Nature*, 436, 504-509, 2005; [5] Wray J.J., et al., 44<sup>th</sup> LPSC, abs. n.3065, 2013; [6] Rogers A.D. & Nekvasil H., *Geophys. Res. Lett.*, 42, 2619-2626, 2015; [7] Carli C. and Sgavetti M., *Icarus*, 211, 1034–1048, 2011; [8] Carli C. et al., *SGL*, doi 10.1144/SP401.19, 2015.

**CURRENT CATEGORY:** Laboratory Research

**CURRENT :** None

**AUTHORS (FIRST NAME, LAST NAME):** Simone De Angelis<sup>1</sup>, Cristian Carli<sup>1</sup>, Paola Manzari<sup>1</sup>, Maria Cristina De Sanctis<sup>1</sup>, Fabrizio Capaccioni<sup>1</sup>

**INSTITUTIONS (ALL):**

1. IAPS, INAF, Rome, Italy.

**Contributing Teams:**