



Publication Year	2016
Acceptance in OA	2020-11-20T16:41:40Z
Title	On the radial distribution of Galactic cosmic rays
Authors	Recchia, S., Blasi, P., MORLINO, GIOVANNI
Publisher's version (DOI)	10.1093/mnrasl/slw136
Handle	http://hdl.handle.net/20.500.12386/28493
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
Volume	462

On the radial distribution of Galactic cosmic rays

S. Recchia,¹★ P. Blasi^{2,1} and G. Morlino^{1,2}

¹INFN – Gran Sasso Science Institute, viale F. Crispi 7, I-67100 L'Aquila, Italy.

²INAF – Osservatorio Astrofisico di Arcetri, L.go E. Fermi, 5, I-50125 Firenze, Italy.

Accepted 2016 July 7. Received 2016 July 7; in original form 2016 April 26

ABSTRACT

The spectrum and morphology of the diffuse Galactic γ -ray emission carries valuable information on cosmic ray (CR) propagation. Recent results obtained by analyzing *Fermi*-LAT data accumulated over 7 yr of observation show a substantial variation of the CR spectrum as a function of the distance from the Galactic Centre. The spatial distribution of the CR density in the outer Galaxy appears to be weakly dependent upon the galactocentric distance, as found in previous studies as well, while the density in the central region of the Galaxy was found to exceed the value measured in the outer Galaxy. At the same time, *Fermi*-LAT data suggest a gradual spectral softening while moving outwards from the centre of the Galaxy to its outskirts. These findings represent a challenge for standard calculations of CR propagation based on assuming a uniform diffusion coefficient within the Galactic volume. Here, we present a model of non-linear CR propagation in which transport is due to particle scattering and advection off self-generated turbulence. We find that for a realistic distribution of CR sources following the spatial distribution of supernova remnants and the space dependence of the magnetic field on galactocentric distance, both the spatial profile of CR density and the spectral softening can easily be accounted for.

Key words: ISM: general – gamma-rays: diffuse background.

1 INTRODUCTION

Among the numerous open questions in cosmic ray (CR) physics there is an old one, known as ‘the radial gradient problem’, concerning the dependence of CR intensity on Galactocentric distance: the CR density as measured from γ -ray emission in the Galactic disc is much more weakly dependent upon galactocentric distance than the spatial distribution of the alleged CR sources, modelled following pulsar and supernova remnant (SNR) catalogues. This result was obtained for the first time from the analysis of the γ -ray emissivity in the Galactic disc derived from the SAS-2 data by Stecker & Jones (1977) and later from the analysis of the COS-B data by Bhat et al. (1986) and Bloemen et al. (1986), then confirmed by work based on EGRET data (Strong & Mattox 1996; Hunter et al. 1997). In more recent years, the existence of a ‘gradient problem’ in the external region of our Galaxy has also been confirmed by data collected by *Fermi*-LAT (Ackermann et al. 2011, 2012). The results of a more detailed analysis of *Fermi*-LAT data, including also the inner part of the Galaxy, was published in two independent papers (Acero et al. 2016; Yang, Aharonian & Evoli 2016) based on data accumulated over 7 yr. This study highlighted a more complex situation: while in the outer Galaxy the density gradient problem has been confirmed once more, the density of CR protons in the inner Galaxy turns out

to be appreciably higher than in the outer regions of the disc. Moreover, this analysis suggests a softening of the CR spectrum with Galactocentric distance, with a slope ranging from 2.6, at a distance of ~ 3 kpc, to 2.9 in the external regions. These are very interesting results, but it is worth keeping in mind that the spatial distribution of CRs in the Galaxy is not the result of a direct measurement, in that it can only be inferred from the observed γ -ray emission through a careful modelling of the spatial distribution of the gas that acts as target for nuclear collisions and by separating the γ -ray contribution due to hadrons from that due to leptonic interactions (bremsstrahlung and inverse Compton scattering).

The scenario that emerges from this analysis is difficult to reconcile with the standard approach to CR propagation, which is based upon solving the transport equation under the assumption that the diffusive properties are the same in the whole propagation volume (see, e.g. Berezhinskii et al. 1990). Within the context of this approach, several proposals have been put forward to explain the radial gradient problem. Among them: (a) assuming a larger halo size or (b) a flatter distribution of sources in the outer Galaxy (Ackermann et al. 2011); (c) accounting for advection effects due to the presence of a Galactic wind (Bloemen et al. 1993); (d) assuming a sharp rise of the CO-to-H₂ ratio in the external Galaxy (Strong et al. 2004); (e) speculating on a possible radial dependence of the injected spectrum (Erlykin, Wolfendale & Dogiel 2016). None of these ideas, taken individually, can simultaneously account for both the spatial gradient and the spectral behaviour of CR protons.

* E-mail: sarah.recchia@gssi.infn.it.

Moreover, many of them have issues in accounting for other observables (see, e.g. the discussion in Evoli et al. 2012).

A different class of solutions invoke the breakdown of the hypothesis of a spatially constant diffusion coefficient. For instance, Evoli et al. (2012) proposed a correlation between the diffusion coefficient parallel to the Galactic plane and the source density in order to account for both the CR density gradient and the small observed anisotropy of CR arrival directions. Gaggero et al. (2015a,b) followed the same lines of thought and showed that a phenomenological scenario where the transport properties (both diffusion and convection) are position-dependent can account for the observed gradient in the CR density. It is however unsatisfactory that these approaches do not provide a convincing physical motivation for the assumed space properties of the transport parameters.

In this paper, we explore the possibility that diffusion and advection in self-generated waves produced by CR-streaming could play a major role in determining the CR radial density and spectrum. The effects of self-generated diffusion has been shown to provide a viable explanation to the hardening of CR proton and helium spectra observed by PAMELA (Adriani et al. 2011) and AMS-02 (Aguilar et al. 2015), supporting the idea that below ~ 100 GeV, particle transport at the Sun's location may be dominated by self-generated turbulence (Blasi, Amato & Serpico 2012; Aloisio, Blasi & Serpico 2015). We suggest that this could be the case everywhere in the Galaxy and explore the implications of this scenario. In the assumption that the sources of Galactic CRs trace the spatial distribution of SNRs and that the magnetic field drops at large galactocentric distances, the density of CRs and their spectrum are well described if CRs are allowed to diffuse and advect in self-produced waves.

The paper is structured as follows. In Section 2, we show the solution of the CR transport in a 1D slab model with constant advection and with purely self-generated diffusion. In Section 3, we discuss the distribution of sources and the behaviour of the magnetic field in the Galactic plane and we compare our results for the CR proton spectrum with the data obtained by *Fermi*-LAT. Finally, we summarize in Section 4.

2 CR TRANSPORT IN SELF-GENERATED TURBULENCE

For simplicity, we assume that for any given galactocentric distance R , diffusion can be described as one dimensional, so that particles diffuse and are advected only along the z -direction. The transport equation for CR protons can then be written as follows:

$$-\frac{\partial}{\partial z} \left[D(z, p) \frac{\partial f}{\partial z} \right] + w \frac{\partial f}{\partial z} - \frac{p}{3} \frac{\partial w}{\partial z} \frac{\partial f}{\partial p} = Q_0(p) \delta(z), \quad (1)$$

where the advection velocity is

$$w(z) = \text{sign}(z) v_A, \quad (2)$$

directed away from the disc, and the Alfvén speed is $v_A = B_0 / \sqrt{4\pi m_p n_i}$. We assume that the ion density n_i is constant everywhere in the halo, which is assumed to extend out to $|z| = H$, while the magnetic field strength is constant along z but can be a function of the Galactocentric distance, R .

The injection of particles occurs only in the Galactic disc (at $z = 0$) and is a power law in momentum:

$$Q_0(p) = \frac{\xi_{\text{inj}} E_{\text{SN}} \mathcal{R}_{\text{SN}}(R)}{4\pi \Lambda c (m_p c)^4} \left(\frac{p}{m_p c} \right)^{-\gamma}. \quad (3)$$

Here, $E_{\text{SN}} = 10^{51}$ erg is the total kinetic energy released by a single supernova explosion, ξ_{inj} is the fraction of such energy channelled into CRs, $\mathcal{R}_{\text{SN}}(R)$ is the SN explosion rate per unit area and the normalization constant is $\Lambda = \int_{p_{\text{min}}}^{p_{\text{max}}} dy y^{2-\gamma} [(y^2 + 1)^{1/2} - 1]$.

A standard technique to solve the transport equation (1) is to integrate between 0^- and 0^+ and between 0^+ and z with the boundary conditions $f(0, p) = f_0(p)$ and $f(H, z) = 0$ (see, e.g. Blasi et al. 2012; Recchia, Blasi & Morlino 2016). One gets the following result for $f(z, p)$:

$$f(z, p) = f_0(p) \frac{1 - e^{-\xi(z, p)}}{1 - e^{-\xi(0, p)}}, \quad (4)$$

where $\xi(z, p) = \int_z^H v_A / D(z', p) dz'$ and the distribution function in the disc is

$$f_0(p) = \int_p^{p_{\text{max}}} \frac{dp'}{p'} \frac{3Q_0(p')}{2v_A} \exp \left[- \int_p^{p'} \left(\frac{d\hat{p}}{\hat{p}} \frac{3}{e^{\xi(0, \hat{p})} - 1} \right) \right]. \quad (5)$$

When magnetic perturbations are weak ($\delta B \ll B_0$), one can use quasi-linear theory to determine the diffusion coefficient, which is usually written as

$$D(z, p) = D_B \left[\frac{1}{\mathcal{F}(k)} \right]_{k=1/r_L}, \quad (6)$$

where $D_B = r_L v / 3$ is the Bohm diffusion coefficient, with r_L the Larmor radius and v the particle's speed. $\mathcal{F}(k)$ is the normalized energy density per unit logarithmic wavenumber k , calculated at the resonant wavenumber $k = 1/r_L$. The local value of \mathcal{F} is determined by the balance between the CR-driven growth of Alfvén waves and their damping. The growth rate due to the CR-streaming instability is (Skilling 1975)

$$\Gamma_{\text{cr}} = \frac{16\pi^2}{3} \frac{v_A}{\mathcal{F}(k) B_0^2} \left[p^4 v(p) \frac{\partial f}{\partial z} \right]_{p=eB_0/kc}, \quad (7)$$

while the dominant damping process in a region where the background gas is totally ionized is the non-linear Landau Damping (NLLD) that occurs at a rate (Ptuskin & Zirakashvili 2003):

$$\Gamma_{\text{nllD}} = (2c_k)^{-3/2} k v_A \mathcal{F}(k)^{1/2}, \quad (8)$$

with $c_k = 3.6$. Since damping is much faster than wave advection at the Alfvén speed, a good approximation for the wave distribution can be obtained by equating Γ_{cr} with Γ_{nllD} , which returns the following implicit form for the wave spectrum:

$$\mathcal{F}(k) = (2c_k)^3 \left[\frac{16\pi^2 p^4}{B_0^2} D \frac{\partial f}{\partial z} \right]^2. \quad (9)$$

Inserting the diffusive flux, $D \partial f / \partial z$ (from equation 4) into equation (9), we can derive the diffusion coefficient using equation (6):

$$D(z, p) = D_H(p) + 2v_A (H - z), \quad (10)$$

where $D_H(p)$ is the diffusion coefficient at $z = H$ and reads

$$D_H(p) = \frac{D_B}{(2c_k)^3} \left[\frac{B_0^2}{16\pi^2 p^4} \frac{1 - e^{-\xi(0, p)}}{v_A f_0(p)} \right]^2. \quad (11)$$

It follows that the diffusion coefficient is maximum at $z = 0$ and decreases linearly with z .

The exact solutions, equations (4)–(5) and (10)–(11), can be written in an explicit form in the two opposite limits of diffusion-dominated and advection-dominated transport. In particular, it is straightforward to verify that in the diffusion-dominated case (i.e. when $v_A H \ll D_H$) the leaky box solution is recovered. In fact

in this limit $e^{-\xi(0,p)} \approx 1 - v_A H/D_H$ and D becomes constant in z , namely

$$D(z, p) \rightarrow D_H(p) \rightarrow D_B^{1/3} \frac{1}{2c_k} \left[\frac{B_0^2 H}{16\pi^2 p^4 f_0(p)} \right]^{2/3}. \quad (12)$$

In the same limit, equation (4) reduces to the well known $f(z, p) = f_0(p)(1 - z/H)$, where $f_0(p) = Q_0 H/(2D_H)$. Replacing this last expression for f_0 into equation (12), we obtain explicit expressions for both D_H and f_0 , which read

$$D_H(p) = D_B \frac{1}{(2c_k)^3} \left[\frac{2B_0^2}{16\pi^2 p^4 Q_0} \right]^2 \propto p^{2\gamma-7} \quad (13)$$

and

$$f_0(p) = \frac{Q_0}{2} \frac{H}{D_H} = \frac{3c_k^3}{r_L v} \left(\frac{16\pi^2 p^4}{B_0^2} \right)^2 H Q_0(p)^3 \propto p^{7-3\gamma}, \quad (14)$$

respectively. In the opposite limit, when $v_A H \gg D_H$, we have that $e^{-\xi} \approx 0$, hence $f(z, p) \rightarrow f_0(p)$ and from equation (5) we recover

$$f_0(p) = \frac{3Q_0(p)}{2v_A \gamma}. \quad (15)$$

On the other hand, we see from equation (10) that the diffusion coefficient in the disc behaves like a constant in momentum, namely $D(z=0, p) \rightarrow 2v_A H$. This happens because for small p , $\mathcal{F} \rightarrow D_B/(2v_A H)$, hence $D = D_B/\mathcal{F} \rightarrow 2v_A H$. Clearly, this dependence is restricted to the momenta for which diffusion becomes comparable to advection, typically below $\sim 10 \text{ GeV } c^{-1}$ (see below). We refer to this regime as advection-dominated regime, although particles never reach a fully advection-dominated transport because diffusion and advection time are of the same order.

We are interested in describing the dependence of the CR spectrum on the Galactocentric distance, which enters the calculation only through the injection term and the magnetic field strength. Comparing equations (14) and (15), we see that the CR density has the following scalings with quantities depending on R :

$$\begin{aligned} f_0(p) &\propto (Q_0/B_0)^3 && \text{(diffusive regime)} \\ f_0(p) &\propto Q_0/B_0 && \text{(advective regime)}. \end{aligned} \quad (16)$$

In a more general case, equations (5), (10) and (11) can be solved iteratively. We start by choosing a guess function for $D_H(p)$ (for instance the expression, equation 13, obtained without advection) and then we iterate until convergence is reached, a procedure which usually requires only few iterations. Notice that the general case of a transport equation (1) where the advection speed may depend on the z -coordinate has been recently discussed by Recchia et al. (2016) and used to describe CR-induced Galactic winds. For the sake of simplicity, and to retain the least number of parameters, here we assume that the advection velocity is simply the Alfvén speed of self-generated waves and we assume that v_A is independent of z .

3 RESULTS

3.1 Fitting the local CR spectrum

The rate of injection of CRs per unit surface can be calibrated to reproduce the energy density and spectrum of CRs as observed at the Earth. In all our calculations, following most of current literature, we choose a size of the halo $H = 4 \text{ kpc}$, while the ion density in the halo is fixed as $n_i = 0.02 \text{ cm}^{-3}$, a value consistent with the density of the warm ionized gas component (see, e.g. Ferrière 2001). The magnetic field B_0 at the location of the Sun is assumed

to be $B_\odot = 1 \mu\text{G}$ (since we are only describing the propagation in the z -direction, this should be considered as the component of the field perpendicular to the disc). Notice that, given B_0 and n_i , the value of the Alfvén speed is also fixed, and this is very important in that it also fixes the momentum where the transition from advection propagation to diffusion-dominated propagation takes place for a given injection spectrum and product of injection efficiency times the local SN explosion rate, $\xi_{\text{inj}} \times \mathcal{R}_{\text{SN}}(R_\odot)$. Following Blasi et al. (2012), Aloisio & Blasi (2013) and Aloisio et al. (2015), we adopted a slope at injection $\gamma = 4.2$. Then, by requiring that the local CR density at $\sim 10\text{--}50 \text{ GeV}$ is equal to the observed one, we get $\xi_{\text{inj}}/0.1 \times \mathcal{R}_{\text{SN}}/(1/30 \text{ yr}) = 0.29$.

It is worth stressing that the CR spectrum in the energy region $\gtrsim 100 \text{ GeV}$ may be heavily affected by either pre-existing turbulence Blasi et al. (2012); Aloisio & Blasi (2013); Aloisio et al. (2015) or a z -dependent diffusion coefficient (Tomassetti 2012). Both possibilities have been proposed to explain the spectral hardening observed in both the protons' and helium spectrum at rigidities above $\sim 200 \text{ GV}$. For this reason, a model including only self-generated diffusion can be considered as reliable only below $\sim 50 \text{ GeV}$. In the following, we limit our attention to CRs that are responsible for the production of γ -rays of energy $\sim 2 \text{ GeV}$, as observed by *Fermi*-LAT, namely protons with energy of the order of $\sim 20 \text{ GeV}$. This threshold is sufficiently low that the slope derived by ignoring the high energy spectral hardening can be considered reliable.

The injection parameters (efficiency and spectrum) found by fitting the CR density and spectrum at the Sun's location are assumed to be the same for the whole Galaxy. As discussed in Section 3.2, the rate of injection of CRs per unit surface is then proportional to the density of SNRs as inferred from observations.

3.2 CR spectrum in the Galactic disc

The SNR distribution is usually inferred based on two possible tracers: radio SNRs and pulsars. Here, we adopt the distribution of SNRs recently obtained by Green (2015) from the analysis of bright radio SNRs. He adopted a cylindrical model for the Galactic surface density of SNRs as a function of the Galactocentric radius, in the form

$$f_{\text{SNR}} \propto \left(\frac{R}{R_\odot} \right)^\alpha \exp \left(-\beta \frac{R - R_\odot}{R_\odot} \right), \quad (17)$$

where the position of the Sun is assumed to be at $R_\odot = 8.5 \text{ kpc}$. For the best-fitting Green (2015) obtained $\alpha = 1.09$ and $\beta = 3.87$, so that the distribution is peaked at $R = 2.4 \text{ kpc}$. However, as noted by Green (2015), it is worth keeping in mind that the best-fitting model is not very well defined, as there is some level of degeneracy between the parameters α and β .

Case & Bhattacharya (1998) also adopted a fitting function as in equation (17) but obtained their best fit for $\alpha = 2.0$ and $\beta = 3.53$, resulting in a distribution peaked at $R = 4.8 \text{ kpc}$ and broader for larger values of R with respect to the one of Green (2015). Case & Bhattacharya (1998) estimated the source distances using the so called ' Σ - D ' relation, that is well known to be affected by large uncertainties. Moreover, Green (2015) argued that the Σ - D used by Case & Bhattacharya (1998) appears to have been derived incorrectly.

An important caveat worth keeping in mind is that the SNR distribution derived in the literature is poorly constrained for large galactocentric radii. For instance, Green (2015) used a sample of 69 bright SNRs but only two of them are located at galactic latitude $l > 160^\circ$. Similarly, Case & Bhattacharya (1998) used a larger sample




Figure 1. CR density at $E > 20$ GeV (Acero et al. 2016) and emissivity per H atom (Yang et al. 2016) as a function of the Galactocentric distance, as labelled. Our predicted CR density at $E > 20$ GeV is shown as a dashed line. The case of exponentially suppressed magnetic field is shown as a solid line. The dotted line shows the distribution of sources (Green 2015).

with 198 SNRs, but only 7 of them are located at $R > 13$ kpc and there are no sources beyond 16 kpc.

The distribution of pulsars is also expected to trace that of SNRs after taking into account the effect of birth kick velocity, that can reach ~ 500 km s^{-1} . These corrections are all but trivial (see, e.g. Faucher-Giguère & Kaspi 2006), hence in what follows we adopt the spatial distribution as inferred by Green (2015).

One last ingredient needed for our calculation is the magnetic field strength, $B_0(R)$, as a function of galactocentric distance R . While there is a general consensus that the magnetic field in the Galactic disc is roughly constant in the inner region, in particular in the so-called ‘molecular ring’, between 3 and 5 kpc (Stanev 1997; Jansson & Farrar 2012), much less is known about what the trend is in the very inner region around the Galactic Centre, and in the outer region, at $R > 5$ kpc. Following the prescription of Jansson & Farrar (2012, see also Stanev 1997), we assume the following radial dependence:

$$\begin{aligned} B_0(R < 5 \text{ kpc}) &= B_\odot R_\odot / 5 \text{ kpc} \\ B_0(R > 5 \text{ kpc}) &= B_\odot R_\odot / R, \end{aligned} \quad (18)$$

where the normalization is fixed at the Sun’s position, that is $B_\odot = 1 \mu\text{G}$. Using this prescription, we calculate the CR spectrum as a function of the Galactocentric distance, as discussed in Section 2. In Fig. 1, we plot the density of CRs with energy $\gtrsim 20$ GeV (dashed line) and compare it with the same quantity as derived from *Fermi*-LAT data. Our results are in remarkably good agreement with data, at least out to a distance of ~ 10 kpc. At larger distances, our predicted CR density drops faster than the one inferred from data, thereby flagging again the well-known CR gradient problem. In fact, the non-linear theory of CR propagation, in its most basic form (dashed line) makes the problem even more severe: where there are more sources, the diffusion coefficient is reduced and CRs are trapped more easily, but where the density of sources is smaller the corresponding diffusion coefficient is larger and the CR density drops. A similar situation can be seen in the trend of the spectral slope as a function of R , plotted in Fig. 2. The dashed line reproduces well the slope inferred from *Fermi*-LAT data out to a distance of ~ 10 kpc, but not in the outer regions where the predicted spectrum




Figure 2. Radial dependence of the power-law index of the proton spectrum as inferred by (Acero et al. 2016, filled circle) and (Yang et al. 2016, filled triangle). Our predicted slope for the basic model is shown as a dashed line, while the solid line illustrates the results for the exponentially suppressed magnetic field.

is steeper than observed. It is important to understand the physical motivation for such a trend: at intermediate values of R , where there is a peak in the source density, the diffusion coefficient is smaller and the momenta for which advection dominates upon diffusion is higher. This implies that the equilibrium CR spectrum is closer to the injection spectrum, $Q(p)$ (harder spectrum). On the other hand, for very small and for large values of R , the smaller source density implies a larger diffusion coefficient and a correspondingly lower momentum where advection dominates upon diffusion. As a consequence the spectrum is steeper, namely closer to $Q(p)/D(p)$. In fact, at distances $R \gtrsim 15$ kpc, the spectrum reaches the full diffusive regime, hence $f_0 \sim p^{7-3\gamma} = p^{-5.6}$, meaning that the slope in Fig. 2 is 3.6. As pointed out in Section 2, the non-linear propagation is quite sensitive to the dependence of the magnetic field on R .

Both the distribution of sources and the magnetic field strength in the outer regions of the Galaxy are poorly known. Hence, we decided to explore the possibility that the strength of the magnetic field may drop faster than $1/R$ at large galactocentric distances. As a working hypothesis, we assumed the following form for the dependence of B_0 on R , at $R \gtrsim 10$ kpc:

$$B_0(R > 10 \text{ kpc}) = \frac{B_\odot R_\odot}{R} \exp\left[-\frac{R - 10 \text{ kpc}}{d}\right], \quad (19)$$

where the scalelength, d , is left as a free parameter. We found that using $d = 3.1$ kpc, both the resulting CR density and spectral slope describe very well the *Fermi*-LAT data in the outer Galaxy. The results of our calculations for this case are shown in Figs 1 and 2 with solid lines.

The diffusion coefficient resulting from the non-linear CR transport in the Galaxy, calculated as in Section 2, is illustrated in Fig. 3, for different galactocentric distances. It is interesting to notice that at all values of R (and especially at the Sun’s position) $D(p)$ is almost momentum independent at $p \lesssim 10$ GeV c^{-1} . This reflects the fact that at those energies the transport is equally contributed by both advection and diffusion, as discussed above. This trend, that comes out as a natural consequence of the calculations, is remarkably similar to the one that in numerical approaches to CR transport is imposed by hand in order to fit observations.




Figure 3. Diffusion coefficient $D(z=0, p)$ as a function of momentum in GeV/c for different Galactocentric distances as labelled.

Contrary to a naive expectation, in the case in which $B_0(R)$ drops exponentially, the diffusion coefficient becomes smaller in the external Galaxy than in the inner part, in spite of the smaller number of sources in the outer Galaxy. This counterintuitive result is due to the fact that $D_H(p) \propto B_0^4/Q_0^2$ (see equation 13) and that both B_0 and Q_0 are assumed to drop exponentially at large R . Clearly, this result loses validity when $\delta B/B_0$ approaches unity and the amplification enters the non-linear regime. Using equation (10), such condition in the disc can be written as $\mathcal{F}(z=0, k) \approx D_B/(2v_A H) \gtrsim 1$ which, for 1 GeV particles occurs for $R \gtrsim 28$ kpc (red-dashed line in Figs 1 and 2). In any case, the density of CRs at large galactocentric distances drops down, as visible in Fig. (1).

4 CONCLUSIONS

The CR density recently inferred from *Fermi*-LAT observations of the diffuse Galactic γ -ray emission, as carried out during the last 7 yr, appears to be all but constant with galactocentric distance R Acero et al. (2016); Yang et al. (2016). In the inner ~ 5 kpc from the Galactic Centre, such density shows a pronounced peak around 3–4 kpc, while it drops with R for $R \gtrsim 5$ kpc, but much slower than what one would expect based on the distribution of SNRs, as possible sources of Galactic CRs. Moreover, the inferred slope of the CR spectrum shows a gradual steepening in the outer regions of the Galaxy. This puzzling CR gradient is hard to accommodate in the standard picture of CR transport.

Here, we showed that both the gradient and the spectral shape can be explained in a simple model of non-linear CR transport: CRs excite waves through streaming instability in the ionized Galactic halo and are advected with such Alfvén waves. In this model, the diffusion coefficient is smaller where the source density is larger and this phenomenon enhances the CR density in the inner Galaxy. In the outer Galaxy, the data can be well explained only by assuming that the background magnetic field drops exponentially at $R \gtrsim 10$ kpc,

with a suppression scale of ~ 3 kpc. This scenario also fits well the spectral slope of the CR spectrum as a function of R , as a result of the fact that at different R the spectrum at a given energy (~ 20 GeV) may be dominated by advection (harder spectrum) or diffusion (softer spectrum). A simple prediction of our calculations is that the spectral hardening should disappear at higher energies, where transport is diffusion dominated at all galactocentric distances.

ACKNOWLEDGEMENTS

The authors acknowledge useful conversations with R. Aloisio and C. Evoli.

REFERENCES

- Acero F. et al., 2016, *ApJS*, 223, 26
Ackermann M. et al., 2011, *ApJ*, 726, 81
Ackermann M. et al., 2012, *A&A*, 538, A71
Adriani O. et al., 2011, *Science*, 332, 69
Aguilar M. et al., 2015, *Phys. Rev. Lett.*, 114, 171103
Aloisio R., Blasi P., 2013, *J. Cosmol. Astropart. Phys.*, 7, 001
Aloisio R., Blasi P., Serpico P. D., 2015, *A&A*, 583, A95
Berezinskii V. S., Bulanov S. V., Dogiel V. A., Ptuskin V. S., 1990, *Astrophysics of Cosmic Rays*. North-Holland, Amsterdam
Bhat C. L., Mayer C. J., Rogers M., Wolfendale A. W., Zan M., 1986, *J. Phys. G: Nucl. Part. Phys.*, 12, 1087
Blasi P., Amato E., Serpico P. D., 2012, *Phys. Rev. Lett.*, 109, 061101
Bloemen J. B., G. M. et al., 1986, *A&A*, 154, 25
Bloemen J. B. G. M., Dogiel V. A., Dorman V. L., Ptuskin V. S., 1993, *A&A*, 267, 372
Case G. L., Bhattacharya D., 1998, *ApJ*, 504, 761
Erlykin A. D., Wolfendale A. W., Dogiel V. A., 2016, *Adv. Space Res.*, 57, 519
Evoli C., Gaggero D., Grasso D., Maccione L., 2012, *Phys. Rev. Lett.*, 108, 211102
Faucher-Giguère C.-A., Kaspi V. M., 2006, *ApJ*, 643, 332
Ferrière K. M., 2001, *Rev. Mod. Phys.*, 73, 1031
Gaggero D., Urbano A., Valli M., Ullio P., 2015a, *Phys. Rev. D*, 91, 083012
Gaggero D., Grasso D., Marinelli A., Urbano A., Valli M., 2015b, *ApJ*, 815, L25
Green D. A., 2015, *MNRAS*, 454, 1517
Hunter S. D. et al., 1997, *ApJ*, 481, 205
Jansson R., Farrar G. R., 2012, *ApJ*, 757, 14
Ptuskin V. S., Zirakashvili V. N., 2003, *A&A*, 403, 1
Recchia S., Blasi P., Morlino G., 2016, preprint ([arXiv:1603.06746](https://arxiv.org/abs/1603.06746))
Skillington J., 1975, *MNRAS*, 173, 255
Stanev T., 1997, *ApJ*, 479, 290
Stecker F. W., Jones F. C., 1977, *ApJ*, 217, 843
Strong A. W., Mattox J. R., 1996, *A&A*, 308, L21
Strong A. W., Moskalenko I. V., Reimer O., Digel S., Diehl R., 2004, *A&A*, 422, L47
Tomassetti N., 2012, *ApJ*, 752, L13
Yang R., Aharonian F., Evoli C., 2016, *Phys. Rev. D*, 93, 123007

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.