



Publication Year	2024
Acceptance in OA	2025-05-16T14:50:41Z
Title	Study of solar brightness profiles in the 18-26 GHz frequency range with INAF radio telescopes. II. Evidence of coronal emission
Authors	MARONGIU, Marco, PELLIZZONI, ALBERTO PAOLO, RIGHINI, Simona, Mulas, S., NESTI, Renzo, Burtovoi, A., Romoli, M., SERRA, Giampaolo, Valente, G., EGRON, Elise Marie Jeanne, Murtas, G., Iacolina, M. N., MELIS, Andrea, GUGLIELMINO, SALVATORE LUIGI, LORU, Sara, Zucca, P., ZANICHELLI, Alessandra, BACHETTI, Matteo, BEMPORAD, Alessandro, BUFFA, Franco, CONCU, Raimondo, DEIANA, Gian Luigi, Karakotia, C., LADU, Adelaide, MACCAFERRI, Andrea, MARONGIU, Pasqualino, MESSEROTTI, Mauro, NAVARRINI, Alessandro, ORFEI, Alessandro, ORTU, Pierluigi, PILI, Mauro, PISANU, Tonino, PUPILLO, Giuseppe, ROMANO, Paolo, SABA, Andrea, SCHIRRU, Luca, Tiburzi, C., Abbo, L., FRASSATI, Federica, Giarrusso, M., JERSE, GIOVANNA, LANDINI, Federico, PANCRAZZI, Maurizio, RUSSANO, Giuliana, SASSO, CLEMENTINA, SUSINO, ROBERTO
Publisher's version (DOI)	10.1051/0004-6361/202348770
Handle	http://hdl.handle.net/20.500.12386/37148
Journal	ASTRONOMY & ASTROPHYSICS
Volume	684

Study of solar brightness profiles in the 18–26 GHz frequency range with INAF radio telescopes

II. Evidence of coronal emission

M. Marongiu¹, A. Pellizzoni¹, S. Righini², S. Mulas¹, R. Nesti³, A. Burtovoi^{4,3}, M. Romoli^{4,3}, G. Serra⁵, G. Valente⁵, E. Egron¹, G. Murtas⁶, M. N. Iacolina⁵, A. Melis¹, S. L. Guglielmino⁷, S. Loru⁷, P. Zucca⁸, A. Zanichelli², M. Bachetti¹, A. Bemporad⁹, F. Buffa¹, R. Concu¹, G. L. Deiana¹, C. Karakotia², A. Ladu¹, A. Maccaferri², P. Marongiu¹, M. Messerotti^{10,11}, A. Navarrini^{1,12}, A. Orfei², P. Ortu¹, M. Pili¹, T. Pisanu¹, G. Pupillo², P. Romano⁷, A. Saba⁵, L. Schirru¹, C. Tiburzi¹, L. Abbo⁹, F. Frassati⁹, M. Giarrusso^{4,7}, G. Jerse¹⁰, F. Landini⁹, M. Pancrazzi⁹, G. Russano¹³, C. Sasso¹³, and R. Susino⁹

¹ INAF – Cagliari Astronomical Observatory, Via della Scienza 5, 09047 Selargius (CA), Italy
e-mail: marco.marongiu@inaf.it

² INAF – Institute of Radio Astronomy, Via Gobetti 101, 40129 Bologna, Italy

³ INAF – Astrophysical Observatory of Arcetri, Largo Enrico Fermi 5, 50125 Firenze, Italy

⁴ University of Florence, Department of Physics and Astronomy, Via Giovanni Sansone 1, 50019 Sesto Fiorentino (FI), Italy

⁵ ASI – c/o Cagliari Astronomical Observatory, Via della Scienza 5, 09047 Selargius (CA), Italy

⁶ Los Alamos National Laboratory, Bikini Atoll Rd, Los Alamos, NM 87545, USA

⁷ INAF – Catania Astrophysical Observatory, Via Santa Sofia 78, 95123 Catania, Italy

⁸ ASTRON – The Netherlands Institute for Radio Astronomy, Oude Hoogeveensedijk 4, 7991 PD Dwingeloo, The Netherlands

⁹ INAF – Turin Astrophysical Observatory, Via Osservatorio 20, 10025 Pino Torinese (TO), Italy

¹⁰ INAF – Trieste Astronomical Observatory, Via Giambattista Tiepolo 11, 34131 Trieste, Italy

¹¹ Department of Physics, University of Trieste, Via Alfonso Valerio 2, 34127 Trieste, Italy

¹² NRAO – Central Development Laboratory, 1180 Boxwood Estate Rd, Charlottesville, VA 22903, USA

¹³ INAF – Astronomical Observatory of Capodimonte, Salita Moiariello 16, 80131 Napoli, Italy

Received 28 November 2023 / Accepted 10 February 2024

ABSTRACT

Context. One of the most important objectives of solar physics is to gain a physical understanding of the solar atmosphere, whose structure can also be described in terms of the density (N) and temperature (T) distributions of the atmospheric matter. Several multi-frequency analyses have shown that the characteristics of these distributions are still under debate, especially for outer coronal emission.

Aims. We aim to constrain the T and N distributions of the solar atmosphere through observations in the centimetric radio domain. We employed single-dish observations from two of the INAF radio telescopes at the K -band frequencies (18–26 GHz). We investigated the origin of the significant brightness temperature (T_B) detected up to the upper corona (at an altitude of ~ 800 Mm with respect to the photospheric solar surface).

Methods. To probe the physical origin of the atmospheric emission and to constrain instrumental biases, we reproduced the solar signal by convolving specific 2D antenna beam models. We performed an analysis of the solar atmosphere by adopting a physical model that assumes the thermal bremsstrahlung as the emission mechanism, with specific T and N distributions. We compared the modelled T_B profiles with those observed by averaging solar maps obtained at 18.3 and 25.8 GHz during the minimum of solar activity (2018–2020).

Results. We probed any possible discrepancies between the T and N distributions assumed from the model and those derived from our measurements. The T and N distributions are compatible (within a 25% of uncertainty) with the model up to ~ 60 Mm and ~ 100 Mm in altitude, respectively.

Conclusions. Our analysis of the role of the antenna beam pattern on our solar maps proves the physical nature of the atmospheric emission in our images up to the coronal tails seen in our T_B profiles. Our results suggest that the modelled T and N distributions are in good agreement (within 25% of uncertainty) with our solar maps up to altitudes of $\lesssim 100$ Mm. A subsequent, more challenging analysis of the coronal radio emission at higher altitudes, together with the data from satellite instruments, will require further multi-frequency measurements.

Key words. methods: data analysis – Sun: atmosphere – Sun: chromosphere – Sun: corona – Sun: photosphere – Sun: radio radiation

1. Introduction

Our understanding of the physical processes that govern mass and energy flow in the solar atmosphere represents one of

the fundamental goals of solar physics (for a review, see e.g. Shibasaki et al. 2011). Analyses of the physical properties characterising the solar atmosphere (e.g. the density and the temperature) are crucial to improving our understanding of these

processes. Multi-frequency observations (from radio to EUV-X domain) of the Sun probe different layers of the solar atmosphere, with respect to the photosphere level of $R_{\odot, \text{opt}} = 695.66 \pm 0.14 \text{ Mm}$ (corresponding to $959.16 \pm 0.19 \text{ arcsec}$ at Sun-Earth distance of 1 AU^1 ; Mamajek et al. 2015; Prša et al. 2016; Haberreiter et al. 2008). Such observations offer insights into the thermal origin (free-free bremsstrahlung radiation) of the atmospheric emission of the quiet Sun. Many atmospheric models are based on optical spectral analysis (e.g. Vernazza et al. 1973, 1976, 1981; Fontenla et al. 1990, 1991, 1993, 2002), allowing us to describe the layers of the solar atmosphere in terms of the temperature and the density distributions.

At radio frequencies, the quiet Sun is characterised by pure free-free emission at local thermal equilibrium (LTE). Under LTE conditions, radio waves cannot propagate outwards through regions below a specific radius, R_{ω} , where the local plasma frequency, ω_p , equals the observing frequency; ω_p is defined as $\sqrt{N_e e^2 / (m_e \epsilon_0)}$, where N_e is the electron number density, e is the elementary charge, m_e is the electron mass, and ϵ_0 is the vacuum permittivity. In this frequency domain (especially in the MHz range), the atmosphere of the Sun (from the photosphere to the upper corona, as well as its structure) was observed and modelled by several authors in the literature over many decades (e.g. Newkirk 1961; Wild 1970; Dulk et al. 1977; Kundu et al. 1983; Wang et al. 1987; Ramesh et al. 1998, 2010; Krissinel' 2005; Zhang et al. 2022). These models play a fundamental role in the interpretation of solar radio emission and in the study of outwardly flowing coronal material into the solar wind (Lallement et al. 1986; Fludra et al. 1999; Meyer-Vernet 2012). In addition, they have indicated variability in the corona, depending upon solar activity (e.g. Saito et al. 1977). Among these models, starting from the observations at 17 GHz with the Nobeyama Radioheliograph (NoRH; Nakajima et al. 1994, 1995) Selhorst et al. (2005, 2008) developed a two-dimensional (2D) temperature and density model to reproduce the observations ranging from 1.4 to 400 GHz, assuming the thermal bremsstrahlung as the emission mechanism. This theoretical model reproduces the brightness temperature of the quiet-Sun, T_{qs} , at the disk centre, the centre-to-limb brightening distribution, the value of the solar radius, R_{\odot} , and it provides an estimation of the local distribution of temperature and density of the solar atmosphere. However, this model and other similar approaches (e.g. Mercier & Chambe 2015; Vocks et al. 2018; Ramesh et al. 2020) are characterised for simplicity by a symmetrical solar disk. This aspect cannot independently explain both the equatorial and the polar limb profiles, thus preventing an accurate 3D description of the density and the temperature distributions of the atmospheric layers of the Sun. The comparison between the theoretical and the observed data is crucial to improve the modelling of the solar corona and, hence, to better understand the atmospheric structures in terms of density and temperature (e.g. Saint-Hilaire et al. 2012; Mercier & Chambe 2015; Vocks et al. 2018; Ramesh et al. 2020).

In this paper, we focus on the study of the atmospheric emission through the analysis of the brightness profiles of about 300 solar images, obtained through single-dish observations from the newly appointed Medicina Gavril Grueff Radio Telescope (hereafter, Grueff Radio Telescope) and the Sardinia Radio Telescope (SRT) at the radio K -band (18.3–25.8 GHz), over about half a solar cycle (from 2018 to mid-2023), in the frame of the SunDish

project aimed at the development of solar imaging observing modes for the INAF radio telescopes (Pellizzoni et al. 2022). Through this project, accurate brightness temperature, T_B , profiles of the solar disk were obtained and the solar size was deeply analysed, finding that the measured solar radii, R_{\odot} – ranging between 959 and 994 arcsec – calculated for each observing frequency and radio telescope, are consistent with the other values reported in the literature (Marongiu et al. 2024). The coverage of the entire solar disk, the low noise (RMS of the solar maps of $<10 \text{ K}$) and the low contribution of the absolute calibration in the uncertainty on T_B ($\leq 3\%$), make our data set valuable for analysing and modelling the solar atmospheric emission in the radio K -band. These aspects have allowed us to detect significant emission up to the weak outer corona contributing to the ‘tails’ extending outside the solar disk in our brightness profiles.

We organise this paper as follows. A brief review of the solar imaging observations with INAF radio telescopes is presented in Sect. 2. The description of techniques employed for analysing the solar atmosphere and related results is reported in Sect. 3. In particular, to corroborate the physical origin of the weak features in our brightness profiles, as seen in the correspondence of the outer corona, we modelled our solar maps by accounting for a specific 2D model (Sect. 3.1); the brightness temperature spectrum profiles of our observed maps are compared with the theoretical estimations (discussed in the literature) in Sect. 3.2. After the discussion of our results in Sect. 4, we give our conclusions in Sect. 5.

2. Observations and data reduction

The data set used in this work was provided by single-dish observation campaigns at the central frequency ranging between 18.1 and 26.1 GHz, with the network of the INAF single-dish radio telescopes². This network includes the Grueff Radio Telescope, the Sardinia Radio Telescope (SRT), and the Noto³ Radio Telescope. The duration of these solar campaigns covers over five years, from 2018 to mid-2023. These solar observations are the core of the ‘SunDish Project’ (PI: A. Pellizzoni)⁴, in collaboration with INAF and ASI (Pellizzoni et al. 2019, 2022; Plainaki et al. 2020), aimed at the imaging and monitoring of the solar atmosphere.

The 32-m Medicina Radio Telescope, located near Bologna at a 25 m elevation, observes and monitors the Sun once a week since February 2018. The solar observations are carried out through the K -band dual-feed receiver operating at central frequency ranging between 18.3 and 25.8 GHz. The beam pattern of this receiver is characterised by a beam size ranging between 1.5 and 2.1 arcmin.

The 64-m SRT is located at 650 m elevation in Sardinia (Italy). Up to date, SRT has observed the Sun mostly at 18.8 and 24.7 GHz once a month through a seven-feed dual-polarization K -band receiver, customised for solar observations (Bolli et al. 2015; Prandoni et al. 2017; Pellizzoni et al. 2022) and characterised by a beam size of 1.0 and 0.8 arcmin, respectively. This radio telescope is currently in the final step of an upgrade phase that includes the installation of new receivers (suitable also for solar observations) operating up to 116 GHz in the context of the National Operative Programme (Programma Operativo Nazionale-PON; Govoni et al. 2021)⁵.

² <https://www.radiotelesopes.inaf.it>

³ Solar observations setup and operations are not yet implemented at Noto.

⁴ <https://sites.google.com/inaf.it/sundish>

⁵ <https://sites.google.com/a/inaf.it/pon-srt/home>

¹ In this paper, we reported the angular solar size normalised to 1 AU in all our conversions from the ‘absolute’ size (expressed in units of Mm), unless stated otherwise.




Fig. 1. Examples of polar (red line) and equatorial (green line) brightness temperature profiles derived from solar images obtained through the *Grueff* Radio Telescope and SRT. Top: profiles derived on September 6, 2020 from the 18.3 GHz (left) and 25.8 GHz (right) *Grueff* Radio Telescope images. Bottom: profiles derived on January 28, 2020 from the 18.8 GHz (left) and 24.7 GHz (right) SRT images. Typical errors in the brightness profiles are in the 5–20 K range, significantly affecting only the outwards tails. Blue dashed lines indicate the solar radius at the photosphere level $R_{\odot, \text{opt}} = 959.16 \pm 0.19$ arcsec (Mamajek et al. 2015; Prša et al. 2016; Haberreiter et al. 2008).

In the time frame between 2018 and mid-2023, most of our solar maps in *K*-band were performed by the *Grueff* Radio Telescope. For this work we employed an extensive data set of 290 maps (273 maps from *Grueff*, and 17 maps from SRT): in particular, this data set is composed of 145 maps at 18.3 GHz (*Grueff*), 128 maps at 25.8 GHz (*Grueff*), 10 maps at 18.8 GHz (SRT), and 7 maps at 24.7 GHz (SRT). Mostly due to high atmospheric opacity or even instrumental errors, some maps had to be discarded ($\sim 10\%$).

The radio signal is processed using the DISCOS antenna control system⁶ through full-stokes spectral-polarimetric ROACH2-based back-end (SARDARA system, Melis et al. 2018) at SRT, and through total-power and intensity back-end configuration at Medicina⁷. The solar maps are reconstructed from on-the-fly (OTF) scans (Prandoni et al. 2017) in right ascension (RA) and declination (Dec), respectively, covering an area in the sky of 4800×4800 arcsec at Medicina, and

⁶ <https://discos.readthedocs.io/en/latest/user/index.html>

⁷ From July 2021 this radio telescope is also equipped by a full-stokes spectral-polarimetric ROACH2-based back-end through the SARDARA system (Mulas et al. 2022). This back-end is available to the scientific community since the observing semester 2023A.

5400×5400 arcsec at SRT. The imaging and calibration procedures are performed using the SRT Single-Dish Imager (SDI; Egron et al. 2017; Pellizzoni et al. 2019, 2022; Loru et al. 2019, 2021; Marongiu et al. 2020, 2022; Mulas et al. 2022). The data analysis is provided by the Python solar pipeline SUNPIT (Marongiu et al. 2022). Flux density and brightness temperature calibrations are achieved through the exploitation of the supernova remnant Cas A (for details see Pellizzoni et al. 2022; Mulas et al. 2022). Both for the *Grueff* Radio Telescope and SRT, details about the mapping techniques, the setup configurations, the observing strategy, and the data processing are available in Pellizzoni et al. (2022). The data set of *Grueff* and SRT is available in the SunDish Archive⁸.

3. Data analysis

In this work, we analyse the properties of the solar atmospheric layers in terms of the temperature and the density distributions also above the solar radius altitude. This kind of analysis originated from the presence of ‘tails’ extending outside the solar

⁸ <https://sites.google.com/inaf.it/sundish/sundish-images-archive/sundish-solar-archive>

disk signal in our brightness profiles, as shown in Fig. 1. These tails are characterised by the following features:

- No significant differences are observed between polar and equatorial brightness temperature, T_B , within the errors (ranging between 10 and 20 K), as well as between *Grueff* and SRT measurements;
- There are no significant fluctuations of T_B over time for each observing frequency, apart for statistical fluctuations compatible with noise;
- The T_B level of the tails at high K -band (~ 25 GHz) is higher than the T_B level at low K -band (~ 18 GHz), suggesting a thermal nature for the tail (coronal) emission (we estimate a spectral index⁹ $\alpha \gtrsim 1.5$);
- There is no correlation between the tails and the elevation δ of the Sun during the observations both at SRT and *Grueff* radio telescopes (δ ranges between 20 and 60 degrees).

To probe the physical origin of these tails (connected with the atmospheric emission) in our solar images and to obtain more robust measurements of the solar radii (Marongiu et al. 2024), we analysed the degradation effects on the solar images due to the convolution of the antenna beam pattern with the solar signal (Sect. 3.1). Moreover, we compared the brightness temperature profiles of our observed solar maps with those obtained through a specific atmospheric model, assuming the thermal bremsstrahlung as the main emission mechanism (Sect. 3.2).

For this work, we used 290 solar maps (273 with *Grueff* and 17 with SRT) to derive the brightness profiles and analyse the atmospheric emission in the range 18–26 GHz. We selected four frequencies, characterised by a uniform epoch coverage from 2018 to date: 18.3 and 25.8 GHz for *Grueff* and 18.8 and 24.7 GHz for SRT. Moreover, we also used two averaged solar maps at 18.3 and 25.8 GHz obtained with the *Grueff* Radio Telescope during the minimum solar activity (2018–2020).

3.1. Prescription for probing the role of the antenna beam pattern in the observed atmospheric emission

One of the most important features that influence solar radio measurements is the degrading effect of the antenna beam pattern on the solar signal (e.g. Giménez de Castro et al. 2020; Menezes et al. 2022). As we show in this work, our analysis of the role of the antenna beam pattern (to evaluate its impact on the solar radii measurements and to assess the radius determination; see more details in Marongiu et al. 2024) also allows us to probe the physical nature of the external coronal emission observed as ‘tails’ in the brightness profiles of our images of the Sun (see Fig. 1).

The *Grueff* and SRT beam patterns were reconstructed using a dedicated software called GRASP¹⁰; except the measurement noise and some residual optical aberration, the beam patterns measured with the *Grueff* and SRT radio telescopes (Prandoni et al. 2017; Egron et al. 2022) are in good agreement with the simulated ones obtained with GRASP. We used the 2D beam patterns obtained through four observing frequencies (18.3 and 25.8 GHz for *Grueff*; 18.8 and 24.7 GHz for SRT), the most observed by the *Grueff* and SRT radio telescopes during solar observations. The peaks of the main and secondary lobes of these beam patterns are reported in Table 1. Adopting the GRASP beam model, the secondary lobes cannot account for the observed outer tails in the brightness profiles, as an artifact

Table 1. Values of the amplitudes of the main (a_0) and secondary (a_2) lobes (in units of dBi) obtained with GRASP for the analysed beam patterns of *Grueff* and SRT radio telescopes.

Frequency (GHz)	Telescope	a_0 (dBi)	a_2 (dBi)	Δa (dBi)	R_{off} (arcmin)
18.3	<i>Grueff</i>	73.5	53.0	–20.5	3.1
25.8	<i>Grueff</i>	76.2	48.0	–28.2	2.7
18.8	SRT	80.4	61.4	–19.0	1.5
24.7	SRT	82.9	60.8	–22.1	1.3

Notes. Δa reports the difference between a_2 and a_0 , and R_{off} indicates the offset (in units of arcmin) between the secondary and the main lobes centroids.

Table 2. Uniform prior distribution adopted for the Bayesian approach in the solar signal models of our analysis.

Model	Parameter	Range	Unit
GBP	\mathcal{A}	$0.1 \leq \mathcal{A} \leq 10$	–
CCBH	R	$R_0 - 100 \leq R_0 \leq R_0 + 100$	arcsec
CCBH	l	$0.5 \leq l_0 \leq 10000$	K

Notes. CCBH indicates the solar signal model, and GPB indicates the model of the generic beam pattern; these models are described in Sect 3.1.

due to instrumental bias. In fact, beam lobes can drive emission from the solar disk to the tail at a level only <100 – 1000 times lower (depending on the observing frequency) than the disk emission itself, while the tail brightness is still $>1\%$ of that at >8 – 10 arcmin from the disk limb (Fig. 1). Thus, our solar maps show a signal from the high solar atmosphere that is incompatible with instrumental biases caused by the degrading effect of the antenna beam pattern on the solar signal.

For the sake of completeness, to properly take into account possible instrumental biases in our images and the effects of hypothetical instabilities of the opto-mechanical performances of the radiotelescopes during solar observations, we provided a parametrized 2D beam model derived from the original obtained by GRASP, along the lines of other similar works (e.g. Iwai et al. 2017). We modelled the generic beam pattern as a function characterised by (1) the instrumental (fixed) amplitude, a_0 , of the main lobe (Table 1) and (2) a variable amplitude $A_{\text{suc}} = \mathcal{A} \times a_{\text{suc}}$ of the successive lobe peaks (Table 2), where a_{suc} is the instrumental (fixed) amplitude of the successive lobes, and \mathcal{A} is the degradation level of the beam pattern (free parameter). We define \mathcal{A} as a dimensionless constant that describes the amplitude of the secondary lobe, accounting for possible unexpected optical deformations and aberrations. The amplitude of the lobes in this model is expressed in units of dBi, and it adheres to the principle of energy conservation, maintaining constant the volume of the generic beam pattern. In this context, \mathcal{A} is taken inversely proportional to the width of the main lobe.

We convolved this parametrized beam model with specific solar signal models (see below) in order to fit our observed solar images. This procedure (hereafter, ‘‘beam fitting procedure’’) provides as output, the best-fit parameters for (1) the generic beam model adopted as input (\mathcal{A} , and hence the modelled secondary lobe amplitude A_2), and (2) the R_\odot values of the adopted solar signal model. The best-fit parameters of the beam fitting procedure are obtained thanks to a Bayesian approach based on Markov chain MonteCarlo (MCMC) simulations (e.g. Sharma 2017).

⁹ For this estimation we considered a preliminary coronal ring defined between 1.1 and $1.7 R_\odot$.

¹⁰ <https://www.ticra.com/software/grasp/>




Fig. 2. Example of modelled equatorial T_B profiles of a solar map obtained adopting the ECB-model as the solar signal for SRT at 24.7 GHz (October 9, 2019), in comparison with the observed brightness profile. These plots differ in the T_B y-axis scale (linear on top, logarithmic on bottom). Blue dotted line indicates the observed equatorial T_B profile of a solar map at this observing frequency with SRT, green solid line indicates the equatorial T_B profile obtained convolving the solar signal model with the GRASP original beam pattern, and red solid line indicates the modelled equatorial T_B profile obtained with a generic beam pattern with the secondary lobe characterised by $A_2 = 1.2 \times a_2$ (where a_2 is the reference value obtained by GRASP).

For this approach we used the Python `emcee` package¹¹ (Foreman-Mackey et al. 2013). The complete uniform prior distribution adopted in our analysis is listed in Table 2, where the parameters labelled with 0 indicate the initial best-fit parameters. All the uncertainties are reported at 68% (1σ). Further details about this approach applied to 2D solar signal models are available in Marongiu et al. (2024).

The level of the “tails” seen outside the solar disk in the brightness profiles of the 2D-convolved solar maps is directly proportional to \mathcal{A} .

Adopting a simple elliptical-based cylindrical box (ECB-model, described in Marongiu et al. 2024) that trivially describes a pure and flat disk emission as the solar signal, it is not possible to fit the observed brightness profile, even in the assumption of significant beam degradation effects (Fig. 2). Furthermore, when the beam degradation level is significant (i.e. $\mathcal{A} \gtrsim 1.05$), the limb of the solar disk in the 2D-convolved map is more smoothed than the observed solar map, and hence the solar signal requires a horned structure in the limb of the solar disk to compensate




Fig. 3. Example of modelled equatorial T_B profile of a solar map obtained adopting the CCBH-model as the solar signal.

for this limb smoothing in the 2D-convolved map. Thus, we empirically introduce this feature through the combination of a circular-based cylindrical box and a horned function, called the CCBH-model (Fig. 3), defined as:

$$f_M(x, y, R, T_c, l, n) = \begin{cases} U_M & \text{if } U_M \leq T_c + l \\ 0 & \text{otherwise} \end{cases}, \quad (1)$$

where R indicates the box radius, T_c , indicates the brightness temperature, T_B , of the centre of the solar disk (quiet-Sun level), and l is the limb height (with respect to T_c); U_M is the horned function, defined as:

$$U_M(x, y, R, T_c, l) = T_c + |[b(U_C - x_C)^\zeta]|, \quad (2)$$

where $b = l[(R - x_C)^{-\zeta}]$, and $\zeta = 2$ (fixed). Also, U_C indicates the circular base of the cylindrical box, defined as:

$$U_C(x, y) = \sqrt{(x - x_C)^2 + (y - y_C)^2}, \quad (3)$$

where x_C and y_C indicate the centre coordinates of the circle (fixed at 0, as defined for the solar maps at SRT and Medicina).

The CCBH-model (Eq. (1) and Fig. 3) allows us to account for the behaviour of the signal observed at the solar limb, where generally the limb brightening and the coronal emission are observed and smoothed by the characteristics of the antenna beam pattern.

To investigate possible instrumental biases affecting the “tails” of the solar disk, we tested the beam fitting procedure using both solar signal models (ECB-model; CCBH-model, Eq. (1)) to obtain the hypothetical beam pattern that would fit the data and to check whether it is physically plausible. In this context, we compared the modelled value of the amplitude A_2 of the secondary lobe with those obtained through GRASP (a_2 , Table 1), to find possible anomalies with respect to the standard GRASP beam model.

In Table 3 we report the intensity difference ΔA between the original GRASP lobe a_2 (reported in Table 1) and the modelled lobe A_2 (using both the ECB- and CCBH-models) of the hypothetical beam patterns needed to fit the observed tails in our images.

In the observing frequency range 18–26 GHz, the large values obtained for ΔA (ranging between 4.1 and 14.2 dBi) are incompatible with hypothetical opto-mechanical deformations

¹¹ <https://emcee.readthedocs.io/en/stable/>

Table 3. Values of the modelled secondary lobe A_2 of the hypothetical beam patterns needed to fit the observed tails in the range 18–26 GHz with *Grueff* and SRT radio telescopes, in the assumption of ECB/CCBH solar models.

Frequency (GHz)	Instrument	Model	R_c (arcsec)	R_{eq} (arcsec)	R_{pol} (arcsec)	A_2 (dBi)	ΔA (dBi)
18.3	<i>Grueff</i>	ECB	–	$991.7^{+6.1}_{-2.9}$	$987.6^{+3.1}_{-2.8}$	$58.8^{+1.0}_{-0.8}$	5.8
18.3	<i>Grueff</i>	CCBH	$982.7^{+2.0}_{-1.8}$	–	–	$60.6^{+0.6}_{-0.6}$	7.6
18.8	SRT	ECB	–	$980.3^{+2.7}_{-0.6}$	$981.8^{+3.5}_{-0.6}$	$65.5^{+0.2}_{-0.4}$	4.1
18.8	SRT	CCBH	$979.3^{+0.7}_{-0.8}$	–	–	$66.5^{+0.2}_{-0.3}$	5.1
24.7	SRT	ECB	–	$978.7^{+0.5}_{-0.4}$	$979.6^{+1.3}_{-1.8}$	$67.0^{+0.3}_{-0.4}$	6.2
24.7	SRT	CCBH	$977.6^{+1.4}_{-2.0}$	–	–	$67.4^{+0.1}_{-0.3}$	6.6
25.8	<i>Grueff</i>	ECB	–	$989.8^{+4.2}_{-2.9}$	$987.4^{+1.3}_{-1.5}$	$60.2^{+0.9}_{-1.0}$	12.2
25.8	<i>Grueff</i>	CCBH	$977.1^{+1.5}_{-1.3}$	–	–	$62.2^{+0.6}_{-0.4}$	14.2

Notes. We report also the difference between the modelled and the real values $\Delta A = A_2 - a_2$, and the values of the mean, equatorial, and polar radii of the Sun (R_c , R_{eq} , and R_{pol}).

both for *Grueff* and SRT radio telescopes. In fact, mean beam aberration effects at the observed elevations are expected $\lesssim 3$ dBi at worst (Egron et al. 2022), thus suggesting a genuinely solar origin of the tail signals, rather than instrumental biases.

3.2. The theoretical radio atmospheric emission from the quiet Sun

In addition to the empirical solar signal models described in Sect. 3.1, we considered existing theoretical atmospheric models that provide brightness profiles that can be compared with our observations after convolving them with our beam pattern functions. In particular, we adopted the atmospheric SSC model (Selhorst et al. 2005, 2019), which describes the full solar atmosphere from the photosphere to the coronal plasma up to an altitude $h_{SSC} = 39.3$ Mm above the solar surface. For our analysis we tried to describe the full solar atmosphere up to an altitude $h_{ext} \sim 2000$ Mm above the solar surface, following a prescription similar to that used by Zhang et al. (2022). Therefore, in this work we named this model “extended SSC model” (eSSC). The altitude h_{ext} corresponds to a distance from the centre of the Sun $D_{\odot} = 3.9 R_{\odot, opt}$ (~ 3700 arcsec), suitable for analysing our solar maps, characterised by a size of about 5000×5000 arcsec (Sect. 2). This model does not take into account the effects of the strong magnetic fields in active regions, the spicules, the special features observed at the polar regions (Selhorst et al. 2019), and the geometry of radio wave refraction within the solar corona (Smerd 1950). As clarified in Selhorst et al. (2005), the main purpose of this model is to reproduce the brightness temperature spectrum of the solar disk centre, in order to model the full quiet Sun atmosphere from the photosphere to the corona.

In the eSSC model, T_B (expressed as a function of the observing frequency) is derived from a model of the density and temperature of the solar atmosphere (basically composed of hydrogen), assuming the thermal bremsstrahlung as the emission mechanism, by numerically integrating the radiation transfer equation along the line of sight:

$$T_B(\nu) = \int T k_{\nu} e^{-\tau_{\nu}} ds \text{ K}, \quad (4)$$

where T is the physical temperature of the medium (in units of K), k_{ν} is the free-free absorption coefficient (or the opacity, in units of cm^{-1} , Zirin 1988; Simões et al. 2017), and $\tau_{\nu} = \int k_{\nu} ds$ is the optical depth. Equation (4) allows us to obtain theoretical brightness

profiles since the information about the position with respect to the solar centre is contained in ds , thanks to simple geometrical transformations. The T distribution, defined as a function of the altitude $h = R - R_{\odot}$ (in units of km), follows the SSC model up to h_{SSC} , with a constant extension at $T = 1.43 \times 10^6$ K (the last value of T in the SSC model) for the outer corona (up to h_{ext}), as there is no rich information for the temperature of the outer corona in the literature. Moreover, k_{ν} is expressed as:

$$k_{\nu} = 3.7 \times 10^8 T^{-1/2} N_e N_p \nu^{-3} g_{ff} (1 - e^{-h\nu/k_b T}) \text{ cm}^{-1}, \quad (5)$$

where N_e and N_p are, respectively, the density of electrons and protons in the solar atmosphere and outer corona, the term $(1 - e^{-h\nu/k_b T})$ is the correction for stimulated emission, where h and k_b are the Planck and Boltzmann constants, respectively, and g_{ff} is the Gaunt factor (van Hoof et al. 2014). The density distributions, as a function of h (in units of km), follow the SSC model up to h_{SSC} , with an extension for the outer corona (up to h_{ext}) as n times (with $n = 1.27$)¹² the density distribution described in a specific model (Saito et al. 1977), to have a continuous connection with the SSC. With the model described above, a brightness temperature spectrum, $T_B(\nu)$, for the solar disk centre can be obtained by numerically integrating Eq. (4) for a series of frequencies, as shown in Fig. 4. As reported by Selhorst et al. (2005), $T_B(\nu)$ calculated through the SSC model is in good accordance with the radio observations from 1.4 to 400 GHz.

Propagation effects (i.e. the refraction and scattering) were not included in the original SSC model. These effects are very significant at lower frequencies and are inversely proportional to the observing frequency. Thus, we adopted a specific power-law term $T'_B(\nu) = T_B(\nu)(1 - \alpha\nu^{\beta})$ – previously used in Zhang et al. (2022) to represent and model the attenuation of T_B due to propagation effects. In this term, $T'_B(\nu)$ and $T_B(\nu)$ are, respectively, the observed and the modelled brightness temperature spectra, and ν is the observing frequency (in units of Hz). The best-fit parameters for α and β are reported in Table 4 considering different data sets. All the quiet-Sun measurements reported in Fig. 4 are well modelled (Table 4), but the best-fit parameters of our modelling (α and β) are different with respect the results obtained by Zhang et al. (2022), probably caused by a wider data set. Moreover, our results indicate that the propagation effects are

¹² We used a different value of n with respect to that used in Zhang et al. (2022; $n = 1.25$) in order to have a perfect numerical connection between SSC and Saito models.




Fig. 4. Brightness temperature spectrum $T_B(\nu)$ for the solar disk centre obtained from the data of the INAF radio telescopes (averaged values $T_{B,av}$ from Medicina, Table 5), and of previous works: Zirin et al. (1991), Subramanian (2004), Ramesh et al. (2006), Mercier & Chambe (2015), Melnik et al. (2018), Kallunki & Tornikoski (2018), Sharma & Oberoi (2020), Zhang et al. (2022). The black continuous line represents the modelled brightness temperature of bremsstrahlung emission obtained by the SSC model (Selhorst et al. 2005, 2019; Zhang et al. 2022); the black dashed line represents the fitted flux accounting for propagation effects; the dark grey shaded area shows the errors in the fit parameters. Green points indicate the measurements obtained from the averaged solar maps of the *Grueff* Radio Telescope ($T_{B,av}$ in Table 5), according to the analysis of Pellizzoni et al. (2022) (P+22) and Mulas et al. (2022) (M+22). This figure is an updated version of the figure reported by Zhang et al. (2022).

already negligible in the range 18–26 GHz and then these effects are not important for the modelling of the coronal tail emission. As reported in Table 4, we point out that the parameters α and β are highly sensitive to the variation of $T_B(\nu)$.

To strengthen the evidence of the physical nature of the coronal emission in the tails of the T_B profiles (Sect. 3.1) and to constrain the temperature and density parameters of the atmospheric emission, we also compared the modelled and the observed T_B profiles along the equatorial and polar diameters of the quiet Sun during the minimum solar activity (2018–2020) and their variations with the observing frequency. The modelled T_B profiles are obtained through the SSC and the eSSC model (Eq. (4)), convolved with the beam pattern of the instrument for that observing frequency. The observed T_B profiles are obtained with the *Grueff* Radio Telescope through two averaged solar maps, one at 18.3 GHz and the other at 25.8 GHz (green dotted and blue dashed lines in Fig. 5). These averaged solar maps have been created using only the medium- and high-quality maps (to avoid undesirable systematic and/or meteorological effects) during the minimum phase of the solar cycle, to minimise the effects of solar activity. We selected these observing frequencies and this radio telescope because we have a moderately uniform time coverage of solar observations from 2018 up to the present day.

From Fig. 5 and Table 5, we can compare the theoretical T_B profiles with the observed ones using the averaged solar maps of the *Grueff* Radio Telescope. In particular, we observe these features: (1) the level of the averaged tails of the T_B profiles (shown as green dot-dashed and blue dotted lines for the equatorial and the polar profiles, respectively) is higher than those predicted by the eSSC model (shown as a red solid line) by a factor 4.5 at 18.3 GHz, and by a factor 13 at 25.8 GHz, suggesting that the

Table 4. Best-fit parameters obtained by the modelling of the power-law term that describes the attenuation of T_B due to propagation effects.

Data	Model	α	β	χ_r^2
LOFAR	eSSC	$(2.92 \pm 0.16) \times 10^2$	$-(0.35 \pm 0.02)$	0.4
LOFAR/SunDish	eSSC	$(3.87 \pm 0.20) \times 10^2$	$-(0.37 \pm 0.02)$	0.6
All	eSSC	$(4.19 \pm 0.19) \times 10^2$	$-(0.37 \pm 0.02)$	1.1

difference between the model and observation increases with the increase in observing frequency; (2) the strong limb brightening (about 10%) predicted by the theory does not emerge from our averaged limbs in the T_B profiles; (3) the T_B values (at the central position of the solar disk) obtained through the averaged solar maps are slightly higher than those obtained in the modelled T_B profiles using the eSSC model, especially at 25.8 GHz, where the discrepancy is $\sim 6\%$ (Table 5). The same behaviour in the comparison between the theoretical model and the averaged T_B profiles occurs also when the SSC model is considered.

These features suggest that (1) the SSC/eSSC model could be too simplified to describe the radio emission from the quiet Sun, spanning from the photosphere to the upper corona (neglecting effects such as gyro-magnetic emission and propagation effects, as mentioned below in this section) and/or (2) the T and N distributions of the SSC/eSSC model could differ from what we are able to derive from our observations. In this context, we performed a modelling of our observed T_B profiles (both equatorial and polar) for each observing frequency (Figs. 6–8). Our brightness profiles cannot univocally constrain the actual physical parameters required for the fit, but as a first approach, we can check the effects of the variation of density and temperature separately. A possible fitting solution for the N_e and N_p distributions can be obtained, while keeping the T distribution constant according to the SSC/eSSC model; conversely, a valid T distribution can be obtained while maintaining constant the original density distributions. The best-fit results are obtained in iterative mode, as long as the value of T (or N) allows us to model $T_B(\nu)$ of Eq. (4), for each solar coordinate of the observed T_B profiles. The computational error connected with the calculation of the integral to obtain $T_B(\nu)$ (Eq. (4)) is conservatively fixed at 1%. In the fitting procedure, we also carefully take into account the errors related to our brightness temperature measurements: uncertainties due to the image RMS (2.5 K at 18.3 GHz and 3.5 K at 25.8 GHz); errors (2.5%) associated with the calibration procedure adopting the supernova remnant Cas A as calibrator source (Mulas et al. 2022; Pellizzoni et al. 2022); effects caused by the background baseline subtraction during the imaging procedure (5 K at 18.3 GHz and 15 K at 25.8 GHz).

By accounting for these uncertainties, we are able to analyse the solar atmosphere up to a specific T_B threshold (10 K at 18.3 GHz and 20 K at 25.8 GHz). This implies, for example, that we can observe the corona with the *Grueff* Radio Telescope up to (1) 2400 arcsec ($\sim 2.5 R_\odot$), with respect to the Sun’s centre at 18.3 GHz, and (2) 2600 arcsec ($\sim 2.7 R_\odot$), with respect to the Sun’s centre at 25.8 GHz. However, due to the high uncertainties starting from $\sim 1.8 R_\odot$ (see Fig. 8), we consider reliable only the data up to 2050 arcsec ($2.15 R_\odot$).

4. Discussion

As described in Sect. 3.1, the pronounced tail signal in the observed brightness profiles shows features that are




Fig. 5. Averaged and modelled T_B profiles obtained at the observing frequencies of 18.3 (left) and 25.8 GHz (right) with the *Grueff* Radio Telescope. These plots differ in the T_B y-axis scale (linear on top, logarithmic on bottom). Red solid line indicates the modelled T_B profile, obtained from the convolution between the eSSC model (Eq. (4)) and the beam pattern of the receiver for each observing frequency. Green dot-dashed and blue dotted lines indicate the equatorial and polar profiles of the averaged solar map, respectively.

Table 5. Measured T_B at the central position of the Sun – obtained at 18.3 and 25.8 GHz with the *Grueff* Radio Telescope.

Frequency (GHz)	$T_{B,qS}$ (K)	$T_{B,av}$ (K)	$T_{B,eSSC}$ (K)
18.3	$10\,130 \pm 253$	$10\,159 \pm 254$	9957 ± 100
25.8	9755 ± 244	9848 ± 246	9270 ± 93

Notes. T_B was obtained from (1) the modelled quiet-Sun level ($T_{B,qS}$) according to the Cas A calibrator source (Mulas et al. 2022; Pellizzoni et al. 2022), (2) the averaged solar maps ($T_{B,av}$), and (3) the modelled value of the eSSC model, convolved with the beam pattern of the instrument ($T_{B,eSSC}$).

incompatible with hypothetical opto-mechanical deformations both for *Grueff* and SRT radio telescopes, suggesting a genuinely solar origin of the tail signals, rather than instrumental biases. Furthermore, the comparison between the observed brightness profiles (averaged maps) and those obtained from the theory (SSC and eSSC models) shows that the modelled brightness profiles (Eq. (4)) differ from those observed (Sect. 3.2). Also, on this occasion, these observed profiles show that the level of the averaged tails (and especially that of the limb) is higher than what is predicted by theory (Fig. 5). The discrepancies between the theoretical models (SSC and eSSC) and the observed brightness profiles (Fig. 5) led us to further investigate the behaviour of T ,

N_e , and N_p in order to properly fit the data according to the prescriptions and limitations described in Sect. 3.2. The obtained best-fit density and temperature distributions (coherent with our data) are shown in Figs. 6 and 7. In each of these figures, the right panel indicates the differences, ΔT (for the temperature) and ΔN (for the density), between the best-fit distributions and the distributions of the theoretical models (SSC and eSSC). Thanks to our analysis, the reconstructed (and modified) T and N distributions of the SSC and eSSC models allow us to fit with a good agreement both the limbs and the coronal tails in our averaged T_B profiles.

As we can see in Figs. 6 and 7 (left panel), we observe that the modelled T and N distributions are independent, within a 1σ error, of the observing frequency. Moreover, the distributions reconstructed from the equatorial profile are compatible with the polar counterpart within 1σ , over the entire range of altitudes and for each observing frequency.

Adopting the SSC model, whose maximum altitude, h_{SSC} , is indicated by the blue vertical line in Figs. 6 and 7, the reconstructed T (Fig. 6, left panel) and N (Fig. 7, left panel) distributions are compatible within 1σ with this model. In particular, the reconstructed n distributions (Fig. 7, left panel) are in good accordance with the literature in the radio domain (e.g. Saint-Hilaire et al. 2012; Ramesh et al. 2020). A coronal analysis of the Sun (Chiuderi Drago et al. 1999) – assuming hydrostatic equilibrium and using EUV and radio data – shows an electron density derived at the base of the corona ($h \lesssim$




Fig. 6. Distribution of T (left) and ΔT (right) as a function of the altitude from the solar surface, defined as the photosphere level $R_{\odot, \text{opt}}$. The respective shaded area indicates the 1σ error of these distributions. The black line indicates the theoretical T distribution of the original eSSC model. The blue vertical line indicates the altitude, h_{SSC} , below which the SSC model is defined. The y-axis of these plots is shown in logarithmic scale. The best-fit T distribution is compatible (within an uncertainty $\leq 25\%$) with the original eSSC model below ~ 100 Mm. Above this altitude, this distribution is characterised by strong degeneracy.




Fig. 7. Distribution of N (electrons N_e and protons N_p , left) and ΔN (right) as a function of the altitude from the solar surface, defined as the photosphere level $R_{\odot, \text{opt}}$. The respective shaded area indicates the 1σ error of these distributions. The black line indicates the theoretical N distributions of the original eSSC model. The blue vertical line indicates the altitude, h_{SSC} , below which the SSC model is defined. The y-axis of these plots is shown in logarithmic scale. The best-fit n distributions are compatible (within an uncertainty $\leq 25\%$) with the original eSSC model below ~ 100 Mm; above this altitude, these distributions are characterised by high uncertainty (up to 70%).

35 Mm) $N_e(0) \approx 3 \times 10^8 \text{ cm}^{-3}$, in good accordance (within 3σ) with our results ($N_e(0) \approx (2.2 \pm 0.4) \times 10^8 \text{ cm}^{-3}$) and the SSC model. As shown in Fig. 6 (left panel), the T distribution reaches a maximum of $(1.5 \pm 0.2) \times 10^6 \text{ K}$ at h_{SSC} , compatible with the finding of other authors in the radio domain (e.g. Newkirk 1961; Gary & Hurford 1987; Mann et al. 1999; Saint-Hilaire et al. 2012). On the other hand, this value is lower, but still compatible within 5σ – than $2.2 \times 10^6 \text{ K}$ obtained from other authors in the MHz range (Mercier & Chambe 2015; Vocks et al. 2018). In the EUV domain, the T distribution in the coronal region falls below 10^6 K when coronal holes are observed in this frequency domain (e.g. Dulk et al. 1977; David et al. 1998; Fludra et al. 1997, 1999).

Using the eSSC model in the analysis, our data can be fitted (compatibility within 1σ) up to (1) ~ 60 Mm of altitude ($1.09 R_{\odot}$) for T , and (2) ~ 960 Mm of altitude ($2.4 R_{\odot}$) for N . The application of Eq. (4) (with only T as free parameter to model the

pronounced tails in our T_B profiles) produces strong degeneracy and high uncertainties for altitudes higher than $\gtrsim 60$ Mm (see Fig. 6, right panel). This suggests that the estimation of the T distribution using the eSSC model is not accurate above 60 Mm. The modelling using N (N_e and N_p) as free parameter produces good results up to ~ 100 Mm ($1.14 R_{\odot}$), where the uncertainty is $< 25\%$ (Fig. 7). Above this altitude, the modelled N distribution decreases over altitude at a much slower rate than the distribution defined by the eSSC model, reaching an uncertainty of $\sim 70\%$ on the measurement of N at ~ 800 Mm ($2.15 R_{\odot}$). Although the excess of our N extrapolations at high altitudes with the eSSC model is characterised by high uncertainty, in this range of altitudes the N distributions are compatible within 1σ level with the measurements obtained between March 22-25, 2022 by the Metis coronagraph (Antonucci et al. 2020; Fineschi et al. 2020), aboard the Solar Orbiter ESA-NASA observatory (e.g. Müller et al. 2020), as shown in Fig. 8. The latter values of N_e




Fig. 8. Distribution of N (left) and ΔN (right) as a function of the altitude from the solar surface. Green and red lines indicate the measured N_e obtained through observations with *Grueff* at 18.3 GHz (green) and 25.8 GHz (red) during a period of minimum solar activity (2018–2020), respectively (see the legend of the right panel in Fig. 7). The shaded area indicates the 1σ error of the distributions. Blue lines indicate the measured N_e obtained through observations with Metis (dashed line: equatorial component; dotted line: polar component) during the ascending phase of the solar cycle (March 2022). The black line indicates the theoretical N_e associated with the original eSSC model. The y-axis of these plots is shown in logarithmic scale. In the right panel, the difference between ΔN_e referred to Metis data is reported in terms of absolute value.

were estimated by applying the inversion procedure described in van de Hulst (1950) to the Metis visible light images of level 2 (L2, data release ‘V01’), as described in De Leo et al. (2023). We note that this comparison is affected by the fact that (1) Metis and radio observations have not simultaneous thus far (coronal density depends on the solar activity) and (2) N_e values inferred from Metis are obtained through a different approach.

The discrepancies of our reconstructed N distributions (right panel of Figs. 7 and 8), especially in the chromospheric and the upper coronal regions, could explain the tails observed in our T_B profiles, more pronounced than those predicted by the theory (Fig. 5). We pointed out that above h_{SSC} (1) the N distributions are characterised by an ever greater uncertainty (up to $\sim 70\%$) and by a slope lower than the distribution defined by the eSSC model (Fig. 8) and (2) the T distribution is characterised by high uncertainty. This discrepancy could be ascribed to the simple assumption of pure free-free emission in LTE conditions in the eSSC model, which allows us to model the solar atmosphere up to ~ 2000 Mm of altitude. This model might lead to an overestimate of T and N under the hypothesis of the likely presence of additional emission components (e.g. gyro-magnetic emission; White & Kundu 1997) or other effects (e.g. weak plasma emission, spicules, and refraction effects), neglected in the SSC/eSSC model (Sect. 3.2). Moreover, this discrepancy could be explained by the fact that the worse angular resolution at low frequencies than the higher frequencies (e.g. visible, EUV, X-rays) overestimates the radio T_B , resulting in a higher value of N in the corona (e.g. Chiuderi Drago et al. 1999). The assessment of this discrepancy should be addressed by multi-frequency campaigns – such as simultaneous observations with SunDish telescope in radio and various currently operating visible light coronagraphs such as Metis/Solar Orbiter, LASCO/SOHO (Brueckner et al. 1995), and COR2/STEREO-A (Howard et al. 2008) – to properly account for the expected coronal variability. This campaign is crucial to improve the modelling of the solar atmosphere and to better understand the behaviour of the T and N distributions. Furthermore, to disentangle these discrepancies and possible polar and equatorial anisotropies of T_B (and then of the density) of the coronal radio tails, deeper exposures would be

required up to two degrees from the solar centroid to improve sensitivity.

In addition to the eSSC model, other literature models have been used to try to describe the quiet solar atmosphere and its radio emission at different frequencies, such as the 3D numerical code known as PAKAL (De La Luz et al. 2008, 2010) and its updated version PAKALMPI (De la Luz et al. 2011). This model includes distributions for T (extended up to 10^5 Mm) and N (extended up to 10^3 Mm) slightly different than those both included in the eSSC model and in our results. In particular, the T distribution of PAKAL is compatible with our T distribution up to h_{SSC} . For altitudes higher than h_{SSC} , the PAKAL distribution reaches a maximum value of $\sim 2 \times 10^6$ K at ~ 300 Mm, where our T distribution shows strong degeneracy. The N distribution of PAKAL is compatible with our N distribution up to ~ 100 Mm. Above this altitude, our N distribution decreases over altitude at a much slower rate than the distribution defined with PAKAL, reaching a difference in density of about one order of magnitude at ~ 300 Mm.

According to the discrepancy of the T_B profiles in the limb regions, a preliminary modelling process in the radio domain (18–26 GHz) using our solar maps convolved with the empirical CCBH model (Sect. 3.1) appears to show the presence of faint limb brightening (whose level ranges between 3% and 7%); this is in agreement with other works (e.g. Selhorst et al. 2003). As stated by several authors (e.g. Selhorst et al. 2005; Menezes et al. 2022), a high limb brightening observed in the solar disk can produce an overestimation in the determination of R_\odot , resulting in an underestimation of the limb brightening itself. This underestimation could be also due to the presence of chromospheric features such as spicules located close to the limb, which are usually not included in the theoretical models (Selhorst et al. 2019). Moreover, since the SSC/eSSC model has an ideal assumption of straight light of sight integral (Eq. (4)) – true for close-to-center regions – refractive effects (e.g. Smerd 1950; Alissandrakis 1994; Vocks et al. 2018) could deviate from a straight line the radio wave propagation (especially at MHz-level, see Fig. 4) in the limb regions, causing the appearance of the limb brightening in the model. A deeper analysis of this

phenomenon is beyond the scope of this paper and it will be the subject of a future work.

Finally, it is worth noting that the analysis of R_{\odot} via our solar maps (Marongiu et al. 2024) showed that the measurements of R_{\odot} (obtained with a specific 2D model¹³) range between the values predicted by the SSC/eSSC models and the most widely used methods for estimating the radio R_{\odot} in the literature (e.g. Costa et al. 1999; Selhorst et al. 2011; Emilio et al. 2012; Menezes & Valio 2017; Alissandrakis et al. 2017; Menezes et al. 2021). This aspect suggests that the methods aimed at the measurements of R_{\odot} , when they are improved by the addition of specific features of the solar disk and the atmosphere (such as the coronal emission level and the limb brightening), allowing us to obtain values of R_{\odot} that are closer to the canonical value $R_{\odot, \text{opt}}$ and more accurate.

5. Summary

In this paper, we focus on the analysis of the brightness profiles of solar images obtained in the frame of the SunDish project. We used 290 single-dish observations over five years – from 2018 to mid-2023 – corresponding to about half solar cycle in the radio K -band (18–26 GHz). For our analysis, we also used two averaged solar maps at 18.3 and 25.8 GHz acquired at Medicina during the minimum solar activity (2018–2020). We analysed the weak tailing emission seen outside the solar disk to probe its physical origin. In this context, we adopted a Bayesian approach to study the role of the antenna beam pattern on our solar maps, through the modelling of the solar signal (the observed brightness profiles) with a horned 2D function (CCBH model) convolved with the beam pattern. Moreover, we constrained the T and N distributions of the atmospheric layers through fitting procedures of the observed brightness profiles, adopting the SSC model and its extended version (eSSC). The results of our analysis are summarised as follows:

- We probed significant emission of physical origin of the weak tailing emission seen outside the solar disk, taking into account all possible biases related to instrumental effects. In particular, analysing the role of the antenna beam pattern on our solar maps allowed us to prove the physical origin of the coronal tail emission in our images since no realistic beam pattern can produce these tails as artefacts. In this context, our images are important not only to improve the existing solar atmospheric models, but also to probe directly different high-lying atmospheric layers of the corona.
- The comparison between the observed brightness profiles (averaged maps) and those obtained from the theory (SSC and eSSC models) shows that the modelled brightness profiles differ from those observed in the coronal tail (where the observation is more pronounced than the model) and especially in the limb of the solar disk (where the model is more pronounced than the observation).
- In particular, the SSC model is in good agreement with our solar maps up to an altitude, h_{SSC} , essentially corresponding to the base of the corona. Above an altitude of ~ 60 Mm for T and ~ 100 Mm for N , our results obtained with the eSSC model reveal a discrepancy with respect to the original T and N distributions, especially in the upper chromospheric and coronal regions. These discrepancies could be also ascribed to (1) the simple assumption of pure free-free emission in

LTE conditions (especially in the eSSC model) and the lack of neglected effects in the eSSC model (e.g. gyro-magnetic emission), along with (2) the worse angular resolution at our radio frequencies than the higher frequencies (e.g. visible, EUV, X-rays), which overestimates N in the corona (e.g. Chiuderi Drago et al. 1999).

Future observations and detailed theoretical analysis with *Grueff* and SRT for longer periods of time and with a multi-frequency approach (also thanks to the new PON receivers operating up to 116 GHz) are crucial for probing different layers of the solar atmosphere over time. Furthermore, simultaneous radio-visible light observations (e.g. with SunDish and Metis) and multi-frequency observations are necessary to properly account for the expected coronal variability, as well as to better understand the behaviour of the N distributions, while also using the same method to extrapolate the density. A detailed theoretical analysis of the coronal model, with the analysis of the distributions of the solar medium in the context of the thermal bremsstrahlung emission mechanism plus gyro-magnetic components, is beyond the scope of this paper and will be the subject of a future work. In addition, these observations are needed to better clarify several aspects, such as (1) the correlation between solar activity and the size of the Sun, (2) the polar and equatorial trends of the solar atmosphere, and (3) the question of limb brightening (especially the presence of the polar limb brightening during the solar minima). A preliminary investigation of the possible presence of limb brightening in our solar maps seems to show the presence of faint limb brightening (whose level ranges between 3% and 7%), in agreement with other works (e.g. Selhorst et al. 2003), but this kind of investigation is beyond the scope of this paper and will be the subject of a future analysis.

Acknowledgements. We thank the anonymous referee for helping us improve the paper. The Medicina radio telescope is funded by the Ministry of University and Research (MUR) and is operated as National Facility by the National Institute for Astrophysics (INAF). The Sardinia Radio Telescope is funded by the Ministry of University and Research (MUR), Italian Space Agency (ASI), and the Autonomous Region of Sardinia (RAS) and is operated as National Facility by the National Institute for Astrophysics (INAF). The Enhancement of the Sardinia Radio Telescope (SRT) for the study of the Universe at high radio frequencies is financially supported by the National Operative Program (Programma Operativo Nazionale – PON) of the Italian Ministry of University and Research “Research and Innovation 2014-2020”, Notice D.D. 424 of 28/02/2018 for the granting of funding aimed at strengthening research infrastructures, in implementation of the Action II.1 – Project Proposal PIR01_00010. Solar Orbiter is a space mission of international collaboration between ESA and NASA, operated by ESA. Metis was built and operated with funding from the Italian Space Agency (ASI), under contracts to the National Institute of Astrophysics (INAF) and industrial partners. Metis was built with hardware contributions from Germany (Bundesministerium für Wirtschaft und Energie through DLR), from the Czech Republic (PRODEX) and from ESA. We acknowledge the Computing Centre at INAF – Istituto di Radioastronomia for providing resources and staff support during the processing of solar data presented in this paper. This work was supported by the Istituto Nazionale di Astrofisica (Bando per il finanziamento della Ricerca Fondamentale 2023 – IDEA-SW project).

References

- Alissandrakis, C. E. 1994, *Adv. Space Res.*, 14, 81
 Alissandrakis, C. E., Patsourakos, S., Nindos, A., & Bastian, T. S. 2017, *A&A*, 605, A78
 Antonucci, E., Romoli, M., Andretta, V., et al. 2020, *A&A*, 642, A10
 Bolli, P., Orlati, A., Stringhetti, L., et al. 2015, *J. Astron. Instrum.*, 04, 1550008
 Brueckner, G. E., Howard, R. A., Koomen, M. J., et al. 1995, *Sol. Phys.*, 162, 357
 Chiuderi Drago, F., Landi, E., Fludra, A., & Kerdran, A. 1999, *A&A*, 348, 261
 Costa, J. E. R., Silva, A. V. R., Makhmutov, V. S., et al. 1999, *ApJ*, 520, L63
 David, C., Gabriel, A. H., Bely-Dubau, F., et al. 1998, *A&A*, 336, L90
 De La Luz, V., Lara, A., Mendoza, E., & Shimojo, M. 2008, *Geofis. Int.*, 47, 197

¹³ In this model, tailored to empirically describe both the solar disk and the atmospheric emission, the solar signal is defined as the combination between the ECB-model and a 2D-Gaussian function (2GECB-model).

- De la Luz, V., Lara, A., Mendoza-Torres, J. E., & Selhorst, C. L. 2010, *ApJS*, **188**, 437
- De la Luz, V., Lara, A., & Raulin, J.-P. 2011, *ApJ*, **737**, 1
- De Leo, Y., Burtovoi, A., Teriaca, L., et al. 2023, *A&A*, **676**, A45
- Dulk, G. A., Sheridan, K. V., Smerd, S. F., & Withbroe, G. L. 1977, *Sol. Phys.*, **52**, 349
- Egron, E., Pellizzoni, A., Iacolina, M. N., et al. 2017, *MNRAS*, **470**, 1329
- Egron, E., Vacca, V., Carboni, G., et al. 2022, SRT performance measurements (2018-2021), Tech. Rep. 178, OA@INAF
- Emilio, M., Kuhn, J. R., Bush, R. I., & Scholl, I. F. 2012, *ApJ*, **750**, 135
- Fineschi, S., Naletto, G., Romoli, M., et al. 2020, *Exp. Astron.*, **49**, 239
- Fludra, A., Brekke, P., Harrison, R. A., et al. 1997, *Sol. Phys.*, **175**, 487
- Fludra, A., Del Zanna, G., Alexander, D., & Bromage, B. J. I. 1999, *J. Geophys. Res.*, **104**, 9709
- Fontenla, J. M., Avrett, E. H., & Loeser, R. 1990, *ApJ*, **355**, 700
- Fontenla, J. M., Avrett, E. H., & Loeser, R. 1991, *ApJ*, **377**, 712
- Fontenla, J. M., Avrett, E. H., & Loeser, R. 1993, *ApJ*, **406**, 319
- Fontenla, J. M., Avrett, E. H., & Loeser, R. 2002, *ApJ*, **572**, 636
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, **125**, 306
- Gary, D. E., & Hurford, G. J. 1987, *ApJ*, **317**, 522
- Giménez de Castro, C. G., Pereira, A. L. G., Valle Silva, J. F., et al. 2020, *ApJ*, **902**, 136
- Govoni, F., Bolli, P., Buffa, F., et al. 2021, in *2021 XXXIVth General Assembly and Scientific Symposium of the International Union of Radio Science (URSI GASS)*, 1-4
- Haberreiter, M., Schmutz, W., & Kosovichev, A. G. 2008, *ApJ*, **675**, L53
- Howard, R. A., Moses, J. D., Vourlidas, A., et al. 2008, *Space Sci. Rev.*, **136**, 67
- Iwai, K., Shimojo, M., Asayama, S., et al. 2017, *Sol. Phys.*, **292**, 22
- Kallunki, J., & Tornikoski, M. 2018, *Sol. Phys.*, **293**, 156
- Krissinel', B. B. 2005, *Astron. Rep.*, **49**, 939
- Kundu, M. R., Gergely, T. E., Turner, P. J., & Howard, R. A. 1983, *ApJ*, **269**, L67
- Lallement, R., Holzer, T. E., & Munro, R. H. 1986, *J. Geophys. Res.*, **91**, 6751
- Loru, S., Pellizzoni, A., Egron, E., et al. 2019, *MNRAS*, **482**, 3857
- Loru, S., Pellizzoni, A., Egron, E., et al. 2021, *MNRAS*, **500**, 5177
- Mamajek, E. E., Prsa, A., Torres, G., et al. 2015, arXiv e-prints [arXiv:1510.07674]
- Mann, G., Jansen, F., MacDowall, R. J., Kaiser, M. L., & Stone, R. G. 1999, *A&A*, **348**, 614
- Marongiu, M., Pellizzoni, A., Egron, E., et al. 2020, *Exp. Astron.*, **49**, 159
- Marongiu, M., Pellizzoni, A., Bachetti, M., et al. 2022, A dedicated pipeline to analyse solar data with INAF Radio Telescopes: SUNPIT (SUNdish Pipeline Tool), Tech. Rep. 137, OA@INAF
- Marongiu, M., Pellizzoni, A., Mulas, S., et al. 2024, *A&A*, **684**, A122 (Paper I)
- Melis, A., Concu, R., Trois, A., et al. 2018, *J. Astron. Instrum.*, **7**, 1850004
- Melnik, V. N., Shepelev, V. A., Poedts, S., et al. 2018, *Sol. Phys.*, **293**, 97
- Menezes, F., & Valio, A. 2017, *Sol. Phys.*, **292**, 195
- Menezes, F., Selhorst, C. L., Giménez de Castro, C. G., & Valio, A. 2021, *ApJ*, **910**, 77
- Menezes, F., Selhorst, C. L., Giménez de Castro, C. G., & Valio, A. 2022, *MNRAS*, **511**, 877
- Mercier, C., & Chambe, G. 2015, *A&A*, **583**, A101
- Meyer-Vernet, N. 2012, *Basics of the Solar Wind* (UK: Cambridge University Press)
- Mulas, S., Pellizzoni, A., Iacolina, M. N., et al. 2022, A New Method for Accurate Calibration of Solar Disk Emission in the Radio Band, Tech. Rep.
- Müller, D., St. Cyr, O. C., Zouganelis, I., et al. 2020, *A&A*, **642**, A1
- Nakajima, H., Nishio, M., Enome, S., et al. 1994, *IEEE Proc.*, **82**, 705
- Nakajima, H., Nishio, M., Enome, S., et al. 1995, *JApAS*, **16**, 437
- Newkirk, G. J., 1961, *ApJ*, **133**, 983
- Pellizzoni, A., Righini, S., Murtas, G., et al. 2019, *Nuovo Cimento C Geophys. Space Phys. C*, **42**, 9
- Pellizzoni, A., Righini, S., Iacolina, M. N., et al. 2022, *Sol. Phys.*, **297**, 86
- Plainaki, C., Antonucci, M., Bemporad, A., et al. 2020, *J. Space Weather Space Clim.*, **10**, 6
- Prandoni, I., Murgia, M., Tarchi, A., et al. 2017, *A&A*, **608**, A40
- Prša, A., Harmanec, P., Torres, G., et al. 2016, *AJ*, **152**, 41
- Ramesh, R., Subramanian, K. R., Sundararajan, M. S., & Sastry, C. V. 1998, *Sol. Phys.*, **181**, 439
- Ramesh, R., Nataraj, H. S., Kathiravan, C., & Sastry, C. V. 2006, *ApJ*, **648**, 707
- Ramesh, R., Kathiravan, C., Barve, I. V., Beeharry, G. K., & Rajasekara, G. N. 2010, *ApJ*, **719**, L41
- Ramesh, R., Kumari, A., Kathiravan, C., et al. 2020, *Geophys. Rev. Lett.*, **47**, e90426
- Saint-Hilaire, P., Hurford, G. J., Keating, G., Bower, G. C., & Gutierrez-Kraybill, C. 2012, *Sol. Phys.*, **277**, 431
- Saito, K., Poland, A. I., & Munro, R. H. 1977, *Sol. Phys.*, **55**, 121
- Selhorst, C. L., Silva, A. V. R., Costa, J. E. R., & Shibasaki, K. 2003, *A&A*, **401**, 1143
- Selhorst, C. L., Silva, A. V. R., & Costa, J. E. R. 2005, *A&A*, **433**, 365
- Selhorst, C. L., Silva-Válio, A., & Costa, J. E. R. 2008, *A&A*, **488**, 1079
- Selhorst, C. L., Giménez de Castro, C. G., Valio, A., Costa, J. E. R., & Shibasaki, K. 2011, *ApJ*, **734**, 64
- Selhorst, C. L., Simões, P. J. A., Brajša, R., et al. 2019, *ApJ*, **871**, 45
- Sharma, S. 2017, *ARA&A*, **55**, 213
- Sharma, R., & Oberoi, D. 2020, *ApJ*, **903**, 126
- Shibasaki, K., Alissandrakis, C. E., & Pohjolainen, S. 2011, *Sol. Phys.*, **273**, 309
- Simões, P. J. A., Kerr, G. S., Fletcher, L., et al. 2017, *A&A*, **605**, A125
- Smerd, S. F. 1950, *Aust. J. Sci. Res. A Phys. Sci.*, **3**, 34
- Subramanian, K. R. 2004, *A&A*, **426**, 329
- van de Hulst, H. C. 1950, *Bull. Astron. Inst. Neth.*, **11**, 135
- van Hoof, P. A. M., Williams, R. J. R., Volk, K., et al. 2014, *MNRAS*, **444**, 420
- Vernazza, J. E., Avrett, E. H., & Loeser, R. 1973, *ApJ*, **184**, 605
- Vernazza, J. E., Avrett, E. H., & Loeser, R. 1976, *ApJS*, **30**, 1
- Vernazza, J. E., Avrett, E. H., & Loeser, R. 1981, *ApJS*, **45**, 635
- Vocks, C., Mann, G., Breiting, F., et al. 2018, *A&A*, **614**, A54
- Wang, Z., Schmahl, E. J., & Kundu, M. R. 1987, *Sol. Phys.*, **111**, 419
- White, S. M., & Kundu, M. R. 1997, *Sol. Phys.*, **174**, 31
- Wild, J. P. 1970, *PASA*, **1**, 365
- Zhang, P., Zucca, P., Kozarev, K., et al. 2022, *ApJ*, **932**, 17
- Zirin, H. 1988, *Astrophysics of the Sun* (Cambridge: University Press)
- Zirin, H., Baumert, B. M., & Hurford, G. J. 1991, *ApJ*, **370**, 779