



<b>Publication Year</b>	2020
<b>Acceptance in OA</b>	2021-08-27T10:10:11Z
<b>Title</b>	A 3D view of the Taurus star-forming region by Gaia and Herschel: multiple populations related to the filamentary molecular cloud
<b>Authors</b>	Roccatagliata, V., FRANCIOSINI, Elena, SACCO, GIUSEPPE GERMANO, RANDICH, Maria Sofia, Sicilia-Aguilar, A.
<b>Publisher's version (DOI)</b>	10.1051/0004-6361/201936401
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/30981">http://hdl.handle.net/20.500.12386/30981</a>
<b>Journal</b>	ASTRONOMY & ASTROPHYSICS
<b>Volume</b>	638

# A 3D view of the Taurus star-forming region by *Gaia* and *Herschel*

## Multiple populations related to the filamentary molecular cloud<sup>★,★★</sup>

V. Roccatagliata<sup>1,2,3</sup>, E. Franciosini<sup>2</sup>, G. G. Sacco<sup>2</sup>, S. Randich<sup>2</sup>, and A. Sicilia-Aguilar<sup>4</sup>

<sup>1</sup> Dipartimento di Fisica “Enrico Fermi”, Università di Pisa, Largo Pontecorvo 3, 56127 Pisa, Italy  
e-mail: [veronica.roccatagliata@unipi.it](mailto:veronica.roccatagliata@unipi.it)

<sup>2</sup> INAF-Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy

<sup>3</sup> INFN, Sezione di Pisa, Largo Bruno Pontecorvo 3, 56127 Pisa, Italy

<sup>4</sup> SUPA, School of Science and Engineering, University of Dundee, Nethergate, DD1 4HN, Dundee, UK

Received 29 July 2019 / Accepted 18 April 2020

### ABSTRACT

**Context.** Taurus represents an ideal region to study the three-dimensional distribution of the young stellar population and relate it to the associated molecular cloud.

**Aims.** The second *Gaia* data release (DR2) enables us to investigate the Taurus complex in three dimensions, starting from a previously defined robust membership. The molecular cloud structured in filaments can be traced in emission using the public far-infrared maps from *Herschel*.

**Methods.** From a compiled catalog of spectroscopically confirmed members, we analyze the 283 sources with reliable parallax and proper motions in the *Gaia* DR2 archive. We fit the distribution of parallaxes and proper motions with multiple populations described by multivariate Gaussians. We compute the cartesian Galactic coordinates ( $X, Y, Z$ ) and, for the populations associated with the main cloud, also the galactic space velocity ( $U, V, W$ ). We discuss the spatial distribution of the populations in relation to the structure of the filamentary molecular cloud traced by *Herschel*.

**Results.** We discover the presence of six populations which are all well defined in parallax and proper motions, with the only exception being Taurus D. The derived distances range between  $\sim 130$  and  $\sim 160$  pc. We do not find a unique relation between stellar population and the associated molecular cloud: while the stellar population seems to be on the cloud surface, both lying at similar distances, this is not the case when the molecular cloud is structured in filaments. Taurus B is probably moving in the direction of Taurus A, while Taurus E appears to be moving towards them.

**Conclusions.** The Taurus region is the result of a complex star formation history which most probably occurred in clumpy and filamentary structures that are evolving independently.

**Key words.** stars: pre-main sequence – open clusters and associations: individual: Taurus – parallaxes – proper motions – radio continuum: ISM

## 1. Introduction

During the early phases of the star and cluster formation process, filamentary structures have not only been theoretically predicted (e.g., Dale et al. 2012), but have also been detected in low- and high-mass star-forming regions, and in regions where no star formation is currently active (e.g., André et al. 2010). In particular, Myers (2009) highlights that near young stellar groups or clusters are associated with a “hub-filament structure”. Filaments traced commonly by extinction maps from infrared observations can now be traced in great detail in emission in the far-infrared thanks to *Herschel*. Within 300 pc we find examples of a main prominent filament in Chamaeleon I or Corona Australis (e.g., Schmalzl et al. 2010; Palmeirim et al. 2013; Sicilia-Aguilar et al. 2011, 2013); Lupus I is instead located along a filament at the converging location of two bubbles (e.g., Gaczkowski et al. 2015, 2017; Krause et al. 2018), while Serpens Main is at the converging point of filaments (e.g., Roccatagliata et al. 2015).

A step further in the study of filamentary molecular clouds is to relate their physical properties to young stellar populations. This can be done by combining the astrometric *Gaia* data with the *Herschel* maps of the molecular cloud in emission. The second release of *Gaia* is revolutionizing the study of star formation and young clusters under different aspects. A 3D density map for early-type and pre-main sequence sources was recently constructed by, for example, Zari et al. (2018), finding that younger stars within 500 pc to the Sun are mostly distributed in dense and compact clumps, while older sources are instead widely distributed. Several studies on large-scale star-forming regions are combining *Gaia* data with photometric surveys (e.g., Vela OB2, Armstrong et al. 2018; Cantat-Gaudin et al. 2019). New members have been found in nearby associations (e.g., Gagné & Faherty 2018), and even new nearby associations have been discovered (e.g., Gagné et al. 2018). Recently, multiple populations were also detected in young clusters (Roccatagliata et al. 2018; Franciosini et al. 2018).

One of the most studied young star forming regions, where both a prominent molecular cloud is structured in a main filament and a young population is present, is the Taurus complex. The discussion of star formation history in space and time started more than 15 yr ago. Palla & Stahler (2002) found a younger

\* Movie associated to Fig. A.1 is available at <http://www.aanda.org>

\*\* Full Table A.1 is only available at the CDS via anonymous ftp to [cdsarc.u-strasbg.fr](ftp://cdsarc.u-strasbg.fr) (130.79.128.5) or via <http://cdsarc.u-strasbg.fr/viz-bin/cat/J/A+A/638/A85>.


population inside the filaments and a more dispersed and older population outside them, and concluded that an age spread was present in the region. In the same year, overlaying the young stellar content to the  $^{12}\text{CO}$  maps, Hartmann (2002) found that most of the sources in “*Taurus Main*” follow three near parallel elongated bands, while only a few of them are localized in widely distributed groups. This led him to organize the sources into different groups according to their spatial distribution, without finding a significant age spread in the K7-M1 spectral type range. Continuum and line maps of the region with increasing spatial resolution have been obtained for Taurus in recent years. A large-scale survey of Taurus in  $^{12}\text{CO}$  and  $^{13}\text{CO}$  was presented by Goldsmith et al. (2008), showing a very complex and highly structured cloud morphology including filaments, cavities, and rings. Analyzing the  $^{13}\text{CO}$  integrated intensities in three different velocity intervals, these latter authors found the highest velocity ( $7\text{--}9\text{ km s}^{-1}$ ) corresponding to L1506 and L1498 and part of L1495, while the other clouds (B213, L1521, B18, L1536) have mainly intermediate velocities ( $5\text{--}7\text{ km s}^{-1}$ ) and only a few points at the borders of those clouds are found at even lower velocities ( $3\text{--}5\text{ km s}^{-1}$ ). A different orientation of the magnetic field was also found: the long filament of L1506 is parallel to the field, while the magnetic field is perpendicular to the extension of the long axis of the B216 and B217 filaments (Goodman et al. 1992).

The main filament in the B213-L1495 region has been extensively studied by Hacar et al. (2013) in high-density tracers such as  $\text{C}^{18}\text{O}$ ,  $\text{N}_2\text{H}^+$ , and  $\text{SO}$ , and by Hacar et al. (2016) in the three main isotopologues of  $^{12}\text{CO}$ ,  $^{13}\text{CO}$ , and  $\text{C}^{18}\text{O}$ . These latter authors suggested that the velocity broadening of the lines could be caused by the effects of different structures in 3D, which are clumpy or tangly on the line of sight projection of the velocity component. A first analysis of the *Gaia* DR2 data of Taurus recently presented by Luhman (2018) and Esplin & Luhman (2019) enabled these authors to revise the membership and to constrain the initial mass function of Taurus. Using the *Gaia* data, Luhman (2018) found that the population older than 10 Myr is not associated to Taurus. The goal of the present paper is to relate the star formation history in the Taurus complex to the filamentary structure of the molecular cloud. To this aim, the new analysis of the *Gaia* DR2 data starts from a reliable membership (presented in Sect. 2) and applies a rigorous approach which takes into account the covariances (not considered in the studies of Luhman 2018; Esplin & Luhman 2019) between parallax and proper motions (available in the *Gaia* Archive). The cluster kinematics and the relation between stellar content and filamentary structures are discussed in Sect. 4, together with the estimate of the distance of the molecular cloud. Our conclusions are presented in Sect. 5.


## 2. Cluster membership and *Gaia* DR2 data

Of the initial catalog of 518 spectroscopically confirmed members of Esplin & Luhman (2019), 443 sources are present in the *Gaia* DR2 archive. From this sample, we selected the sources with good astrometry following Lindegren (2018), who suggested the use of the renormalized unit weight error (RUWE), which is a more reliable and informative goodness-of-fit statistic than the astrometric excess noise commonly used until now by the community. The RUWE is defined as

$$\text{RUWE} = \frac{\sqrt{\chi^2/(n\text{Obs} - \nu)}}{f(G, G_{\text{BP}} - G_{\text{RP}})}, \quad (1)$$



**Fig. 1.** Histogram of the parallaxes of the Taurus members. The dashed red lines indicate the range in parallax commonly adopted in the literature.



**Fig. 2.** Proper motions in  $\alpha$  and  $\delta$  color coded by the parallax of the sources.

where  $n\text{Obs}$  is the number of observations,  $\nu$  is the number of parameters solved, and  $f$  is a renormalization function. Our further analysis is based on the 283 sources with RUWE of less than 1.4.

Figure 1 shows the distribution of the parallaxes of the cluster members. The resulting mean parallax is  $7.01 \pm 0.06$  mas, where the associated error is computed as the standard deviation ( $0.78$  mas) divided by  $\sqrt{N}$ . However, this mean parallax value lies exactly in the middle of a parallax distribution that appears to be multimodal. Furthermore, the distribution in proper motions in Fig. 2 shows that multiple populations might be present in the region. A deeper analysis of the *Gaia* data is therefore necessary to understand the star formation history of the region.

## 3. Analysis

Figures 1 and 2 clearly show that Taurus cannot be described by a single population because both parallaxes and proper motions are not randomly distributed around single mean values. Therefore, we model the distribution of the astrometric parameters using a similar approach to that used by Roccatagliata et al. (2018) to study the two sub-populations in Chamaeleon I. We assume that the Taurus star forming region is composed of multiple populations, each described by a 3D multivariate Gaussian

(as in [Lindegren et al. 2000](#))<sup>1</sup>. The  $j$ -th population is defined by seven parameters: the mean parallax ( $\pi_j$ ), the mean proper motions along right ascension and declination ( $\mu_{\alpha,j}$  and  $\mu_{\delta,j}$ ), the intrinsic dispersion of the parallax ( $\sigma_{\pi,j}$ ), the intrinsic dispersion of proper motions along right ascension and declination ( $\sigma_{\mu_{\alpha,j}}$  and  $\sigma_{\mu_{\delta,j}}$ ), and a normalization factor that takes into account the fraction of stars belonging to that population ( $f_j$ ). We assume that the mean proper motions and the mean parallaxes of each population are not correlated. To fit our distribution we use a maximum likelihood technique as in [Jeffries et al. \(2014\)](#) and [Franciosi et al. \(2018\)](#), adopting the SLSQP minimization method as implemented in `scipy`. The likelihood function of the  $i$ -th star for the  $j$ -th population is given by

$$L_{i,j} = (2\pi)^{-3/2} |C_{i,j}|^{-1/2} \times \exp\left[-\frac{1}{2}(a_i - a_j)' C_{i,j}^{-1} (a_i - a_j)\right], \quad (2)$$

where  $C_{i,j}$  is the covariance matrix,  $|C_{i,j}|$  its determinant, and  $(a_i - a_j)'$  is the transpose of the vector

$$a_i - a_j = \begin{bmatrix} \pi_i - \pi_j \\ \mu_{\alpha,i} - \mu_{\alpha,j} \\ \mu_{\delta,i} - \mu_{\delta,j} \end{bmatrix}. \quad (3)$$

Following [Lindegren et al. \(2000\)](#), the covariance matrix  $C_{i,j}$  is given by:

$$C_i = \begin{bmatrix} \sigma_{\pi,i}^2 + \sigma_{\pi,j}^2 & \sigma_{\pi,i} \sigma_{\mu_{\alpha,i}} \rho_i(\pi, \mu_{\alpha}) & \sigma_{\pi,i} \sigma_{\mu_{\delta,i}} \rho_i(\pi, \mu_{\delta}) \\ \sigma_{\pi,i} \sigma_{\mu_{\alpha,i}} \rho_i(\pi, \mu_{\alpha}) & \sigma_{\mu_{\alpha,i}}^2 + \sigma_{\mu_{\alpha,j}}^2 & \sigma_{\mu_{\alpha,i}} \sigma_{\mu_{\delta,i}} \rho_i(\mu_{\alpha}, \mu_{\delta}) \\ \sigma_{\pi,i} \sigma_{\mu_{\delta,i}} \rho_i(\pi, \mu_{\delta}) & \sigma_{\mu_{\alpha,i}} \sigma_{\mu_{\delta,i}} \rho_i(\mu_{\alpha}, \mu_{\delta}) & \sigma_{\mu_{\delta,i}}^2 + \sigma_{\mu_{\delta,j}}^2 \end{bmatrix}, \quad (4)$$

where  $\rho_i(\pi, \mu_{\alpha})$ ,  $\rho_i(\pi, \mu_{\delta})$ ,  $\rho_i(\mu_{\alpha}, \mu_{\delta})$  are the non-diagonal elements of the covariance matrix of each star provided by the *Gaia* archive.


The total likelihood for the  $i$ -th star is given by:

$$L_i = \sum_{j=1}^n f_j L_{i,j}, \quad (5)$$

where  $n$  is the number of populations and  $\sum_{j=1}^n f_j = 1$ . The probability for each star of belonging to the  $j$ -th population is then

$$P_{i,j} = f_j \frac{L_{i,j}}{L_i}. \quad (6)$$

We fitted our data with a series of models, starting with two populations, and adding one population at a time. We note that each additional population requires that seven new free parameters be fitted (the three centroids of parallax and proper motions and the relative dispersions, plus the fraction of stars). The statistical significance of each model was tested using the Bayesian information criterion (BIC), which takes into account both the maximum-likelihood and the number of free parameters of the model in order to reduce the risk of overfitting. The resulting values are shown in Fig. 3: we see that the BIC decreases up to six populations and then starts to increase again. Therefore, we conclude that the model with six populations is the best fit of our data, given our assumptions. The best fit parameters for the six populations, which we call Taurus A, B, C, D, E and F, are listed in Table 1, while the probability of each star belonging to



**Fig. 3.** Values of the BIC calculated for the fits with different numbers  $n$  of populations.

each population is reported in Table A.1. Between the six probabilities for each source, the highest identifies the population with which that source is associated. We find that 253 of the 283 stars used for the calculations have a probability of higher than 80% of belonging to one of the six populations: 62 to Taurus A, 46 to Taurus B, 36 to Taurus C, 27 to Taurus D, 40 to Taurus E, and 42 to Taurus F. The distribution of parallaxes and proper motions of these 253 stars are shown in Figs. 4 and 5, respectively. In Fig. 6 (see Sect. 4.2) we show their spatial distribution relative to the molecular cloud traced by *Herschel* far-infrared images.

We note that the majority of the members of the six populations that have been selected using only proper motions and parallaxes are also well separated in the plane of the sky. This is independent confirmation of the results of our analysis.

The relative errors on the derived parallaxes are of the order of 1% or less, and therefore the bias to the distance calculated by inverting the parallaxes is negligible ([Luri et al. 2018](#); [Bailey-Jones 2015](#)). We find that three of the populations are associated with *Taurus Main*: Taurus A and B, located at similar distances (130.5<sup>+0.5</sup><sub>-0.5</sub> pc and 131.0<sup>+1.0</sup><sub>-1.0</sub> pc, respectively), and Taurus E further away at a distance of 160.3<sup>+0.6</sup><sub>-0.6</sub> pc. Taurus C and F are located in the clouds outside *Taurus Main*, at 158.4<sup>+0.6</sup><sub>-0.6</sub> pc and 146.0<sup>+0.6</sup><sub>-0.6</sub> pc, respectively. With a mean distance of 162.7<sup>+4.4</sup><sub>-4.1</sub> pc, Taurus D is the least defined population, with a large spread both in astrometric parameters and in the spatial distribution.

All the given errors in the parallaxes and distances include only the statistical uncertainties from the fit. At these distances, the conservative systematic error of 0.1 mas discussed by [Luri et al. \(2018\)](#) corresponds to  $\sim 2$  pc.

## 4. Discussion

We organize the discussion as follows. First we compare the distances of the six populations to the literature values. We then investigate the properties and distance of the molecular cloud in order to be able to relate the 3D distribution of the stellar population to the molecular cloud.

### 4.1. Distances of the Taurus stellar populations

The distance of the Taurus complex reported in the literature (e.g., [Preibisch & Smith 1997](#)) and commonly used so far is consistent with the average parallax of 7.01 mas that we found previously, but this can no longer be considered valid in light of the present results, which show different distances for the six populations, ranging from 130 pc for Taurus A to 160 pc for Taurus C.

<sup>1</sup> Equations (6)–(10).

**Table 1.** Results from the maximum-likelihood fit for the parallaxes and proper motions, and relative dispersions, of Taurus A, B, C, D, E and F.

	Taurus A	Taurus B	Taurus C	Taurus D	Taurus E	Taurus F
$\pi$ [mas]	$7.660 \pm 0.029$	$7.634 \pm 0.058$	$6.314 \pm 0.025$	$6.145 \pm 0.161$	$6.240 \pm 0.024$	$6.850 \pm 0.029$
$\mu_\alpha$ [mas yr <sup>-1</sup> ]	$8.780 \pm 0.144$	$6.883 \pm 0.258$	$4.618 \pm 0.108$	$5.572 \pm 0.454$	$10.490 \pm 0.209$	$12.413 \pm 0.167$
$\mu_\delta$ [mas yr <sup>-1</sup> ]	$-25.373 \pm 0.201$	$-21.380 \pm 0.170$	$-24.842 \pm 0.138$	$-17.368 \pm 0.934$	$-17.172 \pm 0.168$	$-19.077 \pm 0.145$
$\sigma_{\pi,0}$ [mas]	$0.192 \pm 0.025$	$0.271 \pm 0.058$	$0.101 \pm 0.027$	$0.663 \pm 0.101$	$0.088 \pm 0.034$	$0.136 \pm 0.024$
$\sigma_{\mu_\alpha,0}$ [mas yr <sup>-1</sup> ]	$1.073 \pm 0.113$	$1.358 \pm 0.174$	$0.591 \pm 0.079$	$2.298 \pm 0.314$	$1.141 \pm 0.164$	$0.976 \pm 0.147$
$\sigma_{\mu_\delta,0}$ [mas yr <sup>-1</sup> ]	$1.268 \pm 0.159$	$0.847 \pm 0.123$	$0.770 \pm 0.109$	$4.738 \pm 0.612$	$1.006 \pm 0.126$	$0.891 \pm 0.103$
$f$	$0.233 \pm 0.027$	$0.189 \pm 0.028$	$0.129 \pm 0.021$	$0.127 \pm 0.027$	$0.163 \pm 0.024$	$0.159 \pm 0.057$
$d$ [pc]	$130.5^{+0.5}_{-0.5}$	$131.0^{+1.0}_{-1.0}$	$158.4^{+0.6}_{-0.6}$	$162.7^{+4.4}_{-4.1}$	$160.3^{+0.6}_{-0.6}$	$146.0^{+0.6}_{-0.6}$

**Notes.** In the last line we report the distance in parsecs of the six populations computed as described in the text.

Several measurements at different wavelengths and using different methods have been taken in recent decades to investigate the distance of the Taurus complex. Accurate distances derived from VLBI astrometry are available for several stars. Values of  $128.5 \pm 0.6$  pc,  $132.8 \pm 0.5$  pc, and  $132.8 \pm 2.3$  pc were found for Hubble 4, HDE 283572, and the binary V773 Tau, respectively (Torres et al. 2007, 2012), while a distance of  $161.2 \pm 0.9$  pc was found for the HPTau/G2 system (Torres et al. 2009). Galli et al. (2018) derived VLBI distances for an additional 18 Taurus members, finding values ranging from 84 to 162 pc. We note also that Kenyon et al. (1994) already suggested that the canonical distance of  $140 \pm 10$  pc was not representative of most stars in this region since they measured a significant depth effect using optical spectrophotometry of field stars projected on the molecular cloud. On the contrary, Dzib et al. (2018), using the *Gaia* data alone, found that L1495 in Taurus lies at 129 pc, which corresponds to the distance of Taurus A in our analysis.

The presence of multiple populations was also investigated by a recent study of Luhman (2018), who used colors to identify different populations; we report those colours in Table A.1. We find that all the stars that we classify as part of Taurus A and Taurus B belong to the red population of Luhman (2018), while stars in Taurus C belong to his cyan population. The blue population corresponds to Taurus E and F, while Taurus D includes sources from all the populations found by Luhman (2018). This is to be expected since this is the most spatially dispersed population, without a clear peak in the parallax distribution. We conclude that the populations found by Luhman (2018) are only partially consistent with ours. This is because we adopted completely different approaches. In his case, starting from the spatial distribution of sources in correspondence to the different clouds, Luhman (2018) characterized their average parallax and proper motions without any additional statistical analysis able to take into account the errors and correlations between astrometric parameters. Recently, Galli et al. (2019) also studied the Taurus populations, instead applying a hierarchical clustering algorithm and removing the outliers using the minimum covariance determinant. These latter authors found 21 clusters, which are also indicated in Table A.1, most of which, however, have less than eight members. To compare this latter result with the four populations found by Luhman (2018), Galli et al. (2019) defined six groups of clusters, which are consistent with the four populations of Luhman (2018). Comparing our results with the group classification of Galli et al. (2019), we find that Taurus A and B, both red in Luhman (2018), correspond to Group F in Galli et al. (2019), that Taurus C corresponds to Group C (cyan in Luhman 2018), Taurus E mostly corresponds

to Group A, and Taurus F corresponds to Group B (both blue in Luhman 2018).


It is important to highlight the fact that our decision to not start our analysis from the spatial distribution of the sources (as is highly recommended in clusters without a centrally peaked spatial distribution) is motivated by the fact that we wanted to investigate whether or not the molecular cloud, which is mainly structured in filaments, is shaping the spatial distribution of the different populations.

#### 4.2. Taurus molecular clouds and stellar populations


We can now begin to discuss the Taurus star formation history by combining the new *Gaia* results with high-resolution observations of the filaments in Taurus from *Herschel*. A similar approach was used by Großschedl et al. (2018) with Orion A finding that this is a cometary-like cloud and that the straight filamentary cloud is only its projection.

A large-scale continuum map of the Taurus complex was computed from the near-infrared extinction map from Schmalzl et al. (2010). The *Herschel* observations of Taurus were obtained in the context of the Gould Belt survey (Kirk et al. 2013; Palmeirim et al. 2013) allowing point-source extraction and the determination of the temperature and density profile of the main filament in the LDN 1495 region. Figure 6 shows a composite *Herschel* map of the Taurus complex combining all the public SPIRE observations available in the archive<sup>2</sup>. The positions of the Taurus A, B, C, D, E and F populations are also shown, as well as the position of the known clouds in the region. We find that, in the *Taurus Main* cloud, Taurus A and B are at a similar distance, while Taurus E is located 30 pc further away. The greatest concentration of sources of Taurus A spreads over the B209 cloud. In the rest of *Taurus Main*, all populations follow the filamentary structure traced by the *Herschel* maps. Taurus B, in particular, is mostly distributed along the L1524 and L1529 clouds, while Taurus E is projected in the direction of the filamentary clouds L1495 and B213 and on the cloud L1536. The bulk of the population of Taurus C corresponds to the L1517 cloud, while Taurus F is concentrated on L1551. Few sources of Taurus D are close to the L1544 and L1558 clouds, but in general this population is distributed over the entire complex. A possible explanation for this behavior, given also the spread in parallax and proper motions, is that Taurus D is not a single population but the combination of multiple substructures, which our analysis is not able to distinguish; it may also represent a dispersed young population.

<sup>2</sup> <http://archives.esac.esa.int/hsa/whsa/>



**Fig. 4.** Parallax distribution of the most probable members ( $P \geq 80\%$ ) of the Taurus A (red), Taurus B (blue), Taurus C (yellow), Taurus D (green), Taurus E (cyan), and Taurus F (magenta) populations. In each panel the maximum-likelihood parallaxes for the six populations are also shown as vertical dashed lines using the color code of the histograms.



**Fig. 5.** Proper motions in  $\alpha$  and  $\delta$  of the six populations, color coded as in Fig. 4. The filled dots represent sources with probabilities higher than 80%, while the empty circles represent sources with probabilities lower than 80%.

Figure A.1 shows the 3D spatial distribution of the six populations in cartesian galactic coordinates ( $X, Y, Z$ ), where the  $X$  axis points towards the galactic center,  $Y$  towards the local direction of rotation in the plane of the galaxy, and  $Z$  towards the North Galactic Pole. In particular, given the galactic coordinates ( $l, b$ ),  $X, Y, Z$  are defined as:

$$X = d \cos b \cos l \quad (7)$$

$$Y = d \sin l \cos b \quad (8)$$

$$Z = d \sin b. \quad (9)$$

The projection of the 3D distribution of the six populations on the individual planes is shown in Fig. 7. We find that Taurus A and B in the ( $X, Z$ ) projection are compact and overlapped, while in the other two projections they are separated but close; both populations extend for about 20 pc along the  $X$  axis and 10 pc along the  $Y$  and  $Z$  axes. In the ( $X, Y$ ) and ( $X, Z$ ) planes, Taurus C is distributed along three parallel lines separated by a few parsecs and extending for about 10 pc. In the ( $Y, Z$ ) projection, this population has a more compact structure in the center, which is slightly elongated by about 5 pc in the  $Y$  direction. Taurus D is also confirmed to be a spread population in 3D. Taurus E has a compact structure of  $20 \times 10$  pc only in the ( $X, Z$ ) plane; in the other two projections, it shows a compact structure of  $20 \times 5$  pc and  $5 \times 10$  pc in the ( $X, Y$ ) and ( $Y, Z$ ) planes, respectively, which corresponds to the L1536 cloud. This structure is connected by a few sources along 10 pc to a linear distribution of sources of 10 pc in length corresponding to the L1495 filament. The bulk of sources of Taurus F are concentrated within 10 pc in  $X$  and 5 pc in  $Y$  and  $Z$ ; this corresponds to the L1551 cloud. A few additional sources are scattered over 20 pc in  $X$  and  $Z$ , and  $\sim 40$  pc in  $Y$ .

The distance of the molecular cloud can be estimated using the field stars in and outside of the cloud from *Gaia* DR2. This was done recently by Yan et al. (2019), who found a single distance of  $145^{+12}_{-16}$  pc for the entire molecular cloud. A similar approach was adopted by Zucker et al. (2019), who compiled a uniform catalog of distances to local molecular clouds. The methodology of these latter authors, based on the work of Schlafly et al. (2014), infers a joint probability distribution function on distance and reddening for individual stars based on optical and near-infrared photometric surveys and *Gaia* parallaxes, and also on modeling of the cloud as a simple dust screen