



<b>Publication Year</b>	2016
<b>Acceptance in OA</b>	2020-07-14T10:43:41Z
<b>Title</b>	Mesospheric CO <sub>2</sub> ice clouds on Mars observed by Planetary Fourier Spectrometer onboard Mars Express
<b>Authors</b>	Aoki, Shohei, GIURANNA, MARCO, Sato, Yuki, Nakagawa, Hiromu, Sato, Takao M., WOLKENBERG, PAULINA MARIA, Murata, Isao, Kasaba, Yasumasa
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/26447">http://hdl.handle.net/20.500.12386/26447</a>
<b>Journal</b>	GEOPHYSICAL RESEARCH ABSTRACTS



## Mesospheric CO<sub>2</sub> ice clouds on Mars observed by Planetary Fourier Spectrometer onboard Mars Express

Shohei Aoki (1,2), Marco Giuranna (1), Yuki Sato (2), Hiromu Nakagawa (2), Takao M. Sato (3), Paulina Wolkenberg (1,4), Isao Murata (2), and Yasumasa Kasaba (2)

(1) Istituto di Astrofisica e Planetologia Spaziali (IAPS), Istituto Nazionale di Astrofisica (INAF), Roma, Italy (shohei.aoki@iaps.inaf.it), (2) Department of Geophysics, Tohoku University, Sendai, Japan, (3) Institute of Space and Astronautical Science (ISAS), Japan Aerospace Exploration Agency (JAXA), Sagami-hara, Japan, (4) Centrum Badań Kosmicznych Polskiej Akademii Nauk (CBK-PAN), Warszawa, Polska

We investigate mesospheric CO<sub>2</sub> ice clouds on Mars detected by the Planetary Fourier Spectrometer (PFS) onboard Mars Express (MEx). The relatively high spectral resolution of PFS allows firm identification of the clouds' reflection spike. A total of 279 occurrences of the CO<sub>2</sub> ice clouds features has been detected at the bottom of 4.3 μm CO<sub>2</sub> band from the MEx/PFS data during the period from MY27 to MY32. 115 occurrences out of them are also confirmed by simultaneous observations by MEx/OMEGA imaging spectrometer. The spatial and seasonal distributions of the CO<sub>2</sub> ice clouds observed by PFS are consistent with the previous studies: the CO<sub>2</sub> ice clouds are only observed between L<sub>s</sub>=0° and 140° at distinct longitudinal corridors around the equatorial region (±20°N). The CO<sub>2</sub> ice clouds are preferentially detected at local time between 15-17h. The relatively high spectral resolution of PFS allows us to investigate the spectral shape of the CO<sub>2</sub> ice clouds features. The CO<sub>2</sub> ice clouds reflection spike is peaked between 4.24 and 4.29 μm, with no evidence of the secondary peak at 4.32-4.34 μm observed by MEx/OMEGA (Määttä et al., 2010). In most of the cases (about 75%), the peak is present between 4.245 and 4.255 μm. Moreover, small secondary peaks are found around 4.28 μm (about 15 occurrences). These spectral features cannot be reproduced by the synthetic spectra with the assumption of a spherical particle shape in our radiative transfer model (DISORT). This can be due to the fact that the available CO<sub>2</sub> ice reflective indexes are either inaccurate or inappropriate for the mesospheric temperatures, or that the particle shape is not spherical. Accurate measurements of the reflective index depending on temperature and detailed comparison with the model taking into account non-spherical shapes will give a clue to solve this issue.