



<b>Publication Year</b>	2016
<b>Acceptance in OA</b>	2020-05-12T08:59:52Z
<b>Title</b>	GRB 160712A: Swift-XRT observations
<b>Authors</b>	Kennea, J. A., CUSUMANO, GIANCARLO, PERRI, Matteo, Evans, P. A., Osborne, J. P., Burrows, D. N.
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/24725">http://hdl.handle.net/20.500.12386/24725</a>
<b>Journal</b>	GRB Coordinates Network
<b>Volume</b>	19686

TITLE: GCN CIRCULAR  
NUMBER: 19686  
SUBJECT: GRB 160712A: Swift-XRT observations  
DATE: 16/07/12 21:28:03 GMT  
FROM: Phil Evans at U of Leicester <pae9@leicester.ac.uk>

J.A. Kennea (PSU), G. Cusumano (INAF-IASF PA), M. Perri (ASDC), P.A. Evans (U. Leicester), J.P. Osborne (U. Leicester) and D.N. Burrows (PSU) report on behalf of the Swift-XRT team:

The XRT began observing the field of GRB 160712A at 20:47:52.7 UT, 3256.7 seconds after the BAT trigger. The position determined from promptly downlinked data differs significantly from the on-board position, suggesting that the XRT may have centroided on a cosmic ray; the initial XRT position notice should be treated with caution. Using promptly downlinked data we find an uncatalogued X-ray source with an enhanced position: RA, Dec 304.1600, -26.9582 which is equivalent to:

RA(J2000) = 20h 16m 38.40s

Dec(J2000) = -26d 57' 29.4"

with an uncertainty of 2.1 arcseconds (radius, 90% containment). This location is 89 arcseconds from the BAT onboard position, within the BAT error circle. This position may be improved as more data are received; the latest position is available at <http://www.swift.ac.uk/sper>.

A power-law fit to a spectrum formed from promptly downlinked event data gives a column density in excess of the Galactic value ( $8.30 \times 10^{20} \text{ cm}^{-2}$ , Willingale et al. 2013), with an excess column of  $2.7 (+2.97/-2.55) \times 10^{21} \text{ cm}^{-2}$  (90% confidence).