



Publication Year	2015
Acceptance in OA	2020-07-07T09:55:57Z
Title	Dust Measurements in the Coma of Comet 67P/Churyumov-Gerasimenko Inbound to the Sun Between 3.7 and 3.4 AU
Authors	FULLE, Marco, DELLA CORTE, VINCENZO, Rotundi, Alessandra, Accolla, M., FERRARI, MARCO, IVANOVSKI, STAVRO LAMBROV, LUCARELLI, Fabrizio, SORDINI, Roberto, ZAKHAROV, VLADIMIR, MAZZOTTA EPIFANI, Elena, Lopez-Moreno, J. J., Rodriguez, J., COLANGELI, Luigi, Palumbo, P., Bussoletti, E., Crifo, J., ESPOSITO, Francesca, Green, S. F., Gruen, E., Lamy, P. L., McDonnell, J. A. M., MENNELLA, Vito, Molina, A., Morales, R., Moreno, F., Ortiz, J. L., PALOMBA, Ernesto, Perrin, J., Rodrigo, R., Weissman, P., Zarnecki, J. C., Cosi, M., Giovane, F., Gustafson, B., Herranz, M. L., Jeronimo, J. M., Leese, M. R., Lopez-Jimenez, A. C., Altobelli, N., Sierks, H., Agarwal, J., Bertini, I., Fornasier, S., Gutierrez, P. J., Lara, L., Guettler, C., Marzari, F., Oaklay, N., Snodgrass, C., Tubiana, C., Vincenzo, J. B.
Handle	http://hdl.handle.net/20.500.12386/26357

Dust Measurements in the Coma of Comet 67P/Churyumov-Gerasimenko Inbound to the Sun Between 3.7 and 3.4 AU. M. Fulle¹, V. Della Corte², A. Rotundi^{2,3}, M. Accolla^{2,3}, M. Ferrari^{2,3}, S. Ivanovski², F. Lucarelli³, R. Sordini², V. Zakharov⁴, E. Mazzotta Epifani⁵, J. J. López-Moreno⁶, J. Rodríguez⁶, L. Colangeli⁷, P. Palumbo^{2,3}, E. Bussoletti³, J. Crifo⁸, F. Esposito⁵, S. F. Green⁹, E. Grün¹⁰, P. L. Lamy¹¹, J. A. M. McDonnell^{9,12}, V. Mennella⁵, A. Molina¹³, R. Morales⁶, F. Moreno⁶, J. L. Ortiz⁶, E. Palomba², J. Perrin^{8,14}, R. Rodrigo^{15,16}, P. Weissman¹⁷, J. C. Zarnecki¹⁶, M. Cossi¹⁸, F. Giovane¹⁹, B. Gustafson²⁰, M. L. Herranz⁶, J. M. Jerónimo⁶, M. R. Leese⁹, A. C. López-Jiménez⁶, N. Altobelli²¹, H. Sierks²², J. Agarwal²², I. Bertini²³, S. Fornasier⁴, P. J. Gutiérrez⁶, L. Lara⁶, C. Guettler²², F. Marzari²⁴, N. Oklay²², C. Snodgrass^{9,22}, C. Tubiana²², J.B. Vincent²².

1 INAF – Osservatorio Astronomico di Trieste, Via Tiepolo Giambattista, 11, 34131 Trieste, Italy.

2 INAF – Istituto di Astrofisica e Planetologia Spaziali, Via Fosso del Cavaliere, 100, 0133, Rome, Italy.

3 Università degli Studi di Napoli “Parthenope”, Dip. di Scienze e Tecnologie, CDN IC4, 80143, Naples, Italy

4 LESIA, Obs. de Paris, CNRS, Univ Paris 06, Univ. Paris-Diderot, 5 place J. Janssen, 92195, Meudon, France

5 INAF – Osservatorio Astronomico di Capodimonte, Salita Moiariello, 16, 80133, Naples, Italy.

6 Instituto de Astrofísica de Andalucía (CSIC), Glorieta de la Astronomía s/n, 18008 Granada, Spain.

7 ESA - ESTEC, Keplerlaan 1, 2201 AZ Noordwijk, The Netherlands.

8 LATMOS, 11 Bd d'Alembert, F78280 Guyancourt, France.

9 Planetary and Space Sciences, Department of Physical Sciences, The Open University, Milton Keynes, MK7 6AA, UK.

10 Max-Planck-Institut fuer Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany

11 Astrophysics Laboratory of Marseilles, UMR7326 - CNRS-INSU, Université d'Aix-Marseille, Marseilles, France

12 The University of Kent, Canterbury, Kent, CT2 7NZ, UK

13 Departamento de Física Aplicada, Universidad de Granada, Facultad de Ciencias, Avda. Severo Ochoa, s/n, 18071, Granada, Spain.

14 Observatoire de Haute Provence OSU Pythéas UMS 2244 CNRS-AMU, 04870 Saint Michel l'Observatoire, France

15 Centro de Astrobiología (INTA-CSIC), 28691 Villanueva de la Canada, Madrid, Spain

16 International Space Science Institute, Hallerstrasse 6, 3012, Bern, Switzerland

17 Planetary Science Section, Jet Propulsion Laboratory, 4800 Oak Grove Dr, Pasadena, CA 91109, USA

18 Selex-ES, Via A. Einstein, 35, 50013 - Campi Bisenzio - (Firenze), Italy

19 Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA

20 University of Florida, Gainesville, Florida, 32611, USA

21 ESA-ESAC, Camino Bajo del Castillo, s/n., Urb. Villafranca del Castillo, 28692 Villanueva de la Cañada, Madrid, Spagna

22 Max Planck Institute for Solar System Research, Justus-von-Liebig-Weg 3, 37077 Göttingen, Germany

23 Center of Studies and Activities for Space (CISAS), University of Padova,

I-35131 Padova, Italy

24 Department of Physics and Astronomy, University of Padova, I-35131 Padova, Italy

Introduction: Comets are the most primitive bodies in the solar system. They retain a cosmo-chemical record of conditions in the solar nebula when the planets were forming, 4.5 billion years ago. While accurate measurements of the gas loss rate from comets are possible under favorable conditions even from Earth, estimates of the dust loss rate so far have been much more uncertain. Multi-parametric models are needed to extract global dust parameters from the dust features of comets (e.g. coma, tails and trails) observed from ground and Earth orbiting telescopes, and it is often difficult to establish the uniqueness of these model results. Past space missions had on board instruments devoted to the measurement of the dust flux. Since all these missions were fast flybys, it was impossible to disentangle the dust grains coming directly from the nucleus from those reflected back by solar radiation pressure [1,2]. The latter component could explain all or part of the excess of mm-sized particles observed

during flybys at 1P/Halley [3] and at short-period comets 26P/Grigg-Skjellerup, 81P/Wild 2 and 9P/Tempel 1 [4-6].

An even more severe bias could affect all estimates of dust to gas (d/g) ratio obtained so far in comets. The d/g measured in 1P/Halley was close to 2 [5], but this number is valid up to the largest mass of about 1 g observed by the DIDSY detector (actually, this largest mass grain was invoked to explain the spacecraft precession-inducing impact that occurred just before closest approach). We expect that 1P/Halley was then ejecting much larger masses, and the d/g ratio strongly depends on the actual largest grain ejected in the coma. Since it was impossible to fix the size distribution between 1 g and the unknown largest ejected mass, we cannot exclude d/g values higher than 10 or even 50. In this paper we show that for comet 67P/CG we can disentangle the two families of ejected grains (direct and reflected), and extract the dust size distribution up to

the largest ejected grain, obtaining for the first time an accurate estimate of the total dust loss rate from the nucleus.

Methods: Critical measurements for understanding the process of accretion and the refractory to volatiles ratio in the solar nebula are being obtained by the Grain Impact Analyzer and Dust Accumulator (GIADA) experiment onboard ESA's Rosetta spacecraft, now orbiting comet 67P/Churyumov-Gerasimenko (67P/CG). GIADA measures the mass, momentum and velocity of individual grains, providing the dust loss rate over three orders of magnitude in mass for grains from tens to hundreds of microns in diameter. GIADA consists of three subsystems: 1) the Grain Detection System (GDS) to detect dust grains as they pass through a laser curtain, 2) the Impact Sensor (IS) to measure grain momentum derived from the impact on a plate connected to five piezoelectric sensors, and 3) the MicroBalances System (MBS); five quartz crystal microbalances in roughly orthogonal directions providing the cumulative dust flux of grains smaller than 10 microns. GDS provides data on grain speed and its optical cross section. The IS grain momentum measurement, when combined with the GDS detection time, provides a direct measurement of grain speed and mass [7, 8]. These combined measurements characterize single grain dust dynamics in the coma of 67P/CG.

Results: The first grain was detected on 1 August 2014 at 814 km from the comet nucleus. Between then and 13 September 2014 GIADA detected 35 grains ranging in mass from $\sim 5 \times 10^{-10}$ to 8×10^{-8} kg. Including complementary data from the OSIRIS narrow angle camera, the dust mass loss was calculated over an additional three orders of magnitude in mass, extending the ejected dust grain sizes up to 2 cm. Combined with data from the MIRO and the ROSINA instruments onboard Rosetta we find a dust/gas mass ratio of 4 ± 2 averaged over the sunlit nucleus surface. The dust to gas ratio may change as the comet approaches closer to the Sun.

References: [1] M. Fulle, L. Colangeli, V. Menella, A. Rotundi, E. Bussoletti, (1995). *Astron. Astrophys.* 304, 622-630. [2] M. Fulle, A. C. Levasseur-Regourd, N. McBride, E. Hadamcik, (2000) *Astron. J.* 119, 1968-1977 [3] J. A. M. McDonnell, G. S. Pankiewicz, P. N. W. Birchley, S. F. Green, C. H. Perry. In Proc. 20th Lunar and Planet. Sci. Conf. (Houston, TX, Lunar and Planetary Institute, (1990) pp. 373-378. [4] J. A. M. McDonnell et al., (1993) *Nature* 362, 732-734. [5] S. F. Green et al., *J. Geophys. Res.*

109, (E12), E12S04 (2004). [6] T. E. Economou, S. F. Green, D. E. Brownlee, B. C. Clark, *Icarus* 222, Issue 2, 526-539 (2013). [7] V. Della Corte et al., (2014) *Journal of Astronomical Instrumentation* 3, Issue 1, 1350011-11, doi:10.1142/S2251171713500116. [8] L. Colangeli, et al., (2007) *Space Science Reviews* 128, Issue 1-4, 803-821.

Acknowledgments: GIADA was built by a consortium led by the Univ. Napoli "Parthenope" & INAF-Oss. Astr. Capodimonte, in collaboration with the Inst. de Astrofísica de Andalucía, Selex-ES, FI and SENEC. GIADA is presently managed & operated by Ist. di Astrofísica e Planetologia Spaziali-INAf, IT. GIADA was funded and managed by the Agenzia Spaziale Italiana, IT, with the support of the Spanish Ministry of Education and Science MEC, ES. GIADA was developed from a PI proposal from the University of Kent; sci. & tech. contribution were provided by CISAS, IT, Lab. d'Astr. Spat., FR, and Institutions from UK, IT, FR, DE and USA. We thank the RSGS/ESAC, RMOC/ESOC & Rosetta Project/ESTEC for their outstanding work. Science support provided was by NASA through the US Rosetta Project managed by the Jet Propulsion Laboratory/California Institute of Technology. GIADA calibrated data will be available through ESA's PSA web site (www.rssd.esa.int/index.php?project=PSA&page=index). We would like to thank Angioletta Coradini for her contribution as a GIADA Co-I. We thank the MIRO, OSIRIS and ROSINA teams for sharing their early results with us.