



Publication Year	2019
Acceptance in OA	2021-02-26T17:22:56Z
Title	Addressing the circularity problem in the Ep-Eiso correlation of gamma-ray bursts
Authors	AMATI, LORENZO, D'AGOSTINO, Rocco, Luongo, Orlando, Muccino, Marco, TANTALO, MARIA
Publisher's version (DOI)	10.1093/mnrasl/slz056
Handle	http://hdl.handle.net/20.500.12386/30660
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY. LETTERS
Volume	486

Addressing the circularity problem in the $E_p - E_{\text{iso}}$ correlation of gamma-ray bursts

Lorenzo Amati,¹ Rocco D'Agostino,² Orlando Luongo^{3,4*}, Marco Muccino³ and Maria Tantalò⁵

¹INAF, Istituto di Astrofisica Spaziale e Fisica Cosmica, Bologna, Via Gobetti 101, I-40129 Bologna, Italy

²Sezione INFN, Università di Roma 'Tor Vergata', Via della Ricerca Scientifica 1, I-00133 Roma, Italy

³Istituto Nazionale di Fisica Nucleare, Laboratori Nazionali di Frascati, I-00044 Frascati, Italy

⁴NNLOT, Al-Farabi Kazakh National University, Al-Farabi av. 71, 050040 Almaty, Kazakhstan

⁵Dipartimento di Fisica, Università di Roma 'Tor Vergata', Via della Ricerca Scientifica 1, I-00133 Roma, Italy

Accepted 2019 April 15. Received 2019 April 3; in original form 2019 March 14

ABSTRACT

We here propose a new model-independent technique to overcome the circularity problem affecting the use of gamma-ray bursts (GRBs) as distance indicators through the use of $E_p - E_{\text{iso}}$ correlation. We calibrate the $E_p - E_{\text{iso}}$ correlation and find the GRB distance moduli that can be used to constrain dark energy models. We use observational Hubble data to approximate the cosmic evolution through Bézier parametric curve obtained through the linear combination of Bernstein basis polynomials. In doing so, we build up a new data set consisting of 193 GRB distance moduli. We combine this sample with the supernova JLA data set to test the standard Λ CDM model and its w CDM extension. We place observational constraints on the cosmological parameters through Markov Chain Monte Carlo numerical technique. Moreover, we compare the theoretical scenarios by performing the Akaike and Deviance Information statistical criteria. At the 2σ level, while for the w CDM model we obtain $\Omega_m = 0.34^{+0.13}_{-0.15}$ and $w = -0.86^{+0.36}_{-0.38}$ at the 2σ level. Our analysis suggests that Λ CDM model is statistically favoured over the w CDM scenario. No evidence for extension of the Λ CDM model is found.

Key words: cosmology: observations – dark energy – gamma-rays: general.

1 INTRODUCTION

The cosmic speed up is today a consolidated experimental evidence confirmed by several probes (Haridasu et al. 2017). Particularly, Type Ia Supernovae (SNe Ia) have been employed as standard candles (Phillips 1993) to check the onset of cosmic acceleration (Perlmutter et al. 1998, 1999; Riess et al. 1998; Schmidt et al. 1998). Their importance lies in the fact that they may open a window into the nature of the constituents pushing up the universe to accelerate. Even though SNe Ia are considered among the most reliable standard candles, they are detectable at most at redshifts $z \simeq 2$ (Rodney et al. 2015). Thus, at intermediate redshifts the standard cosmological model, dubbed the Λ CDM paradigm, cannot be tested with SNe Ia alone. Consequently, higher redshift distance indicators, such as Baryon Acoustic Oscillations (Percival et al. 2010; Aubourg et al. 2015; Luković, D'Agostino & Vittorio 2016), have been used to alleviate degeneracy among the Λ CDM paradigm and dark energy scenarios. In these respects, a relevant example if

offered by gamma-ray bursts (GRBs), which represent the most powerful cosmic explosions detectable up to $z = 9.4$ (Salvaterra et al. 2009; Tanvir et al. 2009; Cucchiara et al. 2011). Attempts to use GRBs as cosmic rulers led cosmologists to get several correlations between GRB photometric and spectroscopic properties (Amati et al. 2002; Ghirlanda et al. 2004; Schaefer 2007; Amati et al. 2008; Capozziello & Izzo 2008; Dainotti, Cardone & Capozziello 2008; Bernardini et al. 2012; Amati & Della Valle 2013; Wei et al. 2014; Izzo et al. 2015; Demianski et al. 2017a,b). The most investigated correlations involve the rest-frame spectral peak energy E_p , i.e. the rest-frame photon energy at which the νF_ν spectrum of the GRB peaks, and the bolometric isotropic-equivalent radiated energy E_{iso} , or peak luminosity L_p (Amati et al. 2002; Yonetoku et al. 2004; Amati et al. 2008; Amati & Della Valle 2013; Demianski et al. 2017a,b). However, the use of GRBs for cosmology is still affected by some uncertainties due to selection and instrumental effects and the so-called *circularity problem* (see e.g. Kodama et al. 2008). The former issue has been investigated in several studies, with the general, even though still debated, conclusion that these effects should be minor (see e.g. Amati 2006; Ghirlanda, Ghisellini & Firmani 2006; Nava et al. 2012; Amati & Della

* E-mail: luongo@inf.infn.it

Valle 2013; Demianski et al. 2017a). The circularity problem arises from the fact that, given the lack of very low-redshift GRBs, the correlations between radiated energy or luminosity and the spectral properties are established assuming a background cosmology. For example, calibrating GRBs through the standard Λ CDM model, the estimate of cosmological parameters of any dark energy framework inevitably returns an overall agreement with the concordance model.

In this paper, we propose a new model-independent calibration of the E_p – E_{iso} correlation (the *Amati relation*; see e.g. Amati et al. 2008; Amati & Della Valle 2013). We take the most recent values of observational Hubble Data (OHD), consisting of 31 points of Hubble rates got at different redshifts (see Capozziello, D’Agostino & Luongo 2018, and references therein). These data have been obtained through the differential age method applied to pairs of nearby galaxies, providing model-independent measurements (Jimenez & Loeb 2002). We follow the strategy to fit OHD data using a Bézier parametric curve obtained through the linear combination of Bernstein basis polynomials. This treatment is a refined approximated method and reproduces Hubble’s rate at arbitrary redshifts without assuming an a priori cosmological model. We thus use it to calibrate the E_{iso} values by means of a data set made of 193 GRBs (with firmly measured redshift and spectral parameters taken from Demianski et al. 2017a, and references therein), and compute the corresponding GRB distance moduli μ_{GRB} and the 1σ error bars, depending upon the uncertainties on GRB observables. Detailed discussions of possible biases and selection effects can be found, e.g. in Amati & Della Valle (2013), Demianski et al. (2017a), and Dainotti & Amati (2018). From the above model-independent analysis over OHD data, we obtain $H_0 = 67.74 \text{ km s}^{-1} \text{ Mpc}^{-1}$, compatible with the current estimates by the Planck Collaboration VI (2018) and Riess et al. (2018).

As a pure example of fitting procedure, we analyse our data by means of Markov Chain Monte Carlo (MCMC) technique and compare them with the standard cosmological paradigm and its simplest extension, namely the w CDM model. We discuss the limits over our technique in view of the most recent bias and problems related to SN Ia and GRBs. Afterwards, using the above value of H_0 got from our parametric fit analysis, we show that our results are in tension with the concordance paradigm (Planck Collaboration VI 2018) at $\geq 3\sigma$. However, we propose that such results may be affected by systematics and how these limits may be reconsidered in view of future developments. Finally, we compare the statistical performance of the cosmological models through the Bayesian selection criteria.

The paper is divided into four sections. After this Introduction, in Section 2 we describe the main features of our treatment, using OHD data surveys over the Amati relation. In Section 3, we discuss our numerical outcomes concerning the use of our new data set. We thus get constraints over the free parameters of the Λ CDM and w CDM models. In Section 4, we draw conclusions and identify the perspectives of our work.

2 MODEL-INDEPENDENT CALIBRATION OF THE AMATI RELATION

Calibrating the Amati relation represents a challenge due to the problem of circularity (see e.g. Ghirlanda et al. 2004; Ghirlanda et al. 2006; Kodama et al. 2008; Amati & Della Valle 2013). In fact, in the E_p – E_{iso} correlation, the cosmological parameters Ω_i and the Hubble constant H_0 enter in the E_{iso} definition through the luminosity distance d_L , i.e. $E_{\text{iso}}(z, H_0, \Omega_i) \equiv 4\pi d_L^2(z, H_0, \Omega_i) S_{\text{bolo}}/(1+z)$, where S_{bolo} is the observed bolometric GRB fluence and the factor




Figure 1. OHD data (31 black points with the vertical error bars), their best-fitting function (the solid thick blue curve), and its 1σ (the blue curves and the light blue shaded arc) and 3σ (the blue dashed curves) confidence regions.

$(1+z)^{-1}$ transforms the observed GRB duration into the source cosmological rest frame one. The most quoted approach to the calibration of the Amati relation uses the SN Ia Hubble diagram, directly inferred from the observations, and interpolate it to higher redshift using GRBs (see e.g. Kodama et al. 2008; Liang et al. 2008; Demianski et al. 2017a,b). However, this method biases the GRB Hubble diagram by introducing the systematics of the SNe Ia.

Here, we propose an alternative calibration that uses the *differential age method* based on spectroscopic measurements of the age difference Δt and redshift difference Δz of couples of passively evolving galaxies that formed at the same time (Jimenez & Loeb 2002). This method implies that $\Delta z/\Delta t \equiv dz/dt$ and hence the Hubble function can be computed in a cosmology-independent way as $H(z) = -(1+z)^{-1} \Delta z/\Delta t$. The updated sample of 31 OHD (see Capozziello et al. 2018) is shown in Fig. 1. To avoid the circularity problem, we approximate the OHD data by employing a Bézier parametric curve¹ of degree n :

$$H_n(z) = \sum_{d=0}^n \beta_d h_n^d(z), \quad h_n^d(z) \equiv \frac{n!(z/z_m)^d}{d!(n-d)!} \left(1 - \frac{z}{z_m}\right)^{n-d}, \quad (1)$$

where β_d are coefficients of the linear combination of Bernstein basis polynomials $h_n^d(z)$, positive in the range $0 \leq z/z_m \leq 1$, where z_{max} is the maximum z of the OHD data set. For $d=0$ and $z=0$, we easily identify $\beta_0 \equiv H_0$. Besides the simple cases with $n=0$ and $n=1$ leading to a constant value and a linear growth with z of $H(z)$, respectively, the only case providing a monotonic growing function over the limited range in redshift of the OHD data is $n=2$; higher values lead to oscillatory behaviours of the approximating function. Therefore, in the following we use $n=2$ in fitting the OHD data. The best fit with its 1σ and 3σ confidence regions are shown in Fig. 1. The best-fitting parameters are $H_0 = 67.76 \pm 3.68$, $\beta_1 = 103.34 \pm 11.14$, and $\beta_2 = 208.45 \pm 14.29$ (all in units of $\text{km s}^{-1} \text{ Mpc}^{-1}$). The value of H_0 so obtained is compatible with the current estimate of the Planck Collaboration (Planck Collaboration VI 2018) and in agreement at the 1.49σ level with the value measured by Riess et al. (2018).

¹Bézier curves are easy to use in computation, are stable at the lower degrees of control points, and can be rotated and translated by performing the operations on the points.




Figure 2. GRB calibrated distribution in the E_p – $E_{\text{iso}}^{\text{cal}}$ plane (black data points), the best-fitting function (red solid line) and the $1\sigma_{\text{ex}}$ and $3\sigma_{\text{ex}}$ limits (dark-grey and light-grey shaded regions, respectively).

Once the function $H_2(z)$ is extrapolated to redshift $z > z_m$, the luminosity distance is (see e.g. Goobar & Perlmutter 1995)

$$d_L(\Omega_k, z) = \frac{c}{H_0} \frac{(1+z)}{\sqrt{|\Omega_k|}} S_k \left[\sqrt{|\Omega_k|} \int_0^z \frac{H_0 dz'}{H_2(z')} \right], \quad (2)$$

where Ω_k is the curvature parameter, and $S_k(x) = \sinh(x)$ for $\Omega_k > 0$, $S_k(x) = x$ for $\Omega_k = 0$, and $S_k(x) = \sin(x)$ for $\Omega_k < 0$. We note that d_L in equation (2) is not completely independent of cosmological scenarios since it depends upon Ω_k . However, supported by the most recent Planck results (Planck Collaboration VI 2018), which find $\Omega_k = 0.001 \pm 0.002$, we can safely assume $\Omega_k = 0$. In doing so, the dependence upon Ω_k identically vanishes and equation (2) becomes cosmology independent:

$$d_{\text{cal}}(z) = c(1+z) \int_0^z \frac{dz'}{H_2(z')}. \quad (3)$$

We are now in the position to use $d_{\text{cal}}(z)$ to calibrate the isotropic energy $E_{\text{iso}}^{\text{cal}}$ for each GRB fulfilling the Amati relation:²

$$E_{\text{iso}}^{\text{cal}}(z) \equiv 4\pi d_{\text{cal}}^2(z) S_{\text{bolo}} (1+z)^{-1}, \quad (4)$$

where the respective errors $\sigma E_{\text{iso}}^{\text{cal}}$ depend upon the GRB systematics on the observables and the fitting procedure (see confidence regions in Fig. 1). The corresponding E_p – $E_{\text{iso}}^{\text{cal}}$ distribution is displayed in Fig. 2. Following the method by D’Agostini (2005), we fit the calibrated Amati relation using a linear fit $\log(E_p/1\text{keV}) = q + m[\log(E_{\text{iso}}^{\text{cal}}/\text{erg}) - 52]$. We find the best-fitting parameters $q = 2.06 \pm 0.03$, $m = 0.50 \pm 0.02$, and the extra-scatter $\sigma_{\text{ex}} = 0.20 \pm 0.01$ dex (see Fig. 2). The corresponding Spearman’s rank correlation coefficient is $\rho_s = 0.84$ and the p -value from the two-sided Student’s t -distribution is $p = 2.42 \times 10^{-36}$.

We can then compute the GRB distance moduli from the standard definition $\mu_{\text{GRB}} = 25 + 5\log(d_{\text{cal}} \text{Mpc}^{-1})$. Using the fit of the calibrated Amati relation, we obtain

$$\mu_{\text{GRB}} = 25 + \frac{5}{2} \left[\frac{\log E_p - q}{m} - \log \left(\frac{4\pi S_{\text{bolo}}}{1+z} \right) + 52 \right], \quad (5)$$

²Recent works claim that our universe has non-zero curvature and that Ω_k represents at most the 2 percent of the total universe energy density (see e.g. Ooba, Ratra & Sugiyama 2018, and references therein). Relaxing the assumption $\Omega_k = 0$ since its value is still very small, the circularity problem is not completely healed, but it is only restricted to the value of Ω_k since $H(z)$ can be still approximated by the function $H_2(z)$.




Figure 3. *Upper plot:* GRB distance moduli μ_{GRB} distribution compared to the Λ CDM model $\mu_{\Lambda\text{CDM}}$ with $H_0 = 67.36 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3166$, and $\Omega_\Lambda = 0.6847$ as in Planck Collaboration VI (2018; the solid red curve), and two w CDM models with the above Λ CDM parameters and $w = -0.90$ (the dashed blue curve) and $w = -0.75$ (the dot-dashed green curve). *Lower plot:* the deviations of the above three models μ_X from $\mu_{\Lambda\text{CDM}}$ computed as $(\mu_X - \mu_{\Lambda\text{CDM}})/\mu_{\Lambda\text{CDM}}$ (curves retain the same meaning as before).

where now S_{bolo} has been normalized to erg Mpc^{-2} to obtain d_{cal} in the desired units of Mpc. The attached errors on μ_{GRB} take into account the GRB systematics and the statistical errors on q , m , and σ_{ex} . The distribution of μ_{GRB} with z is shown in Fig. 3.

We note that the statistical method adopted for the GRBs calibration may be in principle used also for the analysis of the SN data. This would in fact reduce the propagation errors when the combined fit of both data sets is performed, making the joint sample homogeneous for the cosmological studies. It will be interesting to analyse the impact of such a procedure in a forthcoming study, where the Philips relation of SN will be calibrated in the way we attempted with GRBs prior to performing the cosmological fit.

3 NUMERICAL RESULTS

We here use our sample of GRBs to test cosmological models. In particular, we assume standard barotropic equation of state (EoS). Thus, for each fluid the pressure P_i is a one-to-one function of the density ρ_i : $P_i = w_i \rho_i$. As a consequence of Bianchi’s identity, one gets $\dot{\rho}_i + 3H \rho_i (1 + w_i) = 0$ for each species entering the Einstein equations. Following the standard recipe, we here consider pressureless matter with negligible radiation and define current total density as $\Omega_i = \rho_i/\rho_c$, with $\rho_c \equiv 8\pi G/(3H_0^2)$ is the critical density, one can reformulate the Hubble evolution as

$$H(z) = H_0 \sqrt{\Omega_m (1+z)^3 + \Omega_{\text{DE}} (1+z)^{3(1+w)}}. \quad (6)$$

In the above relation, dark energy takes a net density given by $\Omega_{\text{DE}} = 1 - \Omega_m$ to guarantee that $H(z=0) = H_0$, and w is the dark energy EoS parameter. In particular, equation (6) reduces to the Λ CDM model as $w = -1$, whereas to the w CDM model when w is free to vary. The distance modulus is given by $\mu_{\text{th}}(z) = 25 + 5\log[d_L(z) \text{Mpc}^{-1}]$, where $d_L(z)$ is given by equation (2) with $\Omega_k = 0$. Thus, the likelihood function of the GRB data can be written as

$$\mathcal{L}_{\text{GRB}} = \prod_{i=1}^{N_{\text{GRB}}} \frac{1}{\sqrt{2\pi} \sigma_{\mu_{\text{GRB},i}}} \exp \left[-\frac{1}{2} \left(\frac{\mu_{\text{th}}(z_i) - \mu_{\text{GRB},i}}{\sigma_{\mu_{\text{GRB},i}}} \right)^2 \right], \quad (7)$$

Table 1. Priors used for parameters estimate in the MCMC analysis.

w	Ω_m	M	Δ_M	α	β
$(-0.5, -1.5)$	$(0,1)$	$(-20, -18)$	$(-1, 1)$	$(0,1)$	$(0,5)$

where $N_{\text{GRB}} = 193$ is the number of GRB data points. To obtain more robust observational bounds on cosmological parameters, we consider a complete Hubble diagram by complementing the GRB measurements with the SN JLA sample (Betoule et al. 2014). The latter consists of 740 SN Ia in the redshift range $0.01 < z < 1.3$. The distance modulus of each SN is parametrized as

$$\mu_{\text{SN}} = m_B - M_B + \alpha X_1 - \beta C, \quad (8)$$

where m_B is the B -band apparent magnitude, while C and X_1 are the colour and the stretch factor of the light curve, respectively; M_B is the absolute magnitude defined as

$$M_B = \begin{cases} M & \text{if } M_{\text{host}} < 10^{10} M_{\text{Sun}}, \\ M + \Delta_M & \text{otherwise,} \end{cases} \quad (9)$$

where M_{host} is the host stellar mass, and M , α , and β are nuisance parameters that enter the fits along with cosmological parameters. The likelihood function of the SN data is given as

$$\mathcal{L}_{\text{SN}} = \frac{1}{|2\pi\mathcal{M}|^{1/2}} \exp\left[-\frac{1}{2}(\mu_{\text{th}} - \mu_{\text{SN}})^T \mathcal{M}^{-1}(\mu_{\text{th}} - \mu_{\text{SN}})\right], \quad (10)$$

where \mathcal{M} is the $3N_{\text{SN}} \times N_{\text{SN}} = 2220 \times 2220$ covariance matrix with the statistical and systematic uncertainties on the light-curve parameters given in Betoule et al. (2014).

We thus perform an MCMC integration on the combined likelihood function $\mathcal{L} = \mathcal{L}_{\text{SN}}\mathcal{L}_{\text{GRB}}$ by means of the Metropolis–Hastings algorithm implemented through the MONTE PYTHON code (Audren et al. 2013). In the numerical procedure, we assume uniform priors on the fitting parameters (see Table 1) and we take H_0 as the best-fitting value obtained from the model-independent analysis of the OHD data: $H_0 = 67.74 \text{ km s}^{-1} \text{ Mpc}^{-1}$. We summarize the results for the Λ CDM and w CDM models in Table 2. We show the marginalized 1σ and 2σ confidence contours in Fig. 4. One immediately sees that Ω_m in the Λ CDM model is unusually high compared to previous findings that use SNe Ia and other surveys different from GRBs. In fact, our result is in tension with Planck’s predictions (Planck Collaboration VI 2018) at $\geq 3\sigma$. However, our outcome is well consistent within 1σ with previous analyses that used GRBs (see. e.g. Amati & Della Valle 2013 for a review, and Izzo et al. 2015, Haridasu et al. 2017, and Demianski et al. 2017a,b for recent results). In addition, the tension is reduced as one considers the w CDM model, enabling w to vary. This does not indicate that w CDM is favoured with respect to the standard cosmological model. In fact, we immediately notice that w is consistent within 1σ with the Λ CDM case, i.e. $w = -1$.

We note that the numerical approach using the Metropolis–Hasting algorithm may suffer from some issues related to random walk behaviour. In the case of highly correlated statistical models, the use of more robust integration methods could alleviate many of those issues. Alternative approaches for the multilevel structure of the proper Bayesian model are left for a future study.

3.1 Statistical performances with GRBs

To test the statistical performance of the models under study, we apply the AIC criterion (Akaike 1974):

$$\text{AIC} \equiv 2p - 2 \ln \mathcal{L}_{\text{max}},$$

where p is the number of free parameters in the model and \mathcal{L}_{max} is the maximum probability function calculated at the best-fitting point. The best model is the one that minimizes the AIC value. We also use the DIC criterion (Kunz, Trotta & Parkinson 2006) defined as

$$\text{DIC} \equiv 2p_{\text{eff}} - 2 \ln \mathcal{L}_{\text{max}},$$

where $p_{\text{eff}} = \langle -2 \ln \mathcal{L} \rangle + 2 \ln \mathcal{L}_{\text{max}}$ is the number of parameters that a data set can effectively constrain. Here, the brackets indicate the average over the posterior distribution. Unlike the AIC and BIC criteria, the DIC statistics does not penalize for the total number of free parameters of the model, but only for those that are constrained by the data (Liddle 2007). We thus computed the differences with respect to the reference Λ CDM flat scenario. Both the AIC and DIC results indicate that the Λ CDM model is only slightly favoured with respect to the w CDM model (see Table 2).

4 FINAL OUTLOOKS AND PERSPECTIVES


In this work, we faced out the circularity problem in using GRBs as distance indicators. To do so, we employed the $E_p - E_{\text{iso}}$ (Amati) correlation and we proposed a new technique to build d_L in a model-independent way, using the OHD measurements. In particular, we considered the OHD data points and we approximated the Hubble function by means of a Bézier parametric curve obtained from the linear combinations of Bernstein’s polynomials. Assuming vanishing spatial curvature as suggested by Planck’s results, we were able to calibrate the Amati relation in a model-independent way. We thus obtained a new sample of distance moduli for 193 different GRBs (see Table A1).

We then used the new data sample to constrain two different cosmological scenarios: the concordance Λ CDM model, and the w CDM model, with the dark energy EoS parameter is free to vary. Hence, we performed a Monte Carlo integration through the Metropolis–Hastings algorithm on the joint likelihood function obtained by combining the GRB measurements with the SNe JLA data set. In our numerical analysis, we fixed H_0 to the best-fitting value obtained from the model-independent analysis over OHD data, i.e. $H_0 = 67.74 \text{ km s}^{-1} \text{ Mpc}^{-1}$. Our results for Ω_m and w agree with previous findings using GRBs and our treatment candidates as a severe alternative to calibrate the Amati relation in a model-independent form. Finally, we employed the AIC and BIC selection criteria to compare the statistical performance of the investigated models. We found that the Λ CDM model is preferred with respect to the minimal w CDM extension. Although a pure Λ CDM model is statistically favoured, we note that the values of Ω_m and w for the w CDM model are remarkably in agreement with those obtained by the Dark Energy Survey (Abbott et al. 2018). We can then conclude that no modifications of the standard paradigm are expected as intermediate redshifts are involved.

Future efforts will be dedicated to the use of our new technique to fix refined constraints over dynamical dark energy models. Also, we will compare our outcomes with respect to previous model-independent calibrations.

Table 2. 95 per cent confidence level results of the MCMC analysis for the SN + GRB data. The AIC and DIC differences are intended with respect to the Λ CDM model.

Model	w	Ω_m	M	Δ_M	α	β	Δ AIC	Δ DIC
Λ CDM	-1	$0.397^{+0.040}_{-0.039}$	$-19.090^{+0.037}_{-0.037}$	$-0.055^{+0.043}_{-0.043}$	$0.126^{+0.011}_{-0.012}$	$2.61^{+0.13}_{-0.13}$	0	0
w CDM	$-0.86^{+0.36}_{-0.38}$	$0.34^{+0.13}_{-0.15}$	$-19.079^{+0.046}_{-0.046}$	$-0.055^{+0.042}_{-0.042}$	$0.126^{+0.011}_{-0.012}$	$2.61^{+0.13}_{-0.13}$	1.44	1.24

**Figure 4.** Marginalized 1σ and 2σ contours, and posterior distributions from the MCMC analysis of SN + GRB data for the Λ CDM model (top) and for the w CDM model (bottom).

ACKNOWLEDGEMENTS

The authors are grateful to the anonymous referee for helpful suggestions that helped to improve the quality of the manuscript. RD and OL are grateful to Anna Silvia Baldi and Kuantay Boshkayev for useful discussions. LA acknowledges financial contribution from

the agreement ASI-INAF n.2017-14-H.O. The work was supported in part by the Ministry of Education and Science of the Republic of Kazakhstan, Program 'Fundamental and applied studies in related fields of physics of terrestrial, near-earth and atmospheric processes and their practical application' IRN: BR05236494.

REFERENCES

- DES Collaboration, Abbott T. M. C. DES Collaboration et al., 2018, preprint (arXiv:1811.02375)
- Amati L., 2006, *MNRAS*, 372, 233
- Amati L., Della Valle M., 2013, *Int. J. Mod. Phys. D*, 22, 1330028
- Amati L. et al., 2002, *A&A*, 390, 81
- Amati L., Guidorzi C., Frontera F., Della Valle M., Finelli F., Landi R., Montanari E., 2008, *MNRAS*, 391, 577
- Aubourg É. et al., 2015, *Phys. Rev. D*, 92, 123516
- Audren B., Lesgourgues J., Benabed K., Prunet S., 2013, *J. Cosmol. Astropart. Phys.*, 2, 001
- Bernardini M. G., Margutti R., Zaninoni E., Chincarini G., 2012, *MNRAS*, 425, 1199
- Betoule M. et al., 2014, *A&A*, 568, A22
- Capozziello S., Izzo L., 2008, *A&A*, 490, 31
- Capozziello S., D'Agostino R., Luongo O., 2018, *MNRAS*, 476, 3924
- Cucchiara A. et al., 2011, *ApJ*, 736, 7
- D'Agostini G., 2005, preprint (arXiv:physics/0511182)
- Dainotti M. G., Amati L., 2018, *PASP*, 130, 051001
- Dainotti M. G., Cardone V. F., Capozziello S., 2008, *MNRAS*, 391, L79
- Demianski M., Piedipalumbo E., Sawant D., Amati L., 2017a, *A&A*, 598, A112
- Demianski M., Piedipalumbo E., Sawant D., Amati L., 2017b, *A&A*, 598, A113
- Ghirlanda G., Ghisellini G., Lazzati D., Firmani C., 2004, *ApJ*, 613, L13
- Ghirlanda G., Ghisellini G., Firmani C., 2006, *New J. Phys.*, 8, 123
- Goobar A., Perlmutter S., 1995, *ApJ*, 450, 14
- Haridasu B. S., Luković V. V., D'Agostino R., Vittorio N., 2017, *A&A*, 600, L1
- Izzo L., Muccino M., Zaninoni E., Amati L., Della Valle M., 2015, *A&A*, 582, A115
- Jimenez R., Loeb A., 2002, *ApJ*, 573, 37
- Kodama Y., Yonetoku D., Murakami T., Tanabe S., Tsutsui R., Nakamura T., 2008, *MNRAS*, 391, L1
- Kunz M., Trotta R., Parkinson D., 2006, *Phys. Rev. D*, 74, 023503
- Liang N., Xiao W. K., Liu Y., Zhang S. N., 2008, *ApJ*, 685, 354
- Liddle A. R., 2007, *MNRAS*, 377, L74
- Luković V. V., D'Agostino R., Vittorio N., 2016, *A&A*, 595, A109
- Nava L. et al., 2012, *MNRAS*, 421, 1256
- Ooba J., Ratra B., Sugiyama N., 2018, *ApJ*, 864, 80
- Percival W. J. et al., 2010, *MNRAS*, 401, 2148
- Perlmutter S. et al., 1998, *Nature*, 391, 51
- Perlmutter S. et al., 1999, *ApJ*, 517, 565
- Phillips M. M., 1993, *ApJ*, 413, L105
- Planck Collaboration VI, 2018, preprint (arXiv:1807.06209)
- Riess A. G. et al., 1998, *AJ*, 116, 1009
- Riess A. G. et al., 2018, *ApJ*, 861, 126
- Rodney S. A. et al., 2015, *AJ*, 150, 156
- Salvaterra R. et al., 2009, *Nature*, 461, 1258
- Schaefer B. E., 2007, *ApJ*, 660, 16
- Schmidt B. P. et al., 1998, *ApJ*, 507, 46

Tanvir N. R. et al., 2009, *Nature*, 461, 1254Wei J.-J., Wu X.-F., Melia F., Wei D.-M., Feng L.-L., 2014, *MNRAS*, 439, 3329Yonetoku D., Murakami T., Nakamura T., Yamazaki R., Inoue A. K., Ioka K., 2004, *ApJ*, 609, 935

APPENDIX A:

Table A1. List of the full sample of GRBs used in this work and their redshift z and calibrated μ_{GRB} .

GRB	z	$\mu_{\text{GRB}} \pm \sigma_{\mu, \text{GRB}}$	GRB	z	$\mu_{\text{GRB}} \pm \sigma_{\mu, \text{GRB}}$	GRB	z	$\mu_{\text{GRB}} \pm \sigma_{\mu, \text{GRB}}$	GRB	z	$\mu_{\text{GRB}} \pm \sigma_{\mu, \text{GRB}}$
970228	0.695	43.76 \pm 0.77	051109A	2.346	47.73 \pm 0.89	090323	3.57	47.08 \pm 0.53	120909A	3.93	48.20 \pm 0.93
970508	0.835	44.64 \pm 0.73	060115	3.5328	47.67 \pm 1.07	090328	0.736	45.47 \pm 0.36	121128A	2.2	45.56 \pm 0.30
970828	0.958	43.87 \pm 0.51	060124	2.296	46.47 \pm 0.88	090418A	1.608	48.05 \pm 0.64	130408A	3.758	48.55 \pm 0.46
971214	3.42	47.97 \pm 0.51	060206	4.0559	49.11 \pm 1.22	090423	8.1	50.05 \pm 0.67	130420A	1.297	42.87 \pm 0.29
980613	1.096	46.06 \pm 1.11	060210	3.91	47.52 \pm 0.79	090424	0.544	42.67 \pm 0.30	130427A	0.3399	41.59 \pm 0.41
980703	0.966	45.09 \pm 0.37	060218	0.03351	34.60 \pm 0.42	090516	4.109	47.93 \pm 1.00	130505A	2.27	46.27 \pm 0.39
981226	1.11	44.03 \pm 1.13	060306	3.5	47.48 \pm 1.04	090618	0.54	40.54 \pm 0.30	130518A	2.488	46.31 \pm 0.37
990123	1.6	45.37 \pm 0.70	060418	1.489	45.89 \pm 0.64	090715B	3.	47.01 \pm 0.78	130701A	1.155	44.69 \pm 0.30
990506	1.3	43.74 \pm 0.57	060526	3.22	45.97 \pm 0.50	090812	2.452	48.61 \pm 0.88	130831A	0.4791	41.24 \pm 0.30
990510	1.619	45.14 \pm 0.34	060607A	3.075	47.57 \pm 0.65	090902B	1.822	46.04 \pm 0.40	131011A	1.874	44.68 \pm 0.39
990705	0.842	43.51 \pm 0.73	060614	0.125	38.88 \pm 2.58	090926	2.1062	44.75 \pm 0.35	131030A	1.295	43.82 \pm 0.31
990712	0.434	41.85 \pm 0.44	060707	3.424	47.74 \pm 0.68	090926B	1.24	44.51 \pm 0.30	131105A	1.686	45.26 \pm 0.59
991208	0.706	41.98 \pm 0.31	060729	0.543	42.55 \pm 1.22	091003	0.8969	45.53 \pm 0.52	131108A	2.4	47.08 \pm 0.36
991216	1.02	43.36 \pm 0.52	060814	1.9229	45.63 \pm 0.79	091018	0.971	42.94 \pm 1.05	131117A	4.042	46.99 \pm 0.46
000131	4.5	47.18 \pm 1.04	060908	1.8836	47.14 \pm 1.18	091020	1.71	46.52 \pm 0.39	131231A	0.642	41.55 \pm 0.30
000210	0.846	44.82 \pm 0.34	060927	5.46	47.84 \pm 0.70	091024	1.092	43.92 \pm 0.32	140206A	2.73	46.07 \pm 0.33
000418	1.12	43.97 \pm 0.35	061007	1.262	44.36 \pm 0.41	091029	2.752	46.12 \pm 0.68	140213A	1.2076	43.72 \pm 0.30
000911	1.06	45.76 \pm 0.58	061121	1.314	46.73 \pm 0.40	091127	0.49	39.90 \pm 0.31	140419A	3.956	47.83 \pm 0.84
000926	2.07	44.62 \pm 0.37	061126	1.1588	46.15 \pm 0.76	091208B	1.063	45.18 \pm 0.30	140423A	3.26	47.24 \pm 0.40
010222	1.48	44.52 \pm 0.34	061222A	2.088	46.87 \pm 0.51	100414A	1.368	45.92 \pm 0.37	140506A	0.889	45.78 \pm 0.99
010921	0.45	42.29 \pm 0.49	070125	1.547	45.08 \pm 0.44	100621A	0.542	41.88 \pm 0.41	140508A	1.027	43.69 \pm 0.32
011121	0.36	44.03 \pm 0.70	070521	1.35	45.67 \pm 0.36	100728A	1.567	44.83 \pm 0.34	140512A	0.725	44.31 \pm 1.41
011211	2.14	45.34 \pm 0.35	071003	1.604	47.79 \pm 0.45	100728B	2.106	47.03 \pm 0.39	140515A	6.32	49.32 \pm 0.71
020124	3.198	46.59 \pm 0.79	071010B	0.947	42.48 \pm 0.59	100814A	1.44	43.93 \pm 0.36	140518A	4.707	47.58 \pm 0.46
020405	0.69	43.02 \pm 0.31	071020	2.145	48.45 \pm 0.66	100816A	0.8049	45.58 \pm 0.31	140620A	2.04	45.44 \pm 0.30
020813	1.25	43.73 \pm 0.67	071117	1.331	46.77 \pm 1.30	100906A	1.727	43.93 \pm 0.41	140623A	1.92	48.26 \pm 1.12
020819B	0.41	41.07 \pm 0.75	080207	2.0858	45.41 \pm 1.78	101213A	0.414	43.63 \pm 1.00	140629A	2.275	46.46 \pm 0.50
020903	0.25	39.31 \pm 1.38	080319B	0.937	44.01 \pm 0.36	101219B	0.55	42.89 \pm 0.31	140801A	1.32	44.79 \pm 0.30
021004	2.3	46.86 \pm 1.06	080411	1.03	44.50 \pm 0.38	110106B	0.618	44.31 \pm 0.68	140808A	3.29	48.53 \pm 0.45
021211	1.01	44.21 \pm 0.97	080413A	2.433	47.75 \pm 0.78	110205A	2.22	46.20 \pm 0.98	140907A	1.21	45.20 \pm 0.30
030226	1.98	45.23 \pm 0.55	080413B	1.1	44.63 \pm 0.70	110213A	1.46	44.71 \pm 0.79	141028A	2.33	46.12 \pm 0.35
030323	3.37	48.08 \pm 1.06	080603B	2.69	46.77 \pm 1.02	110213B	1.083	43.79 \pm 0.43	141109A	2.993	47.27 \pm 0.71
030328	1.52	43.58 \pm 0.43	080605	1.64	46.00 \pm 0.65	110422A	1.77	43.76 \pm 0.32	141220A	1.3195	43.78 \pm 0.33
030329	0.1685	38.28 \pm 0.30	080607	3.036	47.24 \pm 0.44	110503A	1.613	45.56 \pm 0.34	141221A	1.452	46.71 \pm 0.47
030429	2.65	46.09 \pm 0.50	080721	2.591	47.33 \pm 0.46	110715A	0.82	43.42 \pm 0.30	141225A	0.915	45.43 \pm 0.41
030528	0.78	41.06 \pm 0.41	080804	2.2045	47.83 \pm 0.35	110731A	2.83	47.76 \pm 0.37	150206A	2.087	45.71 \pm 0.44
040912B	1.563	42.91 \pm 2.16	080913	6.695	50.73 \pm 1.26	110801A	1.858	45.94 \pm 1.06	150301B	1.5169	46.73 \pm 0.52
040924	0.859	43.48 \pm 0.81	080916A	0.689	44.44 \pm 0.30	110818A	3.36	48.79 \pm 0.56	150314A	1.758	45.62 \pm 0.35
041006	0.716	41.64 \pm 0.57	080928	1.6919	43.64 \pm 0.62	111107A	2.893	48.38 \pm 0.72	150323A	0.593	44.32 \pm 0.40
041219	0.31	40.24 \pm 0.65	081007	0.5295	42.92 \pm 0.59	111228A	0.716	40.63 \pm 0.34	150403A	2.06	46.20 \pm 0.42
050318	1.4436	44.22 \pm 0.52	081008	1.9685	45.26 \pm 0.49	120119A	1.728	45.22 \pm 0.34	150413A	3.139	45.92 \pm 0.98
050401	2.8983	46.12 \pm 0.61	081028	3.038	45.51 \pm 0.95	120326A	1.798	45.04 \pm 0.30	150514A	0.807	43.47 \pm 0.43
050416A	0.6535	41.8 \pm 0.54	081118	2.58	45.21 \pm 0.30	120624B	2.1974	45.89 \pm 0.47	150821A	0.755	44.07 \pm 1.20
050525A	0.606	42.13 \pm 0.36	081121	2.512	46.48 \pm 0.34	120711A	1.405	46.00 \pm 0.40	151021A	1.49	43.82 \pm 0.36
050603	2.821	47.76 \pm 0.37	081203A	2.05	47.99 \pm 1.28	120712A	4.1745	48.38 \pm 0.53	151027A	0.81	44.57 \pm 1.31
050820	2.615	47.03 \pm 0.55	081221	2.26	44.55 \pm 0.31	120716A	2.486	45.60 \pm 0.32	151029A	1.423	45.24 \pm 0.51
050904	6.295	50.94 \pm 0.88	081222	2.77	47.00 \pm 0.34	120724A	1.48	44.25 \pm 0.68			
050922C	2.199	47.20 \pm 0.73	090102	1.547	47.01 \pm 0.37	120802A	3.796	46.82 \pm 0.84			
051022	0.809	43.30 \pm 0.82	090205	4.6497	49.78 \pm 0.88	120811C	2.671	45.95 \pm 0.30			

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.