











<b>Publication Year</b>	2021
<b>Acceptance in OA</b>	2023-01-23T13:47:25Z
<b>Title</b>	A PSF-based Approach to TESS High quality data Of Stellar clusters (PATHOS) - IV. Candidate exoplanets around stars in open clusters: frequency and age-planetary radius distribution
<b>Authors</b>	Nardiello, D., Deleuil, M., Mantovan, G., Malavolta, L., Lacedelli, G., Libralato, M., BEDIN, Luigi, BORSATO, LUCA, Granata, V., Piotto, G.
<b>Publisher's version (DOI)</b>	10.1093/mnras/stab1497
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/32992">http://hdl.handle.net/20.500.12386/32992</a>
<b>Journal</b>	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
<b>Volume</b>	505

# A PSF-based Approach to TESS High quality data Of Stellar clusters (PATHOS) – IV. Candidate exoplanets around stars in open clusters: frequency and age–planetary radius distribution

D. Nardiello <sup>1,2★</sup> M. Deleuil,<sup>1</sup> G. Mantovan <sup>2,3</sup> L. Malavolta <sup>2,3</sup> G. Lacedelli <sup>2,3</sup> M. Libralato <sup>4</sup>  
L. R. Bedin <sup>2</sup> L. Borsato <sup>2</sup> V. Granata <sup>2,3</sup> and G. Piotto<sup>2,3</sup>

<sup>1</sup>Aix Marseille Université, CNRS, CNES, LAM, F-13007 Marseille, France

<sup>2</sup>Istituto Nazionale di Astrofisica - Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, I-35122 Padova, Italy

<sup>3</sup>Dipartimento di Fisica e Astronomia 'Galileo Galilei', Università di Padova, Vicolo dell'Osservatorio 3, I-35122 Padova, Italy

<sup>4</sup>AURA for the European Space Agency (ESA), ESA Office, Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

Accepted 2021 May 19. Received 2021 May 18; in original form 2021 February 15

## ABSTRACT

The knowledge of the ages of stars hosting exoplanets allows us to obtain an overview on the evolution of exoplanets and understand the mechanisms affecting their life. The measurement of the ages of stars in the Galaxy is usually affected by large uncertainties. An exception are the stellar clusters: For their coeval members, born from the same molecular cloud, ages can be measured with extreme accuracy. In this context, the project PATHOS is providing candidate exoplanets orbiting members of stellar clusters and associations through the analysis of high-precision light curves obtained with cutting-edge tools. In this work, we exploited the data collected during the second year of the *Transiting Exoplanet Survey Satellite* mission. We extracted, analysed, and modelled the light curves of  $\sim 90\,000$  stars in open clusters located in the Northern ecliptic hemisphere in order to find candidate exoplanets. We measured the frequencies of candidate exoplanets in open clusters for different orbital periods and planetary radii, taking into account the detection efficiency of our pipeline and the false positive probabilities of our candidates. We analysed the age– $R_p$  distribution of candidate and confirmed exoplanets with periods  $< 100$  d and well constrained ages. While no peculiar trends are observed for Jupiter-size and (super-)Earth-size planets, we found that objects with  $4 \lesssim R_p \lesssim 13 R_{\text{Earth}}$  are concentrated at ages  $\lesssim 200$  Myr; different scenarios (atmospheric losses, migration, etc.) are considered to explain the observed age– $R_p$  distribution.

**Key words:** techniques: image processing – techniques: photometric – planets and satellites: general – stars: variables: general – open clusters and associations: general.

## 1 INTRODUCTION

In Summer 2020, the *Transiting Exoplanet Survey Satellite* (TESS; Ricker et al. 2015) concluded its main mission after about 2 yr of observations. In this period, the spacecraft has observed millions of stars in about  $\gtrsim 70$  per cent of the sky with an unprecedented photometric precision and temporal coverage, and new data from the extended mission, characterized by (in part) a new observing strategy, are coming.

Stellar clusters and associations offer the unique opportunity to derive precise stellar parameters (like radius, mass, chemical content, and especially age) for their members simply using theoretical models. During the main mission TESS observed many hundreds stellar (open and globular) clusters and associations in sectors of  $\sim 27$  d. However, the low resolution of the four cameras ( $\sim 21$  arcsec pixel<sup>-1</sup>) makes the extraction of high-precision light curves difficult for the stars located in these dense regions.

The project ‘a PSF-based Approach to TESS High quality data Of Stellar clusters’ (PATHOS; Nardiello et al. 2019, hereafter **Paper I**) was born to exploit the TESS data in order to extract high-precision photometry for members of stellar clusters and associations adopting an innovative approach, based on the use of empirical point spread functions (PSFs) and neighbour subtraction. Scope of the project is the discovery and characterization of new candidate exoplanets in stellar clusters and the analysis of possible correlations between well-measured star properties and candidate exoplanet characteristics. Field stars’ age is usually affected by large uncertainties, but it is also an essential information to constrain the formation and understand the evolution of exoplanets, like for example how and on which temporal scales the mechanisms that bring to the atmosphere evaporation of low-mass close-in exoplanets happen (Lammer et al. 2003; Baraffe et al. 2005; Murray-Clay, Chiang & Murray 2009; Owen & Jackson 2012; Owen & Wu 2017; Wu 2019; Owen 2020). In this context, the PATHOS project is providing interesting candidate exoplanets orbiting stars with well constrained ages. Moreover, the high-precision light curves generated in our project, and publicly available to the astronomical community, allow us to obtain results not only in the research field of exoplanets, but also in other fields like

\* E-mail: domenico.nardiello@lam.fr

asteroseismology (Mackereth et al. 2021), or the analysis of the spin axis orientations of cluster members (Healy & McCullough 2020).

We already successfully applied the PATHOS pipeline in Paper I, when we studied the stars in an extremely crowded region containing the globular cluster 47 Tuc and the Small Magellanic Cloud. In Nardiello et al. (2020, hereafter Paper II), we extracted and analysed the light curves of open cluster members located in the Southern ecliptic hemisphere, finding 33 objects of interest and deriving a first estimate of exoplanet frequency in open clusters. Nardiello (2020, hereafter Paper III) studied the light curves of the members of five young associations, having ages  $\lesssim 10$  Myr; in particular, the author performed a gyrochronological analysis of association members to constrain the age of the stars, analysed the dust in the circumstellar discs of the young members and identified and characterized six strong candidate exoplanets.

In this work, we exploited the *TESS* data collected during Cycle 2 (Sectors 14–26) to obtain high-precision light curves of cluster members in the Northern ecliptic hemisphere by using our cutting-edge tools (Section 2), find and characterize candidate exoplanets in stellar clusters (Section 3), and analyse their frequency and properties as a function of host stars’ characteristics (Sections 4). We summarized and discussed the joined results obtained in this work and in Paper II in Section 5.

## 2 OBSERVATION AND DATA REDUCTION

In this work, we extracted and analysed the light curves of the stars likely members of Northern ecliptic hemisphere open clusters observed by *TESS* during the second year of the mission. The observations used in this work were carried out between 2019 July 18 and 2020 July 4 ( $\sim 352$  d), and are divided into 13 sectors (Sectors 14–26); in Sectors 21, 22, and 23, no open clusters fell in the *TESS* field of view and therefore the final number of analysed sectors is 10.

For the light-curve extraction and correction, we used the PATHOS pipeline described in detail in Papers I and II. We extracted the light curves of stars in a given catalogue from *TESS* full frame images (FFIs) by using the light-curve extractor IMG2LC. This software was developed by Nardiello et al. (2015a, 2016a) for ground-based observations, and it is a versatile tool that can be used with photometric time-series collected also with space-based observatories (see, e.g. Libralato et al. 2016a,b; Nardiello et al. 2016b).

The three main inputs of our PSF-based approach are (i) FFIs, (ii) PSFs, and (iii) input catalogue. For each star in the input catalogue, the light-curve extractor searches for the neighbours within a radius of 20 *TESS* pixels in the *Gaia* DR2 catalogue (Gaia Collaboration et al. 2018), transforms their positions and luminosities in the reference system of the FFI, models them by using a local PSF and then subtracts them from the FFI. Finally, it extracts PSF-fitting and aperture (1-, 2-, 3-, and 4-pixel radius) photometries of the target star from the neighbour-subtracted FFI. This approach has two advantages: (i) It minimizes the dilution effects due to the neighbour contaminants, and (ii) it allows us the extraction of high-precision photometry for stars in the *TESS* faint regime of magnitudes ( $T \gtrsim 15$ ; see, e.g. Apai, Nardiello & Bedin 2021). We corrected the extracted raw light curves for systematic effects by fitting and applying the Cotrending Basis Vectors, as widely discussed in Papers I and II.

As in Paper II, we used as input list the catalogue of cluster members published by Cantat-Gaudin et al. (2018); this catalogue contains the positions, colours, magnitudes, proper motions, parallaxes, and membership probabilities of likely members in 1229

stellar clusters. From this catalogue, we selected all the stars that satisfy these two conditions: (i) magnitude  $G < 17.5$ , because stars with larger magnitude are too faint to be detected by *TESS*; and (ii) ecliptic latitude  $\beta > 4^\circ$ , which corresponds to the part of the Northern ecliptic hemisphere covered by *TESS*.<sup>1</sup>

Fig. 1 shows the 126 372 stars (red points) in the input catalogue overlapped with the *TESS* fields of view (grey points): About 1/3 of them fall outside the *TESS* observations. We extracted 150 216 light curves of 89 858 stars in 411 clusters; about 50.3 per cent, i.e. 45 182 stars, were observed in only one sector, 30 957 stars ( $\sim 34.4$  per cent) were observed in two sectors, 11 844 ( $\sim 13.2$  per cent) in three sectors, and 1 875 ( $\sim 2.1$  per cent) in four or more sectors.

Light curves are released on the Mikulski Archive for Space Telescopes (MAST) as a High Level Science Product (HLSP) under the project PATHOS<sup>2</sup> (DOI: 10.17909/t9-es7m-vw14). A detailed description of the light curves (that are both in `ascii` and `fits` format) is reported in Papers I and II and in the MAST web page of the PATHOS project.

### 2.1 Photometric precision

We explored two different quality parameters, already defined in Papers I, II, and III, to identify for each star the photometric method that gives the best light curve.


The first quality parameter is the simple `rms`, defined as the 68.27th percentile of the  $3.5\sigma$ -clipped sorted residual from the median value. This parameter is sensitive to the (high) variability of some stars, and, for this reason, is not recommended to estimate the photometric precision of the light curve. The mean trends of the `rms` as a function of the *TESS* magnitude  $T$  for the five photometric methods are reported in Fig. 2 (top panel).

The second quality parameter is the `P2P rms`, defined as the 68.27th percentile of the  $3.5\sigma$ -clipped sorted residual from the median value of the vector  $\delta F_j = F_j - F_{j+1}$ , with  $F$  the flux at a given epoch  $j$ . This parameter is not sensitive to the intrinsic luminosity variations of the stars, and we used it to define the interval in which each photometric method works, on average, better than the others. From the mean trends shown in bottom panel of Fig. 2, we found (confirming the results obtained in the previous works) that for stars with  $5.5 \lesssim T \lesssim 7.0$ , the aperture photometry with radius 4-pixel gives the best results; the best photometric methods for the intervals  $7.0 \lesssim T \lesssim 9.0$ ,  $9.0 \lesssim T \lesssim 10.0$ , and  $10.0 \lesssim T \lesssim 13.0$  are 3-pixel, 2-pixel aperture, and PSF-fitting photometries, respectively. For faint stars with  $T \gtrsim 13.0$ , the 1-pixel aperture photometry gives the lower `P2P rms`.


In the following analysis, we used, for each star of magnitude  $T_*$ , the light curve associated with the photometric method that has the lower mean `P2P rms` in  $T_*$ . We excluded from the analysis all the stars whose mean light-curve instrumental magnitude ( $T_{\text{instr}}$ ) is too different from the expected magnitude  $T_{\text{calib}}$ , following this procedure: We extracted the  $\delta T = T_{\text{instr}} - T_{\text{calib}}$  distribution, we calculated its mean ( $\delta \bar{T}$ ) and the standard deviation ( $\sigma_{\delta T}$ ), and we excluded the  $i$ th light curve if  $|\delta T_i - \delta \bar{T}| > 4\sigma_{\delta T}$ . We also excluded all the light curves that have  $< 75$  per cent of well-measured points (i.e. `DQUALITY=0` and `FLUX≠ 0`). The final number of analysed light curves is 138 924 associated with 84 967 stars.

<sup>1</sup>As also shown in Fig. 1, during Sectors 14–16 and 24–26, the *TESS* pointings were modified in order to avoid excessive contamination by stray Earth- and Moon-light in cameras 1 and 2.

<sup>2</sup><https://archive.stsci.edu/hlsp/pathos>.



**Figure 1.** Aitoff projection in ecliptic coordinates of the fields observed by *TESS* in the first 2 yr of mission and of the open cluster members analysed in the PATHOS project: Grey points represent the sources observed in 2-min cadence mode in Sectors 1–26, and blue and red points are the stars in the input list used in Paper II and in this work, respectively.



**Figure 2.** Mean trends of the photometric  $r_{\text{rms}}$  (top-panel) and P2P  $r_{\text{rms}}$  (bottom panel) as a function of the *TESS* magnitude  $T$  for different photometric methods: Black lines are associated with PSF-fitting photometry, and magenta, blue, green, and yellow lines are associated with 1-, 2-, 3-, and 4-pixel aperture photometries, respectively. As an example, the rms distributions obtained with 3-pixel and PSF-fitting photometry are shown in light green and grey crosses, respectively (for clarity, only 10 per cent of the stars are plotted). Red starred symbols represent the saturated stars. The dashed line is the theoretical limit calculated as in Paper II.

### 3 CANDIDATE EXOPLANETS: SEARCHING, VETTING, AND CHARACTERIZATION

We searched for signals of transiting objects among the selected light curves following the procedure described in Papers II and III. Briefly, we removed the intrinsic stellar variability interpolating the light curve with a fifth order spline defined on  $N_{\text{knots}}$  knots. In order to model short- and long-period variability, we considered two different grids of knots, with knots spaced by 6.5 and 13.0 h, respectively. The grids of knots are defined on continuous parts of light curves that does not present ‘breaks’  $>0.5$  d, in order to avoid the introduction of artifacts in the flattened light curve. We also removed from the light curves the photometric points associated with high values of the local sky ( $>5\sigma_{\text{SKY}}$  from the mean local background), with `DQUALITY`  $>0$  and  $4\sigma$  above the median normalized flux. We extracted the transit-fitting least-squares (TLS) periodograms of the flattened light curves adopting the `PYTHON` package `TLS`<sup>3</sup> (Hippke & Heller 2019), and searched for transit signals with period  $0.6 \text{ d} \leq P \leq T_{\text{LC}}$ , with  $T_{\text{LC}}$  being the temporal length of the light curve. We used the output parameters for the first selection of candidate transiting objects, as follows: (i) We selected the stars associated with a depth of the transit  $\delta_t < 10$  per cent and to a significance between odd and even transits  $\sigma_{\text{odd-even}} < 2.5$ ; (ii) we divided the signal detection efficiency (SDE) and the signal-to-noise ratio (SNR) distributions in bins of periods  $\delta P = 0.5$  d, and calculated the  $3.5\sigma$ -clipped mean and standard deviation of SDE ( $\overline{\text{SDE}}$  and  $\sigma_{\text{SDE}}$ ) and SNR ( $\overline{\text{SNR}}$  and  $\sigma_{\text{SNR}}$ ) within each bin. Then we interpolated the binned points  $\overline{\text{SDE}} + 3.5\sigma_{\text{SDE}}$  and  $\overline{\text{SNR}} + 3.5\sigma_{\text{SNR}}$  with a spline, and we selected as good candidates all the stars above these splines. Panels (a) of Fig. 3 show an example of selections based on the output parameters of TLS. We visually inspected the light curves that passed the above selections to exclude false positives due to the presence of artefacts. We applied the above procedure both to the light curves obtained in each single sector, and then to the stacked light curves of stars observed in more than one sector, in order to increase the efficiency of transit detection. The number of stars that passed the first-step selection is 279 ( $\sim 0.3$  per cent).

These candidates were subjected to a series of vetting tests widely described in the previous papers of the PATHOS series. They are as follows: (i) Check for the transit depths in the light curves obtained with different photometric methods; (ii) check for the presence of secondary eclipses in the light curves phased with a period  $0.5 \times P$ ,  $1.0 \times P$ , and  $2.0 \times P$ , with  $P$  the period found by the TLS routine; (iii) check for the transit depths by comparing binned odd/even transits (panels b of Fig. 3); and (iv) check for contamination through the analysis of the in/out-of-transit difference centroid (panel c of Fig. 3). After this second-step selection, 39 transiting objects of interest survived ( $\sim 0.05$  per cent); one of these objects showed only one transit in its light curve.

#### 3.1 TESS Objects of Interest

We cross-matched the *TESS* Objects of Interest (TOI) list<sup>4</sup> with our input catalogue of cluster members. Four candidates, found by the Quick-Look Pipeline (QLP; Huang et al. 2020), are also in our input catalogue, but only one of them (TOI-1535, TIC 420288086) is in our final list of transiting objects of interest. Two of them (TOI-1497, TIC 371673488 and TOI-1321, TIC 195199644) were not detected by our pipeline because no transit signals are present in the light curves we analysed, even if the mean scatter of our

light curves is lower than the scatter of the light curves shown in the QLP data validation report. We checked the notes about these two candidates on ExoFOP<sup>5</sup>: (i) the depth-aperture correlation for TOI-1497.01, reported in a note on ExoFOP, is confirmed by our analysis; (ii) for TOI-1321, a depth-aperture correlation is also reported; moreover, a note associated with a photometric follow-up with MuSCAT2 reports a deep transit signal from a nearby star at  $\sim 1$  arcmin from TOI-1321. Therefore, both these candidates are likely contaminated by neighbour sources. The fourth QLP candidate, TOI-1188 (TIC 15247229), was excluded from our final list after the centroid analysis. Its vetting tests are reported in Fig. 3. Our conclusion is also supported by the notes reported in the photometric follow-up section of the ExoFOP website.

#### 3.2 Stellar parameters

We fitted theoretical models from the last release of BaSTI (‘a Bag of Stellar Tracks and Isochrones’) models (Hidalgo et al. 2018) to the CMDs of the 32 open clusters that host the stars associated with the 39 transiting objects of interest. In this way, we were able to extract primary information (stellar radius, mass, density, effective temperature) of the stars that host candidate transiting objects. Because metallicity measurements are not available for the large part of the clusters and because open clusters have, on average, metallicities similar to that of the Sun, in our fit, we used isochrones with  $[\text{Fe}/\text{H}] = 0.0 \pm 0.3$  as already done in Paper II, and we added the contribution of the uncertainties on the metallicity to the final errors on the stellar parameter estimates.

We transformed the isochrones from the theoretical to the observational plane using the distance modulus of the clusters obtained by Cantat-Gaudin et al. (2018), and the reddening and ages measured by Kharchenko et al. (2016), Röser et al. (2016), and Bossini et al. (2019); since some of the catalogues do not provides error estimates, for homogeneity, we used a conservative error of 10 per cent on the age and reddening values. Gulliver 49 is an open cluster discovered by Cantat-Gaudin et al. (2018), and no age estimate is provided in literature: We followed the technique by Nardiello et al. (2015b) based on the use of the  $\chi^2$ -minimization between isochrones and fiducial lines to derive an estimate of the cluster age. We found an age of  $200 \pm 20$  Myr.

Clusters’ parameters adopted for the isochrone fitting are reported in Table 1. Stellar parameters of the transiting candidates’ hosts obtained from isochrone fitting were used as priors in transit modelling described in the next section, and are reported in Table A1.

#### 3.3 Transit modelling

We modelled the transits of the objects of interest using the `PYTHON` package `PYORBIT`<sup>6</sup> (Malavolta et al. 2016, 2018, see also Benatti et al. 2019; Carleo et al. 2021; Lacedelli et al. 2021), based on the combined use of the package `BATMAN` (Kreidberg 2015), the global optimization algorithm `PYDE`<sup>7</sup> (Storn & Price 1997), and the affine invariant Markov chain Monte Carlo sampler `EMCEE` (Foreman-Mackey et al. 2013).

For the transit modelling, we included the central time of the first transit ( $T_0$ ), the period ( $P$ ), the impact parameter ( $b$ ), the planetary-to-stellar-radius ratio ( $R_p/R_*$ ), the stellar density ( $\rho_*$ ), and the dilution


<sup>3</sup>`TLS` v. 1.0.24; <https://github.com/hippke/tls>.

<sup>4</sup><https://tess.mit.edu/toi-releases/go-to-alerts/>.

<sup>5</sup><https://exofop.ipac.caltech.edu/tess>.

<sup>6</sup><https://github.com/LucaMalavolta/PyORBIT>.

<sup>7</sup><https://github.com/hpparvi/PyDE>.



**Figure 3.** Candidate selection and vetting in the case of the star TIC 15247229 (TOI-1188). Panels (a<sub>1</sub>) and (a<sub>2</sub>) show the TLS SNR and SDE versus detected period, respectively, while panel (a<sub>3</sub>) reports the  $\sigma_{\text{odd-even}}$  as a function of  $\delta_t$ ; grey points are all the analysed stars, orange points the light curves that passed the selection described in the text, and the blue circle is TOI-1188. In panels (b<sub>1</sub>) and (b<sub>2</sub>), we compare the binned odd (azure) and even (green) transits of the candidate. Panel (c) represents the analysis of the in-/out-of-transit difference centroid: The centroid is shifted on a neighbour star that contaminates the target. Panel (d) is the  $G$  versus  $(G_{\text{BP}} - G_{\text{RP}})$  colour–magnitude diagram (CMD) of Roslund 6, the open cluster that hosts TOI-1188 (blue circle). Panel (e) explains the procedure of flattening of the light curve: Grey points form the original, cleaned light curve; red line is the model defined by a spline on knots spaced by 13.0 h; and black points are the flattened light curve. Azure and green triangles indicate the odd and even transits, respectively.

factor ( $df$ ). The latter quantity is included as a free parameter, with a Gaussian prior obtained considering all the stars in the *Gaia* DR2 catalogue that fall in the same pixel of the target,<sup>8</sup> and transforming

<sup>8</sup>In such crowded environments, the completeness of the *Gaia* DR2 catalogue is  $\gtrsim 80$ –90 per cent for stars with  $16 \lesssim G \lesssim 18$  and  $\gtrsim 95$  per cent for brighter stars (see *Gaia* DR2 Documentation release 1.2, Section 10.7.4), and therefore the values of dilution factor obtained in this work represent, within the errors, a good estimate of the real dilution factor. In the case of close binaries with separation  $< 1$ –2 arcsec, Ziegler et al. (2018) demonstrated that *Gaia* satellite is not always able to resolve the components. If two (or more stars) are not resolved by *Gaia*, the result will be a single source whose flux will approximately be equal to the sum of the fluxes coming from the different not-resolved components; this does not affect the subtraction process

their *Gaia* magnitudes in *TESS* magnitudes adopting the equations by Stassun et al. (2019). Host star parameters, like the stellar radius ( $R_*$ ), mass radius ( $M_*$ ), gravity ( $\log g$ ), and effective temperature ( $T_{\text{eff}}$ ) come from the isochrone fits described in the previous section. On the basis of  $\log g$  and  $T_{\text{eff}}$ , we obtained information on the limb-darkening (LD) by using the grid of values published by Claret (2018); we adopted the LD parametrization described by Kipping

in the extraction of the light curve because we have considered the multiple components as a single source whose flux is the sum of components’ fluxes. It might affect the estimate of the dilution factor used for the modelling of the transits, and, for this reason, further AO follow-up of the candidates are mandatory.

**Table 1.** Cluster parameters.

Cluster name	Age (Myr)	Distance (pc)	$E(B - V)$	Reference
ASCC 13	44 ± 4	1078 ± 105	0.22 ± 0.02	(1)
Alessi 37	133 ± 13	707 ± 46	0.25 ± 0.03	(1)
Alessi Teutsch 5	74 ± 7	876 ± 70	0.50 ± 0.05	(2)
Czernik 44	32 ± 3	4696 ± 1500	1.13 ± 0.11	(2)
FSR 0342	376 ± 38	2687 ± 570	0.65 ± 0.07	(2)
Gulliver 49	200 ± 20	1622 ± 425	1.15 ± 0.12	(4)
IC 1396	1 ± 1	913 ± 76	0.42 ± 0.04	(2)
King 5	1230 ± 123	2523 ± 510	0.67 ± 0.07	(2)
King 6	382 ± 38	727 ± 50	0.59 ± 0.06	(1)
King 20	349 ± 35	1093 ± 305	0.67 ± 0.07	(1)
NGC 225	179 ± 18	684 ± 44	0.27 ± 0.03	(3)
NGC 457	24 ± 2	2882 ± 650	0.60 ± 0.06	(2)
NGC 752	1479 ± 148	441 ± 20	0.05 ± 0.01	(1)
NGC 884	16 ± 2	2341 ± 445	0.56 ± 0.06	(2)
NGC 1027	355 ± 35	1097 ± 98	0.45 ± 0.05	(2)
NGC 6811	863 ± 86	1112 ± 101	0.07 ± 0.01	(1)
NGC 6871	10 ± 1	1841 ± 285	0.60 ± 0.06	(2)
NGC 6910	34 ± 3	1350 ± 260	1.20 ± 0.12	(2)
NGC 6940	1023 ± 102	1025 ± 94	0.21 ± 0.02	(1)
NGC 6997	552 ± 55	865 ± 70	0.53 ± 0.05	(1)
NGC 7024	266 ± 27	1182 ± 130	0.63 ± 0.06	(2)
NGC 7086	116 ± 12	1616 ± 225	0.77 ± 0.08	(2)
NGC 7142	1778 ± 178	2376 ± 446	0.45 ± 0.05	(2)
NGC 7209	341 ± 34	1178 ± 125	0.18 ± 0.02	(1)
NGC 7245	355 ± 35	3307 ± 882	0.48 ± 0.05	(2)
NGC 7510	50 ± 5	3177 ± 765	0.95 ± 0.09	(2)
NGC 7654	79 ± 8	1600 ± 220	0.65 ± 0.07	(2)
NGC 7789	1841 ± 184	2074 ± 366	0.22 ± 0.02	(2)
RSG 5	50 ± 5	336 ± 11	0.04 ± 0.00	(3)
RSG 8	316 ± 32	446 ± 20	0.04 ± 0.00	(3)
SAI 25	243 ± 24	2194 ± 498	1.17 ± 0.12	(2)
SAI 149	251 ± 25	3000 ± 690	1.24 ± 0.12	(2)

*References.* (1) Bossini et al. (2019); (2) Kharchenko et al. (2016); (3) Röser, Schilbach & Goldman (2016); (4) this work.

(2013). In the modelling process, the routine takes into account the local variability of the star by fitting a second-degree polynomial to the out-of-transit part of the light curve. The routine modelled the transits with a fixed circular orbital eccentricity ( $e = 0$ ), and taking into account the 30-min cadence of the *TESS* FFIs (Kipping 2010).

The adopted priors on stellar parameters are reported in Table A1. The package `PYORBIT` explored all the parameters in linear space. In the `EMCEE` run, the number of walkers used is 10 times the number of free parameters. We ran, for each model, the sampler for 80 000 steps, excluding the first 15 000 steps as burn-in and using a thinning factor of 100. Fig. 4 shows an example of the modelling process in the case of *PATHOS-53*, a mono-transit object of interest.

In Table 2, we report the results of the transit fitting; Figs A1–A3 give an overview on the main properties of each transiting object of interest (position on the CMD, proper motions, in-/out-of-transit centroid analysis, transit modelling).

## 4 RESULTS

We improved the calculation of the frequencies of exoplanets reported in Paper II, taking into consideration the detection efficiency of our method and the false positive probabilities (FPPs) of our candidates. In this analysis, we considered candidates identified in this work and in Paper II, excluding all the *PATHOS* objects with  $R_p > 2.5R_J$  (because of their doubtful planet nature): 23 candidates survived after this selection (14 and 9 in the Southern and Northern ecliptic hemispheres, respectively).

### 4.1 Detection efficiency


We calculated the detection efficiency of our finding pipeline injecting transit signals of planets having radii  $R_p$  in a sample of light curves extracted randomly from the collection of light curves analysed in Paper II and in this work.

To inject the transit signals in each light curve, we roughly estimated the stellar radius and mass of each the 232 669 stars analysed in Paper II and in this work by using their absolute  $M_{G,0}$  versus  $(G_{BP} - G_{RP})_0$  CMD. To obtain the absolute CMD, we transformed the apparent magnitudes into absolute magnitudes by using the *Gaia* DR2 distances obtained by Bailer-Jones et al. (2018). For each star, we corrected the effects of the extinction on the colour and the magnitude of the stars calculating the reddening value  $[E(B - V)]$  by using the `PYTHON` routine `mw dust`<sup>9</sup> (Bovy et al. 2016) and the `Combined19` dustmap (Drimmel, Cabrera-Lavers & López-Corrodoira 2003; Marshall et al. 2006; Green et al. 2019), and the colour-dependent equation and the coefficients reported by Bossini et al. (2019). Fig. 5 shows the  $M_{G,0}$  versus  $(G_{BP} - G_{RP})_0$  CMD for all the stars in our sample; red starred points indicate the 23 candidates. We selected the likely main-sequence stars as follows: We performed a first guess selection of the main-sequence stars by hand, excluding all the stars that clearly are evolved stars (subgiant and red-giant stars); in the second step, we calculated the fiducial line of the likely main-sequence stars by using the naive estimator (Silverman 1986; see also Nardiello et al. 2015b for the description of the method) and we selected the stars whose colours are within  $2\sigma$  from the mean colour of the fiducial line. We calculated the radius and the mass of each star in our catalogue by using the main-sequence points of the `PARSEC` (Girardi et al. 2002; Bressan et al. 2012; Marigo et al. 2017) isochrones with ages between 10 and 1000 Myr. In Fig. 5, we reported as reference the mean  $M_{G,0}$  absolute magnitude for stars having radius  $R_* = 0.5, 1.0, 1.5,$  and  $2.5 R_\odot$ .

Because our candidate exoplanets orbit stars with  $R_* \lesssim 2.5 R_\odot$ , we first selected the light curves associated with these stars; we divided our sample of light curves on the basis of the *TESS* magnitude of the associated stars, considering bins of size  $\Delta T = 1.0$  in the magnitude range  $6.0 \leq T \leq 18.0$ . In each bin, we randomly caught 350 light curves<sup>10</sup> associated with stars having the previously estimated radii and masses, and, by using the routine `INJECTTRANSIT` implemented in `VARTOOLS` v.1.39 (Hartman & Bakos 2016), we injected in each light curve a periodic transit signal of a planet having radius  $R_p^{\text{inj}}$ , orbital period  $P^{\text{inj}}$ , random inclination  $i^{\text{inj}}$  (with the constraint that there must be a transit), and eccentricity  $e = 0$ . We considered 12 different cases in which  $R_p^{\text{inj}}$  and  $P^{\text{inj}}$  randomly vary between a given minimum and a maximum; (i) we first injected (super-)Earth-size planets having radii  $0.85 \leq R_p^{\text{inj}} \leq 3.9 R_{\text{Earth}}$ ; (ii) the second case we considered was for (super-)Neptune planets having radii  $3.9 < R_p^{\text{inj}} \leq 11.2 R_{\text{Earth}} \sim 1 R_J$ ; finally, we considered giant planets with radii  $1.0 < R_p^{\text{inj}} \leq 2.5 R_J$ . For each  $R_p^{\text{inj}}$  case, we considered random orbital periods in the following intervals: (a) short periods  $0.5 \leq P^{\text{inj}} \leq 2.0$  d; (b) short-medium periods  $2.0 < P^{\text{inj}} \leq 10.0$  d; (c) long-medium periods  $10.0 < P^{\text{inj}} \leq 85.0$  d; and (d) long periods  $85.0 < P^{\text{inj}} \leq 365.0$  d. We have not made any selection on the time length of the light curves, so, on the basis of the associated star observability, the light curve can span randomly between  $\sim 27$  and  $\sim 365$  d.

<sup>9</sup><https://github.com/jobovy/mwdust>.

<sup>10</sup>If the number of light curves in the considered magnitude bin was  $< 350$ , we considered all the light curves belonging to that bin.



**Figure 4.** Overview on the transit modelling of PATHOS-53: The left-hand panel shows the  $G$  versus  $(G_{BP} - G_{RP})$  CMD of the members of the open cluster RSG 5 and the isochrone fit (in magenta,  $t_{AGE} = 50$  Myr) used to derive the stellar parameters of PATHOS-53 (red point). The top right-hand panel shows the light curve of PATHOS-53 collected in Sectors 14 and 15; only one transit is detected (pointed by the green triangle). The bottom right-hand panel shows the model fit (green line) performed by `PYORBIT` on the single transit.

We injected the transit signals in the raw light curves and we followed the same pipeline for the correction of the light curves and the detection of transit signals described in Sections 2 and 3. We considered the injected planet as recovered if it passes the selections illustrated in Section 3 and in Fig. 3 and if  $|P^{inj} - P_{TLS}^{inj}| < 4.0\sigma(P_{TLS}^{inj})$ , where  $P_{TLS}^{inj}$  is the period obtained by the extraction of the TLS periodogram, and  $\sigma(P_{TLS}^{inj})$  its error. We also considered the planet as recovered if  $P_{TLS}^{inj}$  is equal to  $0.5 \times P^{inj}$  or  $2.0 \times P^{inj}$ , within  $4.0\sigma(P_{TLS}^{inj})$ . On a sample of 100 light curves selected randomly from those that passed the selections, we performed the vetting tests described in Section 3<sup>11</sup>: all the selected objects passed the tests. We finally calculated the detection efficiency in each magnitude interval  $\Delta T$  as the ratio  $N_{rec}/N_{inj}$ , where  $N_{rec}$  is the number of simulated planets recovered, and  $N_{inj}$  the number of light curves in which we injected transit signals. In Fig. 6 and Table 3, we reported the detection efficiency in the  $P^{inj}$  versus  $T$  grid for the three different planetary radius intervals: For Earth-size planets with  $R_p \lesssim 3.9R_{Earth}$ , the detection efficiency is low ( $\lesssim 40$  per cent), mainly because the large part of low-mass stars, for which the detection of small size planets is easier, is concentrated at magnitudes  $T \gtrsim 11$  (see fig. 8 of Paper II); for short period Neptune-size planets ( $P < 10$  d) orbiting stars with magnitude  $T \lesssim 12$ , the detection efficiency is  $\gtrsim 30$ –50 per cent, and decreases at fainter magnitude to 20–30 per cent; we obtained a detection efficiency always  $\gtrsim 30$  per cent for giant planets with periods  $\lesssim 10$  d. For periods  $P \gtrsim 10$  d, the detection efficiencies are always  $\lesssim 20$  per cent: This is an effect caused by the fact that only  $\sim 15$  per cent of the stars in our sample are observed in

more than two sectors, the detection of long-period planets difficult. Finally, even if we used different grids to flat the light curves to take into account the different kinds of variability, the detection efficiency can be lower for very active stars in young stellar clusters.

## 4.2 FPP estimation


We used the tool `VESPA` v. 0.6<sup>12</sup> (Morton 2012, 2015) to estimate the FPP in a Bayesian framework of each of the 23 candidate exoplanets previously selected. This software estimates the probability that the transit signal detected in a light curve is due to a real planet or to a different source. Different scenarios are taken into consideration to explain the signal: (i) simple eclipsing binary (EB, single/double period); (ii) a hierarchical triple system where two components eclipse (HEB, single/double period); (iii) a blended eclipsing binary inside the photometric aperture of the target (BEB, single/double period); and (iv) a transiting planet on the target star (pl). We want to highlight that `VESPA` does not consider the scenario in which transit signals are due to non-astrophysical sources, even if in our case the probability it happens is low, because the analysed candidates passed a series of vetting tests that allow to exclude signals due to systematic effects. For each configuration, `VESPA` simulates a representative stellar population, constrained by the information we have from the isochrone fitting and the light-curve modelling; in particular, we put constraints on the  $G$ ,  $G_{BP}$ ,  $G_{RP}$ ,  $J_{2MASS}$ ,  $H_{2MASS}$ ,  $K_{2MASS}$  magnitudes of the target star, the  $(\alpha, \delta)$  coordinates of the target, the effective temperature  $T_{eff}$ , the density  $\rho_*$ , and the surface gravity  $\log(g)$  of the star, the orbital period  $P$  of the candidate exoplanet, the planet-to-star radius ratio  $R_p/R_*$ . Moreover, we gave as an input the light

<sup>11</sup>The analysis of the centroid was excluded from these vetting tests because it is obtained analysing the images, where there are no signals for the simulated transits.

<sup>12</sup><https://github.com/timothydmorton/VESPA>.

**Table 2.** Results of transit modelling.

TIC	PATHOS	Cluster	$P$ (d)	$T_0$ (BTJD)	$R_p/R_*$	$b$	$a$ (AU)	$\rho_*$ ( $\rho_\odot$ )	$i$ ( $^\circ$ )	$R_p$ ( $R_J$ )	$R_p$ ( $R_\oplus$ )	Note
0013416465	44	NGC 6910	5.5776 <sup>+0.0587</sup> <sub>-0.0524</sub>	1688.665 <sup>+0.105</sup> <sub>-0.117</sub>	0.282 <sup>+0.144</sup> <sub>-0.081</sub>	1.00 <sup>+0.18</sup> <sub>-0.13</sub>	0.0860 <sup>+0.0006</sup> <sub>-0.0005</sub>	0.41 <sup>+0.01</sup> <sub>-0.01</sub>	84.2 <sup>+0.8</sup> <sub>-1.1</sub>	5.16 <sup>+2.64</sup> <sub>-1.49</sub>	57.79 <sup>+29.6</sup> <sub>-16.7</sub>	
0013866376	45	NGC 6910	9.3972 <sup>+0.0032</sup> <sub>-0.0031</sub>	1686.476 <sup>+0.005</sup> <sub>-0.005</sub>	0.075 <sup>+0.014</sup> <sub>-0.006</sub>	0.93 <sup>+0.03</sup> <sub>-0.02</sub>	0.1575 <sup>+0.0018</sup> <sub>-0.0018</sub>	0.12 <sup>+0.01</sup> <sub>-0.01</sub>	84.2 <sup>+0.2</sup> <sub>-0.2</sub>	2.68 <sup>+0.52</sup> <sub>-0.22</sub>	30.06 <sup>+5.8</sup> <sub>-2.5</sub>	
0013875852	46	NGC 6910	8.3300 <sup>+0.0014</sup> <sub>-0.0014</sub>	1684.099 <sup>+0.005</sup> <sub>-0.005</sub>	0.255 <sup>+0.166</sup> <sub>-0.063</sub>	0.98 <sup>+0.21</sup> <sub>-0.10</sub>	0.1110 <sup>+0.0003</sup> <sub>-0.0003</sub>	0.43 <sup>+0.01</sup> <sub>-0.01</sub>	85.7 <sup>+0.5</sup> <sub>-0.9</sub>	4.54 <sup>+2.95</sup> <sub>-1.13</sub>	50.91 <sup>+33.1</sup> <sub>-12.7</sub>	
0050361536	47	NGC 1027	12.9725 <sup>+0.0088</sup> <sub>-0.0086</sub>	1793.465 <sup>+0.006</sup> <sub>-0.006</sub>	0.327 <sup>+0.106</sup> <sub>-0.100</sub>	1.22 <sup>+0.11</sup> <sub>-0.11</sub>	0.1482 <sup>+0.0013</sup> <sub>-0.0014</sub>	0.15 <sup>+0.01</sup> <sub>-0.01</sub>	84.3 <sup>+0.5</sup> <sub>-0.5</sub>	8.28 <sup>+2.68</sup> <sub>-2.54</sub>	92.75 <sup>+30.0</sup> <sub>-28.4</sub>	
0051022999	48	NGC 1027	7.7336 <sup>+0.0020</sup> <sub>-0.0019</sub>	1794.506 <sup>+0.003</sup> <sub>-0.003</sub>	0.219 <sup>+0.006</sup> <sub>-0.006</sub>	0.08 <sup>+0.008</sup> <sub>-0.005</sub>	0.0736 <sup>+0.0003</sup> <sub>-0.0003</sub>	1.77 <sup>+0.01</sup> <sub>-0.01</sub>	89.8 <sup>+0.2</sup> <sub>-0.2</sub>	1.69 <sup>+0.04</sup> <sub>-0.04</sub>	18.99 <sup>+0.5</sup> <sub>-0.5</sub>	
0065557265	49	NGC 7789	1.6828 <sup>+0.0002</sup> <sub>-0.0002</sub>	1785.339 <sup>+0.011</sup> <sub>-0.063</sub>	0.114 <sup>+0.138</sup> <sub>-0.094</sub>	0.75 <sup>+0.22</sup> <sub>-0.11</sub>	0.0318 <sup>+0.0004</sup> <sub>-0.0013</sub>	0.55 <sup>+0.32</sup> <sub>-0.34</sub>	79.8 <sup>+2.7</sup> <sub>-1.8</sub>	1.42 <sup>+3.29</sup> <sub>-1.18</sub>	15.93 <sup>+36.9</sup> <sub>-13.2</sub>	
0067424670	50	NGC 752	1.1328 <sup>+0.0001</sup> <sub>-0.0001</sub>	1790.941 <sup>+0.001</sup> <sub>-0.001</sub>	0.188 <sup>+0.006</sup> <sub>-0.006</sub>	0.71 <sup>+0.03</sup> <sub>-0.03</sub>	0.0200 <sup>+0.0001</sup> <sub>-0.0001</sub>	1.92 <sup>+0.01</sup> <sub>-0.01</sub>	82.8 <sup>+0.4</sup> <sub>-0.3</sub>	1.39 <sup>+0.05</sup> <sub>-0.04</sub>	15.53 <sup>+0.5</sup> <sub>-0.5</sub>	
0106235729	51	NGC 6871	4.0550 <sup>+0.0003</sup> <sub>-0.0003</sub>	1684.418 <sup>+0.002</sup> <sub>-0.002</sub>	0.366 <sup>+0.060</sup> <sub>-0.051</sub>	1.11 <sup>+0.07</sup> <sub>-0.06</sub>	0.0730 <sup>+0.0003</sup> <sub>-0.0003</sub>	0.40 <sup>+0.01</sup> <sub>-0.01</sub>	81.9 <sup>+0.5</sup> <sub>-0.5</sub>	7.08 <sup>+1.16</sup> <sub>-0.99</sub>	79.41 <sup>+13.0</sup> <sub>-11.1</sub>	
0154304816	52	Alessi 37	3.8552 <sup>+0.0009</sup> <sub>-0.0009</sub>	1742.164 <sup>+0.002</sup> <sub>-0.002</sub>	0.259 <sup>+0.162</sup> <sub>-0.119</sub>	1.12 <sup>+0.17</sup> <sub>-0.15</sub>	0.0537 <sup>+0.0001</sup> <sub>-0.0001</sub>	0.57 <sup>+0.01</sup> <sub>-0.01</sub>	82.5 <sup>+1.0</sup> <sub>-1.2</sub>	3.39 <sup>+2.12</sup> <sub>-1.57</sub>	38.02 <sup>+23.8</sup> <sub>-17.6</sub>	
0185779182	53	RSG 5	70.0427 <sup>+19.4350</sup> <sub>-19.5648</sub>	1711.871 <sup>+0.004</sup> <sub>-0.004</sub>	0.228 <sup>+0.012</sup> <sub>-0.011</sub>	0.49 <sup>+0.25</sup> <sub>-0.25</sub>	0.3046 <sup>+0.0541</sup> <sub>-0.0598</sub>	1.74 <sup>+0.17</sup> <sub>-0.17</sub>	89.7 <sup>+0.1</sup> <sub>-0.10</sub>	1.69 <sup>+0.12</sup> <sub>-0.10</sub>	18.99 <sup>+1.1</sup> <sub>-1.1</sub>	
0251494772	54	SAI 25	5.3034 <sup>+0.0024</sup> <sub>-0.0023</sub>	1795.088 <sup>+0.005</sup> <sub>-0.006</sub>	0.121 <sup>+0.004</sup> <sub>-0.004</sub>	0.72 <sup>+0.04</sup> <sub>-0.04</sub>	0.0894 <sup>+0.0014</sup> <sub>-0.0014</sub>	0.05 <sup>+0.01</sup> <sub>-0.01</sub>	81.4 <sup>+0.8</sup> <sub>-0.9</sub>	4.70 <sup>+3.4</sup> <sub>-0.29</sub>	52.64 <sup>+3.8</sup> <sub>-3.3</sub>	
0251975224	55	King 20	3.5618 <sup>+0.0008</sup> <sub>-0.0008</sub>	1956.781 <sup>+0.003</sup> <sub>-0.004</sub>	0.187 <sup>+0.004</sup> <sub>-0.004</sub>	0.08 <sup>+0.008</sup> <sub>-0.006</sub>	0.0531 <sup>+0.0001</sup> <sub>-0.0001</sub>	0.46 <sup>+0.01</sup> <sub>-0.01</sub>	89.4 <sup>+0.4</sup> <sub>-0.6</sub>	2.73 <sup>+0.06</sup> <sub>-0.06</sub>	30.65 <sup>+0.7</sup> <sub>-0.7</sub>	
0260167199	56	IC 1396	17.6348 <sup>+0.0078</sup> <sub>-0.0077</sub>	1741.326 <sup>+0.005</sup> <sub>-0.005</sub>	0.312 <sup>+0.120</sup> <sub>-0.106</sub>	1.13 <sup>+0.13</sup> <sub>-0.13</sub>	0.1494 <sup>+0.0004</sup> <sub>-0.0004</sub>	0.46 <sup>+0.01</sup> <sub>-0.01</sub>	87.1 <sup>+0.3</sup> <sub>-0.3</sub>	4.44 <sup>+1.70</sup> <sub>-1.51</sub>	49.72 <sup>+19.0</sup> <sub>-16.9</sub>	
0269519402	57	Gulliver 49	3.3736 <sup>+0.0000</sup> <sub>-0.0000</sub>	1806.271 <sup>+0.001</sup> <sub>-0.001</sub>	0.120 <sup>+0.002</sup> <sub>-0.002</sub>	0.39 <sup>+0.05</sup> <sub>-0.07</sub>	0.0630 <sup>+0.0005</sup> <sub>-0.0005</sub>	0.18 <sup>+0.01</sup> <sub>-0.01</sub>	85.9 <sup>+0.8</sup> <sub>-0.7</sub>	2.95 <sup>+0.09</sup> <sub>-0.09</sub>	33.07 <sup>+1.0</sup> <sub>-1.0</sub>	
0270022396	58	NGC 7654	3.7786 <sup>+0.0000</sup> <sub>-0.0000</sub>	1807.429 <sup>+0.001</sup> <sub>-0.001</sub>	0.347 <sup>+0.055</sup> <sub>-0.056</sub>	1.12 <sup>+0.07</sup> <sub>-0.07</sub>	0.0666 <sup>+0.0002</sup> <sub>-0.0002</sub>	0.36 <sup>+0.01</sup> <sub>-0.01</sub>	81.1 <sup>+0.6</sup> <sub>-0.5</sub>	6.66 <sup>+1.05</sup> <sub>-1.08</sub>	74.71 <sup>+11.8</sup> <sub>-12.1</sub>	
0270618239	59	NGC 6811	2.6377 <sup>+0.0002</sup> <sub>-0.0002</sub>	1683.548 <sup>+0.002</sup> <sub>-0.002</sub>	0.217 <sup>+0.003</sup> <sub>-0.003</sub>	0.03 <sup>+0.03</sup> <sub>-0.02</sub>	0.0381 <sup>+0.0001</sup> <sub>-0.0001</sub>	1.08 <sup>+0.01</sup> <sub>-0.01</sub>	89.8 <sup>+0.1</sup> <sub>-0.2</sub>	2.10 <sup>+0.03</sup> <sub>-0.03</sub>	23.56 <sup>+0.4</sup> <sub>-0.3</sub>	
0270920839	60	Czemik 44	5.2236 <sup>+0.0001</sup> <sub>-0.0001</sub>	1768.090 <sup>+0.002</sup> <sub>-0.002</sub>	0.151 <sup>+0.008</sup> <sub>-0.010</sub>	0.81 <sup>+0.02</sup> <sub>-0.03</sub>	0.1088 <sup>+0.0011</sup> <sub>-0.0012</sub>	0.12 <sup>+0.02</sup> <sub>-0.02</sub>	82.6 <sup>+0.6</sup> <sub>-0.6</sub>	5.42 <sup>+0.39</sup> <sub>-0.35</sub>	60.80 <sup>+4.3</sup> <sub>-3.9</sub>	
0271443321	61	SAI 149	5.9718 <sup>+0.0001</sup> <sub>-0.0001</sub>	1765.415 <sup>+0.002</sup> <sub>-0.002</sub>	0.393 <sup>+0.030</sup> <sub>-0.026</sub>	0.89 <sup>+0.05</sup> <sub>-0.07</sub>	0.0970 <sup>+0.0019</sup> <sub>-0.0019</sub>	0.07 <sup>+0.01</sup> <sub>-0.01</sub>	80.9 <sup>+0.4</sup> <sub>-0.5</sub>	14.17 <sup>+1.15</sup> <sub>-1.00</sub>	158.79 <sup>+12.9</sup> <sub>-11.2</sub>	
0285249796	62	ASCC 13	4.7520 <sup>+0.0009</sup> <sub>-0.0009</sub>	1816.836 <sup>+0.002</sup> <sub>-0.002</sub>	0.257 <sup>+0.008</sup> <sub>-0.008</sub>	0.64 <sup>+0.03</sup> <sub>-0.04</sub>	0.0664 <sup>+0.0001</sup> <sub>-0.0001</sub>	0.52 <sup>+0.01</sup> <sub>-0.01</sub>	86.2 <sup>+0.2</sup> <sub>-0.2</sub>	3.72 <sup>+0.12</sup> <sub>-0.12</sub>	41.70 <sup>+1.4</sup> <sub>-1.3</sub>	
0298292983	63	NGC 6940	1.2816 <sup>+0.0001</sup> <sub>-0.0131</sub>	1701.051 <sup>+0.002</sup> <sub>-0.052</sub>	0.146 <sup>+0.005</sup> <sub>-0.016</sub>	0.76 <sup>+0.05</sup> <sub>-0.08</sub>	0.0267 <sup>+0.0001</sup> <sub>-0.0002</sub>	0.33 <sup>+0.02</sup> <sub>-0.02</sub>	77.2 <sup>+1.5</sup> <sub>-1.1</sub>	2.37 <sup>+0.14</sup> <sub>-0.28</sub>	26.60 <sup>+3.2</sup> <sub>-3.1</sub>	
0316246231	64	King 6	8.5751 <sup>+0.0020</sup> <sub>-0.0020</sub>	1975.982 <sup>+0.002</sup> <sub>-0.002</sub>	0.360 <sup>+0.073</sup> <sub>-0.065</sub>	1.12 <sup>+0.09</sup> <sub>-0.08</sub>	0.0992 <sup>+0.0002</sup> <sub>-0.0002</sub>	0.43 <sup>+0.01</sup> <sub>-0.01</sub>	85.2 <sup>+0.4</sup> <sub>-0.5</sub>	5.63 <sup>+1.01</sup> <sub>-1.01</sub>	63.14 <sup>+12.8</sup> <sub>-11.3</sub>	
0323717669	65	RSG 8	4.2378 <sup>+0.0018</sup> <sub>-0.0015</sub>	1955.852 <sup>+0.006</sup> <sub>-0.007</sub>	0.169 <sup>+0.008</sup> <sub>-0.007</sub>	0.14 <sup>+0.14</sup> <sub>-0.10</sub>	0.0453 <sup>+0.0002</sup> <sub>-0.0002</sub>	2.66 <sup>+0.01</sup> <sub>-0.01</sub>	89.5 <sup>+0.4</sup> <sub>-0.5</sub>	1.05 <sup>+0.05</sup> <sub>-0.05</sub>	11.77 <sup>+0.5</sup> <sub>-0.6</sub>	
0326483210	66	FSR 0342	3.8130 <sup>+0.0004</sup> <sub>-0.0004</sub>	1741.329 <sup>+0.003</sup> <sub>-0.003</sub>	0.200 <sup>+0.007</sup> <sub>-0.010</sub>	0.35 <sup>+0.10</sup> <sub>-0.10</sub>	0.0603 <sup>+0.0005</sup> <sub>-0.0005</sub>	0.33 <sup>+0.01</sup> <sub>-0.01</sub>	87.2 <sup>+0.6</sup> <sub>-0.6</sub>	3.55 <sup>+0.14</sup> <sub>-0.17</sub>	39.80 <sup>+1.6</sup> <sub>-2.0</sub>	
0332258412	67	NGC 457	7.1465 <sup>+0.0035</sup> <sub>-0.0035</sub>	1794.905 <sup>+0.004</sup> <sub>-0.004</sub>	0.371 <sup>+0.074</sup> <sub>-0.068</sub>	1.10 <sup>+0.09</sup> <sub>-0.09</sub>	0.1235 <sup>+0.0005</sup> <sub>-0.0005</sub>	0.22 <sup>+0.01</sup> <sub>-0.01</sub>	83.3 <sup>+0.6</sup> <sub>-0.5</sub>	10.10 <sup>+2.03</sup> <sub>-1.88</sub>	114.24 <sup>+22.8</sup> <sub>-21.0</sub>	
0334949878	68	Alessi Teutsch 5	19.4784 <sup>+0.0002</sup> <sub>-0.0002</sub>	1741.392 <sup>+0.001</sup> <sub>-0.001</sub>	0.349 <sup>+0.055</sup> <sub>-0.048</sub>	1.26 <sup>+0.06</sup> <sub>-0.05</sub>	0.2126 <sup>+0.0010</sup> <sub>-0.0011</sub>	0.28 <sup>+0.01</sup> <sub>-0.01</sub>	86.4 <sup>+0.2</sup> <sub>-0.2</sub>	7.76 <sup>+1.22</sup> <sub>-1.06</sub>	87.01 <sup>+13.7</sup> <sub>-11.9</sub>	
0348608380	69	NGC 884	16.3026 <sup>+0.5559</sup> <sub>-0.2298</sub>	1806.922 <sup>+0.003</sup> <sub>-0.003</sub>	0.380 <sup>+0.064</sup> <sub>-0.058</sub>	1.11 <sup>+0.08</sup> <sub>-0.07</sub>	0.2476 <sup>+0.0056</sup> <sub>-0.0024</sub>	0.19 <sup>+0.02</sup> <sub>-0.02</sub>	85.9 <sup>+0.3</sup> <sub>-0.3</sub>	12.60 <sup>+2.22</sup> <sub>-1.93</sub>	141.26 <sup>+24.9</sup> <sub>-21.6</sub>	
0356973763	70	NGC 6997	10.0250 <sup>+0.0035</sup> <sub>-0.4778</sub>	1720.499 <sup>+0.015</sup> <sub>-0.328</sub>	0.104 <sup>+0.113</sup> <sub>-0.017</sub>	0.92 <sup>+0.09</sup> <sub>-0.18</sub>	0.1182 <sup>+0.0014</sup> <sub>-0.0029</sub>	0.13 <sup>+0.02</sup> <sub>-0.02</sub>	84.5 <sup>+1.1</sup> <sub>-0.9</sub>	2.71 <sup>+2.94</sup> <sub>-0.58</sub>	30.34 <sup>+33.0</sup> <sub>-6.5</sub>	
0377619148	71	NGC 7510	19.8963 <sup>+0.0004</sup> <sub>-0.1048</sub>	1784.408 <sup>+0.002</sup> <sub>-0.002</sub>	0.373 <sup>+0.077</sup> <sub>-0.077</sub>	1.17 <sup>+0.09</sup> <sub>-0.09</sub>	0.2713 <sup>+0.0062</sup> <sub>-0.0065</sub>	0.05 <sup>+0.01</sup> <sub>-0.01</sub>	84.1 <sup>+0.5</sup> <sub>-0.4</sub>	18.65 <sup>+3.81</sup> <sub>-3.85</sub>	209.06 <sup>+42.7</sup> <sub>-43.2</sub>	
0408094816	72	NGC 7142	5.4037 <sup>+0.0001</sup> <sub>-0.0001</sub>	1743.980 <sup>+0.002</sup> <sub>-0.002</sub>	0.172 <sup>+0.005</sup> <sub>-0.005</sub>	0.52 <sup>+0.08</sup> <sub>-0.13</sub>	0.0711 <sup>+0.0003</sup> <sub>-0.0003</sub>	0.09 <sup>+0.02</sup> <sub>-0.02</sub>	84.8 <sup>+1.5</sup> <sub>-1.2</sub>	4.41 <sup>+0.38</sup> <sub>-0.37</sub>	49.40 <sup>+4.2</sup> <sub>-4.2</sub>	
0408358709	73	NGC 7142	6.3320 <sup>+0.0001</sup> <sub>-0.0001</sub>	1741.434 <sup>+0.002</sup> <sub>-0.090</sub>	0.337 <sup>+0.108</sup> <sub>-0.090</sub>	1.03 <sup>+0.13</sup> <sub>-0.13</sub>	0.0778 <sup>+0.0003</sup> <sub>-0.0003</sub>	0.14 <sup>+0.02</sup> <sub>-0.01</sub>	82.1 <sup>+1.2</sup> <sub>-1.0</sub>	7.40 <sup>+2.29</sup> <sub>-2.12</sub>	82.92 <sup>+25.7</sup> <sub>-23.8</sub>	
0417058223	74	NGC 7086	5.4767 <sup>+0.0004</sup> <sub>-0.0005</sub>	1712.263 <sup>+0.002</sup> <sub>-0.002</sub>	0.347 <sup>+0.086</sup> <sub>-0.078</sub>	1.18 <sup>+0.10</sup> <sub>-0.09</sub>	0.0853 <sup>+0.0003</sup> <sub>-0.0003</sub>	0.31 <sup>+0.01</sup> <sub>-0.01</sub>	82.3 <sup>+0.6</sup> <sub>-0.6</sub>	7.04 <sup>+1.74</sup> <sub>-1.57</sub>	78.89 <sup>+19.5</sup> <sub>-17.6</sub>	
0420288086	75	NGC 225	6.5343 <sup>+0.0002</sup> <sub>-0.0002</sub>	1769.585 <sup>+0.006</sup> <sub>-0.005</sub>	0.043 <sup>+0.002</sup> <sub>-0.002</sub>	0.85 <sup>+0.02</sup> <sub>-0.02</sub>	0.0950 <sup>+0.0006</sup> <sub>-0.0006</sub>	0.25 <sup>+0.01</sup> <sub>-0.01</sub>	84.7 <sup>+0.2</sup> <sub>-0.1</sub>	0.93 <sup>+0.05</sup> <sub>-0.05</sub>	10.46 <sup>+0.5</sup> <sub>-0.5</sub>	(1)
0421630760	76	IC 1396	4.1869 <sup>+0.0008</sup> <sub>-0.0008</sub>	1742.859 <sup>+0.005</sup> <sub>-0.004</sub>	0.095 <sup>+0.003</sup> <sub>-0.003</sub>	0.10 <sup>+0.10</sup> <sub>-0.07</sub>	0.0556 <sup>+0.0001</sup> <sub>-0.0001</sub>	0.47 <sup>+0.01</sup> <sub>-0.01</sub>	89.4 <sup>+0.5</sup> <sub>-0.7</sub>	1.30 <sup>+0.05</sup> <sub>-0.05</sub>	14.61 <sup>+0.5</sup> <sub>-0.5</sub>	
0427943483	77	NGC 7209	6.2981 <sup>+0.0007</sup> <sub>-0.0007</sub>	1738.788 <sup>+0.001</sup> <sub>-0.001</sub>	0.238 <sup>+0.004</sup> <sub>-0.004</sub>	0.70 <sup>+0.02</sup> <sub>-0.02</sub>	0.0741 <sup>+0.0005</sup> <sub>-0.0005</sub>	0.55 <sup>+0.01</sup> <sub>-0.01</sub>	86.6 <sup>+0.1</sup> <sub>-0.1</sub>	3.14 <sup>+0.07</sup> <sub>-0.06</sub>	35.21 <sup>+0.8</sup> <sub>-0.7</sub>	
0602870459	78	NGC 457	3.1259 <sup>+0.0010</sup> <sub>-0.0010</sub>	1792.949 <sup>+0.004</sup> <sub>-0.004</sub>	0.333 <sup>+0.109</sup> <sub>-0.097</sub>	1.09 <sup>+0.13</sup> <sub>-0.13</sub>	0.0469 <sup>+0.0002</sup> <sub>-0.0002</sub>	0.55 <sup>+0.02</sup> <sub>-0.02</sub>	81.5 <sup>+1.0</sup> <sub>-1.0</sub>	4.43 <sup>+1.45</sup> <sub>-1.29</sub>	49.62 <sup>+16.3</sup> <sub>-14.4</sub>	
0645455722	79	King 5	2.7663 <sup>+0.0004</sup> <sub>-0.0004</sub>	1794.690 <sup>+0.001</sup> <sub>-0.001</sub>	0.190 <sup>+0.009</sup> <sub>-0.007</sub>	0.35 <sup>+0.20</sup> <sub>-0.24</sub>	0.0484 <sup>+0.0011</sup> <sub>-0.0012</sub>	0.25 <sup>+0.01</sup> <sub>-0.06</sub>	86.2 <sup>+0.2</sup> <sub>-2.7</sub>	3.65 <sup>+0.52</sup> <sub>-0.27</sub>	40.88 <sup>+5.8</sup> <sub>-3.1</sub>	
0645713782	80	King 6	6.8572 <sup>+0.0287</sup> <sub>-0.3488</sub>	1790.975 <sup>+0.006</sup> <sub>-0.009</sub>	0.317 <sup>+0.124</sup> <sub>-0.126</sub>	1.09 <sup>+0.15</sup> <sub>-0.17</sub>	0.0715 <sup>+0.0004</sup> <sub>-0.0023</sub>	1.23 <sup>+0.01</sup> <sub>-0.01</sub>	86.1 <sup>+0.6</sup> <sub>-0.6</sub>	2.92 <sup>+1.14</sup> <sub>-1.16</sub>	32.76 <sup>+12.8</sup> <sub>-13.0</sub>	
1961935435	81	NGC 7024	2.7707 <sup>+0.0005</sup> <sub>-0.0005</sub>	1713.398 <sup>+0.002</sup> <sub>-0.002</sub>	0.202 <sup>+0.012</sup> <sub>-0.008</sub>	0.77 <sup>+0.04</sup> <sub>-0.03</sub>	0.0430 <sup>+0.0001</sup> <sub>-0.0001</sub>	0.55 <sup>+0.01</sup> <sub>-0.01</sub>	83.5			



**Figure 5.** The  $M_{G,0}$  versus  $(G_{BP} - G_{RP})_0$  CMD of all the stars analysed in Paper II and in this work. Black points are the main-sequence stars with  $R_* \leq 2.5R_\odot$ , and grey points are the stars excluded from the analysis of candidate exoplanets’ frequency (see the text for details). For clarity, only 50 per cent of the stars are plotted. Red starred symbols indicate the positions on the CMD of the 23 candidate transiting exoplanets identified in stellar clusters from *TESS* data.

$P_{\max}$ , we calculated three frequencies associated with orbital periods  $P_{\min}$ ,  $P_{\max}$ , and  $P_{\text{mid}} = 0.5(P_{\min} + P_{\max})$ . To calculate the frequency of candidate exoplanets in open clusters, we used the following formula:

$$f_*(R_p, P) = \frac{\sum_{i=1}^{N_{\text{cand}}} [1 - \text{FPP}_c^i(R_p, P)]}{\sum_{j=1}^{N_{\text{stars}}} [\text{de}^j(R_p, P) \times \text{Pr}_{\text{transit}}^j(P)]}, \quad (1)$$

where  $N_{\text{cand}}$  and  $N_{\text{stars}}$  are associated with the candidate exoplanets and the stars selected in the previous sections;  $\text{FPP}_c^i(R_p, P)$  are the corrected FPPs of the candidate exoplanets having period  $P$  and radius  $R_p$  in the considered  $(P, R_p)$  bin;  $\text{de}^j$  is the detection efficiency associated with  $j$ th star having *TESS* magnitude  $T_j$ , obtained interpolating the values reported in Fig. 6 for each specific interval  $(P, R_p)$ ; and  $\text{Pr}_{\text{transit}}^j(P) \simeq R_*/a(P)$  is the transit probability associated with the  $j$ th star having radius  $R_*$ , with  $a$  the orbital semimajor axis


calculated on the basis of the third law of Kepler and using the three different periods  $P_{\min}$ ,  $P_{\text{mid}}$ , and  $P_{\max}$ . When the numerator is  $< 1$ , we calculated an upper limit of the frequency considering the numerator equal to 1. The frequencies  $f_*^k$ , with  $k = \text{min, mid, max}$  for the different scenarios  $(P, R_p)$  are reported in Table 5. This is a new approach to calculate the frequencies of exoplanets around a sample of stars, in which we directly used the FPP values obtained with *VESPA*, including also those candidates with a not clear solution (FPPs  $\sim 0.3$ – $0.7$ ). For this kind of candidates, independent statistical validation methods (like that proposed by Armstrong, Gamper & Damoulas 2020) are mandatory.

#### 4.4 Age–planetary radius distribution

Fig. 7 shows the stellar age versus planetary radius distribution for the candidate transiting exoplanets in open clusters identified in this work and in Papers II, and reported as ‘pl’ in Table 4, and the candidates around stars in young associations identified in Paper III; the colours indicate the insolation flux  $S$  calculated on the basis of the stellar radius and effective temperature and of the planetary semimajor axis. Moreover, we included confirmed and candidate exoplanets from literature, orbiting stars in stellar clusters and associations, i.e. with a well-constrained age (coloured triangles). The list of literature objects plotted in Fig. 7 and the corresponding references are reported in Table 6. Finally, we also added objects (coloured squares, Nardiello et al. in preparation) that are under investigation in the context of the ‘GAPS Young Objects’ (GAPS-YO) project (Carleo et al. 2020), aimed at the monitoring of young and intermediate-age stars for the discovery and characterization of young planets. All the objects showed in Fig. 7 have orbital periods  $< 100$  d, i.e. semimajor axis  $a \lesssim 0.5$  AU.

We found that Jupiters with  $R_p \gtrsim 1R_J$  are distributed randomly between  $\sim 10$  Myr and  $\sim 10$  Gyr. Objects having Neptune-sizes or smaller (Earths/super-Earths) are concentrated at ages  $> 200$ – $300$  Myr; anyway the lack of this kind of objects around young stars might be an observational bias due to the difficulty of detecting their transits in the highly variable light curves of active young stars. Objects having a planetary radius  $4 \lesssim R_p \lesssim 10R_{\text{Earth}}$  are concentrated at ages  $< 100$ – $200$  Myr. Is it an observational bias, or the lack of super-Neptune/sub-Jovian-size planets orbiting (on short periods) stars with (well-measured) ages  $\gtrsim 200$  Myr is due to an evolutionary effect of the planets?

The majority of low-mass close-in exoplanets (with  $R_p = 1$ – $4R_{\text{Earth}}$ ) detected until today orbit field stars with ages  $\gtrsim 1$ – $3$  Gyr. Some of these planets are almost totally rocky (e.g. Kepler-93; Dressing et al. 2015), others have low densities that can be explained by the presence of an extended H/He atmosphere. Among them, the Kepler-36 system is particular because formed by two planets with very similar semimajor axis, but totally different densities (Carter et al. 2012), with the inner planet less massive than the external planet. Lopez & Fortney (2013) proved that both the exoplanets in the Kepler-36 system were born with H/He atmospheres and were more massive in the early stages of their life, and that, given the lower core mass of the inner planet, the latter has lost large part of its atmosphere, despite the outer planet that was able to retain about half of its initial atmosphere. As explained in detail in the review by Owen (2019, see also references therein), there is strong evidence that atmospheric escape is the mechanism that prevails in the first stages of a low-mass close-in exoplanet’s evolution, and it depends (in a first approximation) on the characteristics of the host star (and its high-energy emissions), the distance of the planet from the star, and its core mass. During their formation, these planets accreted



**Figure 6.** Transit detection efficiency (normalized to 1) of our pipeline in the injected period  $P^{\text{inj}}$  versus  $T$  magnitude plane for different size simulated planets: Earth and super-Earth-size planets (top panel), Neptune-size planets (middle panel), and Jupiter-size planets (bottom panel).

large amount of H/He into their expanded atmospheres, inflating their radius at  $5\text{--}13R_{\text{Earth}}$ ; in few hundreds Myr, if irradiated strongly enough, these planets lose the large part of their atmosphere, resulting in (super-)Earth/sub-Neptune-size planets (Owen et al. 2020). Even if Fig. 7 does not show any particular dependence on the insolation flux, the distribution of short-period (candidate) exoplanets with radii  $113 \lesssim R_p \lesssim 13R_{\text{Earth}}$  seems to confirm the idea of atmospheric

escape on timescales of  $\lesssim 100\text{--}200$  Myr; anyway, as demonstrated by Owen (2020), even if challenging, for many of these candidates, mass measurements are mandatory in order to understand the mechanisms and constraint the time-scales of planetary atmospheric evolution.

Other possible explanations for the lack of planets older than  $\sim 200$  Myr in the interval  $4 \lesssim R_p \lesssim 10R_{\text{Earth}}$  may be linked to the

**Table 3.** Detection efficiencies of the PATHOS pipeline.

$T$ (mag)	Detection efficiency ( per cent)			
	0.5– 2.0 d	$0.85 < R_p \leq 3.9R_{\text{Earth}}$		85.0– 365.0 d
		2.0– 10.0 d	10.0– 85.0 d	
6–7	40.00 ± 23.66	10.00 ± 10.00	<0.01	<0.01
7–8	40.38 ± 7.38	14.42 ± 3.98	5.21 ± 2.39	3.85 ± 1.96
8–9	24.79 ± 3.64	10.26 ± 2.20	2.75 ± 1.14	2.14 ± 0.97
9–10	16.86 ± 2.37	8.00 ± 1.57	4.95 ± 1.27	2.86 ± 0.92
10–11	14.29 ± 2.16	7.43 ± 1.51	4.62 ± 1.22	4.86 ± 1.21
11–12	10.29 ± 1.80	4.29 ± 1.13	5.79 ± 1.37	3.43 ± 1.01
12–13	6.00 ± 1.35	9.71 ± 1.75	4.10 ± 1.16	4.57 ± 1.17
13–14	6.00 ± 1.35	6.57 ± 1.41	4.13 ± 1.17	2.00 ± 0.76
14–15	5.43 ± 1.28	4.29 ± 1.13	2.41 ± 0.92	1.14 ± 0.57
15–16	2.86 ± 0.92	3.14 ± 0.96	0.35 ± 0.35	1.43 ± 0.64
16–17	2.29 ± 0.82	2.00 ± 0.76	0.69 ± 0.49	0.86 ± 0.50
17–18	1.91 ± 0.97	0.96 ± 0.68	<0.01	<0.01
		$3.9 < R_p \leq 11.2R_{\text{Earth}}$		
6–7	80.00 ± 37.95	60.00 ± 30.98	10.00 ± 10.00	<0.01
7–8	82.69 ± 12.05	56.73 ± 9.25	6.73 ± 2.63	3.85 ± 1.96
8–9	75.64 ± 7.54	52.56 ± 5.85	5.13 ± 1.52	2.56 ± 1.06
9–10	73.14 ± 6.02	40.57 ± 4.04	5.43 ± 1.28	3.14 ± 0.96
10–11	64.00 ± 5.48	40.86 ± 4.05	2.86 ± 0.92	5.71 ± 1.31
11–12	52.29 ± 4.77	30.37 ± 3.37	8.00 ± 1.57	3.43 ± 1.01
12–13	47.14 ± 4.45	22.57 ± 2.81	3.71 ± 1.05	2.86 ± 0.92
13–14	44.86 ± 4.31	28.57 ± 3.24	4.29 ± 1.13	3.43 ± 1.01
14–15	45.43 ± 4.34	21.14 ± 2.71	6.00 ± 1.35	1.14 ± 0.57
15–16	38.00 ± 3.87	18.29 ± 2.49	2.57 ± 0.87	2.00 ± 0.76
16–17	24.29 ± 2.94	11.43 ± 1.91	0.86 ± 0.50	1.43 ± 0.64
17–18	8.13 ± 2.05	1.91 ± 0.97	<0.01	<0.01
		$1.0 < R_p \leq 2.5R_J$		
6–7	80.00 ± 37.95	70.00 ± 34.50	<0.01	<0.01
7–8	97.12 ± 13.57	87.50 ± 12.56	11.54 ± 3.52	2.88 ± 1.69
8–9	90.17 ± 8.56	78.21 ± 7.72	11.54 ± 2.35	2.14 ± 0.97
9–10	90.00 ± 6.99	78.00 ± 6.30	12.29 ± 1.99	4.00 ± 1.09
10–11	90.00 ± 6.99	69.71 ± 5.81	7.71 ± 1.54	3.71 ± 1.05
11–12	84.57 ± 6.68	70.57 ± 5.86	8.29 ± 1.60	6.29 ± 1.38
12–13	85.14 ± 6.71	68.86 ± 5.76	6.29 ± 1.38	4.86 ± 1.21
13–14	78.86 ± 6.35	69.14 ± 5.78	8.57 ± 1.63	3.14 ± 0.96
14–15	69.43 ± 5.80	61.43 ± 5.32	6.57 ± 1.41	2.57 ± 0.87
15–16	51.14 ± 4.70	42.86 ± 4.18	3.71 ± 1.05	1.43 ± 0.64
16–17	53.71 ± 4.86	29.71 ± 3.32	3.71 ± 1.05	0.57 ± 0.41
17–18	46.41 ± 5.70	28.23 ± 4.16	<0.01	<0.01

dynamical evolution of this kind of planets after their formation, such as migration due to interaction with the protoplanetary disc or planet-planet scattering (Fabrycky & Tremaine 2007; Chatterjee et al. 2008; Kley & Nelson 2012). These phenomena are expected to modify the orbital characteristics of the planets, such as inclination, eccentricity, and semimajor axis, making the detection of a possible transit of these planets more difficult. These phenomena occur on scales of a few tens of Myr or less, and may not yet have come into play for the objects reported in this analysis.

## 5 SUMMARY AND CONCLUSION

The aim of the PATHOS project is the discovery and first characterization of transiting objects around stars in stellar clusters and associations observed by *TESS*. Stellar clusters and associations offer the rare opportunity to obtain precise measurements of the ages of the stars (usually affected by large uncertainties), in addition to stars' physical parameters such as radius, mass, effective temperature, etc., and analyse planet characteristics as a function of its host's properties. Because of the low resolution of *TESS* cameras, stellar clusters appear as very crowded regions on FFIs, and appropriate tools are necessary

**Table 4.** FPPs and most likely scenarios for PATHOS candidate exoplanets.

TIC	PATHOS	FPP	Scenario (prob.)	FPP <sub>c</sub>	RUWE
Southern ecliptic hemisphere					
0039291805	3	0.92	BEB (0.51)	0.87	1.023
0088977253	6	0.76	EB (0.47)	0.72	1.079
0125414447	9	0.98	BEB (0.79)	0.93	0.929
0126600730	10	0.97	BEB (0.97)	0.92	0.971
0159059181	16	0.99	BEB (0.52)	0.94	1.230
0306385801	20	0.51	EB (0.51)	0.48	0.851
0308538095	21	0.00	pl (1.00)	0.00	0.895
0372913337	23	0.74	BEB (0.73)	0.70	1.133
0410450228	25	0.46	pl (0.54)	0.44	0.901
0432564189	28	0.96	BEB (0.92)	0.91	1.078
0460205581	30	0.21	pl (0.79)	0.20	0.985
0460950389	31	0.22	pl (0.78)	0.21	0.891
0748919024	33	0.00	pl (1.00)	0.00	2.638
1036769612	34	0.63	BEB (0.63)	0.60	1.062
Northern ecliptic hemisphere					
0051022999	48	0.95	BEB (0.88)	0.90	0.947
0065557265	49	1.00	BEB (0.41)	0.95	1.066
0067424670	50	0.53	pl (0.47)	0.50	0.989
0185779182	53	0.36	pl (0.64)	0.34	1.086
0270618239	59	0.28	pl (0.72)	0.27	1.226
0298292983	63	1.00	EB (0.67)	0.95	1.112
0323717669	65	0.98	BEB (0.96)	0.93	1.893
0420288086	75	0.00	pl (1.00)	0.00	0.971
0421630760	76	1.00	BEB (0.93)	0.95	1.029


to obtain high-precision light curves for cluster members. In order to obtain the best light curves, we developed a cutting-edge technique for the extraction of high-precision photometry of stars in crowded fields, based on the use of empirical PSFs and neighbour subtraction, that allows us to minimize neighbour contamination and extract light curves for very faint objects.

In this work, the fourth of the PATHOS series, we extracted and corrected 150 216 light curves of 89 858 open cluster members listed in the Cantat-Gaudin et al. (2018)'s catalogue and observed during the Cycle 2 (Sectors 14–26) of the *TESS* mission. By using the pipeline already tested in the previous papers of the series, we searched for transit signals among the light curves, finding 39 transiting objects of interest, which are added to the 33 objects orbiting open cluster members identified in Paper II. We modelled their light curves to extract planet parameters. From the two lists of objects of interest, we isolated 23 candidates with planetary radius  $R_p \lesssim 2.5R_J$ , and we calculated their FPPs to be a planet, finding that about 3/5 of them are likely false positives.

Taking into account of the transit detection efficiency of our pipeline and of the rate of false positives, we calculated the frequencies of candidate exoplanets in open clusters ( $f_*$ ) for different (Earth/Neptune/Jupiter) sizes planets in different orbital periods intervals (between 0.5 and 365 d). Because we did not detect any strong candidate with orbital period  $< 2$  d, we calculated the frequency upper limit, finding  $f_* \lesssim 0.06$ ,  $\lesssim 0.007$ , and  $\lesssim 0.004$  per cent for Earth-, Neptune-, and Jupiter-size candidate exoplanets, respectively. For candidate exoplanets with periods  $2.0 < P < 10.0$  d, we found for Earth-size objects, an upper limit  $f_* \lesssim 0.2$  per cent, while for candidate planets with  $R_p \gtrsim 3.9R_{\text{Earth}}$ , the frequency is  $\sim 0.06$  per cent; in the range  $10.0 < P < 85.0$  d, we obtained that the frequency of Earth and Neptune-size candidates is  $\lesssim 2$ . per cent, while the measured frequency for giant planets is  $\sim 1.6$  per cent; finally, for long period planets ( $> 85$  d), we found that the frequency

**Table 5.** Candidate exoplanets' frequencies.

Period (d)	$0.85 < R_P \leq 3.9R_{\text{Earth}}$			$3.9 < R_P \leq 11.2R_{\text{Earth}}$			$1.0 < R_P \leq 2.5R_J$		
	$f_{\star}^{\text{min}}$ (per cent)	$f_{\star}^{\text{mid}}$ (per cent)	$f_{\star}^{\text{max}}$ (per cent)	$f_{\star}^{\text{min}}$ (per cent)	$f_{\star}^{\text{mid}}$ (per cent)	$f_{\star}^{\text{max}}$ (per cent)	$f_{\star}^{\text{min}}$ (per cent)	$f_{\star}^{\text{mid}}$ (per cent)	$f_{\star}^{\text{max}}$ (per cent)
0.5–2.0	<0.034	<0.063	<0.086	<0.004	<0.007	<0.009	<0.002	<0.004	<0.006
2.0–10.0	<0.092	<0.191	<0.269	$0.034 \pm 0.025$	$0.070 \pm 0.052$	$0.098 \pm 0.073$	$0.023 \pm 0.013$	$0.047 \pm 0.027$	$0.066 \pm 0.038$
10.0–85.0	<0.533	<1.505	<2.218	<0.327	<0.925	<1.364	$0.555 \pm 0.336$	$1.568 \pm 0.955$	$2.312 \pm 1.413$
85.0–365.0	<2.716	<5.200	<7.174	<2.197	<4.205	<5.806	<2.102	<4.022	<5.553



**Figure 7.** Stellar age versus planetary radius  $R_P$  for candidate exoplanets identified in Papers II, III, and in this work (circles), for candidates and confirmed exoplanets from literature (triangles), and for the objects under investigation in the GAPS-YO programme (squares). Coloured points represent the exoplanets orbiting stars with well-constrained ages (from isochrone fitting and/or gyrochronological analysis of cluster/association members that host the stars). Different colours are associated with different insolation fluxes on the basis of the bar shown on the top.

of candidate exoplanets is  $< 5\text{--}7$  per cent. The large part of the measured and upper limit frequencies are lower than the values reported by Fressin et al. (2013) in the same period intervals for the corresponding size exoplanets around field stars (even if bias can be introduced by the poor statistic of our sample); an exception are the giant planets with 10–85 d period: We found a mean frequency of  $1.6 \pm 0.9$  per cent, in agreement with the value obtained by Fressin et al. (2013,  $1.5 \pm 0.2$  per cent), but also by Santerne et al.

(2016,  $f_{\star} = 0.90 \pm 0.26$  per cent) and Deleuil et al. (2018,  $f_{\star} = 1.86 \pm 0.68$  per cent).

We investigated the stellar age versus planetary radius distribution by using the results obtained in the PATHOS project and the results obtained in other works for exoplanets orbiting stars with well constrained ages. We only used planets with orbital periods  $< 100$  d and we divided the distribution on the basis of the planetary radius  $R_P$ : (1) for (candidate) exoplanets with  $R_P \gtrsim 1R_J$ , we found no

Table 6. Confirmed and candidate transiting exoplanets from literature.

Object	Cluster/association	Age (Myr)	Period (d)	$R_p$ ( $R_{\text{Earth}}$ )	Reference	Object	Cluster/association	Age (Myr)	Period (d)	$R_p$ ( $R_{\text{Earth}}$ )	Reference
K-66b	NGC 6811	863 $^{+30}_{-30}$	17.815 815 $^{+0.000075}_{-0.000075}$	2.80 $^{+0.16}_{-0.16}$	Meibom et al. (2013)	K2-33b	Upp-Sco	9.3 $^{+1.1}_{-1.1}$	5.424 865 $^{+0.000035}_{-0.000031}$	5.04 $^{+0.34}_{-0.37}$	Mann et al. (2016a)
K-67b	NGC 6811	863 $^{+30}_{-30}$	15.725 90 $^{+0.00011}_{-0.00011}$	2.94 $^{+0.16}_{-0.16}$	Meibom et al. (2013)	TOI-2000b	Tuc-Hor	40 $^{+5}_{-5}$	8.1387 $^{+0.0005}_{-0.0005}$	5.63 $^{+0.22}_{-0.21}$	Benatti et al. (2019)
HD 283869b	Hyades	728 $^{+71}_{-71}$	106 $^{+74.0}_{-25.0}$	1.96 $^{+0.13}_{-0.16}$	Vanderburg et al. (2018)	TOI-451b	Psc-Eri	134 $^{+6.5}_{-6.5}$	1.858 703 $^{+0.00025}_{-0.00035}$	1.91 $^{+0.12}_{-0.12}$	Newton et al. (2021)
K2-25b	Hyades	728 $^{+71}_{-71}$	3.484 552 $^{+0.000031}_{-0.000037}$	3.43 $^{+0.95}_{-0.31}$	Mann et al. (2016b)	TOI-451c	Psc-Eri	134 $^{+6.5}_{-6.5}$	9.192 522 $^{+0.00006}_{-0.00010}$	3.10 $^{+0.13}_{-0.13}$	Newton et al. (2021)
K2-136Ab	Hyades	728 $^{+71}_{-71}$	7.975 292 $^{+0.00070}_{-0.00070}$	0.99 $^{+0.04}_{-0.04}$	Mann et al. (2018)	TOI-451d	Psc-Eri	134 $^{+6.5}_{-6.5}$	16.364 988 $^{+0.00044}_{-0.00044}$	4.07 $^{+0.15}_{-0.15}$	Newton et al. (2021)
K2-136Ac	Hyades	728 $^{+71}_{-71}$	17.307 137 $^{+0.000252}_{-0.000284}$	2.91 $^{+0.11}_{-0.10}$	Mann et al. (2018)	TOI-1098b	Melange-I	250 $^{+50}_{-70}$	10.182 71 $^{+0.00004}_{-0.00004}$	3.2 $^{+0.1}_{-0.1}$	Tofflemire et al. (2021)
K2-136Ad	Hyades	728 $^{+71}_{-71}$	25.575 065 $^{+0.000418}_{-0.000357}$	1.45 $^{+0.11}_{-0.08}$	Mann et al. (2018)	HD 63433b	UMa	414 $^{+23}_{-23}$	7.108 01 $^{+0.00036}_{-0.00034}$	2.15 $^{+0.10}_{-0.10}$	Mann et al. (2020)
K2-95b	Praesepe	670 $^{+67}_{-67}$	10.135 091 $^{+0.000495}_{-0.000488}$	3.7 $^{+0.2}_{-0.2}$	Mann et al. (2017)	HD 63433c	UMa	414 $^{+23}_{-23}$	20.5455 $^{+0.00011}_{-0.00011}$	2.64 $^{+0.12}_{-0.12}$	Mann et al. (2020)
K2-100b	Praesepe	670 $^{+67}_{-67}$	1.673 915 $^{+0.000011}_{-0.000011}$	3.5 $^{+0.2}_{-0.2}$	Mann et al. (2017)	HIP 67522b	Sco-Cen	17 $^{+2}_{-2}$	6.959 503 $^{+0.000016}_{-0.000015}$	10.07 $^{+0.47}_{-0.47}$	Rizzuto et al. (2020)
K2-101b	Praesepe	670 $^{+67}_{-67}$	14.677 303 $^{+0.000824}_{-0.000809}$	3.0 $^{+0.1}_{-0.1}$	Mann et al. (2017)	HIP 67522c	Sco-Cen	17 $^{+2}_{-2}$	54.0 $^{+70.0}_{-24.0}$	8.01 $^{+0.75}_{-0.71}$	Rizzuto et al. (2020)
K2-102b	Praesepe	670 $^{+67}_{-67}$	9.915 651 $^{+0.000194}_{-0.000175}$	1.3 $^{+0.1}_{-0.1}$	Mann et al. (2017)	AU Mic b	AU Mic	22 $^{+3}_{-3}$	8.463 21 $^{+0.00004}_{-0.00004}$	4.20 $^{+0.20}_{-0.20}$	Plavchan et al. (2020)
K2-103b	Praesepe	670 $^{+67}_{-67}$	21.169 687 $^{+0.000636}_{-0.000655}$	2.2 $^{+0.2}_{-0.1}$	Mann et al. (2017)	AU Mic c	AU Mic	22 $^{+3}_{-3}$	30.0 $^{+6.0}_{-6.0}$	2.35 $^{+0.67}_{-0.67}$	Plavchan et al. (2020)
K2-104b	Praesepe	670 $^{+67}_{-67}$	1.974 189 $^{+0.000110}_{-0.000109}$	1.9 $^{+0.2}_{-0.1}$	Mann et al. (2017)	V 1298c	Tau	23 $^{+4}_{-4}$	8.249 58 $^{+0.00072}_{-0.00072}$	5.59 $^{+0.36}_{-0.32}$	David et al. (2019)
K2-264b	Praesepe	670 $^{+67}_{-67}$	5.839 770 $^{+0.000063}_{-0.000063}$	2.27 $^{+0.20}_{-0.16}$	Rizzuto et al. (2018)	V 1298d	Tau	23 $^{+4}_{-4}$	12.4032 $^{+0.00015}_{-0.00015}$	6.41 $^{+0.45}_{-0.40}$	David et al. (2019)
K2-264c	Praesepe	670 $^{+67}_{-67}$	19.663 650 $^{+0.000303}_{-0.000306}$	2.77 $^{+0.20}_{-0.18}$	Rizzuto et al. (2018)	V 1298b	Tau	23 $^{+4}_{-4}$	24.1396 $^{+0.00018}_{-0.00018}$	10.27 $^{+0.58}_{-0.53}$	David et al. (2019)
K2-231b	Ruprecht 147	3000 $^{+250}_{-250}$	13.841 901 $^{+0.000352}_{-0.000352}$	2.5 $^{+0.2}_{-0.2}$	Curtis et al. (2018)	V 1298e	Tau	23 $^{+4}_{-4}$	60 $^{+60}_{-18}$	8.74 $^{+0.84}_{-0.72}$	David et al. (2019)
K2-284b	Cas-Tau	120 $^{+640}_{-20}$	4.795 07 $^{+0.00012}_{-0.00012}$	2.77 $^{+0.12}_{-0.12}$	David et al. (2018)						

particular trends, and the planets are randomly distributed on over the range of ages; (2) objects with  $R_p \approx 4R_{\text{Earth}}$  are concentrated at ages  $>100$ – $200$  Myr, but it might be an observational bias due to the difficulty to detect (super-)Earth transits in the light curve of young active stars; and (3) in the range  $4 \lesssim R_p \lesssim 13R_{\text{Earth}}$ , there is a concentration of objects around young stars with ages  $<100$  Myr. A possible explanation of such concentration is that these objects are young planets with rocky cores that, in the early stages of their formation, have accreted large amount of hydrogen and helium in their atmospheres, inflating their radius. On the time-scales of  $\sim 100$ – $200$  Myr, they lose large part of their atmosphere mainly because of the strong irradiation of the host star (see Owen 2019 for a review of all the mechanisms that contribute to the atmosphere escape), and within few hundreds Myr, their radius decreases to that typical of (super-)Earth/Neptune planets. Other explanations can be related to the dynamical evolution of these exoplanets (planet-planet scattering, migration, etc.).

For this reason, the analysis of light curves of members of young associations (ages  $\lesssim 200$ – $300$  Myr), the subject of the next PATHOS works, will be essential to shed light on the mechanisms of planet formation and evolution of close-in sub-Jovian planets.

## ACKNOWLEDGEMENTS

DN acknowledges the support from the French Centre National d'Etudes Spatiales (CNES). GL acknowledges support by CARIPARO Foundation, according to the agreement CARIPARO-Università degli Studi di Padova (Pratica n. 2018/0098). LB acknowledges the funding support from Italian Space Agency (ASI) regulated by 'Accordo ASI-INAF n. 2013-016-R.0 del 9 luglio 2013 e integrazione del 9 luglio 2015 CHEOPS Fasi A/B/C'. The authors thank the anonymous referee for carefully reading this paper and for the useful suggestions that improved this work. This paper includes data collected by the *TESS* mission. Funding for the *TESS* mission is provided by the NASA Explorer Program. This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC; <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement. Some tasks of the data analysis have been carried out using *VARTOOLS* v 1.39 (Hartman & Bakos 2016) and *TLSPYTHON* routine (Hippke & Heller 2019).

## DATA AVAILABILITY

The data underlying this paper are available in MAST at [doi:10.17909/t9-es7m-vw14](https://doi.org/10.17909/t9-es7m-vw14) and <https://archive.stsci.edu/hlsp/pathos>.

## REFERENCES

- Apai D., Nardiello D., Bedin L. R., 2021, *ApJ*, 906, 64  
 Armstrong D. J., Gampert J., Damoulas T., 2020, *MNRAS*, 504, 5327  
 Bailler-Jones C. A. L., Rybizki J., Founesneau M., Mantelet G., Andrae R., 2018, *AJ*, 156, 58  
 Baraffe I., Chabrier G., Barman T. S., Selsis F., Allard F., Hauschildt P. H., 2005, *A&A*, 436, L47  
 Belokurov V. et al., 2020, *MNRAS*, 496, 1922  
 Benatti S. et al., 2019, *A&A*, 630, A81  
 Bossini D. et al., 2019, *A&A*, 623, A108  
 Bovy J., Rix H.-W., Green G. M., Schlafly E. F., Finkbeiner D. P., 2016, *ApJ*, 818, 130

- Bressan A., Marigo P., Girardi L., Salasnich B., Dal Cero C., Rubele S., Nanni A., 2012, *MNRAS*, 427, 127
- Cantat-Gaudin T. et al., 2018, *A&A*, 618, A93
- Carleo I. et al., 2020, *A&A*, 638, A5
- Carleo I. et al., 2021, *A&A*, 645, A71
- Carter J. A. et al., 2012, *Science*, 337, 556
- Chatterjee S., Ford E. B., Matsumura S., Rasio F. A., 2008, *ApJ*, 686, 580
- Claret A., 2018, *A&A*, 618, A20
- Curtis J. L., et al., 2018, *AJ*, 155, 173
- David T. J. et al., 2018, *AJ*, 156, 302
- David T. J., Petigura E. A., Luger R., Foreman-Mackey D., Livingston J. H., Mamajek E. E., Hillenbrand L. A., 2019, *ApJ*, 885, L12
- Deleuil M. et al., 2018, *A&A*, 619, A97
- Dressing C. D. et al., 2015, *ApJ*, 800, 135
- Drimmel R., Cabrera-Lavers A., López-Corredoira M., 2003, *A&A*, 409, 205
- Fabrycky D., Tremaine S., 2007, *ApJ*, 669, 1298
- Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, *PASP*, 125, 306
- Fressin F. et al., 2013, *ApJ*, 766, 81
- Gaia Collaboration et al., 2018, *A&A*, 616, A1
- Girardi L., Bertelli G., Bressan A., Chiosi C., Groenewegen M. A. T., Marigo P., Salasnich B., Weiss A., 2002, *A&A*, 391, 195
- Green G. M., Schlafly E., Zucker C., Speagle J. S., Finkbeiner D., 2019, *ApJ*, 887, 93
- Hartman J. D., Bakos G. Á., 2016, *Astron. Comput.*, 17, 1
- Healy B. F., McCullough P. R., 2020, *ApJ*, 903, 99
- Hidalgo S. L. et al., 2018, *ApJ*, 856, 125
- Hipke M., Heller R., 2019, *A&A*, 623, A39
- Huang C. X. et al., 2020, *Res. Notes AAS*, 4, 204
- Kharchenko N. V., Piskunov A. E., Schilbach E., Röser S., Scholz R. D., 2016, *A&A*, 585, A101
- Kipping D. M., 2010, *MNRAS*, 408, 1758
- Kipping D. M., 2013, *MNRAS*, 435, 2152
- Kley W., Nelson R. P., 2012, *ARA&A*, 50, 211
- Kreidberg L., 2015, *PASP*, 127, 1161
- Lacedelli G. et al., 2021, *MNRAS*, 501, 4148
- Lammer H., Selsis F., Ribas I., Guinan E. F., Bauer S. J., Weiss W. W., 2003, *ApJ*, 598, L121
- Libralato M., Bedin L. R., Nardiello D., Piotto G., 2016a, *MNRAS*, 456, 1137
- Libralato M. et al., 2016b, *MNRAS*, 463, 1780
- Lopez E. D., Fortney J. J., 2013, *ApJ*, 776, 2
- Mackereth J. T. et al., 2021, *MNRAS*, 502, 1947
- Malavolta L. et al., 2016, *A&A*, 588, A118
- Malavolta L. et al., 2018, *AJ*, 155, 107
- Mann A. W. et al., 2016a, *AJ*, 152, 61
- Mann A. W. et al., 2016b, *ApJ*, 818, 46
- Mann A. W. et al., 2017, *AJ*, 153, 64
- Mann A. W. et al., 2018, *AJ*, 155, 4
- Mann A. W. et al., 2020, *AJ*, 160, 179
- Marigo P. et al., 2017, *ApJ*, 835, 77
- Marshall D. J., Robin A. C., Reylé C., Schultheis M., Picaud S., 2006, *A&A*, 453, 635
- Meibom S. et al., 2013, *Nature*, 499, 55
- Morton T. D., 2012, *ApJ*, 761, 6
- Morton T. D., 2015, *Astrophysics Source Code Library*, record ascl:1503.011)
- Murray-Clay R. A., Chiang E. I., Murray N., 2009, *ApJ*, 693, 23
- Nardiello D. et al., 2015a, *MNRAS*, 447, 3536
- Nardiello D. et al., 2015b, *MNRAS*, 451, 312
- Nardiello D. et al., 2019, *MNRAS*, 490, 3806 (Paper I)
- Nardiello D. et al., 2020, *MNRAS*, 495, 4924 (Paper II)
- Nardiello D., 2020, *MNRAS*, 498, 5972 (Paper III)
- Nardiello D., Libralato M., Bedin L. R., Piotto G., Ochner P., Cunial A., Borsato L., Granata V., 2016a, *MNRAS*, 455, 2337
- Nardiello D., Libralato M., Bedin L. R., Piotto G., Borsato L., Granata V., Malavolta L., Nascimbeni V., 2016b, *MNRAS*, 463, 1831
- Newton E. R. et al., 2021, *AJ*, 161, 65
- Owen J. E., 2019, *Annu. Rev. Earth Planet. Sci.*, 47, 67
- Owen J. E., 2020, *MNRAS*, 498, 5030
- Owen J. E., Jackson A. P., 2012, *MNRAS*, 425, 2931
- Owen J. E., Wu Y., 2017, *ApJ*, 847, 29
- Owen J. E., Shaikhislamov I. F., Lammer H., Fossati L., Khodachenko M. L., 2020, *Space Sci. Rev.*, 216, 129
- Plavchan P. et al., 2020, *Nature*, 582, 497
- Ricker G. R. et al., 2015, *J. Astron. Telesc. Instrum. Syst.*, 1, 014003
- Rizzuto A. C. et al., 2020, *AJ*, 160, 33
- Rizzuto A. C., Vanderburg A., Mann A. W., Kraus A. L., Dressing C. D., Agüeros M. A., Douglas S. T., Krolkowski D. M., 2018, *AJ*, 156, 195
- Röser S., Schilbach E., Goldman B., 2016, *A&A*, 595, A22
- Santerne A. et al., 2016, *A&A*, 587, A64
- Silverman B. W., 1986, *Density Estimation for Statistics and Data Analysis*, Chapman and Hall, London
- Stassun K. G. et al., 2019, *AJ*, 158, 138
- Storn R., Price K., 1997, *J. Glob. Optim.*, 11, 341
- Tofflemire B. M. et al., 2021, *AJ*, 161, 32
- Vanderburg A. et al., 2018, *AJ*, 156, 46
- Wu Y., 2019, *ApJ*, 874, 91
- Ziegler C. et al., 2018, *AJ*, 156, 259

## SUPPORTING INFORMATION


Supplementary data are available at [https://archive.stsci.edu/hlsp/p\\_athos](https://archive.stsci.edu/hlsp/p_athos) online.

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

## APPENDIX A: LIGHT-CURVE MODELLING

**Table A1.** Star parameters and priors for the modelling.

TIC	PATHOS	Cluster	$\alpha$ ( $^{\circ}$ )	$\delta$ ( $^{\circ}$ )	$T$ (mag)	$R_*$ ( $R_{\odot}$ )	$M_*$ ( $M_{\odot}$ )	Period (d)	$T_0$ (BTJD)	$LD_{c1}$	$LD_{c2}$	$df$
0013416465	44	NGC 6910	305.4104	40.6532	14.0	$1.88 \pm 0.05$	$2.74 \pm 0.05$	$\mathcal{U}(5.5, 6.0)$	$\mathcal{U}(1688, 1689)$	$0.15 \pm 0.10$	$0.18 \pm 0.10$	$0.02 \pm 0.05$
0013866376	45	NGC 6910	305.6858	40.5015	11.6	$3.68 \pm 0.08$	$5.90 \pm 0.19$	$\mathcal{U}(9, 9.5)$	$\mathcal{U}(1686.0, 1687.0)$	$0.06 \pm 0.10$	$0.10 \pm 0.10$	$0.04 \pm 0.01$
0013875852	46	NGC 6910	305.7604	40.9818	13.9	$1.84 \pm 0.05$	$2.64 \pm 0.05$	$\mathcal{U}(8.0, 8.5)$	$\mathcal{U}(1683.5, 1684.5)$	$0.15 \pm 0.10$	$0.18 \pm 0.10$	$0.14 \pm 0.05$
0050361536	47	NGC 1027	40.4690	61.7671	11.6	$2.64 \pm 0.12$	$2.58 \pm 0.07$	$\mathcal{U}(12.6, 13.4)$	$\mathcal{U}(1793, 1794)$	$0.15 \pm 0.10$	$0.20 \pm 0.10$	$0.01 \pm 0.05$
0051022999	48	NGC 1027	41.9287	61.6554	16.2	$0.80 \pm 0.03$	$0.89 \pm 0.00$	$\mathcal{U}(7.5, 8.0)$	$\mathcal{U}(1794.0, 1795.0)$	$0.37 \pm 0.10$	$0.33 \pm 0.10$	$0.92 \pm 0.05$
0065557265	49	NGC 7789	359.1359	57.0475	14.0	$2.06 \pm 0.67$	$1.54 \pm 0.03$	$\mathcal{U}(1.5, 2.0)$	$\mathcal{U}(1785.2, 1785.5)$	$0.28 \pm 0.10$	$0.28 \pm 0.10$	$0.01 \pm 0.05$
0067424670	50	NGC 752	29.0476	37.9182	13.6	$0.76 \pm 0.03$	$0.83 \pm 0.05$	$\mathcal{U}(0.9, 1.3)$	$\mathcal{U}(1790.5, 1791.5)$	$0.39 \pm 0.10$	$0.35 \pm 0.10$	$0.08 \pm 0.05$
0106235729	51	NGC 6871	301.0821	35.7889	13.0	$2.01 \pm 0.04$	$3.16 \pm 0.06$	$\mathcal{U}(3.7, 4.3)$	$\mathcal{U}(1684, 1685)$	$0.13 \pm 0.10$	$0.16 \pm 0.10$	$0.03 \pm 0.05$
0154304816	52	Alessi 37	342.4106	46.3500	12.7	$1.35 \pm 0.03$	$1.39 \pm 0.05$	$\mathcal{U}(3.5, 4.0)$	$\mathcal{U}(1742.0, 1743.0)$	$0.27 \pm 0.10$	$0.28 \pm 0.10$	$0.02 \pm 0.05$
0185779182	53	RSG 5	302.5300	45.0274	13.3	$0.76 \pm 0.05$	$0.77 \pm 0.07$	$\mathcal{U}(0.1, 99)$	$\mathcal{U}(1711, 1712)$	$0.42 \pm 0.10$	$0.36 \pm 0.10$	$0.02 \pm 0.03$
0251494772	54	SAI 25	45.3145	57.1103	13.3	$4.42 \pm 0.35$	$3.39 \pm 0.16$	$\mathcal{U}(5.0, 5.5)$	$\mathcal{U}(1795, 1796)$	$0.15 \pm 0.10$	$0.19 \pm 0.10$	$0.01 \pm 0.05$
0251975224	55	King 20	353.1928	58.3744	14.0	$1.49 \pm 0.04$	$1.57 \pm 0.06$	$\mathcal{U}(3.3, 3.9)$	$\mathcal{U}(1956, 1957)$	$0.23 \pm 0.10$	$0.26 \pm 0.10$	$0.04 \pm 0.02$
0260167199	56	IC 1396	325.3457	57.1089	13.5	$1.43 \pm 0.03$	$1.34 \pm 0.05$	$\mathcal{U}(17.3, 18.0)$	$\mathcal{U}(1741.0, 1742.0)$	$0.33 \pm 0.10$	$0.31 \pm 0.10$	$0.10 \pm 0.05$
0269519402	57	Gulliver 49	350.6974	61.9366	13.7	$2.54 \pm 0.07$	$2.93 \pm 0.07$	$\mathcal{U}(3.0, 3.5)$	$\mathcal{U}(1806.0, 1806.5)$	$0.15 \pm 0.10$	$0.18 \pm 0.10$	$0.12 \pm 0.05$
0270022396	58	NGC 7654	351.2741	61.6596	13.1	$2.00 \pm 0.05$	$2.76 \pm 0.06$	$\mathcal{U}(3.6, 3.9)$	$\mathcal{U}(1807.3, 1807.5)$	$0.15 \pm 0.10$	$0.18 \pm 0.10$	$0.18 \pm 0.05$
0270618239	59	NGC 6811	293.6701	46.2956	14.5	$0.99 \pm 0.03$	$1.07 \pm 0.05$	$\mathcal{U}(2.2, 3.0)$	$\mathcal{U}(1683.0, 1684.0)$	$0.32 \pm 0.30$	$0.30 \pm 0.30$	$0.05 \pm 0.05$
0270920839	60	Czernik 44	353.5841	62.0029	14.3	$3.94 \pm 0.09$	$6.30 \pm 0.22$	$\mathcal{U}(5.0, 5.5)$	$\mathcal{U}(1767.5, 1768.5)$	$0.05 \pm 0.10$	$0.08 \pm 0.10$	$0.70 \pm 0.05$
0271443321	61	SAI 149	354.4998	60.5393	13.9	$5.00 \pm 0.32$	$3.41 \pm 0.17$	$\mathcal{U}(5.7, 6.2)$	$\mathcal{U}(1765.0, 1766.0)$	$0.15 \pm 0.10$	$0.21 \pm 0.10$	$0.04 \pm 0.05$
0285249796	62	ASCC 13	78.2136	44.5317	12.7	$1.49 \pm 0.03$	$1.73 \pm 0.05$	$\mathcal{U}(4.5, 5.0)$	$\mathcal{U}(1816, 1817)$	$0.20 \pm 0.10$	$0.24 \pm 0.10$	$0.31 \pm 0.05$
0298292983	63	NGC 6940	308.6667	27.8038	12.7	$1.67 \pm 0.05$	$1.56 \pm 0.05$	$\mathcal{U}(1.2, 1.3)$	$\mathcal{U}(1700.9, 1701.2)$	$0.24 \pm 0.10$	$0.26 \pm 0.10$	$0.13 \pm 0.05$
0316246231	64	King 6	51.9527	56.5553	12.4	$1.62 \pm 0.03$	$1.77 \pm 0.05$	$\mathcal{U}(8.3, 8.7)$	$\mathcal{U}(1795.5, 1796.5)$	$0.20 \pm 0.10$	$0.24 \pm 0.10$	$0.01 \pm 0.05$
0323717669	65	RSG 8	348.0553	59.4878	14.6	$0.64 \pm 0.03$	$0.69 \pm 0.05$	$\mathcal{U}(4.0, 4.5)$	$\mathcal{U}(1955.0, 1956.0)$	$0.45 \pm 0.10$	$0.36 \pm 0.10$	$0.70 \pm 0.05$
0326483210	66	FSR 0342	331.8775	53.3684	15.0	$1.83 \pm 0.05$	$2.01 \pm 0.07$	$\mathcal{U}(3.5, 4.0)$	$\mathcal{U}(1741.0, 1742.0)$	$0.17 \pm 0.10$	$0.22 \pm 0.10$	$0.57 \pm 0.05$
0332258412	67	NGC 457	19.5199	58.7259	13.0	$2.86 \pm 0.06$	$4.93 \pm 0.06$	$\mathcal{U}(7.0, 7.5)$	$\mathcal{U}(1794.5, 1795.5)$	$0.07 \pm 0.10$	$0.12 \pm 0.10$	$0.02 \pm 0.05$
0334949878	68	Alessi Teutsch 5	331.7424	60.5581	11.0	$2.38 \pm 0.06$	$3.38 \pm 0.05$	$\mathcal{U}(19.0, 20.0)$	$\mathcal{U}(1741.0, 1742.0)$	$0.13 \pm 0.10$	$0.16 \pm 0.10$	$0.01 \pm 0.05$
0348608380	69	NGC 884	35.5582	57.3225	11.0	$4.13 \pm 0.08$	$7.63 \pm 0.06$	$\mathcal{U}(16, 99)$	$\mathcal{U}(1806.0, 1807.5)$	$0.02 \pm 0.10$	$0.01 \pm 0.10$	$0.01 \pm 0.05$
0356973763	70	NGC 6997	314.4065	44.5798	11.6	$2.51 \pm 0.15$	$2.24 \pm 0.06$	$\mathcal{U}(9.5, 10.3)$	$\mathcal{U}(1720, 1721)$	$0.17 \pm 0.10$	$0.22 \pm 0.10$	$0.03 \pm 0.05$
0377619148	71	NGC 7510	348.0732	60.4962	11.6	$7.15 \pm 0.46$	$6.77 \pm 0.47$	$\mathcal{U}(19.5, 20.0)$	$\mathcal{U}(1784, 1785)$	$0.06 \pm 0.10$	$0.11 \pm 0.10$	$0.01 \pm 0.05$
0408094816	72	NGC 7142	326.3290	65.7443	14.5	$2.57 \pm 0.26$	$1.64 \pm 0.07$	$\mathcal{U}(5.3, 5.9)$	$\mathcal{U}(1743.5, 1744.5)$	$0.28 \pm 0.10$	$0.28 \pm 0.10$	$0.04 \pm 0.05$
0408358709	73	NGC 7142	326.6784	65.7431	14.7	$2.17 \pm 0.32$	$1.57 \pm 0.06$	$\mathcal{U}(6.0, 6.5)$	$\mathcal{U}(1741.0, 1742.0)$	$0.27 \pm 0.10$	$0.28 \pm 0.10$	$0.02 \pm 0.05$
0417058223	74	NGC 7086	322.8993	51.7531	13.2	$2.09 \pm 0.05$	$2.76 \pm 0.07$	$\mathcal{U}(5.2, 5.8)$	$\mathcal{U}(1712.0, 1713.0)$	$0.15 \pm 0.10$	$0.18 \pm 0.10$	$0.44 \pm 0.05$
0420288086	75	NGC 225	10.9627	61.8356	10.4	$2.20 \pm 0.06$	$2.69 \pm 0.06$	$\mathcal{U}(6.3, 6.9)$	$\mathcal{U}(1769.0, 1770.0)$	$0.15 \pm 0.10$	$0.19 \pm 0.10$	$0.01 \pm 0.05$
0421630760	76	IC 1396	324.6780	57.8705	13.7	$1.41 \pm 0.03$	$1.32 \pm 0.05$	$\mathcal{U}(4.0, 4.5)$	$\mathcal{U}(1742.0, 1743.0)$	$0.33 \pm 0.10$	$0.32 \pm 0.10$	$0.10 \pm 0.05$
0427943483	77	NGC 7209	331.5198	46.8716	13.7	$1.35 \pm 0.03$	$1.37 \pm 0.05$	$\mathcal{U}(6.0, 6.5)$	$\mathcal{U}(1738, 1739)$	$0.27 \pm 0.10$	$0.28 \pm 0.10$	$0.01 \pm 0.05$
0602870459	78	NGC 457	19.2565	58.4526	16.7	$1.37 \pm 0.05$	$1.41 \pm 0.07$	$\mathcal{U}(3.0, 3.5)$	$\mathcal{U}(1792.5, 1793.5)$	$0.27 \pm 0.10$	$0.28 \pm 0.10$	$0.95 \pm 0.05$
0645455722	79	King 5	48.2841	52.2956	13.0	$18.44 \pm 0.58$	$1.98 \pm 0.14$	$\mathcal{U}(2.5, 3.0)$	$\mathcal{U}(1794.5, 1795.0)$	$0.43 \pm 0.30$	$0.38 \pm 0.30$	$0.65 \pm 0.05$
0645713782	80	King 6	52.1609	56.5984	15.2	$0.94 \pm 0.03$	$1.04 \pm 0.05$	$\mathcal{U}(6.3, 6.9)$	$\mathcal{U}(1790.5, 1791.5)$	$0.32 \pm 0.10$	$0.30 \pm 0.10$	$0.75 \pm 0.05$
1961935435	81	NGC 7024	316.4611	41.5474	14.9	$1.35 \pm 0.03$	$1.38 \pm 0.05$	$\mathcal{U}(2.5, 3.0)$	$\mathcal{U}(1713.0, 1714.0)$	$0.27 \pm 0.10$	$0.28 \pm 0.10$	$0.78 \pm 0.05$
2015243161	82	NGC 7245	333.8243	54.2961	14.9	$1.85 \pm 0.05$	$2.06 \pm 0.07$	$\mathcal{U}(11.2, 11.8)$	$\mathcal{U}(1745.5, 1746.5)$	$0.17 \pm 0.10$	$0.22 \pm 0.10$	$0.46 \pm 0.05$



**Figure A1.** Overview on the candidate exoplanets PATHOS-44–PATHOS-58. On the left-hand, the  $G$  versus  $G_{BP} - G_{RP}$  CMD of the cluster that hosts the target star (red star or dashed line when  $G_{BP} - G_{RP}$  colour is not available) and the isochrone (blue) fitted with the cluster parameters listed in Table 1. The top right-hand panel shows the folded light curve (grey points) of the candidate and the model (in red) found with PYORBIT; the middle panel shows the difference between the observed points and the model. The bottom left-hand panel shows the  $95 \times 95$  arcsec<sup>2</sup> finding chart centred on the target star; red circle is the aperture adopted to extract photometry, crosses are the in-/out-of-transit difference centroid. The bottom right-hand panel shows the vector-point diagram, centred on the target star, for all the stars that are within 10 arcmin from the target star; black points are the cluster members listed in the catalogue by Cantat-Gaudin et al. (2018).




Figure A2. As in Fig. A1, but for PATHOS-59–PATHOS-73.




Figure A3. As in Fig. A1, but for PATHOS-74–PATHOS-82.

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.