



| | |
|-------------------------|---|
| Publication Year | 2018 |
| Acceptance in OA | 2024-03-14T16:04:26Z |
| Title | IR spectroscopy of ammoniated phyllosilicates at low pressure/high temperature conditions |
| Authors | DE ANGELIS, Simone, FERRARI, MARCO, DE SANCTIS, MARIA CRISTINA, Ammannito, Eleonora |
| Handle | http://hdl.handle.net/20.500.12386/34968 |

IR spectroscopy of ammoniated phyllosilicates at low pressure/high temperature conditions

S. De Angelis (1), M. Ferrari (1), M.C. De Sanctis (1), E. Ammannito (2)

(1) Institute for Space Astrophysics and Planetology, IAPS-INAF, Rome Italy, (simone.deangelis@iaps.inaf.it) (2) Italian Space Agency – ASI, Rome, Italy

Introduction

Ammonium phyllosilicates are thought to be among the constituents of dwarf planet (1) Ceres surface, based on ground-based telescopic [1] and VIR-Dawn spectral data [2]. Following these findings, several works are currently trying to reproduce in the laboratory the Ceres surface composition, in terms of multi-component mineral mixtures [3,4]. Thus it is of interest to investigate the behavior and stability of ammonium compounds, when measured at pressure-temperature conditions that are different from standard laboratory values. Here we describe Visible-Infrared spectroscopic measurements of an ensemble of ammonium phyllosilicates: spectra have been acquired at various pressure-temperature conditions, by means of a P-T cell realized at INAF-IAPS laboratory.

1. Samples and experimental setup

Ammoniated phyllosilicates were produced in the laboratory starting from natural samples, following a procedure described in a series of works [e.g. 6]. Infrared spectra of five ammoniated samples (montmorillonite, Sca-3, two nontronites, NAu-1 and NAu-2, illite-smectite, ISCz-1 and hectorite, SHCa-1) were then acquired in the spectral range 0.35-2.5 μm , by using an ASD FieldSpec Pro 4 spectro-photometer equipped with a QTH lamp. The instrument is characterized by a spectral resolution of about 3-10 nm in the whole range; the spatial resolution of the setup was about 5 mm on the sample. All samples were analyzed in the form of powder, with grain size $d < 36 \mu\text{m}$.

In order to acquire reflectance spectra at varying conditions, the samples were placed inside a P-T environmental cell, developed at INAF-IAPS [5]. The measurements strategy was the following: (i) acquisition at room P-T, (ii) sequence of acquisitions

at room T during pumping; in this stage the pumping was first performed with only primary diaphragm pump (down to a limit of 3-4 mbar) and then also with turbo-molecular pump (down to vacuum pressure of 10^{-4} - 10^{-5} mbar); (iii) acquisitions in vacuum at higher temperatures.

2. IR Spectral measurements.

Here we report, as an example, on the analyses performed on one sample, nontronite (NAu1). The spectra of NH_4 -Nontronite are shown in fig.1. Data were acquired in three stages. The first spectrum (fig.1, A) is at room pressure and temperature. In panel B spectra were acquired at room T during primary pumping from ambient to 10^{-4} mbar. Finally in panel C spectra acquired in vacuum at different temperatures (50-240°C) are shown.

3. Results and Conclusions

Spectra of nontronite are characterized by Fe^{2+} - Fe^{3+} bands at 0.7-1 μm , and by OH/ H_2O bands at 1.4 and 1.9 μm [7]. The feature at 2.3 μm is Fe-OH [7]. NH_4^+ absorption is visible at 2.12 μm in the room P-T spectrum. After pumping the adsorbed water is removed and NH_4^+ features become evident also at 1.55 and 2.01 μm : at room P-T these two features are shoulders in the 1.4 and 1.9 μm bands. After heating up to 240°C all NH_4^+ bands are visible and quite separate from hydration bands.

We can see that NH_4^+ bands remain quite unaltered both by the process of pumping, at least down to a vacuum of about 10^{-4} mbar, and by the process of heating up to 240°C. For higher temperatures the nontronite sample is subject to structural changes: all the water is removed, then ammonia and finally dehydroxylation occurs (fig. 1C).

NH₄-NONTRONITE




Fig.1. NH₄-Nontronite (NAu-1). A): ambient pressure and temperature; B): ambient temperature, during pumping; C): higher temperatures, in vacuum. Vertical lines indicate NH₄⁺ absorptions.

Acknowledgements

The experiment is funded by ASI.

References

- [1] King T.V.V. et al., Science, vol.255, 1551-1553, 1992
- [2] De Sanctis M.C., et al., Nature, vol.528, 241-244, 2015.
- [3] De Angelis S. et al., EPSC abstract n.830, vol.11, 2017
- [4] Ehlmann B.L. et al., MAPS, 10.1111/maps.13103, 2018
- [5] De Angelis S. et al., 49th LPSC, abstract n.1428, 2018
- [6] Ferrari M. et al., 49th LPSC, abstract n.2413, 2018
- [7] Clark R.N. et al., JGR, vol.95, B8, 12,653-12,680, 1990