



<b>Publication Year</b>	2024
<b>Acceptance in OA</b>	2025-04-11T14:43:17Z
<b>Title</b>	Jupiter's Hotspots as observed by JIRAM-Juno: limb darkening in thermal infrared
<b>Authors</b>	GRASSI, Davide, MURA, Alessandro, ADRIANI, Alberto, Sindoni, G, Atreya, S K, Fletcher, L N, Orton, G S, PLAINAKI, CHRISTINA, TOSI, Federico, BIAGIOTTI, Francesco, Olivieri, A, CASTAGNOLI, Chiara, D'AVERSA, EMILIANO, MIGLIORINI, Alessandra, MOIRANO, ALESSANDRO, NOSCHESI, Raffaella, PICCIONI, GIUSEPPE, SORDINI, Roberto, Bolton, S
<b>Publisher's version (DOI)</b>	10.1093/mnras/stae1858
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/37053">http://hdl.handle.net/20.500.12386/37053</a>
<b>Journal</b>	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
<b>Volume</b>	533

# Jupiter's Hot Spots as observed by JIRAM-Juno: limb-darkening in thermal infrared

D. Grassi<sup>1\*</sup>, A. Mura<sup>1</sup>, A. Adriani<sup>1</sup>, G. Sindoni<sup>2</sup>, S. K. Atreya<sup>3</sup>, L. N. Fletcher<sup>4</sup>, G. S. Orton<sup>5</sup>, C. Plainaki<sup>2</sup>, F. Tosi<sup>1</sup>, F. Biagiotti<sup>6,1</sup>, A. Olivieri<sup>2</sup>, C. Castagnoli<sup>1,7,8</sup>, E. D'Aversa<sup>1</sup>, A. Migliorini<sup>1</sup>, A. Moirano<sup>1,6</sup>, R. Noschese<sup>1</sup>, G. Piccioni<sup>1</sup>, R. Sordini<sup>1</sup>, S. Bolton<sup>9</sup>

<sup>1</sup> Istituto Nazionale di Astrofisica, Istituto di Astrofisica e Planetologia Spaziali, Rome, Italy

<sup>2</sup> Agenzia Spaziale Italiana, Rome, Italy

<sup>3</sup> Department of Climate and Space Sciences and Engineering, Univ. of Michigan, Ann Arbor MI, USA

<sup>4</sup> School of Physics and Astronomy, University of Leicester, UK

<sup>5</sup> Jet Propulsion Laboratory, California Institute of Technology, Pasadena CA, USA

<sup>6</sup> Università degli Studi di Roma "Sapienza", Rome, Italy

<sup>7</sup> Università degli Studi di Roma "Tor Vergata", Rome, Italy

<sup>8</sup> Consiglio Nazionale delle Ricerche, Istituto di Scienze dell'Atmosfera e del Clima, Bologna, Italy

<sup>9</sup> Southwest Research Institute, San Antonio TX, USA

\*Corresponding author: [davide.grassi@inaf.it](mailto:davide.grassi@inaf.it)

## Abstract

The Jupiter InfraRed Auroral Mapper (JIRAM) instrument on board the Juno spacecraft performed repeated observations of Jupiter's North Equatorial Belt (NEB) around the time of 12<sup>th</sup> Juno pericenter passage on April 1<sup>st</sup> 2018. The data consist of thermal infrared images and show, among other atmospheric features, two bright Hot Spots on the boundary between the NEB and the Equatorial zone. Night-time images of the same areas at different emission angles were used to constrain the trend of the limb-darkening function. Comparison with simulated observations, computed for different emission angles, total opacities, single scattering albedo  $\omega_0$  and asymmetry parameter  $g$  suggest that  $\omega_0 \sim 0.90 \pm 0.05$  and  $g \sim 0.37 \pm 0.15$  provide best match with data. Subsequently, we computed the  $\omega_0$  and  $g$  resulting from different size distributions, taking into account the complex refractive indices of ammonium hydrosulfide (NH<sub>4</sub>SH) by Howett et al. [2007] and Ferraro et al., [1980]. Only the former dataset is marginally consistent with JIRAM observations. Similarly, ammonia and hydrazine barely reproduce the experimental data. Tholin, although not usually considered a realistic component for Jupiter's aerosols, provides a better match for particle radii between 0.7 and 1  $\mu\text{m}$ , both as a pure material as well as a thick coating over NH<sub>4</sub>SH cores. Notably, this radius range is consistent with the mean radius of aerosols as estimated by Regent et al., [1998] on the basis of Galileo Entry Probe data. Comparison with literature suggests that similar results can be achieved by a large variety of contaminants bearing C-N bounds.

Keywords: planets and satellites: atmospheres, techniques: photometric, planets and satellites: individual: Jupiter

## 1. Introduction

The infrared emission emerging from Jupiter's atmosphere at wavelengths around 5  $\mu\text{m}$  has been measured both by ground-based instruments [e.g., Fletcher et al., 2016; Giles et al., 2016; Orton et al., 2017] as well as space-borne ones [Carlson et al., 1993; Roos-Serote et al., 1998, 1999; Giles et al., 2015]. The brightness at 5  $\mu\text{m}$  is highly time-variable, as shown by Fletcher et al., [2017] and Antuñano et al. [2018]. An extensive dataset, covering all latitudes, has been produced since 2016 by the Jupiter InfraRed Auroral Mapper (JIRAM) on board the Juno spacecraft [see Adriani et al., 2014, for an instrument description and Adriani et al., 2018; Grassi et al., 2018, 2020 for selected results].

Arguably, the most remarkable feature of these observations is the strong radiance contrast between different regions, exceeding a factor of 100 at low and intermediate latitudes (50°S-40°N), with marked latitudinal trends (e.g., Fig. 4b in Grassi et al., [2020]). Most latitudes (especially those associated with bright 'zones' in the visible spectral range) display low infrared emission, with equivalent brightness temperatures down to 170 K ~~below 200K~~. Globally-averaged equilibrium thermodynamic models [Atreya et al., 1999] ~~that~~ expect the occurrence of at least three distinct cloud decks composed of water, ammonium hydrosulfide, and ammonia, from bottom to top. Optically thick clouds prevent most of the thermal radiation from deeper (warmer) levels from reaching the observer, and the measured signal essentially originates from the thermal emission of the atmosphere at the cloud tops. However, some subtleties should be taken into account. The optical thickness of the  $\text{NH}_3$  ice cloud should be low enough to allow the emergence of some radiation from warmer regions at lower altitudes, at least in the equatorial region [Drossart et al., 1998]. Similarly, Irwin et al. [2001] demonstrated that variations in the 5  $\mu\text{m}$  signal are correlated with opacity between 1 and 2 bars (the expected location of ammonium hydrosulfide clouds), rather than the thickness of the higher ammonia cloud. The brightness temperatures observed in low emission regions are therefore consistent with a relatively thin upper cloud of ammonia located above an optically thick ammonium hydrosulfide cloud deck.

Other latitudes (notably those associated with dark 'belts' in the visible spectral range) exhibit higher infrared emission, with brightness temperatures exceeding 260K. Under these conditions, the observer records photons thermally emitted at pressures of a few bars, with  $\text{H}_2$  collision-induced absorption being the primary source of opacity at these wavelengths [Sromovsky & Fry, 2018]. These photons experience only minor absorption from the thin cloud layers found in these regions.

At low latitudes, the higher infrared signal is observed in the so-called 'Hot Spots,' which are large bright areas elongated in longitude (typically spanning a few thousand kilometers) found between the Equatorial Zone (EZ) and the North Equatorial Belt (NEB). These areas are associated, in the visible spectral range, with grey 'festoons', bluish 'dark projections' and 'dark formations' (see Rogers [1995] for an historical review and Choi et al. [2013]). The low cloud opacity of these areas was anticipated from ground-based data [Terrile & Westphal, 1977] before being confirmed by direct on-site measurements by the Galileo Entry Probe (GEP) [Ragent et al., 1998]. There is consensus around the interpretation of these features as the result of trapped Rossby waves [Ortiz et al., 1998; Hueso 1999; Showman and Dowling, 2000; Friedson 2005; Arregi et al., 2006]. Several studies [Irwin et al., 1998; Roos-Serote et al., 1998; Nixon et al., 2001] investigated the Hot Spots using spectra at wavelengths around 5  $\mu\text{m}$  acquired by the Near Infrared Mapping Spectrometer (NIMS) on board the Galileo mission. More recently, we presented two studies [Grassi et al., 2017a, 2020] based on JIRAM spectral data, detailing the atmospheric composition within the Hot Spots. Properties of Hot Spots from thermal infrared and microwave data is discussed in Fletcher et al. [2020], along with their association with visible dark formations.

Our previous investigations were, however, based on a very simplified model for properties of residual aerosols over these features. Physical conditions in one Hot Spot have been measured *in situ* by the GEP during its descent on December 7<sup>th</sup>, 1995: the probe entered at 6.5°N 4.9°W, at the southern rim of a 5- $\mu\text{m}$  Hot Spot [Orton et al., 1998]. The main aerosol layer encountered by the probe extended approximately between +7 and -8 km around the reference 1-bar isobaric surface, with a mean radius of particles between 0.8 and 1.1  $\mu\text{m}$  [Ragent et al., 1998]. While these observations provide the only direct measurements of the aerosol characteristics, it is not apparent whether they can be generalized to all Hot Spots, especially considering that the Galileo probe entered one of the largest and long-lived hot spots on Jupiter.

In this work, we discuss the observations of two Jupiter's Hot Spots at 4.7  $\mu\text{m}$  performed by JIRAM during the 12<sup>th</sup> Juno perijove passage (PJ) on April 1<sup>st</sup> 2018. In this passage, JIRAM made repeated observations of the same large area of the NEB (including Hot Spots) at a variety of emission angles  $\epsilon$ . The variations of observed signal along  $\epsilon$  (the so-called “limb darkening”) are used to constrain the scattering properties of aerosols (namely, the single scattering albedo  $\omega_0$  and asymmetry parameter  $g$ ) and, indirectly, the size distribution. Several factors combine to make the Hot Spots ideal targets for this type of analysis: 1) they are believed to have the simplest cloud structure that can be found over the disk of Jupiter, with essentially one relatively thin layer of aerosol [West et al., 2004]; 2) their intrinsically strong infrared signal reduces the impact of possible residual calibration artefacts on our analysis; 3) their large spatial extent and smooth spatial variations minimize the ambiguities related to variable spatial resolution between different images.

## 2. Dataset

The JIRAM instrument includes a spectrometer and two imaging channels operating in different spectral regions. In this paper, we will focus our attention solely on one of these channels (the so called M-filter), where images are acquired by integrating the incoming radiance over the 4.54-5.03  $\mu\text{m}$  range. Images have a size of 432 x 128 pixels. All image pixels are acquired simultaneously, i.e.: the bi-dimensional image is formed directly on the focal plane and not by the stacking of consecutive lines in a push-broom scan. The field of view of individual pixels is about 240  $\mu\text{rad}$  [Adriani et al., 2014]. The random noise in calibrated M-filter images is by far dominated by shot noise. Data included in this analysis have a SNR between 20 and 80.

Juno is a spin-stabilized spacecraft and JIRAM acquires one image every Juno spacecraft rotation (2 rpm). JIRAM pointing has only one degree of freedom, and the pointing can be set only along the maximum circle orthogonal to the spin axis. Notably, such an arrangement allows JIRAM to compensate for Juno rotation during exposure. Acquisition of images often (but not necessarily always) occurs in sequences, i.e.: a set of consecutive images where the initial pointing angle is progressively increased (or decreased) in order to achieve, as far as possible, a continuous spatial coverage (i.e. a mosaic). Few examples of mosaics derived from such sequences are presented in Fig. 1.

During the 12<sup>th</sup> Juno perijove passage on April 1<sup>st</sup> 2018, JIRAM acquired a set of more than 210 individual images (arranged in 16 sequences) over Jupiter’s equatorial region, within a time span of 3.5h. The pixel resolution in the range from 5°N to 10°N varies between 395 and 83km at the reference 1-bar level because of the high eccentricity of Juno’s orbit. The same data have emission angles between 30° and 70°.

Table I lists the initial and final image that define each sequence selected for this study. The original calibrated data (Adriani et al., 2019) can be found at this Planetary Data System URL:

[https://atmos.nmsu.edu/PDS/data/PDS4/juno\\_jiram\\_bundle/data\\_calibrated/orbit12/](https://atmos.nmsu.edu/PDS/data/PDS4/juno_jiram_bundle/data_calibrated/orbit12/)

Sequence name <i>First image – last image</i>	Pixel resolution (min-max), km	Emission angle (min-max), degrees
180401_034801 <i>2018091T034404 - 2018091T035216</i>	395-281	70-62
180401_040758 <i>2018091T040401 - 2018091T041213</i>	331-201	67-52
180401_041812 <i>2018091T041415 - 2018091T042227</i>	285-180	64-48
180401_042826 <i>2018091T042429 - 2018091T043240</i>	298-163	66-45
180401_043809 <i>2018091T043412 - 2018091T044224</i>	314-151	68-42

180401_044823 2018091T044426 - 2018091T045237	261-139	64-39
180401_050820 2018091T050423 - 2018091T051235	292-124	68-34
180401_052848 2018091T052450 - 2018091T053302	252-112	66-32
180401_054845 2018091T054447 - 2018091T055259	222-103	64-30
180401_055859 2018091T055501 - 2018091T060313	230- 99	66-31
180401_060913 2018091T060515 - 2018091T061327	170- 96	59-31
180401_062909 2018091T062512 - 2018091T063324	126-91	51-33
180401_064937 2018091T064540 - 2018091T065351	107-87	48-37
180401_070934 2018091T070536 - 2018091T071348	102-84	50-41
180401_071948 2018091T071550 - 2018091T072402	105-83	53-44
180401_073001 2018091T072604 - 2018091T072808	110-83	57-46
<p>Table I. JIRAM data considered in this study. Sequences are series of consecutive images acquired by the instrument, around the time given by sequence name according the scheme yymmdd_hhmmss. While sequence names were defined for the purposes of this work, the names of initial and last image corresponds to the nomenclature adopted in JIRAM data repository at NASA PDS. For each sequence, ranges of spatial resolution and emission angle for pixels in the latitude range 5N-10N are provided.</p>		

130 Geometric parameters of each image (geographic locations of pixel corners and centers, solar zenith, emission and phase angles, slant distance, etc.) are computed by means of the SPICE Toolkit [Acton, 1996, Acton et al., 2017] on the basis of the re-constructed kernel files available at <https://naif.jpl.nasa.gov/pub/naif/JUNO/kernels/>.

135 In this work we consider JIRAM imaging data rather than JIRAM spectra, given the much larger number of pixels available. Moreover, only images acquired on the Jupiter's night side are included, to avoid the potential residual contribution to total signal from reflection of solar radiation.

### 3. Methods

140 Our study used a series of spectral forward models to assess the trends of JIRAM signal  $I$  (as measured in M-filter images) versus emission angle  $\epsilon$ , for different values of scattering parameters and cloud total opacity  $\tau$ . As scattering parameters we consider 1) the single scattering albedo  $\omega_0$ , i.e.: the ratio between the scattering cross section and the total extinction cross section of a particle; 2) the asymmetry parameter  $g$ , i.e.: the average value of the cosine of scattering angle.  $g = 0$  implies isotropic scattering, while values close to 1 imply strong forward scattering.

145 We consider, for simplicity, the model of Hot Spots described in Grassi et al. [2017b]. Namely, this model assumes just one cloud, extended between -7 and +7 km (with respect to the 1 bar level) and the simplified

representation of the gas mixing ratios profiles described in Irwin et al., [1998]. Mixing ratios assumed for minor gases are the averages values observed in Hot Spots, as presented in Grassi et al. [2020].

The forward code adopted to compute spectra [Ignatiev et al., 2005] has been modified to exploit the correlated-k technique (as described in Irwin et al., [2008]). The multiple scattering is modelled by means of DISORT algorithm [Stamnes et al, 1998], in the assumption of a Henyey-Greenstein phase function (developed in Legendre polynomials up to 64 streams). Resulting simulated spectra are multiplied by the M-filter response function to obtain theoretical  $I$  values. We fit the resulting simulated limb-darkening functions with a Minnaret's function, i.e.:

$$\mu \stackrel{\text{def}}{=} \cos(\epsilon) \quad (1)$$

$$I(\mu) = I_0 \mu^k$$

where  $I_0$  and  $k$  are the free parameters of the fit. However, after initial tests, it became quickly evident that, even for a fixed pair of  $\omega_0$  and  $g$  values, 1)  $k$  - i.e.: the slope of the limb darkening curve in the  $\log(I)$ - $\log(\mu)$  space - is not constant with  $\mu$ , therefore making the Minnaret' function inadequate for our purposes, and, more important, 2) the  $k$  parameter at a fixed  $\mu$  varies substantially with the optical depth  $\tau$  (Fig.2 and Fig.3). Notably, the dependency of limb darkening upon opacity requires to include explicitly  $\tau$  in whatever empirical function adopted to model the data, even when considering more sophisticated functions of  $\mu$  and  $\epsilon$  than Minnaret's one. Therefore, the inference of  $\omega_0$  and  $g$  from the limb-darkening trend *in a specific location* is necessarily limited by the ambiguity of the  $\tau$  value to be assumed.

The method presented below circumvents this issue on the basis of several assumptions. First, we *assume* that  $\omega_0$  and  $g$  retain the same values over areas with different opacities inside the Hot Spot. Then, we exploit the monotonic decrease of signal with increasing  $\tau$  *at a fixed emission angle* for a given  $(\omega_0, g)$  pair<sup>1</sup>. In the *further assumption* that observed signal variations are dominated by variations of  $\tau$  - rather than by variations in composition, in vertical temperature profile or in cloud structure - the monotonic trend implies that the signal *at a given reference emission angle*  $I_{ref}$  - as inferred from the *observed* trend of  $I$  versus  $\epsilon$  in a specific location - can be adopted as an effective proxy for  $\tau$  there, in comparison with nearby areas. Potentially, this implies that repeated JIRAM observations of the same area at different emission angles allow us to sample the function  $I=f_0(I_{ref}, \epsilon)$  that, in turn, derives from a function  $I=f_1(\tau, \epsilon)$ . The theoretical shape of  $f_0$  can be computed for any given pair of  $\omega_0$  and  $g$  values from simulated observations and compared directly with the observed shape.

In this preliminary phase of the study, a quick data inspection made clear the difficulties related to the study of more opaque regions, where the JIRAM signal becomes extremely low and it can be affected considerably by residual calibration artefacts (such as imperfect removal of thermal background). Moreover, Braude et al. [2020] demonstrated that in regions of very high opacity (notably, the adjacent Equatorial Zone) the simplistic Hot-Spot model of a single, thin, cloud is no longer appropriate and multiple cloud layers must be included.

In practical terms, the data analysis proceeds along the following steps:

1. Each image in each of the sequences listed in Table I is mapped on the plane according to an azimuthal projection centered at 0°N, 180°W. The projection is performed for each of the four corners that define the IFOV of each individual JIRAM pixel. Before the projection, the longitude of each corner needs to be compensated for zonal motions occurring during the 3.5h acquisition period. This is done by considering the wind profile by Porco et al. [2003] and adjusting each longitude to the same common time (06.30 UT April 1<sup>st</sup>, 2018). Choi et al., [2013] pointed out the slight difference in speed between the NEB jetstreams (~114 m/s) and the dark formations associated to Hot Spots (~103 m/s). Direct tests have shown that this difference in speed has no impact on our final estimates of  $\omega_0$  and  $g$ , nor on their corresponding uncertainties.
2. Over the projection plane considered above, we define a uniform, orthogonal sampling grid. The step of the grid is equal to 183km at 0°N, 180°W. This grid defines the pixels for the re-sampled images produced at the next step (step 3).

<sup>1</sup>This holds true as long as clouds are located in a region of the atmosphere where temperature decreases with altitude, as is the case on Jupiter, where clouds are found well below the tropopause.

- 200 3. For each projected image defined at step 1, we produce a re-sampled version over the grid defined at step 2. Each pixel of the original images contributes to the value of a pixel on the re-sampled image with a weight proportional to its spatial overlap. This step is required to compare different JIRAM images consistently. The value of the emission angle at the center of each original pixel is also re-sampled in an analogous manner. Eventually, in each pixel of re-sampled images (now made spatially consistent with each other), we achieve a pair of values (signal  $I$ , emission angle  $\epsilon$ ) from each original image covering the area. *Hereafter, the term pixel will refer always to the re-sampled image.*
- 205 4. These pairs allow one to estimate, by interpolation on their linear fit, the limb-darkening function for each pixel. Notably, this interpolation is performed for the *sole* purpose of estimating the radiance expected - for each pixel - at the reference emission angle of  $\epsilon = 45^\circ$  ( $I_{45}$  hereafter) rather than to find a complete limb-darkening function such as those presented in Fig. 2. To reduce interpolation errors, the analysis is performed only for pixels where at least six pairs are available and the range of emission angles in JIRAM data encompasses the reference emission angle of  $45^\circ$  (i.e.: extrapolation is not allowed). Given the spread in original JIRAM data and the limited range of  $\epsilon$ , interpolation is performed assuming a linear fit, separately for each pixel (see examples in Fig. 4). Results of the fit are accepted (and the derived  $I_{45}$  included in the subsequent analysis) once the resulting intercept and slope have a value at least 10 times greater than corresponding uncertainties, as estimated from measurements errors and data spread. Eventually, uncertainties on linear fit parameters allow one to compute the uncertainty on the  $I_{45}$  estimated for each pixel. These latter uncertainties have a mean value of  $0.006 \text{ W}/(\text{m}^2 \text{ sr})$ , to be compared against a mean signal of  $0.15 \text{ W}/(\text{m}^2 \text{ sr})$ . Fig. 4 5 presents the  $I_{45}$  map derived in this analysis and corresponding uncertainties. Fig. 5b shows that uncertainties on  $I_{45}$  estimates are dominated by the spread of JIRAM data from different sequences – more evident in overlap regions - rather than by the uncertainty of signal in individual JIRAM images. This spread is due to residual calibration artefacts, most likely to be attributed to minor drifts in the instrument’s temperature and amplifiers gains. The value of  $\epsilon = 45^\circ$  is chosen as intermediate for the range of JIRAM observations (see Table I).
- 210 215 220 225 230 235 5. At this step of the analysis, for each pixel, the original set of pairs derived in step 3 become a set of triplets ( $I, I_{45}, \epsilon$ ),  $I_{45}$  being identical for every triplet from the same pixel. These triplets can be seen as direct samples ( $z, x, y$ ) of the  $z=f_0(x, y)$  surface (i.e.: the  $I=f_0(I_{\text{ref}}, \epsilon)$  function described above) and represent the final experimental constraint offered by JIRAM data to be considered for comparison against theoretical simulations. The triplets considered in this study are presented in Fig. 6. Notably, the study excludes all the pixels where  $I_{45} < 0.02 \text{ W}/(\text{m}^2 \text{ sr})$ , to reduce the possible effect of calibration artifacts, as discussed above. The overall distribution of the original pairs ( $I, \epsilon$ ) as derived in step 3 can also be inferred from Fig. 6, as the distribution of points in the horizontal ( $\epsilon$ ) and out-of-the-page ( $I$ , color-encoded) directions. To better clarify the experimental trends from JIRAM data, points in Fig. 6 were binned in the ( $I_{45}, \mu=\cos(\epsilon)$ ) space; mean and standard deviation of  $I$  inside each bin are presented in Fig. 7.

Considering now the theoretical simulations:

- 240 245 250 1. A set of simulated JIRAM measurements is computed for each possible combination of the following inputs.  $\epsilon$ : from  $32.5^\circ$  to  $62.5^\circ$  (with steps of  $5^\circ$ ),  $\tau = [0.01, 0.031, 0.1, 0.31, 1., 3.1, 10.]$ ,  $\omega_0 = [0.75, 0.8, 0.85, 0.9, 0.95]$ ,  $g = [0.22, 0.27, 0.32, 0.37, 0.42, 0.47, 0.52, 0.57, 0.62]$ . Spectra are computed considering the same model and the same methods described for the preliminary simulations discussed at the beginning of this section, then multiplied by the M-filter response function to estimate the JIRAM measurements  $I(\epsilon, \tau, \omega_0, g)$
2. For each possible fixed pair of ( $\omega_0, g$ ) considered in simulations - and separately for each possible value of opacity  $\tau$  - we interpolate the trend of signal  $I$  versus  $\epsilon$  to estimate signal at the reference emission angle of  $\epsilon = 45^\circ$ .
3. For the given pair of ( $\omega_0, g$ ), each simulation now provides a triplet ( $I, I_{45}, \epsilon$ ), being  $I_{45}$  the same for every simulation with the same  $\tau$ . The triplets for all  $\tau$  are used to perform a polynomial fit of the  $f_{0,\text{simul}}$  surface pertinent for the current pair of  $\omega_0$  and  $g$ .

For each pair of ( $\omega_0, g$ ), the parameters of the polynomial fit allow one to compute the value of  $f_{\text{simul}}$  at the ( $I_{45}, \epsilon$ ) positions of observed triplets described at point 5 of data analysis above, and to compute the total deviation against data as the sum of  $|f_{\text{simul}} - f_0|$  over the entire population of experimental triplets. This ensures that the comparison is indeed performed in regions of  $f_0$  actually covered by data. More explicitly,

being:  $j$  an index to label individual triplets,  $(I_{45}, \epsilon)_j$  the  $I_{45}$  value and emission angle respectively for the  $j$ -th triplet,  $f_{O,j}$  the observed signal  $I$  for the  $j$ -th triplet and  $f_{simul}(I_{45}, \epsilon)_j$  the simulated signal at the  $(I_{45}, \epsilon)_j$  position for a given  $(\omega_0, g)$  pair, we can compute

$$C(\omega_0, g) = \sum_{j=1}^N \frac{[f_{simul}(I_{45}, \epsilon)_j - f_{O,j}]^2}{S_j^2} / N \quad (2)$$

where  $S_j$  is an effective value of data spread in  $(I_{45}, \epsilon)$  position of the  $j$ -th triplet, as estimated from figure 7b, and  $N=133,653$  the total number of triplets considered.

This dimensionless quantity represents the cost function  $C$  to be minimized by varying  $(\omega_0, g)$ .

#### 4. Results

Fig. 8 presents the value of cost function  $C$  in the  $\omega_0 - g$  space. The lowest  $C$  value implies the best agreement with data. Despite the coarse sampling of  $\omega_0 - g$  space by our simulations (made necessary by the high computation times required by DISORT usage), a minimum is observed for  $\omega_0=0.90$  and  $g=0.37$ . In order to assess more directly the quality of the fit, it is useful to compare the trend of  $f_{simul}$  in the assumption of  $\omega_0=0.90$  and  $g=0.37$  with the mean experimental JIRAM values (Fig. 7a). Figure 9a presents this  $f_{simul}(I_{45}, \epsilon)$  over the same region of  $I_{45} - \epsilon$  space sampled by JIRAM data, as well as its difference with respect to figure 7a. The difference between theoretical and experimental surfaces (Fig. 9b) is, in general, well within the spread of experimental data as given in Fig. 7b (see Fig. 9c for the ratio between these quantities), especially in the regions better covered by experimental data (compare Fig. 9c and 7c).

Given the large number of steps in the analysis, it is difficult to perform a rigorous computation of the uncertainties on our  $\omega_0$  and  $g$  estimate.

To perform at least a rough assessment, we evaluated the variations of  $C$  associated with random fluctuations according to a simple bootstrap scheme. First, we computed the uncertainties  $\sigma_z$  associated with the  $I$  value of each triplet: this was done considering the uncertainties on the original JIRAM measurements and propagating them through the averaging process implicit in spatial resampling. Then, for each value of  $\omega_0$  and  $g$ , we performed one thousand trials  $t_i$ , by adding random fluctuations to the  $I$  value in each triplet (with zero mean and standard deviation equal to the specific  $\sigma_z$  pertinent to that triplet). At the same time, we also introduced random fluctuations on the  $I_{45}$  values of each triplet. These random fluctuations are statistically independent from those introduced on  $I$  and have standard deviations equal to the uncertainties on  $I_{45}$ , as estimated from the fit procedure described in previous section 3, step 4. Finally, we recomputed the corresponding value of  $C(\omega_0, g, t_i)$ . The standard deviation of this 1000-element population (with elements from different  $t_i$ ) can be seen as an effective uncertainty on the  $C$  values presented in Fig. 8b. The level curve presented there corresponds to the minimum of  $C$  plus three times its uncertainty. Our estimate can therefore be better described as  $\omega_0=0.90 \pm 0.05$  and  $g=0.37 \pm 0.15$ . These values can be compared against the estimates of Giles et al. [2015] in the same spectral region, who adopted  $\omega_0=0.9$  and  $g=0.7$  in their analysis of Cassini VIMS data. However, their work is based on data acquired “in the equatorial region of the planet ( $-2.5^\circ$  to  $2.5^\circ$ )” and therefore in very different physical conditions that those of the Hot Spots considered here.

In order to address the dependency of our results on the simulation assumptions, we repeated the fit procedure for the search of  $g$  and  $\omega_0$  best values, considering the following variations.

1. The entire set of simulated spectra computations was repeated by shifting, for all cases, the residual cloud from the original altitude  $[-7; +7]$  km down to  $[-14; -7]$  km. While the Galileo Entry Probe nephelometer measurements are indeed consistent with our original assumption (see summary table 5.1 in West et al., 2004, with the main cloud located between 0.75 and 1.3 bars), lower clouds in Hot Spots between 1.3 and 1.6 bars (altitudes between  $-7$  and  $-14$  km) are predicted by the theoretical model of Hueso et al. [1999]. Lower clouds lead to  $g$  and  $\omega_0$  values of 0.42 and 0.90 respectively, consistent - within uncertainties - with our estimate at the original cloud altitude.
2. The entire set of simulated spectra computations was repeated by increasing, for all cases, the air temperatures of  $+14$ K at all pressure levels. While the original  $T(p)$  profile is the same measured by the Galileo Entry Probe [Ragent et al., 1998], the theoretical model by Hueso et al. [1999] predicts temperature deficiencies in the Hot Spots up to 6K with respect to the surrounding

environment (Fig. 3, panel E there). Despite air temperatures highly exceeding expected variations, the resulting  $g$  and  $\omega_0$  values (0.42 and 0.85 respectively) are again consistent - within uncertainties - with our estimate from original assumptions.

3. The entire set of simulated spectra computations was repeated by expanding, for all cases, the residual cloud from the original extension [-7;+7] km to [-14,+7] km. Broader clouds lead to  $g$  and  $\omega_0$  values of 0.32 and 0.90 respectively, consistent - within uncertainties - with our estimate for the original extension assumption.

With a further assumption on cloud composition, the scattering parameters  $\omega_0$  and  $g$  derived from our analysis can provide constraints on the size distribution of cloud particles. Ammonium hydrosulfide ( $\text{NH}_4\text{SH}$ ) ice remains the most likely candidate as the *main* component of residual clouds over Hot Spots [Atreya et al., 1999], although our analysis of the solar-dominated short-wavelength spectral region presented in Grassi et al. [2021] suggests that contamination from other components may be important in other low-latitude regions of the planet. The coating of clouds with complex haze produced photochemically was originally proposed and extensively discussed by Atreya et al. [2005]. Complex refractive indices of solid ammonium hydrosulfide have been inferred from experimental transmissivity measurements described in the two independent studies by Ferraro et al., [1980] and Howett et al., [2007]. However, Ferraro et al., [1980] did not publish any final  $(n,k)$  table and the derived refractive indices have been circulated only informally in the science community since then. For the purposes of this study, we considered  $n=1.38$  and  $k=0.020$  as the Ferraro et al. [1980] effective values for the JIRAM M-filter. With the simplifying assumption of spherical particles (indeed, a rather strong assumption for ice particles) and for a given set of refractive indices, Mie theory allows us to compute the expected  $(\omega_0, g)$  for each possible pair of parameters effective radius ( $r_{\text{eff}}$ ) and variance ( $v$ ) describing a size distribution.

Fig. 10 presents the results of these Mie computations. Namely, we show the regions in the  $r_{\text{eff}} - v$  space centred around the contour lines corresponding to the  $\omega_0$  and  $g$  values inferred from JIRAM and bounded by their respective uncertainties. We define these regions as acceptance regions. Overlaps in acceptance regions of  $\omega_0$  and  $g$  shall be considered consistent with data. The Ferraro et al. [1980] set (Fig. 10a) is *only marginally* not consistent with JIRAM-derived estimates, mostly because of the difficulty in producing the required high  $\omega_0$  with high values of  $k$  (i.e.: absorption coefficients). The simulations based on Howett et al. [2007] data (Fig. 10b) produce the required combination of  $\omega_0$  and  $g$  in two distinct regions of  $r_{\text{eff}} - v$  space. The first region starts at  $v=0.2$  and  $r_{\text{eff}} = 0.3 \mu\text{m}$  and extend to smaller radii and higher variances. The second region develops between 2 and 4  $\mu\text{m}$  in a large range of variances. Notably, in neither of the two areas we have a crossing of the curve levels corresponding to  $\omega_0=0.90$  and  $g=0.37$ . *Pure ammonia ice* (as modelled by the refractive indices in Howett et al. [2007], Fig. 10c) produces a marginally consistent region, developing from  $v=0.6$  and  $r_{\text{eff}} = 0.2 \mu\text{m}$  and extending to smaller radii and higher variances. More convincing agreement is provided by hydrazine (as given in Clapp and Miller [1996], Fig. 10d), where the consistent area starts at  $v=0.1$  and  $r_{\text{eff}} = 0.2 \mu\text{m}$ . We also considered the tholin material [Imanaka et al., 2012] adopted in our previous paper Grassi et al., [2021], there chosen for its capability to reproduce the shape of the reflectance spectral maximum seen in JIRAM spectra at 2.74  $\mu\text{m}$ . While this material cannot be considered as a realistic candidate for the main component of Jupiter clouds (tholin are simulants of the products of  $\text{N}_2\text{-CH}_4$  photochemistry on Titan), it can nonetheless reproduce an external coating by a poorly-characterized compound that includes N-H bounds responsible for the absorption band observed in Jupiter spectra around 3  $\mu\text{m}$  (Sromovsky and Fry [2010]). Fig. 10e presents the results of Mie computation for pure tholin. The overlap of acceptance regions for  $\omega_0$  and  $g$  is achieved for  $0.4 < r_{\text{eff}} < 1 \mu\text{m}$  and a rather large range of  $v$ , including the very low values previously considered in literature (e.g.: 0.05 in Irwin et al., [1998]). Notably, among pure materials, tholin is the only one capable to produce crossing of the curve levels corresponding to  $\omega_0=0.90$  and  $g=0.37$ , achieved for  $v=0.3$  and  $r_{\text{eff}} = 0.45 \mu\text{m}$ .

For sake of completeness, we investigated the space  $r_{\text{eff}}\text{-}n\text{-}k$  (being  $n,k$  the real and imaginary parts of the refractive index at 4.8  $\mu\text{m}$ ) to determine which combinations are consistent with JIRAM data. The results are presented in Fig. 11. As previously observed, both  $\text{NH}_4\text{SH}$  (according Howett et al.) and tholin can produce consistent results, while other pure materials considered have imaginary components of refractive index either too low (hydrazine and ammonia) or too high ( $\text{NH}_4\text{SH}$  according Ferraro). From the panels on Fig. 10 we can also observe that the acceptance region where  $0.85 < \omega_0 < 0.95$  moves consistently toward low  $r_{\text{eff}}$  while  $k$  decreases.

We also considered the case of coated particles, formed by a  $\text{NH}_4\text{SH}$  core (as modeled by Howett et al., [2007] indices) and an external coating (i.e.: an outer layer) formed by ammonia, hydrazine or tholin. The scattering properties of these spherical coated particles were computed by mean of the BART code

[Quirantes, 2020]. Results are shown in Fig. 12, where we considered relative core-to-particle radius ratios of 0.85 and 0.15. These two values are intended to represent cases dominated by the core or by the coating, respectively. The coated particles considered in our simulations seem not adequate to produce a significant improve of overlap regions once compared to pure materials. Coated particles dominated by the  $\text{NH}_4\text{SH}$  core (Fig. 12a,c,e) retain a first region consistent with JIRAM estimates for  $2 < r_{\text{eff}} < 3 \mu\text{m}$  and second for  $r_{\text{eff}} < 0.4 \mu\text{m}$  and high  $v$  (very small for ammonia, more pronounced for tholin). In no case, we observe a crossing of the curve levels corresponding to  $\omega_0=0.90$  and  $g=0.37$ . Particles dominated by coatings (Fig. 12b,d,f) largely reproduce the properties previously observed for the corresponding pure materials. Tholin coating provides the most convincing match with JIRAM-derived values of  $\omega_0$  and  $g$ , for radii between 0.7 and 1  $\mu\text{m}$ . These values can be compared against the average particle radius of 0.8-1.1  $\mu\text{m}$  reported by Regent et al., [1998] for the residual cloud in Hot Spots measured by the Galileo Entry Probe nephelometer.

## 5. Discussion

The results presented in the previous section depends on a number of assumptions, most notably, on the hypothesis that brightness variations inferred from JIRAM data at the reference emission angle are mostly driven by variations in the optical thickness of clouds. Unfortunately, such an assumption cannot be circumvented with repeated observations of the same region, since such an observation does not – alone – address the dependence of limb-darkening function upon cloud opacity shown in Fig. 3. Longitudinal variations in the amounts of water vapour and ammonia inside Hot Spots, such as those presented in Grassi et al. [2020] (Fig. 7c and 8c there), and in Fletcher et al., [2020] are probably the main limit of our analysis. JIRAM estimates are consistent - within uncertainties - either with  $\text{NH}_4\text{SH}$ -dominated particles with radii between 2 and 3  $\mu\text{m}$  or, alternatively, with smaller particles (0.7 – 1  $\mu\text{m}$ ) dominated by contaminants (modelled by tholin). The latter scenario appears more convincing, given a. the best correspondence with the JIRAM estimates b. the consistency with independent size measurements by Galileo Entry Probe.

We noticed also that the refractive indices of the organic samples produced by ion bombardment of ices ( $\text{N}_2:\text{CH}_4:\text{CO}$ ) described in Baratta et al. [2015] have a value of [1.6, 0.07], also consistent with JIRAM data, as demonstrated by Fig. 11. Both in the case of Imanaka et al., [2012] tholin as well as for the organic samples of Baratta et al. [2015], the value of  $k$  in the band of JIRAM images is substantially risen by a sharp absorption feature located at about  $2180 \text{ cm}^{-1}$  ( $\sim 4.58 \mu\text{m}$ ), see Fig. 3,4,5 and 6 in Imanaka et al. and Fig. 11 in Baratta et al. The latter paper interprets this feature as caused by  $\text{C}\equiv\text{N}$  bounds in the sample. Furthermore, Carlson et al. [2016] describe the optical properties of a chromophore from photolyzed ammonia reacting with acetylene, to simulate the specific conditions of Jupiter atmosphere. Although these authors do not provide refractive indices values for the spectral range of our interest, they present an absorption spectrum of the material, with a clear sharp feature at  $2056 \text{ cm}^{-1}$  ( $\sim 4.86 \mu\text{m}$ ) interpreted as due to diazo hydrocarbons ( $\text{C}=\text{N}=\text{N}$ ). Therefore, it appears that intermediate  $k$  values ( $\sim 0.01$ ) along with low  $n$  values ( $\sim 1.6$ ) are common among the by-products of degradation of carbon- and nitrogen-bearing species. Results from Braude et al [2020] (Fig. 13d there) suggests a substantial increase of  $r_{\text{eff}}$  for the equatorial clouds considered by Giles et al. [2015]. Both Fig. 10 and 12 point toward a general increase of the asymmetry parameter  $g$  for increasing  $r_{\text{eff}}$ , and the discrepancy in the  $g$  estimates provided here and in Giles et al. [2015] can probably be reconciled considering the different particle sizes in the areas considered in the two studies. Notably,  $r_{\text{eff}}$  variations produce much less pronounced variations on  $\omega_0$ , that is indeed consistent in the two studies.

The model of Hot Spots presented by Friedson [2005] foresees that air parcels, flowing from west to east, descend from altitudes well above the 0.6 bar isosurface down to the 1.3-bar surface (base of residual cloud in our model) when they encounter the minimum of the Rossby wave (corresponding to brighter regions in the infrared), moving along the isolevels of potential temperature  $\Theta$  (Fig. 8 there). Substantial downdraft in the brightest regions of Hot Spots was also expected in the model of Hueso [1999]. This scenario is consistent with the occurrence of substantial amounts of contaminants produced by the combination of hydrocarbon haze particles falling from Jupiter's stratosphere and hydrazine created by photodissociation of ammonia in the upper troposphere (Atreya et al. [2005]) at the location of residual clouds over Hot Spots.

## Conclusions

415 JIRAM M-filter images acquired over Jupiter's 5- $\mu\text{m}$ -bright Hot Spots during the 12<sup>th</sup> Juno perijove  
passage have been analyzed to constrain, through the study of limb-darkening, the scattering properties of  
residual clouds. Our results point toward a relatively small particle size ( $r_{\text{eff}} \sim 1.0 \mu\text{m}$ ) for these clouds and  
the occurrence of contaminants, possibly over the putative cores of ammonium hydrosulfide. The presence  
of contaminants is consistent with the downward air motions expected by theoretical models over brightest  
420 regions of Hot Spots. Further theoretical work will be required to compare the typical life times of  $\text{NH}_4\text{SH}$   
cloud particles in Hot Spots (where they are subject to periodic sublimation/condensation associated to the  
vertical air motions associated to the Rossby wave) against the times required to accumulate an amount of  
contaminants on their surface adequate to mask their optical properties.

These results allow a refinement of previous analysis of JIRAM spectra, with improved treatment of  
425 scattering by residual clouds in computation of theoretical spectra. The information content of limb-  
darkening measurements demonstrated by this study strongly support the inclusion joint spectral/limb  
darkening measurements in the planning of instruments capable of acquiring extended spectral datasets at  
high spatial resolution, notably MAJIS on board the ESA's JUICE mission.

## Acknowledgements

430 This work was supported by the Italian Space Agency through ASI-INAF contract I/010/10/0 and 2014-  
050-R.0. SKA acknowledge support from NASA through the Juno Project. LNF is supported by a European  
Research Council Consolidator Grant (under the European Union's Horizon 2020 research and innovation  
programme, grant agreement No 723890) at the University of Leicester. GSO acknowledges support from  
435 NASA through funds that were distributed to the Jet Propulsion Laboratory, California Institute of  
Technology.

We thank Nikolay Ignatiev, Space Research Institute of the Russian Academy of Sciences, Moscow, Russia,  
for the permission to use his ARS code for forward simulations and his crucial suggestions.

440 We thank Jonathan Lunine, Cornell University, Ithaca (NY), USA, for his valuable comments and  
revisions.

We thank Riccardo Urso, Osservatorio Astrofisico di Catania, Italy, for his suggestions on optical properties  
of organic materials derived from ion bombardment of ices.

445 The JIRAM instrument was conceived and brought to reality by our late collaborator and institute Director  
Dr. Angioletta Coradini (1946-2011).

## Data availability

The original calibrated data (Adriani et al., 2019) used in this study can be found at this Planetary Data  
System URL: [https://atmos.nmsu.edu/PDS/data/PDS4/juno\\_jiram\\_bundle/data\\_calibrated/orbit12/](https://atmos.nmsu.edu/PDS/data/PDS4/juno_jiram_bundle/data_calibrated/orbit12/)

450 Geometric parameters of each image (geographic locations of pixel corners and centers, solar zenith,  
emission and phase angles, slant distance, etc.) were computed by means of the SPICE Toolkit [Acton,  
1996, Acton et al., 2017] on the basis of the re-constructed kernel files available at  
<https://naif.jpl.nasa.gov/pub/naif/JUNO/kernels/>

## References

- 455 Acton, C. H. (1996) *Ancillary Data Services of NASA's Navigation and Ancillary Information Facility*, Planetary and Space Science, 44(1), doi: 10.1016/0032-0633(95)00107-7
- Acton, C. H., et al. (2017) *A look toward the future in the handling of space science mission geometry*, Planetary and Space Science, doi: 10.1016/j.pss.2017.02.013
- 460 Adriani, A. et al. (2014) *JIRAM, the Jovian Infrared Auroral Mapper*, Space Sci Rev, doi:10.1007/s11214-014-0094-y
- Adriani, A. et al. (2018), *Geometric cyclonic patterns in Jovian Polar Regions*, Nature, doi:10.1038/nature25491
- Adriani, A., R. Noschese, and L. Huber (2019), *Juno JIRAM Bundle*, PDS Atmospheres (ATM) Node, <https://doi.org/10.17189/1518967>.
- 465 Antuñano, A., et al. (2018). *Infrared characterization of Jupiter's equatorial disturbance cycle*, Geophysical Research Letters, 45, 10,987–10,995. doi:10.1029/2018GL080382
- Arregi, J., et al. (2006) *Phase dispersion relation of the 5-micron hot spot wave from a long-term study of Jupiter in the visible*, Journal of Geophysical Research (Planets), vol. 111, p.9010- doi:10.1029/2005JE002653
- 470 Atreya, S. K., et al. (1999) *Comparison of the atmospheres of Jupiter and Saturn: deep atmospheric composition, cloud structure, vertical mixing, and origin*. Planet. Space Sci., 47, 1243-62. doi:10.1016/S0032-0633(99)00047-1
- Atreya, S. K., et al. (2005) *Jupiter's ammonia clouds—localized or ubiquitous?* Planet. Space Sci., 53 (5), 498-507, doi:10.1016/j.pss.2004.04.002
- 475 Baratta, G. A., et al. (2015) *Organic samples produced by ion bombardment of ices for the EXPOSE-R2 mission on the International Space Station*, Planetary & Space Sciences, 118, doi: 10.1016/j.pss.2015.08.011
- Braude, A. S., et al. (2020) *Colour and tropospheric cloud structure of Jupiter from MUSE/VLT: Retrieving a universal chromophore*, Icarus, 338, doi: 10.1016/j.icarus.2019.113589.
- 480 Carlson, B. E. et al. (1993) *Tropospheric gas composition and cloud structure of the Jovian north equatorial belt*, J. Geophys. Res., 98(E3), 5251–5290, doi:10.1029/92JE02737
- Carlson, R. W, et al. (2016) *Chromophores from photolyzed ammonia reacting with acetylene: Application to Jupiter's Great Red Spot*, Icarus, 274, doi: 10.1016/j.icarus.2016.03.008
- Choi, D., et al. (2013) *Meteorology of Jupiter's equatorial hot spot and plumes from Cassini*. Icarus, 223, 832-843, doi: 10.1016/j.icarus.2013.02.001
- 485 Clapp, M. L., and Miller, R. E. (1996) *Complex Refractive Indices of Crystalline Hydrazine from Aerosol Extinction Spectra*, Icarus, 123 (2), 396-403, doi: 10.1006/icar.1996.0166
- Drossart, P., et al. (1998) *The solar reflected component in Jupiter's 5- $\mu$ m spectra from NIMS/Galileo observations*, J. Geophys. Res., 103(E10), 23043–23049, doi:10.1029/98JE01899
- 490 Ferraro, J. R., et al. (1980) *Infrared Intensity Measurements of Cryodeposited Thin Films of NH<sub>3</sub>, NH<sub>4</sub>HS, H<sub>2</sub>S, and Assignments of Absorption Bands*, Appl. Spectrosc., 34, 525-533, doi: doi:10.1366/0003702804731339
- Fletcher, L. N. et al. (2016) *Mid-infrared mapping of Jupiter's temperatures, aerosol opacity and chemical distributions with IRTF/TEXES*, Icarus, 278, 128-161, doi:10.1016/j.icarus.2016.06.008.

- 495 Fletcher, L. N., et al. (2017), *Jupiter's North Equatorial Belt expansion and thermal wave activity ahead of Juno's arrival*, *Geophys. Res. Lett.*, 44, 7140–7148, doi:10.1002/2017GL073383.
- Fletcher, L. N., et al. (2020). *Jupiter's equatorial plumes and hot spots: Spectral mapping from Gemini/TEXES and Juno/MWR*. *Journal of Geophysical Research: Planets*, 125, e2020JE006399, doi: 10.1029/2020JE006399
- 500 Friedson, A. J. (2005) *Water, ammonia, and H<sub>2</sub>S mixing ratios in Jupiter's five-micron hot spots: A dynamical model*, *Icarus*, 177, p. 1-17, doi: 10.1016/j.icarus.2005.03.004
- Giles, R. S., et al. (2015) *Cloud structure and composition of Jupiter's troposphere from 5-m Cassini VIMS spectroscopy*, *Icarus*, 257, 457-470, doi:10.1016/j.icarus.2015.05.030.
- 505 Giles, R. S., et al. (2016) *Latitudinal variability in Jupiter's tropospheric disequilibrium species: GeH<sub>4</sub>, AsH<sub>3</sub> and PH<sub>3</sub>*, *Icarus*, doi:10.1016/j.icarus.2016.10.023
- Grassi, D., et al. (2017a), *Preliminary results on the composition of Jupiter's troposphere in hot spot regions from the JIRAM/Juno instrument*, *Geophys. Res. Lett.*, 44, 4615– 4624, doi:10.1002/2017GL072841.
- 510 Grassi et al. (2017b) *Analysis of IR-bright regions of Jupiter in JIRAM-Juno data: Methods and validation of algorithms*, *Journal of Quantitative Spectroscopy and Radiative Transfer*, doi:10.1016/j.jqsrt.2017.08.008
- Grassi, D. et al. (2018) *First Estimate of Wind Fields in the Jupiter Polar Regions From JIRAM-Juno Images* *J. Geophys. Res. (Planets)*, doi:10.1029/2018JE005555
- 515 Grassi, D. et al., (2020) *On the spatial distribution of minor species in Jupiter's troposphere as inferred from Juno JIRAM data*, *J. Geophys. Res. (Planets)*, doi: 10.1029/2019JE006206
- Grassi, D. et al (2021), *On the clouds and ammonia in Jupiter's upper troposphere from Juno JIRAM reflectivity observations*, *Monthly Notices of the Royal Astronomical Society*, 503 (4), doi: 10.1093/mnras/stab740
- Howett, C. J. A., et al. (2007) *Optical constants of ammonium hydrosulfide ice and ammonia ice* *J. Opt. Soc. Am. B* 24, 126-136, doi: 10.1364/JOSAB.24.000126
- 520 Hueso, R., et al. (1999) *A theoretical study of parcel stability and cloud distribution in a Jovian hot spot*, *Planetary & Space Sci.*, 47, p. 1263-1275, doi: 10.1016/S0032-0633(99)00049-5
- Ignatiev, N. I., et al., (2005) *Planetary Fourier spectrometer data analysis: Fast radiative transfer models*, *Planetary and Space Science*, 53 (10), 1035-1042, doi: 10.1016/j.pss.2004.12.009.
- 525 Imanaka, H., et al (2012), *Optical constants of Titan tholins at mid-infrared wavelengths (2.5–25 μm) and the possible chemical nature of Titan's haze particles*, *Icarus*, 218, p. 247-261, doi: 10.1016/j.icarus.2011.11.018
- Irwin, P. G. J., et al. (1998) *Cloud structure and atmospheric composition of Jupiter retrieved from Galileo near-infrared mapping spectrometer real-time spectra*, *J. Geophys. Res.*, 103(E10), 23001–23021, doi:10.1029/98JE00948.
- 530 Irwin, P. G. J., et al. (2001) *The Origin of Belt/Zone Contrasts in the Atmosphere of Jupiter and Their Correlation with 5-μm Opacity*, *Icarus*, 149, 397-415, doi:10.1006/icar.2000.6542.
- Irwin, P. G. J., et al. (2008), *The NEMESIS planetary atmosphere radiative transfer and retrieval tool*, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 109 (6), 1136-1150, doi: 10.1016/j.jqsrt.2007.11.006.

- 535 Nixon, C.A. et al., (2001) *Atmospheric Composition and Cloud Structure in Jovian 5- $\mu$ m Hotspots from Analysis of Galileo NIMS Measurements*, Icarus, 150, 48-68, doi:10.1006/icar.2000.6561.
- Ortiz, J. L., et al. (1998) *Evolution and persistence of 5- $\mu$ m hot spots at the Galileo probe entry latitude*, Journal of Geophysical Research, vol. 103, p.23051-23069, doi:10.1029/98JE00696
- 540 Orton et al., (1998) *Characteristics of the Galileo Probe entry site from earth-based remote sensing observations*. J. Geophys. Res., **103**, 22791-22814, doi:10.1029/98JE02380
- Orton, G. S., et al. (2017), *Multiple-wavelength sensing of Jupiter during the Juno mission's first perijove passage*, Geophys. Res. Lett., 44, 4607– 4614, doi:10.1002/2017GL073019.
- Porco, C., et al., (2003), *Cassini imaging of Jupiter's atmosphere, satellites, and rings*. Science, 299 (5612), 1541-1547, doi: 10.1126/science.1079462
- 545 Quirantes, A. (2020) *BART code source file*, retrieved at <https://www.ugr.es/~aquiran/codigos.htm>
- Ragent et al. (1998), *The clouds of Jupiter: Results of the Galileo Jupiter Mission Probe Nephelometer Experiment*, J. Geophys. Res., 103( E10), 22891– 22909, doi:10.1029/98JE00353.
- Rogers, J.H. (1995) *The Giant Planet Jupiter*, University Press, Cambridge
- 550 Roos-Serote, M., et al. (1998) *Analysis of Jupiter north equatorial belt hot spots in the 4–5  $\mu$ m range from Galileo/near-infrared mapping spectrometer observations: Measurements of cloud opacity, water, and ammonia*, J. Geophys. Res., 103(E10), 23023–23041, doi:10.1029/98JE01049.
- Roos-Serote, M., et al., (1999) *Constraints on the Tropospheric Cloud Structure of Jupiter from Spectroscopy in the 5- $\mu$ m Region: A Comparison between Voyager/IRIS, Galileo/NIMS, and ISO/SWS Spectra*, Icarus, 137, 315-340, doi:10.1006/icar.1998.6043.
- 555 Showman, A. P., Dowling, T. E. (2000) *Nonlinear simulations of Jupiter's 5-micron hot spots*. Science 289:1737-1740,doi:10.1126/science.289.5485.1737
- Sromovsky, L.A., and P.M. Fry, P.M. (2010) *The source of widespread 3- $\mu$ m absorption in Jupiter's clouds: Constraints from 2000 Cassini VIMS observations*, Icarus, 210 (1), 230-257, doi:10.1016/j.icarus.2010.06.039
- 560 Stamnes, K., et al (1988) *Numerically stable algorithm for discrete-ordinate-method radiative transfer in multiple scattering and emitting layered media*, Appl. Opt. 27, doi: 10.1364/AO.27.002502
- 565 Terrile, R. J., Westphal, J. A. (1977) *The vertical cloud structure of Jupiter from 5 micron measurements*, Icarus, vol. 30, p.274-281, doi:10.1016/0019-1035(77)90159-2
- West et al., (2004), *Jovian clouds and haze in Jupiter: the planet, satellites and magnetosphere*, by F. Bagenal, T.E. Dowling, and W.B. McKinnon (eds.). Cambridge planetary science series. Cambridge, UK: Cambridge University Press




Fig. 1 Mosaics of JIRAM M-band images acquired during the 12<sup>th</sup> Juno perijove passage. Each mosaic is derived from a single JIRAM sequence (see text for a definition). The title of each panel provides the central acquisition time according the scheme yymmdd\_hhmmss. Note the progressive signal increase at (7°N, 162°W) in panels a.-e. caused by decreasing emission angles (see Table I for further details). Cylindrical projection is adopted.




Fig. 2 Examples of theoretical limb darkening functions for typical Hot-Spot conditions, for different cloud opacities  $\tau$ .  $\mu$  is the cosine of emission angle  $\epsilon$ . Note that both axes are in logarithmic form, to highlight deviations from the Minnaret's formulation (see text).







Fig. 3 Examples of  $k$  values for Minnaret functions modeling theoretical limb darkening in typical Hot-Spot conditions. The parameter depends both on  $\mu$  as well as on  $\tau$ .




a.



b.




c.



d.

Fig. 4. JIRAM data from a given location acquired at different emission angles are used to infer, by interpolation of a linear fit, the value at the reference emission angle of  $45^\circ$ . Panels a.-d. present four examples at different absolute signal levels.



a.

b.

Fig. 5. Estimated signal at  $\epsilon = 45^\circ$  for the Hot-Spot region considered in this study (panel a.) and corresponding uncertainties (panel b.). Note the different color scales of the two panels.

Two distinct spatially-adjacent Hot Spots are covered by our data, being the one on the right ( $7^\circ\text{N}$ ,  $162^\circ\text{W}$ ) the same shown in Fig. 1, panels a.-e., and the one on the left ( $7^\circ\text{N}$ ,  $192^\circ\text{W}$ ) the same shown in Fig. 1, panels e.-f. Gaps in coverage are caused mostly by poor interpolation quality, often related to low intrinsic signal.






Fig. 6. Experimental sampling of the surface  $I = f_0(I_{45}, \mu)$ , as derived from JIRAM observations. Individual points represent individual triplets, as described in the text.



a



b

c

Fig. 7. Data points from Fig. 6, once binned in the ( $I_{45}$ ,  $\mu = \cos(\epsilon)$ ) space. Bins have widths of 0.005  $W/(m^2 sr)$  and 0.005 (dimensionless) in the two directions. a. Mean value of JIRAM signal ( $I$ ) inside each bin, b. Standard deviation of  $I$  inside each bin (note the different scale with respect to panel a.), c. Number of data points (triplets) inside each bin.







Fig. 8. The dimensionless cost function  $C$  – measuring the discrepancy between the observed and theoretical  $f_0$  surfaces (see text for definition) - for different values of  $\omega_0$  and  $g$ . The white level curve indicates the  $3\sigma$  uncertainty in the determination of minimum.



a



b

c

Fig. 9. a. Simulated  $I = f_{\text{simul}}(I_{45}, \epsilon)$  for  $\omega_0 = 0.90$  and  $g = 0.37$ , best-fit values inferred from Fig. 8 b. Difference between experimental data and the best-fit surface (i.e.: between Fig. 7a and Fig. 9a.) c. Ratio between fit discrepancies (Fig. 9a) and spread of experimental data (Fig. 7b).




Fig. 10. Contour lines for  $\omega_0$  (solid curves) and  $g$  (dashed curves) in the  $r_{\text{eff}}$  – variance space. Blue curves gives the values estimated from JIRAM data, red curves the acceptance region bounded by corresponding uncertainties as estimated from Fig. 6. a.  $\text{NH}_4\text{SH}$ , with complex refractive indices from Ferraro et al. b.  $\text{NH}_4\text{SH}$ , from Howett et al. c.  $\text{NH}_3$ , from Howett et al. d.  $\text{N}_2\text{H}_4$ , from Clapp and Miller, e. tholin type 2 from Imanaka et al.




Fig. 11. The plots display with black triangles the regions of the  $n$ - $k$  space (real and imaginary parts of refractive index respectively) that satisfy the two constraints  $0.85 < \omega_0 < 0.95$  and  $0.22 < g < 0.52$  for any  $r_{\text{eff}}$  in the range  $0.3$ - $3.0 \mu\text{m}$ . Red squares provide, for reference, the refractive indices at  $4.8 \mu\text{m}$  for different putative components of Jupiter's clouds. *Hy*:  $\text{N}_2\text{H}_4$ , from Clapp and Miller, *Am*:  $\text{NH}_3$ , from Howett et al., *Fe*:  $\text{NH}_4\text{SH}$ , from Ferraro et al., *Ho*:  $\text{NH}_4\text{SH}$ , from Howett et al., *To*: tholin type 2 from Imanaka et al. a. For a size distribution variance of 0.01 b. for a size distribution variance of 0.05.




Fig. 12. As fig. 10, for coated particles formed by a  $\text{NH}_4\text{SH}$  core (Howett et al.) and different coatings. a.-b.  $\text{NH}_3$  coating. c.-d.  $\text{N}_2\text{H}_4$  coating e.-f. tholin coating. 0.85 and 0.15 labels provide the core-to-particle radii ratio (0.85 large core, 0.15 small core)