




Publication Year	2015
Acceptance in OA	2020-03-07T10:47:45Z
Title	Enhanced detection of terrestrial gamma-ray flashes by AGILE
Authors	MARISALDI, MARTINO, ARGAN, ANDREA, URSI, ALESSANDRO, Gjesteland, T., FUSCHINO, FABIO, LABANTI, CLAUDIO, Galli, M., TAVANI, MARCO, PITTORI, Carlotta, VERRECCHIA, Francesco, D'Amico, F., Østgaard, N., MEREGHETTI, Sandro, CAMPANA, RICCARDO, Cattaneo, P. W., BULGARELLI, ANDREA, Colafrancesco, S., Dietrich, S., Longo, F., GIANOTTI, FULVIO, Giommi, P., Rappoldi, A., TRIFOGLIO, MASSIMO, TROIS, ALESSIO
Publisher's version (DOI)	10.1002/2015GL066100
Handle	http://hdl.handle.net/20.500.12386/23157
Journal	GEOPHYSICAL RESEARCH LETTERS
Volume	42



RESEARCH LETTER

10.1002/2015GL066100

Key Points:

- Tenfold improvement in TGF detection rate by AGILE is obtained after a major configuration change
- The sample is dead time-free, with TGF durations as short as 20 μs
- The relationship between TGF duration and likelihood of association with lightning is confirmed

Correspondence to:

M. Marisaldi,
marisaldi@iasfbo.inaf.it

Citation:

Marisaldi, M., et al. (2015), Enhanced detection of terrestrial gamma-ray flashes by AGILE, *Geophys. Res. Lett.*, 42, doi:10.1002/2015GL066100.

Received 7 SEP 2015

Accepted 19 OCT 2015

Accepted article online 22 OCT 2015

Enhanced detection of terrestrial gamma-ray flashes by AGILE

M. Marisaldi^{1,2}, A. Argan³, A. Ursi^{3,4}, T. Gjesteland^{2,5}, F. Fuschino^{1,6}, C. Labanti¹, M. Galli⁷, M. Tavani^{3,4}, C. Pittori^{8,9}, F. Verrecchia^{8,9}, F. D'Amico¹⁰, N. Østgaard², S. Mereghetti¹¹, R. Campana¹, P.W. Cattaneo¹², A. Bulgarelli¹, S. Colafrancesco^{9,13}, S. Dietrich¹⁴, F. Longo^{15,16}, F. Gianotti¹, P. Giommi⁸, A. Rappoldi¹², M. Trifoglio¹, and A. Trois¹⁷

¹INAF-IASF, National Institute for Astrophysics, Bologna, Italy, ²Birkeland Centre for Space Science, Department of Physics and Technology, University of Bergen, Norway, ³INAF-IAPS Roma, Rome, Italy, ⁴Dipartimento di Fisica, Università Tor Vergata, Rome, Italy, ⁵Department of Engineering Sciences, University of Agder, Norway, ⁶Dipartimento di Fisica e Astronomia, Università di Bologna, Bologna, Italy, ⁷ENEA, Bologna, Italy, ⁸ASI Science Data Center, Rome, Italy, ⁹INAF-OAR, Monteporzio Catone, Rome, Italy, ¹⁰Italian Space Agency, Rome, Italy, ¹¹INAF-IASF Milano, Milano, Italy, ¹²INFN Pavia, Pavia, Italy, ¹³School of Physics, University of the Witwatersrand, Johannesburg, South Africa, ¹⁴CNR-ISAC Roma, Rome, Italy, ¹⁵Dipartimento di Fisica, Università di Trieste, Trieste, Italy, ¹⁶INFN Trieste, Trieste, Italy, ¹⁷INAF-Osservatorio Astronomico di Cagliari, Capoterra, Italy

Abstract At the end of March 2015 the onboard software configuration of the Astrorivelatore Gamma a Immagini Leggero (AGILE) satellite was modified in order to disable the veto signal of the anticoincidence shield for the minicalorimeter instrument. The motivation for such a change was the understanding that the dead time induced by the anticoincidence prevented the detection of a large fraction of Terrestrial Gamma-Ray Flashes (TGFs). The configuration change was highly successful resulting in an increase of one order of magnitude in TGF detection rate. As expected, the largest fraction of the new events has short duration ($<100 \mu\text{s}$), and part of them has simultaneous association with lightning sferics detected by the World Wide Lightning Location Network. The new configuration provides the largest TGF detection rate surface density (TGFs/km²/yr) to date, opening prospects for improved correlation studies with lightning and atmospheric parameters on short spatial and temporal scales along the equatorial region.

1. Introduction

Terrestrial Gamma-ray Flashes (TGFs) are submillisecond bursts of gamma rays associated to lightning and thunderstorm activity and typically observed from space. They represent the observable manifestation of thunderstorm systems as the most energetic natural particle accelerators on Earth [Dwyer *et al.*, 2012]. Although many low-Earth orbiting satellites equipped with gamma-ray detectors exist, TGF observations were routinely reported by only four of them: the Burst And Transient Source Experiment on board the Compton Gamma-ray Observatory [Fishman *et al.*, 1994], the Reuven Ramaty High-Energy Solar Spectroscopic Imager [Smith *et al.*, 2005], the Gamma-ray Burst Monitor (GBM) on board the *Fermi* Gamma-ray Space Telescope [Briggs *et al.*, 2010], and the Astrorivelatore Gamma a Immagini Leggero (AGILE) mission [Marisaldi *et al.*, 2010], the last three currently operative. The reason for this relies on the typical time scale of this phenomenon ($\approx 100 \mu\text{s}$ average duration) which puts strong requirements on satellite data acquisition strategies and eventually on onboard trigger logic. In addition, the high TGF average fluence at satellite altitudes ($\approx 0.1 \text{ cm}^{-2}$ at 500–600 km) combined with their short duration makes all TGF detectors significantly affected by dead time and pileup effects [Grefenstette *et al.*, 2009; Gjesteland *et al.*, 2010; Briggs *et al.*, 2010; Marisaldi *et al.*, 2014]. These effects are instrument dependent and must be carefully modeled and accounted for when trying to derive general unbiased properties of the observed TGF population.

Marisaldi *et al.* [2014] showed that AGILE TGF detections by the minicalorimeter (MCAL) instrument [Labanti *et al.*, 2009] were heavily affected by the dead time induced by the anticoincidence (AC) shield designed to reject signals due to charged particles. Dead time, in fact, prevented the detection of events with duration shorter than $\approx 100 \mu\text{s}$, biasing the duration distribution toward larger values than observed by other spacecrafts. In addition, no precise matches with radio signals located by the World Wide Lightning Location Network (WWLLN) were obtained, consistently with this chance being inversely proportional to the TGF duration [Connaughton *et al.*, 2013; Dwyer and Cummer, 2013]. Moreover, since the AC is a paralyzable detector,

©2015. The Authors.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.




Figure 1. (a) Cumulative TGF rate, (b) longitude, (c) local time, and (d) number of counts distributions for the TGF samples in standard (blue) and enhanced (red hatches) configurations. Red-filled histograms are the corresponding distributions for events with a simultaneous WWLLN match.

the same set of observations can be due to events of different intrinsic fluence and duration; therefore, no one-to-one dead time correction can be applied. As a result the TGF intensity distribution evaluated in [Marisaldi *et al.*, 2014], distorted by dead time as well, was corrected by means of a forward folding approach based on assumptions on the intrinsic duration and fluence distributions.

Based on the understanding that AC-induced dead time was significantly affecting the satellite TGF detection capabilities, the AGILE science team, in agreement with the Italian Space Agency (ASI) and in collaboration with the industrial partners and the ASI Science Data Center (ASDC), has undertaken the necessary steps to modify the onboard software configuration in order to inhibit the AC veto signal on the MCAL detector. Starting from 23 March 2015, the new configuration with AC veto disabled on MCAL has been steadily active on board AGILE. The trigger logic parameters described in Marisaldi *et al.* [2014] were left unchanged except for the threshold on the 16 ms search time window, set from 22 to 41 counts to cope with the background rate increase. However, since all TGFs triggered on shorter time windows (293 μ s or 1 ms) this change does not affect the TGF trigger performance. In addition, the burst detection software is now not active during passage through the South Atlantic Anomaly (SAA). The following sections report the results on TGF detection after 3 months of operations in the new configuration.

2. Experimental Results

The same selection criteria on triggered data described in Marisaldi *et al.* [2014] were applied for TGF selection, thus allowing an unbiased comparison between the TGF samples obtained with the new (enhanced) and previous (standard) configurations. Between 23 March and 24 June 2015, a total of 279 TGFs have been recorded in the enhanced configuration. Figure 1 shows the cumulative detection rate and the distributions of longitude, local time, and counts for the samples in both configurations. The average daily TGF detection rate increased by 1 order of magnitude from 0.3 to 3 TGFs/d. Longitude and local time distributions clearly show the three-continental lightning chimneys and the early morning/afternoon peaks, respectively, suggesting that the number of false events in the sample is low. To support this statement, we consider the ratio between the number of TGFs detected above a TGF-active region and a control region with low lightning activity and consequently expected low TGF detection rate, following the approach outlined in Briggs *et al.* [2013]. TGF-active region is defined as the three-continental longitude bands (Central America: $[-90^\circ, -60^\circ]$;




Figure 2. Normalized duration (t_{50}) distribution for the enhanced (red hatches) and standard (blue line) TGF samples. Red filled histogram: t_{50} distribution of the enhanced TGFs with a simultaneous WWLLN match normalized to the total number of the enhanced sample. Cyan histogram: Fermi GBM t_{50} distribution calculated for counts above 300 keV, from *Connaughton et al.* [2013], Figure 3.

extends in the Southeast Pacific where lightning activity is very small [*Christian et al., 2003*], AGILE equatorial control region is close to the Intertropical Convergence Zone (ITCZ) where thunderstorm and lightning activity occur, especially during the observation period (March to May) [*Christian et al., 2003*].

The time series of each TGF has been fitted by a Gaussian model superimposed to a constant background by means of the maximum likelihood technique in order to avoid the loss of information due to arbitrary time binning, as described in *Marisaldi et al.* [2014]. The duration and intensity of each TGF are then extracted by the model fit parameters. The TGF duration is calculated as $t_{50} = 1.349\sigma$ and $t_{90} = 3.290\sigma$, where t_{50} and t_{90} are defined as the central time intervals including the 50% and 90% of the counts, respectively, and σ is the standard deviation of the Gaussian model. When the paucity of counts does not allow a coherent identification of the start and end points of a transient, t_{50} and t_{90} are convenient duration proxies. After close examination of the events light curves and fit results, we decided to exclude four events with poor convergence of the maximum likelihood procedure and seven events with closely spaced multiple peaks from the plots shown in Figure 1d as well as in subsequent figures. Figure 2 shows the duration distributions for the AGILE enhanced and standard samples and for Fermi GBM events




Figure 3. Black hatches histogram: distribution of the time difference ΔT between the WWLLN detection closest in time to a TGF and the TGF peak time corrected for the light propagation time between the WWLLN location and the satellite. Cyan line: same distribution for Fermi events as published in *Connaughton et al.* [2013], Figure 1, normalized to the number of AGILE events for visualization purposes.

Africa: $[-10^\circ, +30^\circ]$; and Maritime Continent: $[+100^\circ, +150^\circ]$). The control region is defined as the equatorial Pacific Ocean longitude band $[-140^\circ, -110^\circ]$ and has been chosen to be as close as possible to the southeast Pacific control region defined in *Briggs et al.* [2013]. Since in the case of AGILE the exposure on these regions is proportional to the longitude extent, we divide the number of observed TGFs in each region by this value. We obtain a TGF detection rate ratio of 20 between TGF-active and control regions, while *Briggs et al.* [2013] report a value 70 for the equivalent parameter. However, a direct comparison between these numbers cannot be done, mostly because of the different orbital inclination of the AGILE (2.5°) and Fermi (25.6°) spacecraft. In fact, while Fermi control region well

extends in the Southeast Pacific where lightning activity is very small [*Christian et al., 2003*], AGILE equatorial control region is close to the Intertropical Convergence Zone (ITCZ) where thunderstorm and lightning activity occur, especially during the observation period (March to May) [*Christian et al., 2003*].

The time series of each TGF has been fitted by a Gaussian model superimposed to a constant background by means of the maximum likelihood technique in order to avoid the loss of information due to arbitrary time binning, as described in *Marisaldi et al.* [2014]. The duration and intensity of each TGF are then extracted by the model fit parameters. The TGF duration is calculated as $t_{50} = 1.349\sigma$ and $t_{90} = 3.290\sigma$, where t_{50} and t_{90} are defined as the central time intervals including the 50% and 90% of the counts, respectively, and σ is the standard deviation of the Gaussian model. When the paucity of counts does not allow a coherent identification of the start and end points of a transient, t_{50} and t_{90} are convenient duration proxies. After close examination of the events light curves and fit results, we decided to exclude four events with poor convergence of the maximum likelihood procedure and seven events with closely spaced multiple peaks from the plots shown in Figure 1d as well as in subsequent figures. Figure 2 shows the duration distributions for the AGILE enhanced and standard samples and for Fermi GBM events when only counts with energy larger than 300 keV are considered [*Connaughton et al., 2013*], to match the MCAL energy threshold. The duration distribution for TGFs with a close WWLLN match is also included, as discussed in the following. The median of the t_{50} distribution is $86 \mu\text{s}$ and $290 \mu\text{s}$ for the enhanced and standard configurations, respectively, clearly indicating that the enhanced configuration allows the detection of much shorter events than before, as expected.

The database of lightning detected by the World Wide Lightning Location Network (WWLLN) [*Rodger et al., 2009*] has been searched for correlation with the enhanced TGF sample. Figure 3 shows the distribution of the time difference ΔT between the WWLLN detection closest in time to a TGF and the TGF peak time corrected for the light propagation time between the WWLLN location and the satellite, assuming a source

production altitude of 15 km [Dwyer and Smith, 2005]. Given the low number of counts, the peak time of a TGF is defined by the centroid of the Gaussian model fit described above. A total number of 39 sferics within 200 μ s from the TGF peak time has been observed, hereafter defined as simultaneous sferics according to the definition in Connaughton et al. [2013], corresponding to 14% of the sample. The same histogram obtained for Fermi data and shown in Figure 1 of Connaughton et al. [2013], normalized to the number of AGILE events, is also shown for reference.

3. Discussion

The inhibition of the AC veto for the AGILE MCAL instrument has clearly resulted in the detection of a much larger number of TGFs than with the standard configuration, enhancing the sensitivity for events lasting less than 100 μ s. These results validate the data interpretation provided in Marisaldi et al. [2014] regarding the role of dead time in biasing the observed duration and intensity distributions and the lack of detection of simultaneous sferics.

Once the contribution by the AC has been removed, the major source of dead time in the enhanced configuration is the data readout of the scintillation bars comprising MCAL. Each of the 30 bars (26 were active at the time the data presented here were collected) acts as an independent nonparalyzable detector, requiring 20 μ s for a photon acquisition after the discriminator has fired. Any detector trigger occurring during this fixed time is rejected. Although the dead time per count is larger than that reported for Fermi GBM (2.6 μ s [Briggs et al., 2013]), the key point to overall dead time effects mitigation is the MCAL spatial segmentation. Since each of the MCAL bars is independent from the others, two consecutive photons separated in time less than 20 μ s are promptly recorded, provided that they hit two different bars ready for data acquisition. We addressed the issue of MCAL-induced dead time by means of dedicated Monte Carlo simulations, described in details in Marisaldi et al. [2014], using the full AGILE mass model and a typical TGF spectrum as reported in Dwyer and Smith [2005]. We considered a TGF as described by two observables: its fluence at satellite altitude F and its duration expressed in terms of t_{50} . We then consider the combined effect of detector effective area ($A_{\text{eff}} \approx 220 \text{ cm}^2$ for a typical TGF spectrum incident at 60° off-axis angle) and dead time as a function f that maps the $(F \times t_{50})$ into the $(N^{\text{obs}} \times t_{50}^{\text{obs}})$ space, where N^{obs} and t_{50}^{obs} are the measured number of counts and duration. The effective area for 60° off-axis angle has been chosen as an average value for all the possible incoming off-axis angles. The maximum effective area, corresponding to the photon beam hitting orthogonal to the MCAL plane is just 15% larger. We note that, since the detectors are nonparalyzable, the function f can be inverted to extract the expected true fluence and duration from the observed counts and duration; i.e., each TGF can be individually corrected for dead time. This situation is radically different from the standard configuration with the AC acting as a paralyzable detector. In order to build the function f we generated 10^7 simulated TGFs uniformly distributed in the parameter space given by $(0.01 \text{ cm}^{-2} < F < 4 \text{ cm}^{-2}) \times (0.002 \text{ ms} < t_{50} < 0.4 \text{ ms})$. For each simulated TGF the expected number of counts in MCAL was defined according to F and the average effective area given by simulation results; the time series of the counts were randomly extracted according to a Gaussian time profile with $\sigma = 0.74 t_{50}$; and finally, each count was assigned to a detector bar based on a uniform random distribution. We then apply to the MCAL time series the effect of the dead time induced by the MCAL detector processing time, rejecting all counts within a 20 μ s time interval following a count on the same scintillating bar. For each simulated TGF we then count the observed number of counts N^{obs} and evaluate the observed duration t_{50}^{obs} by calculating the standard deviation of the counts time series, multiplied by factor 1.349 to convert from standard deviation to t_{50} . With respect to the simulations described in Marisaldi et al. [2014], we modified the code to account for a 50% fraction of double counts, i.e., counts involving two bars, mostly due to Compton scattering of photons between neighboring bars, which affects the number of active bars and can eventually increase the dead time fraction. The chosen double counts fraction is a worst case estimate, the typical fraction for TGFs being of order of $\approx 30\%$. Although this change, we do not observe a significant variation with respect to previous simulations. Figure 4 presents the enhanced TGF sample in the $(N^{\text{obs}} \times t_{50}^{\text{obs}})$ parameter space. The lines corresponding to true fluence and duration values are superimposed. As a rule of thumb, dead time is significantly affecting the sample in those regions where the lines deviate significantly from a parallel grid. The effect of dead time is that the observed number of photons in a TGF is less than the true number of photons that hit the detector. Also, the estimated duration of the TGF gets longer as the losses due to dead time increase. With the current settings the fluence at detection threshold is $\approx 0.05 \text{ cm}^{-2}$ as evidenced by the TGF population shown in Figure 4.




Figure 4. Enhanced sample (black circles) in the observed ($N^{obs} \times t_{50}^{obs}$), i.e., counts \times duration, parameter space. The color contour highlights the region with the highest detection rate (five logarithmic bins per decade in both coordinates). Magenta dots are the events with a simultaneous WWLLN match. Red and black contours mark points corresponding to the same value of t_{50} (ms) and fluence F (cm^{-2}), respectively, according to the MCAL dead time model for 60° off-axis angle. To obtain N^{obs} , F must be multiplied times the effective area of $\approx 220 \text{ cm}^2$.

The median of the t_{50} distribution for the enhanced sample is $86 \mu\text{s}$, significantly shorter than the $290 \mu\text{s}$ obtained for the standard sample. This median value is also shorter than the $100 \mu\text{s}$ reported for the Fermi GBM sample [Briggs *et al.*, 2013], but a greater evidence for the difference in the duration distribution for the two samples is obtained if we consider the fraction of events with $t_{50} \leq 50 \mu\text{s}$, as shown in Figure 2. Briggs *et al.* [2013] noted the lack of very short events in the Fermi sample suggesting it was a consequence of dead time. According to our dead time model the observed AGILE enhanced sample is in a region where the effect of dead time is very low and substantially negligible with respect to the errors due to counting statistics, as shown in Figure 4. We note from Figure 4 that high-fluence TGFs tend to have a longer duration than dimmer ones, e.g. no TGF with $F \geq 0.1 \text{ cm}^{-2}$ is found with $t_{50} \leq 50 \mu\text{s}$. According to our dead time model, this is not an observational bias due to dead time, but a physical feature of the TGF population that could be a test bench for production models. For example, Dwyer [2012] reports that, in the Relativistic Feedback (RF) model longer TGFs produce less runaway electrons than shorter ones, for a given electric field configuration. This is apparently in contrast with our observations. Conversely, the hypothesis that longer TGFs are the result of closely spaced shorter events can be ruled out based on the work by Connaughton *et al.* [2013], where an inverse proportionality between TGF duration and the likelihood of association with WWLLN sferics is firmly established. Whereas the detection of low fluence events is limited by the detector effective area, there is no instrumental issue preventing the detection of very short events with duration of $\approx 20 \mu\text{s}$ or less. However, we cannot neglect the contribution of Compton scattering of photons through the atmosphere to the observed duration of TGFs at satellite altitude [Østgaard *et al.*, 2008; Grefenstette *et al.*, 2008; Celestin and Pasko, 2012; Fitzpatrick *et al.*, 2014]. Although this effect is more evident for low energy photons below MCAL threshold, Celestin and Pasko [2012] showed that also instantaneously produced 1 MeV photons can be spread due to Compton scattering over a time interval as long as $50 \mu\text{s}$ at satellite altitude. Therefore, the observed $\approx 20 \mu\text{s}$ lower limit in TGF duration may be a measure of the minimum incompressible time spread due to Compton scattering rather than the intrinsic lower limit of TGF duration.

Concerning the correlation with WWLLN sferics, we first note that no simultaneous sferics was observed for the standard sample [Marisaldi *et al.*, 2014], and the reason for that was correctly identified in the bias toward longer events due to dead time suppression of short events, the latter of which are more likely associated to lightning sferics according to Connaughton *et al.* [2013]. The characteristics of the sample with simultaneous WWLLN sferics follow the general pattern discussed in Connaughton *et al.* [2013], with few significant

quantitative differences. The overall fraction of TGFs with simultaneous sferics is 14% while it is 30% in the Fermi sample. This fraction rises to 33% for TGF duration shorter than 50 μs , to be compared to the 60% value reported by *Connaughton et al.* [2013]. The anticorrelation between TGF duration and the probability of being associated to a simultaneous sferics is confirmed, as can be seen in Figure 2. The difference in overall simultaneous sferics detection probability may reside in the different orbital inclination of the AGILE and Fermi missions that make the two spacecrafts cover regions where the WWLLN detection efficiency is remarkably different. In particular, the WWLLN detection efficiency over equatorial Africa, where $\approx 40\%$ of the AGILE TGFs are observed, can be on average the 20% of the efficiency over North America and the Pacific region [*Hutchins et al.*, 2012], i.e., about 25% of the efficiency for the other low latitude regions. This scenario is supported by the low number of simultaneous detections observed over Africa, as shown in Figure 1. If we assume the WWLLN detection efficiency on the remaining equatorial regions covered by AGILE orbit is $\approx 80\%$ the average value for Fermi, which is reasonable considering the plots in *Hutchins et al.* [2012], we can estimate the ratio between the number of TGFs with WWLLN simultaneous match for AGILE and Fermi to be $R_{AF} = (0.4 \times 0.25 + 0.6) \times 0.8 = 0.56$. Although very simplified, this estimate is remarkably consistent with the observed value $0.33/0.60 = 0.55$ obtained for TGF with duration shorter than 50 μs . However, WWLLN efficiency significantly changes as a function of local time and generally improves over time as long as new stations are added to the network; therefore, a more detailed comparison between AGILE and Fermi association rate is difficult.

The WWLLN-TGF time difference Δt distribution shown in Figure 3 could appear to be bimodal, although we found no correlation between Δt and any TGF parameter. We therefore regard this feature as due to the limited statistics. The root-mean-square (RMS) of the Δt distribution within $\pm 200 \mu\text{s}$ is 63 μs . This error value can be regarded as the quadrature sum of several independent timing error contributions, namely, the WWLLN accuracy ($\approx 15 \mu\text{s}$ [*Hutchins et al.*, 2012]), the error on the TGF peak determination by the maximum likelihood procedure ($\approx 10 \mu\text{s}$), and the uncertainty on TGF production height ($\sim 10 \text{ km} \approx 30 \mu\text{s}$). The uncertainty on the GPS-provided AGILE position and the MCAL timing accuracy ($\approx 2 \mu\text{s}$) add negligible contributions. An additional timing uncertainty term of order of $\approx 50 \mu\text{s}$ is still missing to justify the observed Δt distribution RMS if we assume that the TGF-producing electrons are responsible also for the sferics radio pulses [*Connaughton et al.*, 2013; *Dwyer and Cummer*, 2013]. However, *Cummer et al.* [2011] showed that the electromagnetic waveform associated to TGFs can be very complex, containing several fast pulses, and that the pulse corresponding to the localization by ground-based lightning location networks may not be the closest in time to the TGF. In fact, this may be the source of the missing term in the uncertainty analysis discussed above. Therefore, even if we can narrow down to $\pm 100 \mu\text{s}$ the definition for simultaneity, provided the distribution of Figure 3 is confirmed with higher statistics, it will be difficult to further improve this value to gather more information on the relative timing between lightning and TGF.

4. Conclusion

The enhanced configuration, which has been running on board AGILE since 23 March 2015, has been highly successful resulting in a TGF detection rate increase of about 1 order of magnitude and opening up unique scientific opportunities for the understanding of the relationship between TGFs and lightning activity. For the first time, we present a TGF sample unbiased by dead time effects, which may serve as a test bench for production models. We also provide an independent confirmation of the anticorrelation between TGF duration and the likelihood of association with simultaneous WWLLN lightning sferics reported in *Connaughton et al.* [2013]. The large number of expected events, $\approx 1000/\text{yr}$, larger than that of Fermi GBM in continuous Time-Tagged Event data acquisition mode [*Briggs et al.*, 2013], concentrated in a narrow latitude band across the equator thanks to the AGILE orbital inclination of 2.5° , provides an unprecedentedly high TGF detection rate surface density, i.e., number of TGFs observed per unit area per unit time. In turn, this should allow enough counting statistics for correlation studies with lightning activity on small spatial and time scales. Up to now, only correlation studies on timescales of several years have been reported [*Smith et al.*, 2010; *Fuschino et al.*, 2011], so any seasonal variability giving possible clues on the TGF/lightning relation has been smeared out.

We recall here that the selection criteria used in this work for TGF identification are the same as those previously used in *Marisaldi et al.* [2014], including the requirement for the maximum photon energy not to be greater than 30 MeV. The availability of a sample of events with a firm association to WWLLN sferics, therefore surely associated to a TGF process, will allow to relax all selection criteria including the cut on the maximum photon energy, possibly shedding light on the long-standing issue of the highest photon energy in TGFs [*Tavani et al.*, 2011].

Acknowledgments

AGILE is a mission of the Italian Space Agency (ASI), with coparticipation of INAF (Istituto Nazionale di Astrofisica) and INFN (Istituto Nazionale di Fisica Nucleare). This work was carried out in the frame of the ASI-INAF agreement I/028/12/0. This study was supported by the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement 320839 and the Research Council of Norway under contracts 208028/F50, 216872/F50, and 223252/F50 (CoE). The authors wish to thank the World Wide Lightning Location Network (<http://wwlln.net>), a collaboration among over 50 universities and institutions, for providing the lightning location data used in this paper. The properties of the TGF sample presented in this work are publicly available at the ASI Science Data Center (ASDC) website: <http://www.asdc.asi.it/mcaletgfcats/>. All other MCAL data used in this work are available upon request from M. Marisaldi (marisaldi@iasfbo.inaf.it). The authors thank the AGILE industrial team at Compagnia Generale per lo Spazio (CGS) and Telespazio for their support during the configuration change. The authors also thank Valerie Connaughton for her helpful support and two anonymous reviewers for their useful suggestions. This work is dedicated to the memory of Paolo Sabatini, AGILE Program Manager at CGS, whose effort largely contributed to the success of the AGILE mission.

References

- Briggs, M. S., et al. (2010), First results on terrestrial gamma ray flashes from the Fermi Gamma-ray Burst Monitor, *J. Geophys. Res.*, *115*, A07323, doi:10.1029/2009JA015242.
- Briggs, M. S., et al. (2013), Terrestrial gamma-ray flashes in the Fermi era: Improved observations and analysis methods, *J. Geophys. Res. Space Physics*, *118*, 3805–3830, doi:10.1002/jgra.50205.
- Celestin, S., and V. P. Pasko (2012), Compton scattering effects on the duration of terrestrial gamma-ray flashes, *Geophys. Res. Lett.*, *39*, L02802, doi:10.1029/2011GL050342.
- Christian, H. J., et al. (2003), Global frequency and distribution of lightning as observed from space by the Optical Transient Detector, *J. Geophys. Res.*, *108*(D1), 4005, doi:10.1029/2002JD002347.
- Connaughton, V., et al. (2013), Radio signals from electron beams in terrestrial gamma ray flashes, *J. Geophys. Res. Space Physics*, *118*, 2313–2320, doi:10.1029/2012JA018288.
- Cummer, S. A., G. Lu, M. S. Briggs, V. Connaughton, S. Xiong, G. J. Fishman, and J. R. Dwyer (2011), The lightning-TGF relationship on microsecond timescales, *Geophys. Res. Lett.*, *38*, L14810, doi:10.1029/2011GL048099.
- Dwyer, J. R. (2012), The relativistic feedback discharge model of terrestrial gamma ray flashes, *J. Geophys. Res.*, *117*, A02308, doi:10.1029/2011JA017160.
- Dwyer, J. R., and S. A. Cummer (2013), Radio emissions from terrestrial gamma-ray flashes, *J. Geophys. Res. Space Physics*, *118*, 3769–3790, doi:10.1002/jgra.50188.
- Dwyer, J. R., and D. M. Smith (2005), A comparison between Monte Carlo simulations of runaway breakdown and terrestrial gamma-ray flash observations, *Geophys. Res. Lett.*, *32*, L22804, doi:10.1029/2005GL023848.
- Dwyer, J. R., D. M. Smith, and S. A. Cummer (2012), High-energy atmospheric physics: Terrestrial gamma-ray flashes and related phenomena, *Space Sci. Rev.*, *173*, 133–196, doi:10.1007/s11214-012-9894-0.
- Fishman, G. J., et al. (1994), Discovery of intense gamma-ray flashes of atmospheric origin, *Science*, *264*, 1313–1316.
- Fitzpatrick, G., et al. (2014), Compton scattering in terrestrial gamma-ray flashes detected with the Fermi gamma-ray burst monitor, *Phys. Rev. D*, *90*(4), 043008, doi:10.1103/PhysRevD.90.043008.
- Fuschino, F., et al. (2011), High spatial resolution correlation of AGILE TGFs and global lightning activity above the equatorial belt, *Geophys. Res. Lett.*, *38*, L14806, doi:10.1029/2011GL047817.
- Gjesteland, T., N. Østgaard, P. H. Connell, J. Stadsnes, and G. J. Fishman (2010), Effects of dead time losses on terrestrial gamma ray flash measurements with the Burst and Transient Source Experiment, *J. Geophys. Res.*, *115*, A00E21, doi:10.1029/2009JA014578.
- Grefenstette, B. W., D. M. Smith, J. R. Dwyer, and G. J. Fishman (2008), Time evolution of terrestrial gamma ray flashes, *Geophys. Res. Lett.*, *35*, L06802, doi:10.1029/2007GL032922.
- Grefenstette, B. W., D. M. Smith, B. J. Hazelton, and L. I. Lopez (2009), First RHESSI terrestrial gamma ray flash catalog, *J. Geophys. Res.*, *114*, A02314, doi:10.1029/2008JA013721.
- Hutchins, M. L., R. H. Holzworth, J. B. Brundell, and C. J. Rodger (2012), Relative detection efficiency of the World Wide Lightning Location Network, *Radio Sci.*, *47*, RS6005, doi:10.1029/2012RS005049.
- Labanti, C., et al. (2009), Design and construction of the Mini-Calorimeter of the AGILE satellite, *Nucl. Instrum. Methods Phys. Res. A*, *598*, 470–479, doi:10.1016/j.nima.2008.09.021.
- Marisaldi, M., et al. (2010), Detection of terrestrial gamma ray flashes up to 40 MeV by the AGILE satellite, *J. Geophys. Res.*, *115*, A00E13, doi:10.1029/2009JA014502.
- Marisaldi, M., et al. (2014), Properties of terrestrial gamma ray flashes detected by AGILE MCAL below 30 MeV, *J. Geophys. Res. Space Physics*, *119*, 1337–1355, doi:10.1002/2013JA019301.
- Østgaard, N., T. Gjesteland, J. Stadsnes, P. H. Connell, and B. Carlson (2008), Production altitude and time delays of the terrestrial gamma flashes: Revisiting the burst and transient source experiment spectra, *J. Geophys. Res.*, *113*, A02307, doi:10.1029/2007JA012618.
- Rodger, C. J., J. B. Brundell, R. H. Holzworth, and E. H. Lay (2009), Growing detection efficiency of the world wide lightning location network, *AIP Conf. Proc.*, *1118*, 15–20, doi:10.1063/1.3137706.
- Smith, D. M., B. J. Hazelton, B. W. Grefenstette, J. R. Dwyer, R. H. Holzworth, and E. H. Lay (2010), Terrestrial gamma ray flashes correlated to storm phase and tropopause height, *J. Geophys. Res.*, *115*, A00E49, doi:10.1029/2009JA014853.
- Smith, D. M., et al. (2005), Terrestrial gamma-ray flashes observed up to 20 MeV, *Science*, *307*, 1085–1088.
- Tavani, M., et al. (2011), Terrestrial gamma-ray flashes as powerful particle accelerators, *Phys. Rev. Lett.*, *106*(1), 018501, doi:10.1103/PhysRevLett.106.018501.