



Publication Year	2021
Acceptance in OA	2022-06-14T15:41:15Z
Title	AMBITION - comet nucleus cryogenic sample return
Authors	Bockelée-Morvan, D., FILACCHIONE, GIANRICO, Altwegg, Kathrin, BIANCHI, ELEONORA, Bizzarro, Martin, Blum, Jürgen, Bonal, Lydie, CAPACCIONI, FABRIZIO, Choukroun, Mathieu, CODELLA, CLAUDIO, Cottin, Hervé, Davidsson, Björn, DE SANCTIS, MARIA CRISTINA, Drozdovskaya, Maria N., Engrand, Cécile, Galand, Marina, Güttler, Carsten, Henri, Pierre, Herique, Alain, Ivanovski, Stavro, Kokotanekova, Rosita, Levasseur-Regourd, Anny-Chantal, Miller, Kelly E., Rotundi, Alessandra, Schönbächler, Maria, Snodgrass, Colin, Thomas, Nicolas, TUBIANA, Cecilia, Ulamec, Stephan, Vincent, Jean-Baptiste
Publisher's version (DOI)	10.1007/s10686-021-09770-4
Handle	http://hdl.handle.net/20.500.12386/32319
Journal	EXPERIMENTAL ASTRONOMY
Volume	54

AMBITION – Comet Nucleus Cryogenic Sample Return

D. Bockelée-Morvan · Gianrico Filacchione ·
Kathrin Altwegg · Eleonora Bianchi · Martin
Bizzarro · Jürgen Blum · Lydie Bonal ·
Fabrizio Capaccioni · Mathieu Choukroun ·
Claudio Codella · Hervé Cottin · Björn
Davidsson · Maria Cristina De Sanctis ·
Maria Drozdovskaya · Cécile Engrand ·
Marina Galand · Carsten Güttler · Pierre
Henri · Alain Herique · Stavro Ivanovski ·
Rosita Kokotanekova · Anny-Chantal
Levasseur-Regourd · Kelly E. Miller ·
Alessandra Rotundi · Maria Schönbächler ·
Colin Snodgrass · Nicolas Thomas · Cecilia
Tubiana · Stephan Ulamec · Jean-Baptiste
Vincent

Received: date / Accepted: date

D. Bockelée-Morvan

LESIA, Observatory of Paris, PSL University, CNRS, Sorbonne University, University of Paris, 5 place Jules Janssen, 92195, Meudon Cedex, France E-mail: dominique.bockelee@obspm.fr

G. Filacchione · F. Capaccioni · M.C. De Sanctis

Istituto di Astrofisica e Planetologia Spaziali, Istituto Nazionale di Astrofisica, via del Fosso del Cavaliere 100, 00133 Rome, Italy

K. Altwegg

Physikalisches Institut der Universität Bern, Sidlerstr. 5, 3012 Bern, Switzerland

E. Bianchi · C. Codella · L. Bonal · A. Herique

Univ. Grenoble Alpes, Institut de Planétologie et d'Astrophysique de Grenoble, 38000 Grenoble, France

M. Bizzarro

Centre for Star and Planet Formation and Natural History Museum of Denmark, University of Copenhagen, DK-1350 Copenhagen, Denmark

J. Blum

Institut für Geophysik und extraterrestrische Physik, Technische Universität Braunschweig, Mendelssohnstr. 3, 38106 Braunschweig, Germany

M. Choukroun · B. Davidsson

Jet Propulsion Laboratory/California Institute of Technology, 4800 Oak Grove Dr., Pasadena, California, 91109, USA

C. Codella

INAF, Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125, Firenze, Italy

Keywords Space missions – Solar System – Comets

H. Cottin

Laboratoire Interuniversitaire des Systèmes Atmosphériques (LISA), UMR CNRS 7583, Université Paris Est Créteil, Université de Paris, Institut Pierre Simon Laplace, 94000 Créteil, France

M. Drozdovskaya

Center for Space and Habitability (CSH), Universität Bern Gesellschaftsstrasse 6, 3012 Bern, Switzerland

C. Engrand

Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France

M. Galand

Department of Physics, Imperial College London, Prince Consort Road, London SW7 2AZ, UK

C. Güttler

Max Planck Institute for Solar System Research, Justus-von-Liebig-Weg 3, 37077 Göttingen, Germany

P. Henri

LPC2E, CNRS, Université d'Orléans, 3 Avenue de la Recherche Scientifique, 45071 Orléans, France

Laboratoire Lagrange, Observatoire de la Côte d'Azur, CNRS, UCA, 96 Boulevard de l'Observatoire, 06300 Nice, France

S. Ivanovski

INAF, Osservatorio Astronomico di Trieste, Via G.B. Tiepolo, 11 I-34143 Trieste, Italy

R. Kokotanekova

European Southern Observatory, Karl-Schwarzschild Str. 2, 85748 Garching bei München, Germany

A. C. Levasseur-Regourd

Sorbonne Université, LATMOS, CNRS, CNES, IPSL, BC 102, 4 place Jussieu, 75005 Paris, France

K. E. Miller

Southwest Research Institute, 6220 Culebra Rd, San Antonio, TX 78238, USA

M. Schönbachler

Department of Earth Sciences, ETH Zurich, Clausiusstrasse 25, 8092 Zurich, Switzerland

C. Snodgrass

Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK

N. Thomas

Physikalisches Institut, University of Bern, Sidlerstr. 5, 3012, Bern, Switzerland

C. Tubiana

Max Planck Institute for Solar System Research, Justus-von-Liebig-Weg 3, 37077 Göttingen, Germany

Istituto di Astrofisica e Planetologia Spaziali, Istituto Nazionale di Astrofisica, via del Fosso del Cavaliere 100, 00133 Rome, Italy

S. Ulamec

DLR, Space Operations and Astronaut Training, Linder Höhe, 51147 Cologne, Germany

J.-B. Vincent

Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institut für Planetenforschung, Rutherfordstraße 2, 12489 Berlin, Germany

Abstract We describe the *AMBITION* project, a mission to return the first-ever cryogenically-stored sample of a cometary nucleus, that has been proposed for the ESA Science Programme Voyage 2050. Comets are the leftover building blocks of giant planet cores and other planetary bodies, and fingerprints of Solar System’s formation processes. We summarise some of the most important questions still open in cometary science and Solar System formation after the successful Rosetta mission. We show that many of these scientific questions require sample analysis using techniques that are only possible in laboratories on Earth. We summarize measurements, instrumentation and mission scenarios that can address these questions. We emphasize the need for returning a sample collected at depth or, still more challenging, at cryogenic temperatures while preserving the stratigraphy of the comet nucleus surface layers. We provide requirements for the next generation of landers, for cryogenic sample acquisition and storage during the return to Earth. Rendezvous missions to the main belt comets and Centaurs, expanding our knowledge by exploring new classes of comets, are also discussed. The *AMBITION* project is discussed in the international context of comet and asteroid space exploration.

1 Executive Summary

‘Wise men know the comets come back’ (V. Nabokov 1899-1977)

The *Giotto* mission was the first interplanetary probe ever flown by the European Space Agency (ESA). Selected in 1980 and flown in 1985, *Giotto* was the most daring of an international fleet of five spaceprobes, which triumphantly visited the comet of all comets, 1P/Halley, in March 1986. Earlier on, in 1984, ESA and NASA established a Comet Nucleus Sample Return Science Definition Team; their work led in 1993 to the selection of the *Rosetta* mission as the Planetary Cornerstone of ESA’s long-term programme Horizon 2000. *Rosetta* was the first mission ever to land an automated laboratory (*Philae*) on the surface of a cometary nucleus and to accompany an errant body in its active phase through the Solar System. Launched in 2004, *Rosetta* ended its operational life on the 30th of September 2016.

Giotto and *Rosetta* have completely transformed our understanding of comets and have contributed hugely to give Europe a well-deserved leadership in the field of cometary studies. Selecting *Giotto* and *Rosetta*, Europe demonstrated the vision and the ambition to take the lead in the worldwide effort to learn about our origins. ESA has recently selected *Comet Interceptor*, a much smaller-scale mission (F-class) to encounter a yet-to-be-discovered Oort cloud comet, following a novel approach to expand our horizons on a limited budget. Europe has the opportunity to confirm and reinforce this leadership with an even more daring program: *AMBITION*, a mission to return the first-ever cryogenically-stored sample of a cometary nucleus to Earth. The international context is very favourable as, after the recent selection by NASA of the next New Frontiers class candidate mission, no international agency (NASA, CNSA, JAXA) is presently planning a sample return mission from a cometary nucleus

in the next decade. NASA's New Frontiers candidate *CAESAR*, the unique direct competitor of *AMBITION*, was not selected and, even if *CAESAR* could still be a potential candidate for the next New Frontiers call, NASA's present timeline gives Europe a competitive edge with the *AMBITION* endeavour.

The numerous discoveries of the *Rosetta* mission are of great relevance not only for their scientific significance, but also because they provided essential insight into performing critical operations and measurements. This information provides better focus for the next set of fundamental questions that can be answered by the *AMBITION* project. This White Paper presents compelling evidence that a mission capable of returning a cryogenic sample of a cometary nucleus to be analysed on Earth, and of studying the comet both remotely and in-situ, will be able to dig deeper into our past.

2 Introduction

Comets and primitive asteroids are the leftover building blocks of giant planet cores and other planetary bodies, and fingerprints of Solar System's formation processes. They are composed of materials that formed in the earliest stages of Solar System history. These objects have also preserved materials that predate the formation of the protoplanetary disc. Asteroids are mostly found in the inner Solar System, inside the orbit of Jupiter, where they are believed to have formed. Conversely, comets, which are stored in two main reservoirs in the outer Solar System, the Oort Cloud and the Kuiper Belt, are classically thought as ice-rich bodies formed beyond the snow line (Podolak and Zucker, 2004) in the outer Solar System. The distinction between comets and asteroids has recently been blurred by the discovery of both ice-rich asteroids and active asteroids in the asteroid main belt (Snodgrass et al., 2017). From the analysis of cometary dust particles using *Stardust* samples returned from comet 81P/Wild 2 or particles of 67P/Churyumov-Gerasimenko (thereafter, 67P) studied in situ by *Rosetta*, it is also now clear that comets and asteroids constitute part of a continuum in composition, pointing to an extensive transport of inner Solar System material into the outer regions in the early history of the Solar System (Brownlee et al., 2006; Fray et al., 2016). The exploration of the different classes of minor Solar System bodies, including main belt comets and transition objects such as centaurs, is of paramount interest to address Solar System formation and history, and the various processes which have altered these bodies since their formation.

Comets are of great scientific value because their ices sublime when they approach the Sun, allowing to access the primitive volatile component of the solar nebula (the Solar System protoplanetary disc). Besides water, these ices are composed of both simple and complex molecules, and appear to be common to star-forming regions, suggesting a formation in the presolar cloud (the molecular cloud precursor of the Solar System; Mumma and Charnley (2011); Altwegg et al. (2017a)). Unlike asteroids, comets do not seem to have been thermally or aqueously altered after accretion (Capaccioni et al., 2015; Quirico

et al., 2016). In addition, comets have near-solar elemental abundances and a large amount of organic matter found in the refractory material (Bardyn et al., 2017). Comets likely constitute the most primitive material still available in the Solar System.

The space exploration of comets started in the 80's with the flybys of comet 1P/Halley. With its extensive payload, the ESA *Giotto* spacecraft revealed properties of comets previously unimaginable, e.g., the incredibly dark surface of cometary nuclei and the large amount of dust particles made of C, H, O, N atoms (Keller et al., 1986; Kissel et al., 1986). NASA missions to comets 19P/Borrelly, 81P/Wild 2, 9P/Tempel 1, and 103P/Hartley 2 demonstrated an amazing diversity in the surface geology and yielded new insights into the bulk properties of cometary nuclei, the nature of cometary dust, and the processes involved in cometary activity (Soderblom et al., 2002; A'Hearn et al., 2005; Brownlee et al., 2006; A'Hearn et al., 2011). Analysis of the samples of comet 81P/Wild 2 collected and delivered to Earth by the *Stardust* spacecraft led to major findings, though the sample collection at 6.1 km/s left little chance for the survival of volatiles and organic matter (Brownlee et al., 2006).

ESA's *Rosetta* mission, the third cornerstone mission of the ESA programme Horizon 2000, is the most ambitious and sophisticated cometary space project so far (Taylor et al., 2017; Boehnhardt et al., 2017). During its 2 years (August 2014–September 2016) escorting comet 67P, the 10 instruments aboard the spacecraft monitored and performed both in situ and remote analyses of the nucleus, atmosphere, and solar-wind interaction continuously. *Rosetta* enhanced its science return by releasing *Philae* which landed on the nucleus surface: despite the non-nominal landing, the instruments aboard the lander were able to complete a subset of the planned measurements aimed at characterizing the comet nucleus's surface, subsurface, and local environment (Boehnhardt et al., 2017). *Rosetta* has provided many important and often unexpected results, as demonstrated by the number of publications (~ 1200 in early 2021), with many in *Science* and *Nature* journals. A wealth of new knowledge was obtained on:

- the bulk and internal properties of the nucleus, with the demonstration that comets have very low density ($\sim 500 \text{ kg/m}^3$) and high porosity ($> 70\%$) (Kofman et al., 2015; Pätzold et al., 2019), and the evidence for layering in the uppermost parts of the nucleus which may be related to primordial or post-collisional accretion (Massironi et al., 2015; Jutzi et al., 2017); the nucleus interior is very homogeneous down to few-meter scales (Herique et al., 2019);
- the surface morphology, showing an unexpected diversity with large areas of the surface covered by airfall particles originating from the southern hemisphere which receives maximum solar flux during perihelion passage (Thomas et al., 2015; Keller et al., 2017). There are clear indications that no single process dominates the evolution of the surface morphology;
- the distribution of water ice on the surface, and its diurnal and seasonal evolution (e.g., De Sanctis et al., 2015; Fornasier et al., 2016; El-Maarry

- et al., 2019). The refractory-to-ice ratio in the nucleus may exceed 3, inconsistent with the paradigm that comets are very ice-rich (e.g., Pätzold et al., 2019; Fulle et al., 2019; Choukroun et al., 2020);
- the thermal and mechanical properties of the surface and subsurface layers; the very low bulk tensile strength is consistent with a primordial rubble-pile formed in the solar nebula (Attree et al., 2018);
 - the physical properties of dust particles (Levasseur-Regourd et al., 2018), with the discovery of both highly porous or even fractal-like and more compact aggregates made of ~ 100 nm subunits consistent with dust growth starting by low-velocity hierarchical accretion (Fulle et al., 2015; Mannel et al., 2016; Fulle et al., 2016b);
 - the solid organic matter ($\sim 50\%$ in mass) in dust particles which is bound in very large macromolecular compounds and is analogous to the insoluble organic matter in carbonaceous chondrites (Fray et al., 2016; Herique et al., 2016);
 - the volatile composition, with the detection of a wealth of simple and complex species, e.g., O_2 , noble gases, heavy hydrocarbons, alcohols, and glycine (Altwegg et al., 2016; Bieler et al., 2015). Isotopic measurements for H, C, O, Si, S, Ar, and Xe, and similarities and differences with primitive carbonaceous meteorites shed light on the presolar versus solar-nebula origin of cometary ices (Hoppe et al., 2018). Xenon isotopic abundances provide evidence that comets contributed to atmospheric Xe on Earth, or more generally to atmospheric noble gases (Marty et al., 2017);
 - the importance of diurnal and seasonal illumination conditions on the different regions of the nucleus in driving the actual activity (Fulle et al., 2016a);
 - the evolution of the interaction of a comet with its surrounding space environment, from the birth of an induced magnetosphere and a nearly collisionless cometary plasma, to a chemically-active, colder, unmagnetized cometary plasma near perihelion (Engelhardt et al., 2018; Goetz et al., 2017; Heritier et al., 2018; Nilsson et al., 2017).

Although the *Rosetta* mission brought cometary science to a new level of maturity and improved our understanding of the early Solar System, a number of fundamental questions that can only be achieved by space missions remain unresolved.

- How did cometary materials assemble? What are the building blocks of cometary nuclei?
- Which post-planetesimal evolutionary paths need to be considered? Are comets collisional fragments or primordial planetesimals?
- Which cometary components predate the formation of the Solar System? What is the nature of the refractory organic materials? How and where did these components form?
- Are there differences in physical/chemical properties in the comet populations? What are the commonalities and differences between comets and

primitive asteroids? Are these distinct populations? Do comets, main belt comets and active asteroids have different primordial reservoirs?

- How does comet activity work? How do surface and coma observations reconnect with the pristine, deep interior?
- How are the dusty coma, the surrounding plasma, and the nucleus interacting together? Do interactions with the solar wind influence the activity and evolution of comets?
- What contributions did comets make to the reservoirs of volatiles and prebiotic compounds on early Earth?

The holy grail of cometary spacecraft missions is the return to Earth of a cryogenic sample extracted from deep within the nucleus. It is worth mentioning that the original *Rosetta* mission was a comet-nucleus sample return (CNSR) and that significant important studies were carried out towards these goals. CNSR missions allow the ultimate level of detailed study of extraterrestrial samples, although they are technically challenging and expensive. In situ studies with static or mobile landers, equipped with a sophisticated payload and drilling systems, remain a necessary complement to sample return because of inherent problems with maintaining sample integrity, in addition to enhance the science return from the sample by providing the full context. We present the major scientific questions resolvable by future comet missions in Section 3, and propose a few missions for the ESA Voyage 2050 long-term plan in Section 4. The international context of comet space and Earth-based exploration is presented in Section 5.

3 Top Level scientific questions

3.1 How and where did cometary materials get assembled? Which post-planetesimal evolution paths need to be considered?

It is undisputed that the formation of planetary bodies in the solar nebula started with the coagulation of fine dust particles induced by low-velocity sticking collisions (Weidenschilling, 1977). Recent numerical models of the physical agglomeration process, taking into account the latest findings from laboratory experiments (e.g., Güttler et al., 2010), show that mm- to cm-sized agglomerates (hereafter, termed pebbles in accordance with the literature) form within a few hundred to a few thousand orbital timescales (Lorek et al., 2018; Zsom et al., 2010). Due to the systematic increase in collision velocity with increasing agglomerate mass, larger agglomerates no longer simply stick after a collision. However, some of the collisions may still lead to a further gain in mass, even though the sticking threshold has been overcome, e.g., by mass transfer in high-velocity collisions (see Güttler et al., 2010, and references therein). Davidsson et al. (2016) described how cometary nuclei might have formed in such a scenario and compared it to the findings of *Rosetta*. An alternative scenario was presented by Blum et al. (2017) who collected evidence from *Rosetta* data that the comet nucleus might still consist of the primordial pebbles, with fractal




Fig. 1: *Left*: Processes in protoplanetary discs giving rise to gas streaming instabilities and the formation of pebbles and planetesimals (Armitage, 2019); *Right* (67P building blocks): pebbles on the surface imaged by the Comet Infrared and Visible Analyser (CIVA) onboard *Philae* (top) (Poulet et al., 2016) and 3D rendered images obtained with the Micro-Imaging Dust Analysis System (MIDAS) (bottom) (Mannel et al., 2019).

dust aggregates captured between the pebbles (Fulle and Blum, 2017), which indicates that it formed by the gentle gravitational collapse of a concentrated cloud of pebbles (Johansen et al., 2007) (Fig. 1).

Without further collisional evolution, these two formation scenarios predict very different physical properties of the resulting planetesimals, e.g., porosity and tensile strength (Blum, 2018). Morbidelli and Rickman (2015) argued that the planetesimals in the outer Solar System underwent one or more (sub-)catastrophic collisions during the past 4.5 billion years. In this case, the primordial planetesimal properties might have been overwritten by the fragmentation and re-accretion events. However, major parts of the reassembled cometary nucleus might still bear the initial morphology so that not necessarily all information about the underlying formation mechanism might have been lost (Schwartz et al., 2018). Additional difficulties arise when trying to identify the properties and structures that may have evolved via other forms of secondary processing (including, but not limited to, distortion or fragmentation caused by spin-up and/or planetary tides; solar and/or radiogenic heating and associated vapor diffusion, re-condensation, and irreversible phase transitions; gravitational compaction). Indeed, *Rosetta* showed a wide variety of different structures and textures on 67P's nucleus surface (Thomas et al., 2015) and complex surface changes due to activity (El-Maarry et al., 2017; Fornasier et al., 2017). It is thus mandatory to isolate features indicative of the initial

growth process of the planetesimal precursors of cometary nuclei. For example, the *Deep Impact* and *Rosetta* missions demonstrated that stratigraphic layering is an important structural component in some comet nuclei (Belton et al., 2007; Massironi et al., 2015), but it is unknown whether layering forms during the growth of the body or at later stages.

The complex interplay between formation and evolutionary processes requires both advanced modeling and dedicated laboratory work to arrive at realistic predictions on the physical and structural properties of bodies formed in different ways. Processes such as the transition from fractal to non-fractal (but porous) aggregate growth domains, the dependence of growth barriers on physical and structural dust particle properties, the interplay between erosional and growth processes in moderate-velocity impacts, or advanced modeling of collisions among porous and hierarchically structured bodies must be the focus of future studies. An anticipated comet mission with the aim of revealing the formation and evolution processes of planetesimals can only be successful if all the above-mentioned aspects are properly taken into account. Only a holistic approach to revealing the structural properties of the entire comet nucleus (or at least considerable parts of it) can lead to the desired results. The ability to investigate the interior of the nucleus down to a depth of several tens to hundreds of meters with a resolution in the mm-cm range may be the essence of this investigation. It is also important to understand whether structures seen on such size scales are representative for the body as a whole, or if regional variability is substantial. In this respect, a successful mission will need to measure the morphological and compositional properties (particle sizes, dust/ice/organic mixing, arrangement on different scales) of a sample that never experienced modification by cometary activity. This implies that material from the deep interior needs to be studied on a wider range of size scales and with a composition-probing capability. A careful assessment of the relative merit of remote-sensing instruments (e.g., ground-penetrating radar), as well as in situ investigations (drill cores, mole probes, penetrators, impact excavation) is required. The possibility of exploiting the exposure of nucleus interiors following comet fragmentation or splitting events, as well as large-scale outbursts, should be considered.

3.2 What is the presolar heritage of cometary materials, versus a solar nebula origin?

Through their mineralogical, chemical, and isotopic composition, comet nuclei document environmental conditions and processes occurring from the protostellar collapse phase to the protoplanetary disc phase. The protostellar collapse phase includes the prestellar core formation from the parent molecular cloud, its collapse, and the bulk solar accretion phase as the protosun evolved through the protostellar Class 0 and Class I stages (Shu, 1977; André and Montmerle, 1994). Physical processes such as gas compression and associated heating (e.g., in shocks), inward accretion flows, disc winds, and bipolar flows,

take place (Bachiller, 1996; Banerjee and Pudritz, 2006; Machida and Matsumoto, 2011). Icy dust particles feeding the disc are affected by thermal evaporation, photodesorption, irradiation, and sputtering in strong shocks. Complex organic molecules have been observed as early on as in the prestellar cores with their formation stemming from grain-surface chemistry (e.g., Bacmann et al., 2012). Physico-chemical models suggest that their formation is further enhanced during protostellar collapse and continues within the protoplanetary disc midplane (Walsh et al., 2014; Drozdovskaya et al., 2016). To what extent does molecular complexity develop in the protoplanetary disc phase versus being inherited from prestellar ices? How do nebular processes, such as stellar UV, X-ray, and cosmic rays, chemically alter volatiles and dust particles? The chemical inventory of the dust particles entering the protoplanetary disc depends on several parameters: how long did the prestellar core spend in the cold-dense phase, what was the temperature in the prestellar core, and what are the characteristic fluences of FUV and cosmic rays? What was the chemical composition of the prestellar core of the Solar System?

3.2.1 Cometary ices

Over the last few years, thanks to large programs on astrochemistry (using, e.g., the Atacama Large Millimeter Array, ALMA), a chemical census of the gas-phase has been obtained (and modelled) along the disc evolution from the protostellar phase to planet-forming disc. At the same time, the Rosetta Spectrometer for Ion and Neutral Analysis (ROSINA) onboard *Rosetta* identified many new cometary volatiles in comet 67P, including both complex organics (e.g., alcohols, long-chain hydrocarbons, glycine (Altwegg et al., 2016)) and unexpected simple species such as O₂ (Taquet et al., 2018). A comparative study of 67P with the closest Solar-like system that is still in its infant embedded phase of formation, namely the Class 0 low-mass protostar IRAS 16293-2422 B, shows striking correlations for CHO-, N- and S-bearing species suggesting inheritance from the presolar phase (Bockelée-Morvan and Biver, 2017; Drozdovskaya et al., 2019) (Fig. 2). A similar correlation is observed with the Class I hot-corino SVS13-A ($\sim 10^5$ yrs old) for complex species such as NH₂CHO and HCOOCH₃ (Bianchi et al., 2019). The higher O₂ content in 67P with respect to methanol in comparison to IRAS 16293-2422 B suggests a higher temperature in the prestellar core of the Solar System, with respect to the core from which IRAS 16293-2422 B formed (Taquet et al., 2018). These findings call for further investigations. Especially important for a future mission will be the measurement of isotopic ratios, as key diagnostics of formation conditions. These have been obtained in 67P and other comets with significant accuracy only for a small number of species, e.g., D/H in H₂O, HCN and H₂S, ¹⁸O/¹⁶O in H₂O and CO₂ (Bockelée-Morvan et al., 2015; Hässig et al., 2017; Schroeder et al., 2019), ³³S/³²S and ³⁴S/³²S in H₂S, CS₂ and OCS (Altwegg et al., 2015; Calmonte et al., 2017; Altwegg et al., 2017b; Hoppe et al., 2018). Whereas the high HDO/H₂O and D₂O/HDO ratios in 67P suggest a presolar




Fig. 2: *Left*: Abundance of CHO bearing molecules relative to methanol in IRAS 16293-2422 B versus that measured in comet 67P (arrows indicate upper limits) (Drozdovskaya et al., 2019). *Right*: the Rho-Ophiuchi star-forming region harbouring IRAS 16293-2422 B (ESO/Digitized Sky Survey 2/L. Calçada).

heritage, this has to be confirmed in the light of recent findings on the D/H diversity in comets (Lis et al., 2019).

3.2.2 Refractory organics

Cometary dust is rich in refractory organic matter (50% in mass) (Bardyn et al., 2017). The identification and elemental characterization of CHON-rich dust particles were first performed in the coma of 1P/Halley (Jessberger et al., 1988; Lawler and Brownlee, 1992). The *Rosetta* mission, and in particular instruments such as mass and IR spectrometers (ROSINA, Cometary Secondary Ion Mass Analyser (COSIMA), Visible and Infrared Thermal Imaging Spectrometer (VIRTIS)), allowed further characterization of this organic matter made of aliphatic and aromatic moieties (Raponi et al., 2020). Cometary organics share some similarities with chondritic insoluble organic matter, although they are not identical (e.g., Fray et al., 2016; Quirico et al., 2016). Cosmic dust particles collected on Earth (mainly in the stratosphere and in the polar regions) contain material of probable cometary origin in the form of ‘chondritic porous interplanetary dust particles’ (CP-IDPs, e.g., Ishii et al. (2008)), and ‘ultracarbonaceous Antarctic micrometeorites’ (UCAMMs, e.g., Duprat et al. (2010)) (Fig. 3). CP-IDPs and UCAMMs also contain > 50% in mass of organics.

Further chemical and isotopic characterization of cometary organics in the laboratory is necessary to establish a deeper comparison, and to decipher the origin(s) of cometary and chondritic organics, which are highly debated in

the community. Different scenarios are advocated for chondritic organics: interstellar heritage, formation in the solar nebula, and parent body formation (e.g., Alexander et al., 2017). Isotopic fractionation measured in comets and chondrites are lower than those found in some species in interstellar clouds. This shows that primitive organics in the Solar System are either a mixture of interstellar matter and materials processed in situ, or produced in the outer regions of the solar nebula (e.g., Aikawa and Herbst, 1999). It has been suggested that the large organic molecules responsible for the Diffuse Interstellar Bands (DIBs) could be preserved in comets (Bertaux and Lallement, 2017). Determining the interstellar vs. solar nebula origin of cometary organic matter requires knowledge of the exact H, C, and N isotopic compositions of cometary organics, as well as the understanding of the evolution of organics from dense molecular clouds to protoplanetary discs. The direct comparison between cometary and chondritic organics (e.g., elemental and isotopic compositions, functional chemistry, structure, and texture) would also allow to address the question of either a unique or multiple reservoir(s) of organics in the Solar System (at least 2 types of organic matter have been identified in UCAMMs, Engrand et al. (2018)). This comparison would, in turn, enable better understanding of the asteroid-comet continuum and potentially of the dynamics of a protoplanetary disc by constraining exchanges between the inner and outer Solar System. This latter point could also be addressed through the search for and characterization of high-temperature components, such as chondrules and CAIs, in cometary samples (Sect. 3.3).

For comet 67P analyzed by *Rosetta*, glycine was identified in the coma by ROSINA (Altwegg et al., 2016), whereas low-molecular weight organic compounds were not found in the COSIMA spectra (Fray et al., 2016). The absence of identification by COSIMA of organics comparable to the so-called soluble organic matter (SOM) in chondrites might indicate the presence of such soluble organics at a lower abundance in comparison to that of chondrites. Although its origin is debated (interstellar, solar, parent body origin), the SOM is more abundant in the chondrites having experienced more intense aqueous alteration (e.g., Pizzarello et al., 2006). Comets have escaped such secondary processes, as suggested by the lack of evidence for hydrated minerals in cometary dust particles (Quirico et al., 2016), although they were experienced by asteroids. The characterization of SOM in comets will give new constraints on the origin of chondritic soluble organics.

Although comets are rich in volatiles and solid organics, they are depleted in nitrogen in comparison to the solar nebula (e.g., Cochran et al., 2000). However, the recent identification of ammonium salts at the surface and in the dust of comet 67P (Poch et al., 2020; ?) sheds new light on this issue: these salts could provide the heretofore-missing link between comets and the parent interstellar cloud. While this potential new nitrogen reservoir was proposed from *Rosetta* data, no precise characterization of this reservoir is currently possible (such as the relative abundance of the different salts, and their isotopic composition). This would necessarily require the sampling of the cometary surface and its study in laboratories.

3.2.3 Presolar grains

Presolar grains in primitive extraterrestrial materials are identified by large isotopic anomalies compared to the solar composition (e.g., Clayton and Nittler, 2004). The nature of presolar grains gives access to the specific types of stellar sources (e.g., supernovae, red giant stars) in whose mass outflows the grains condensed, and their residence time in the interstellar medium. They also yield insights into stellar nucleosynthesis at an unprecedented level of detail (Clayton and Nittler, 2004). Their relative abundance in meteorites and comets gives insights into processes potentially acting both in the inner and outer regions of the protoplanetary disc. Presolar grains are found at a level of 500 ppm in meteorites (Zinner et al., 2005), and up to 1.5% in IDPs collected during the Grigg-Skjellerup dust stream encounter with Earth in 2003 (Busemann et al., 2009). In comet 81P/Wild 2 samples returned by the *Stardust* mission, the abundance of presolar grains, after correction for destruction upon impact in the aerogel at 6.1 km/s, is 650 to 900 ppm, being largely dominated (>90%) by O-rich grains (Floss et al., 2013). This low level (rather comparable to that of meteorites) was found in contrast with measurements of silicon isotopes from comet 67P dust particles by ROSINA, which show an overall large depletion of heavy Si isotopes in the dust, suggesting a large fraction of presolar Si-bearing minerals (Rubin et al., 2017).

Further characterisation of the composition of cometary dust, both for the organic and mineral part, is clearly needed. Remote sensing and in situ analyses suffer from a lack of measurement context, and are not as precise and accurate as laboratory analyses. *Stardust* samples returned from comet 81P/Wild 2 suffered severely from their hypervelocity collection, hampering meaningful understanding of the composition and structure of the matrix and organic matter, because only refractory components were retrieved. CP-IDPs and UCAMMs may have suffered from heating during atmospheric entry, although their porous structure could have limited their alteration. A cometary sample return would give the only opportunity to retrieve the fragile and low temperature components of cometary matter that are so far unknown in collections of primitive extraterrestrial matter, and to investigate a representative amount of material, giving access to the least abundant species (soluble organics, noble gases). A cryogenic sample return would also allow preservation of semi-volatiles, e.g., salts.

3.3 What do comets tell us about large-scale mixing and dynamical processes in the early Solar System?

During the protostellar collapse phase, dust particles located near the protostar experience harsh conditions. Amorphous silicates from the pre-stellar core anneal and crystallize, some solids evaporate and recondense entirely, and evolve into the granular mixtures of Calcium-Aluminum-Rich Inclusions (CAIs), agglomeratic olivines, amoeboid olivine aggregates, chondrules, and matrix ma-




Fig. 3: From left to right: 1) dust particle of 67P collected by COSIMA/Rosetta (credit: ESA/Rosetta/MPS); 2) 81P 2- μm particle collected by *Stardust* made of Mg-rich crystalline olivine (credit JPL/NASA); 3) transmission electron micrograph of GEMS (Glass with Embedded Metals and Sulfides) embedded in their carbonaceous matrix from a potential cometary particle collected on Earth (CP-IDP) (Bradley, 1994); these glassy inclusions could have a presolar origin; 4) backscattered electron micrograph of an Ultracarbonaceous Antarctic MicroMeteorite (UCAMM) from the Concordia Collection (CSNSM, France); these particles are dominated by organic matter (dark phase) with minor mineral components (small and light phases). UCAMMs most probably have a cometary origin.

terial that are well-known from chondritic meteorites (Krot et al., 2004; Jones, 2005; MacPherson et al., 2005; Ruzicka et al., 2012). Infrared cometary spectra and analyses of *Stardust* samples of comet 81P/Wild 2 showed that such particles together with metals and sulfides were transported to large heliocentric distances (Wooden et al., 2000; Brownlee et al., 2006), potentially accreting processed organics and ices on their way. Once there, they mixed with relatively unprocessed amorphous presolar minerals, organics, and ices. Indeed, the mineral components of comet 81P/Wild 2 samples share many similarities with that of primitive carbonaceous chondrites that are thought to originate from dark asteroids, including CAIs and chondrules (Brownlee, 2014). Cometary dust particles and cosmic dust particles of probable cometary origin that are collected on Earth (CP-IDPs and UCAMMs) contain both crystalline silicates and amorphous silicates of possible presolar origin (Wooden et al., 2000) (Fig. 3). The presence of crystalline and high-temperature minerals in cometary matter, sharing similarities with their counterparts in meteorites, has led to the notion of a comet–asteroid continuum (Gounelle, 2011). Cometary dust, on the other hand, contains much more carbon than carbonaceous meteorites (50% carbon in mass, and even more in CP-IDPs and UCAMMs, versus 5% at most in meteorites) (Bardyn et al., 2017), in a form that is close to (but not identical to) insoluble organic matter in meteorites (Fray et al., 2017a; ?,b; Dartois et al., 2018).

Multiple mechanisms have been proposed to explain the outward transport of solids in the solar nebula prior to being accreted into planetesimals. Two leading theories are: i) solids were launched above the disc by winds that are able to decouple them from the gas flow, and rained back down onto the nebula at different radial locations (Shu et al., 1996; Ciesla, 2009); ii)

processes operating within the nebular gas (turbulence, photophoresis, gravitational torques) allowed solids to migrate outwards through the solar nebula (Bockelée-Morvan et al., 2002; Boss, 2008; Mousis et al., 2007) (Fig. 1). Alternatively, high-temperature materials formed on the disc surface far from the star during episodic accretion bursts (Ábrahám et al., 2009). Depending on the transport mechanism, dust particles were exposed to very different environments during their transfer to the outer disc. Timescales for incorporation into planetesimals also differ. Determining which of the proposed mechanisms was dominant requires information on the degree of silicate crystallinity, the relative content of high-temperature material, their variation among different comet populations, and among primitive Solar System bodies formed at different places in the solar nebula. Also important for constraining transport mechanisms is determining how and at what scales low- and high-temperature materials are heterogeneously or homogeneously mixed together.

The search for extinct ^{26}Al in cometary samples could allow (in the chronological interpretation) a precise dating of the crystallization time of cometary minerals. The absence of ^{26}Al excess in *Stardust* samples measured so far could be interpreted as a late formation of minerals in comet 81P/Wild 2 dust (Matzel et al., 2010; Ogliore et al., 2012; Nakashima et al., 2015). Other dating systems (U-Pb, Rb-Sr, Nd-Sm,...) could provide information about the formation ages and timing of possible later processes occurring on the comet. Establishing a precise chronology in the early Solar System is only possible through analysis of pristine samples in the laboratory.

Comets have historically been distinguished from asteroids by the extended coma and tail that they develop when approaching the Sun. This distinction was blurred in 2006 by the discovery of a significant population of asteroids with comet-like activity in the outer main belt (the so-called *Main Belt Comets* (MBCs) (Hsieh and Jewitt, 2006)). The most recent estimate gives a population of 140–150 active bodies, more than could have found their way from known comet reservoirs by complex gravitational interactions (Hsieh et al., 2015). In addition, ice signatures have been identified on the surface of several asteroids (e.g., Campins et al., 2010; Combe et al., 2019), and water vapor was detected around the ice-rich asteroid Ceres (Küppers et al., 2014). The implied presence of water in the main belt within the snow line in the protoplanetary disc indicates that significant migration of the planetesimals has taken place early in the Solar System. This is in line with dynamical models aimed at understanding the formation of the terrestrial planets and the present architecture of the Solar System. Because of interactions with gas and/or planetesimals, giant planets underwent episodes of migration, scattering bodies by many astronomical units from the location of their formation. The Grand Tack model, describing the inward then backward migration of Jupiter and Saturn in the gaseous nebula, not only helps in explaining the properties of the terrestrial planets, but also the structure of the asteroid belt and the presence of S- and C-type asteroids associated with ordinary and primitive carbonaceous chondrites, respectively (Walsh et al., 2011; DeMeo and Carry, 2014). The formation of the two main comet reservoirs, the Oort Cloud and the Scat-

tered Disc, sources of the long-period and short-period dynamical families, respectively, is believed to have resulted from the gravitational interaction of the three outermost external planets with a massive disc of planetesimals extending out to 30 AU, which made them move outwards, dispersing the disc (so-called Nice model, Tsiganis et al., 2005). This means that the diverse comet and asteroid reservoirs probably include bodies formed at different locations in the Solar System. The suggestion that volatile-rich carbonaceous chondrites originate from the outer Solar System or even from comets is an interesting hypothesis that can be tested by a comet nucleus sample return (Gounelle et al., 2008).

The characterization of asteroid-comet continuum objects, like MBCs, is of prime importance to test Solar System evolution scenarios and for constraining the distribution of water in the early Solar System. Because these objects are weakly active and small, their investigation requires space exploration. The detection of water vapor and its sources would attest to the presence of buried ice, and constrain the activity-triggering mechanism. The measurement of the D/H ratio in water is a prime objective for constraining their formation region, together with physical and chemical properties that can be compared to asteroids and comets properties.

3.4 How does comet activity work? How do surface and coma observations reconnect with the pristine, deep interior?

Comets show several types of activity depending on the nature of the gas and dust involved, flux and distribution of the sources (localized or diffuse) that are related to the thermophysical and chemical properties of the comet nucleus. *Rosetta* and previous missions greatly improved our understanding of how comets work, but there are still open questions. Cometary activity can originate from the immediate surface and from the subsurface, and the associated phenomena are different. Also, the mechanisms driving the gas and dust activity can be different and can operate at different times and heliocentric distances.

Different types of local activity have been observed, including i) diffuse activity from the nucleus and ii) localized activity (jets, outbursts; Fig. 4). Jets seem to be related to surface topography (e.g., sinkholes or fractures), but also to sudden changes in the thermophysical condition of the surface, such as shadow boundaries (e.g., the terminator line, shadows from topographic reliefs) (De Sanctis et al., 2015; Vincent et al., 2016a,b; Kramer and Noack, 2016; Shi et al., 2016)). Diffuse activity is more difficult to trace, because it is not clearly linked to surface features, but is more related to the overall nucleus thermochemical state.

Several mechanisms have been recognized to play a major role in the activity of comets. On a global scale, solar input and seasonal and diurnal variations act on the nucleus, driving the gas activity, which starts with the sublimation of the most volatile ices at large heliocentric distance, increasing close to peri-




Fig. 4: *Left and center*: RGB maps of 67P’s Anhur region obtained with the Optical, Spectroscopic and Infrared Remote Imaging System (OSIRIS), showing exposed water ice (blue patches) on the dark surface, and jet activity (Fornasier et al., 2017). *Right*: VIRTIS spectrum showing the transient signature of surface CO₂ ice on Anhur (dashed vertical lines) (Filacchione et al., 2016b). A combination of water ice, organics, and salts is responsible for the broad 3.2 μm band (Capaccioni et al., 2015; Quirico et al., 2016; Poch et al., 2020; Raponi et al., 2020).

helion with the release of the less volatile species (Hässig et al., 2015; Migliorini et al., 2016; Hansen et al., 2016; Biver et al., 2019). On a local scale, activity is driven by the surface and subsurface conditions, such as the accumulation of gas pockets and subsurface/surface stresses, the volatile ice content, morphological and topographic features that change the thermophysical condition locally (i.e., sinkholes, shadows).

The activity is able to alter the nucleus shape and to modify the internal structure. Clear morphological changes have been observed on the nucleus, related to global diffuse and local activity (El-Maarry et al., 2019). Diffuse, solar-driven activity on a global scale can explain non-gravitational forces in torque and orbit (Attree et al., 2019; Kramer and Lauter, 2019). Specific morphology changes seem to follow orbital cycles triggering the displacement of surface regolith (Keller et al., 2017). Other changes (i.e., landslides, fissures, sinkhole collapses, honeycombs, dunes (El-Maarry et al., 2017; Hu et al., 2017)) are more related to the local activity and subject to the diurnal variations. Some morphologic regions are deemed to be more active than others, e.g., the Anhur and Bes southern regions of 67P (Fornasier et al., 2017), due to higher erosion rates during the perihelion passage when the surface is exposed to a higher solar flux. As a result of surface dust removal, more pristine ice-rich layers are exposed (Filacchione et al., 2016c; Fornasier et al., 2017).

A major deficiency in *Rosetta* data is the lack of constraints on the physical properties of the surface and subsurface layers, and the relationship of volatiles and refractories at depth. This information forms an important input to our understanding of cometary activity, and is necessary to relate coma properties, such as chemical abundances and the dust-to-gas ratio, to intrinsic

nucleus properties. Thermophysical parameters strongly depend on microphysical details such as, the material composition, the morphology and structure of the material (arrangement of the material, coordination number, void spaces) and the mixing of the different components (mineral phases, organics and ices) (Filacchione et al., 2019). One can envisage ices surrounding the dust component, ice matrices within dust matrices, and ices as a component isolated from the refractories. How minor species are mixed with water ice is unknown (e.g., trapped in amorphous ice, clathrate hydrates, pure ices), whereas this affects the conditions under which they are released. Microphysical properties affect the macroscopic behaviour of the cometary surface, such as tensile strength (Gundlach et al., 2018) and compressive strength (Lorek et al., 2016). The (somewhat disputed) high compressive strength measured from *Philae* just below the surface of 67P suggests the presence of sintering processes, driven by sublimation and recondensation of water (Spohn et al., 2015).

Values for the thermal inertia (very low, 10–50 SI; Gulkis et al., 2015; Marshall et al., 2018) and porosity ($> 70\%$; Herique et al., 2016; Pätzold et al., 2019) have been obtained, but the thermal conductivity and heat capacity are not fully constrained, and neither are the gas regime transport and recondensation. The refractory-to-ice ratio in the nucleus remains a subject of lively debate (Choukroun et al., 2020). The relative amounts of volatile species (e.g., CO, noble gases) with respect to water measured in the coma certainly do not reflect their abundance in pristine ices, since the depth of the sublimation fronts are different. The seasonal production patterns of minor species fall in two groups, following either H_2O or CO_2 , and are not correlated with sublimation temperatures (Luspay-Kuti et al., 2015; Gasc et al., 2017). This can be interpreted as a consequence of two different ice phases, H_2O or CO_2 ice-dominated, in which the minor species are embedded in different relative abundances. Whether this is a primordial heterogeneity or due to evolution and differentiation is unclear.

A primary objective for a future comet mission should be improved knowledge of the thermophysical parameters and microphysical properties of the comet surface and subsurface. Properties to be studied as a function of depth include composition, particle size distribution and shapes, morphology and bulk properties of the material, mixing of the different components, location of the sublimation fronts, ice-phase (e.g., whether amorphous or crystalline in the deep interior), and relationships between near-nucleus coma and sub-surface properties. Different mission scenarios with varying levels of complexity can be envisaged, from in situ (lander-based) measurements to a cryogenic sample return. Drilling at depths of > 1 m in regions deprived of dust airfall (which is technically challenging) may permit access to volatile-rich layers and amorphous water ice. The detection of CO_2 ice at the surface of 67P shows that volatiles can condense on the surface during the winter season when they are experiencing a years-long night (Filacchione et al., 2016b) (Fig. 4).

3.5 What was the role of comets in the delivery of volatiles and prebiotic compounds to early Earth?

Since comets are reservoirs of a large number of ingredients necessary for the origin of life, like water and organic molecules, they are targets of prime interest for any astrobiology investigation (Cottin et al., 2017; Oro and Cosmovici, 1997).

The link between comets and the origin of water on Earth has always been puzzling and debated. The water D/H ratio measured in comets is quite diverse: ranging from the terrestrial value in comets 46P and 103P ($\sim 1.5 \times 10^{-4}$) (Hartogh et al., 2011; Lis et al., 2019), to more than 3 times this number in comet 67P (Altwegg et al., 2015). No relation between the D/H of water and the dynamical family of the comet has been established so far (Altwegg et al., 2015). However, it has been recently proposed that comets may all have the same D/H in their water ice and that the variation in the observed ratio could be due to fractionation effects during sublimation (Lis et al., 2019). Deciphering the origin of water on Earth requires not only measuring the D/H in water in more comets to further assess D/H variability among comets (which can be done, albeit with difficulties, from Earth-based observatories or space telescopes), but also to measure it in different phases (gas and icy particles) within a single comet. The last point requires in-situ measurements and/or cryogenic sample return. Beyond water, the contribution of comets to the Earth budget of other volatile compounds (i.e., to the formation of the atmosphere) can also be further constrained by measuring isotopes of noble gases, such as Xe and Ar (Marty et al., 2016, 2017). Xe and Ar isotopes have only been measured in comet 67P and they bring strong constraints on the budget of volatiles brought to Earth (20% of atmospheric Xe is of cometary origin). Their relative abundances have to be confirmed beyond the case of 67P (to test their consistency in comets in general).

For a long time, comets have been considered as a source of organic compounds which could have played a key role in the chemistry leading to the origin of life (Oró, 1961). Key constituents for prebiotic chemistry have been identified (Fig. 5), e.g., HCN, formamide (NH_2CHO), glycolaldehyde (HCOCH_2OH), the simplest sugar and an important intermediate on the path towards forming more complex biologically relevant molecules (Biver et al., 2015), and glycine, the simplest of the amino acids, which are the building blocks of proteins (Altwegg et al., 2016). Besides organic compounds, phosphorus, a key element in the structure of nucleotides, the building blocks of DNA and RNA, has also been detected in the coma of 67P (Altwegg et al., 2016). Furthermore, ammonium salts, also detected in 67P (Poch et al., 2020; ?), are known to be key precursors in the syntheses of amino acids and nucleobases. Further characterization of the chemical nature of the cometary organics will bring strong constraints to the list of chemical ingredients delivered to the early Earth, potentially enabling the chemical evolution that led to the origin of life on our planet, and maybe elsewhere.




Fig. 5: The formation of a protocell from precursors detected in comets and/or carbonaceous meteorites. Amino acids are the building block of proteins, nucleobases; ribose and phosphate are the building blocks of nucleotides (which are the building blocks of RNA and DNA), and amphiphilic molecules are known to spontaneously self-assemble into vesicles in water (i.e., into primitive cell membranes). From Cottin et al. (2017).

The origin of chirality is also a very important question in prebiotic chemistry. Is the asymmetry of life already ‘written’ in the organic material brought from space, or is it the result of a stereospecific mechanism on the early Earth, or a pure random selection followed by amplification while life developed? Measurements in carbonaceous chondrites show a few percent excess of the L configuration for amino acids (Cronin and Pizzarello, 1997). Chiral measurements in comets are necessary to assess the extent to which enantiomeric excesses are widely distributed among primitive small bodies of the Solar System and among families of chemical compounds within a given comet. Such observations would support a scenario in which the Solar System was formed in a region where ice-rich particles were irradiated during the protoplanetary phase by an external source of circularly polarized light, inducing stereospecific photochemistry (Modica et al., 2014). The determination of chirality will be possible by in situ or by sample return measurements (see planned measurements from *Philae*, Goesmann et al., 2014).

3.6 How do the dust coma, the surrounding plasma, and the nucleus interact?

The expanding coma around a cometary nucleus is ionised by solar Extreme UltraViolet (EUV) radiation and energetic electrons (Cravens, 1987; Heritier et al., 2018). This forms the cometary ionosphere, composed of free electrons and cations (e.g., H_2O^+ and H_3O^+). The cometary ionosphere is surrounded by the solar wind plasma, made of free electrons, protons and alpha particles,

which interacts with the cometary coma, e.g., through charge exchange (Simon Wedlund et al., 2019).

Dust–plasma. How are cometary dust dynamics and properties affected by charging processes? What is the feedback on a cometary plasma? Comets are natural dusty plasma laboratories which evolve along their elliptical orbits due to changes in dust properties (density, size, porosity) and outgassing. In the presence of a gas-phase plasma (here, the cometary ionosphere) and close enough to the nucleus, the dust particles become negatively charged (as free electrons stick to the surface of the particles), attract gas-phase low-energy cations, and grow in size. The dust charging is expected to influence the dust dynamics and spatial distribution (through electromagnetic forces), the dust physical characteristics (e.g., erosion) and the gas-phase plasma properties (e.g., charge imbalance, electron cooling) (e.g., Mendis et al., 2013). Charged nanograins were first detected at a comet during the *Rosetta* mission (Burch et al., 2015). They are accelerated outward by gas drag and, away from the nucleus, by the solar-wind induced electric field (Gombosi et al., 2015). Nanograins produced by dust disintegration in the coma may be responsible for the attenuation, near perihelion, of the EUV solar flux measured at *Rosetta*'s location (Johansson et al., 2017). Modelling predicts that dust charging affects particles via electrostatic disruption and erosion, and depletes the plasma of electrons with respect to cations (Vigren et al., 2015). Observation of electron depletion may have been hindered at comet 67P due to the relatively low dust abundance and *Rosetta*'s very negative potential, but it has been observed in space in other dusty plasmas. Understanding dusty plasma properties is of interest to nuclear fusion for energy production. Though dust has been observed in space in other dusty plasmas (protoplanetary discs, Matthews et al. (2012), Titan's atmosphere, Lavvas et al. (2013), and Enceladus' plume, Morooka et al. (2011)) the associated dynamical properties of dusty plasma are not yet understood because of lack of dedicated observations. To visit a comet with instrumentation optimised to detect and assess dusty plasma would not only address how cometary dust interacts with ionospheric plasma but also contribute to increase our understanding of dusty plasma for the benefit of the planetary, astrophysical, and plasma communities.

Neutral gas–plasma. How does partially collisional plasma behave? How does it influence the large-scale structure (e.g., diamagnetic cavity)? How is it affected by transient events? Comets offer a unique neutral-plasma environment which evolves from collisional to collisionless regimes with changes in outgassing and cometocentric distance, similar to the partially ionized neutral-plasma environment expected in protoplanetary discs. It is critical to assess the nature of the plasma fine structure, as it plays a critical role in shaping large-scale structures. At comet 67P, the diamagnetic cavity boundary has been linked to instabilities between the unmagnetised, collisional and magnetised, collisionless plasmas (Henri et al., 2017). As the outgassing increases, ion-neutral chemistry occurs, driving ion composition (Heritier et al., 2017). The collisional regime may be affected by solar (coronal mass ejections, corotative interactive regions) and cometary (outbursts) transient events (Hajra

et al., 2017; Goetz et al., 2018). While *Rosetta* was first to unveil the evolution of the coma and its interaction with the solar wind, observations of the plasma dynamics and the boundary location were hampered by the energy resolution of the ion sensors, *Rosetta*'s very negative potential, and the time resolution of the plasma and field sensors. Quasi-collisionless environments are found in planetary exospheres and comae, and are difficult to model as they are at the frontier between kinetic and fluid approaches. It is not clear how collisional and electromagnetic processes are linked together and which role the ion-neutral friction plays. It is hence critical to acquire observational constraints, especially in the absence of satisfying theoretical approaches. With an expanding coma, neutral densities are higher at comets for a given level of collisionality, which makes measurements easier. Comparing the nature of the diamagnetic cavity present in different levels of collisionality and magnetisation (e.g., at comets, after supernovae explosion, or in laser experiments, Winske et al. (2019)) is of high relevance to different communities.

Nucleus–plasma. How is the nucleus affected by the plasma environment, including solar extreme events? Do interactions with the solar wind influence the activity and evolution of comets? At low cometary activity, solar wind sputtering on dust particles residing on the nucleus surface releases non-volatile species (Wurz et al., 2015). Cometary ion sputtering may be a source of neutrals (Yao and Giapis, 2017), though not the main one for O₂ (Heritier et al., 2018). Bombarded by the solar wind, the nucleus is predicted to become electrically charged; the fine dust on its surface could be electrostatically levitated and ejected away (Nordheim et al., 2015). The plasma influence on the nucleus surface during extreme solar events and over time is not yet understood, but may be vital for the assessment of cometary activity and evolution.




Fig. 6: Roadmap of comet space exploration.

4 Missions and technological requirements

In Section 3, we have shown that a large number of major questions need to be addressed by future comet space missions in order to give final answers on the composition and evolution of these bodies and their relationships with other

Scientific goal	Measurement	Instrument	Mission†	Target‡			
				JFC	CEN	DC	LP
How and where did cometary materials get assembled? Which post-planetesimal evolution paths need to be considered?							
Nucleus Internal structure	Internal mass distribution Deep internal structure	Radio Science Low-frequency radar	O O+L,O	X	X	X	X
Nucleus Surface Properties layering	Morphology 3D shape model Shallow subsurface structure, layering Temperature and thermal inertia	High-Res colour camera Medium-Res camera, Altimeter High-frequency radar MIR Radiometer, IR spectrometer	O, L O O O, L	X	X	X	X
Comet building blocks	Pebbles, dust, particles morphology size distribution, porosity, composition	Dust analysers, AFM High Res cameras Low-frequency radar	O, L O, L O+L, O, L	X	X	X	X
Petrologic assemblage phase relationships between minerals, between organics and minerals	Optical properties, mineralogy, crystallinity, molecular composition of bulk and grains	microscopy, SEM, TEM, Raman, IR spectroscopy, XRD	L, SR	X			
What is the presolar heritage of cometary materials? What do comets tell us about large-scale mixing and dynamical processes in the early Solar System?							
Timescales for formation of cometary matter	radiometric ages from short- and long-lived radio isotopes	SIMS, TIMS, ICPMS	SR	X			
Composition of organic matter up to masses of 1000 Da	Mass spectra, infrared spectra	TOF-SIMS, XANES spectroscopy (IOM), GC/LC-MS and FT-ICR/Orbitrap (SOM) MS/IR Spect.	L, SR O SR, CSR	X		X	X
Origin of organic matter	C, N, H isotopic composition	IRMS,GC-IRMS NanoSIMS	CSR	X			
Water ice origin	D/H, ¹⁸ O/ ¹⁶ O, ¹⁷ O/ ¹⁶ O (within 0.1% for CSR) Form (crystalline versus amorphous)	IRMS, NanoSIMS, Laser Spect. Mass spectrometer Thermal probes	O L	X	X	X	X
Volatile presolar heritage Semi-volatiles (e.g., salts)	Molecular and isotopic, abundances, noble gases	Mass Spectrometer Gas chromatography	CSR O, L	X	X	X	X
Composition of mineral phases	Mineralogy, spectroscopy	microscopy, SEM, TEM, Raman, IR spect., APXS, VIS-IR, mid-IR spect.	SR O, L	X		X	X
Elemental abundances asteroids/planets/Sun relationships	Intensity of mass spectral lines	HR-ICP-MS, ICP-OES MS, Orbitrap	SR O, L	X	X	X	X
Presolar grains	Isotope ratios of major elements, with 1% accuracy, mineralogy	NanoSIMS, SIMS, IRMS, TEM	SR	X			

Continued on the next page.

Continued from previous page.

Scientific goal	Measurement	Instrument	Mission†	Target‡				
				JFC	CEN	DC	LP	MBC
How does comet activity work? How do surface and coma observations reconnect with the pristine, deep interior?								
Nucleus subsurface/surface thermal properties	Temperature and thermal inertia	MIR/nm Radiometer, IR spectrometer	O, L	X	X	X	X	X
Active layers properties	Temperature depth profile	Penetrometer/Thermal sensors	L	X	X	X	X	X
Subsurface mechanical properties	Dielectric properties	High-frequency Radar	O	X	X	X	X	X
Comet activity diurnal/seasonal variations	Material strength and layering	Static/dynamic penetrometer	L	X	X	X	X	X
	Coma dust and gas distributions	Colour Camera/IR spectroscopy	O	X	X	X	X	X
	night-side activity	Dust-impact analysers	O, L	X	X	X	X	X
		Mass spectrometer	O, L	X	X	X	X	X
Activity-induced surface changes	Morphology, colour composition, ice distribution	Medium Res Colour Camera	O, L	X	X	X	X	X
	Mass transfer, erosion	VIS-IR spectrometer	O, L	X	X	X	X	X
		High-freq radar, Altimeter	O	X	X	X	X	X
What was the role of comets in the delivery of volatiles and prebiotic compounds to early Earth?								
Prebiotic compounds	Abundance of amino acids and sugar-related compounds	LC-MS, GC-MS, LD-MS (Orbitrap)	CSR, SR, L	X				
Chirality	Enantiomeric proportions	LC-MS, GC-MS, CDS	SR, L	X				
Earth volatiles	Noble gases abundance	MS	SR, CSR	X			X	
Earth formation	Highly siderophile elements abundances (Os, Ir, Ru, Rh, ...)	ICP-MS, TIMS	O, L	X				
			SR	X				
How do the dust coma, the surrounding plasma, and the nucleus interact?								
Dust-plasma interaction	Electron and positive ion densities	see note 1 in caption	O	X	X	X	X	X
	Electric field	Mutual Impedance/Langmuir probes		X	X	X	X	X
	Negative ion density	Negative Ion Composition Analyser		X	X	X	X	X
	Dust grain flux & size distribution (nano to micro)	Dust Analysers		X	X	X	X	X
Coma-plasma interaction (collisionality)	EUV/FUV brightness (Sun)	Langmuir probe, UV Spect.		X	X	X	X	X
	Ion and neutral composition and number density	Ion and Neutral Mass Spectrometer		X	X	X	X	X
	e^- number density/temperature, ion bulk velocity	Pressure gauge		X	X	X	X	X
	Energetic electron/ion flux	Mutual Impedance/Langmuir probes		X	X	X	X	X
	Magnetic field magnitude & components	e^- & ion composition analyser		X	X	X	X	X
		Flux Gate Magnetometer		X	X	X	X	X
Nucleus-plasma interaction (including solar events)	Same as above box	Same as above box	O, L	X	X	X	X	X
	FUV emission brightness	FUV spectrometer		X	X	X	X	X
	Fluxes of major neutral species	IR/Submm High-Res spectr.		X	X	X	X	X

Table 1: Traceability Matrix. See Sect. 4 and Fig. 7. Type of mission[†]: O=Orbiter, L=Lander, **SR**=Sample Return, **CSR**=Cryogenic Sample Return. Target[‡]: JFC=Jupiter Family Comet, CEN=Centaur, DC=Dormant Comet, LP=Long Period comet, DN= Dynamically New Comet, MBC=Main Belt Comet. Instruments Acronyms: Atomic Force Microscope (AFM), Alpha Particle X-Ray Spectrometer (APXS), Circular Dichroism Spectroscopy (CDS), Fourier-Transform Ion-Cyclotron Resonance (FT-ICR), Gas Chromatography (GC), High Resolution Inductively Coupled Plasma Mass Spectroscopy (HR-ICP-MS), Inductively Coupled Plasma Mass Spectrometry (ICPMS), Inductivity Coupled Plasma-Optical Emission Spectroscopy (ICP-OES), Insoluble Organic Matter (IOM), Isotopic Ratio Mass Spectroscopy (IRMS), Liquid Chromatography/Laser Desorption Mass Spectrometry (LC/LD-MS), Scanning Electron Microscopy (SEM), Secondary Ion Mass Spectrometry (SIMS), Soluble Organic Matter (SOM), Time-Of-Flight (TOF), Transmission Spectroscopy (TEM), Thermal ionization Spectroscopy (TIMS), X-Ray Powder Diffraction (XRD), X-ray Absorption Near-Edge Structure (XANES). *Note 1*: to be developed from Langmuir Probe/Mutual Independence Probe.

primitive Solar System bodies, with important implications for understanding Solar System formation and evolution, and planet habitability.

For the ESA Voyage 2050 long-term (2035-2050) plan, *AMBITION* and associated technical challenges should set the bar (see the prospective for future comet missions in Thomas et al. (2019)). Also to be considered is the international context of comet space exploration (see roadmap in Fig. 6 and details in Sect. 5), in particular the pre-selection of the NASA New Frontiers Comet Astrobiology Exploration Sample Return (*CAESAR*) mission for a Phase A study. Though *CAESAR* was not finally selected, it is a strong candidate for the next New Frontiers call (likely to be issued in 2021 or 2022). Future cometary exploration can be performed by both ESA's L and/or M-class mission budgetary envelopes. We are proposing to adopt the **L-class** mission scenario for a mission including an **Orbiting spacecraft (O)** with a **Lander (L)** (Sect. 4.1) and **(Cryogenic) Sample Return (CSR-SR)** (Sect. 4.2). Conversely, the **M-class** mission scenario, limited to an orbiter, seems more suitable for targeting MBCs or Centaurs (Sect. 4.3). In the Traceability Matrix Table 1, we present the appropriate instrumentation (though the list is not exhaustive), including that for the analysis of returned samples, and the associated mission type for different classes of comet families.

Returning a sample collected at depth, or, still more challenging, at cryogenic temperatures and preserving the stratigraphy of the surface layers, is the next step of comet exploration, addressing questions related to the thermo-physical and microphysical properties of subsurface layers (Sects 3.1, 3.4), in addition to the nature and formation of comet materials (Sects 3.2, 3.3, 3.5). Laboratory analyses of returned samples achieve high precision, high resolu-

tion measurements that cannot be performed in space (e.g., Rotundi et al., 2008).

Despite the paramount importance of a cryogenic sample return approach, several key measurements (Table 1) still need instruments operating on orbiter and lander platforms. These include the investigation of the deep interior and active layers to be performed by low-frequency (monostatic or bistatic) and high-frequency radars (Sect. 3.1), isotopic ratios in trace volatiles (Sect. 3.2), and plasma science (Sect. 3.6). Also, the understanding of diffuse activity requires combining sub-surface thermophysical and near-surface coma measurements, necessitating a lander (Sect. 3.4). A landing mission with drills and possibly mobility (or multiple static landers) is attractive in this respect. Instrumentation could include e.g., microscopic imaging, thermal and permittivity probes, and analytical capabilities such as miniaturized mass spectrometers (see Table 1). We provide details in Sect. 4.1 about next-generation landers for the exploration of small Solar System bodies.

To achieve safe landing and sampling, the orbital spacecraft requires at least a narrow angle camera for site selection. To achieve an optimal scientific choice of landing site and understand the context around it, the payload should also include wide angle cameras and IR spectro-imagers, along with dust-impact and gas analysers. Key surface properties such as temperature, thermal inertia, albedo, and ice content, can be measured using VIS-IR spectrometers and MIR/submm spectro/radiometers.

Rosetta explored a Jupiter-family comet (JFC) originating from the Kuiper Belt. This population of comets has perihelia of typically 1 AU and aphelia near the orbit of Jupiter, and is accessible by rendezvous, landing, and sample return missions. These comets are the most suitable targets for sample return. An attractive subset of this population are the hyperactive comets, which are releasing large amounts of icy aggregates and show a terrestrial D/H ratio in water, unlike 67P (Lis et al., 2019). Prototypes are comet 103P/Hartley 2, encountered by the NASA *EPOXI* mission, and 46P/Wirtanen, the initial *Rosetta* target. The mechanisms driving hyperactivity are unknown, and could be related to CO₂ driven-activity, as observed for 103P/Hartley 2 (A’Hearn et al., 2011), to their small size (Lis et al., 2019), with the possibility that these comets are particularly ice-rich. A landing mission to a hyperactive comet (likely an L-class mission) would provide important insights about activity processes and comet diversity.

Some populations of comets are still completely unexplored. We discuss in Sect. 4.3 a variety of different sized missions towards extinct/dormant comets (DC), MBCs, long-period (LP) and dynamically new comets (DN), Centaurs (CEN) and interstellar objects (ISOs). In the Traceability Matrix Table 1, we assumed rendezvous missions for DC, MBC, CEN and LP, and flyby/orbiter science for DN (appropriate also for ISOs). Approximate mission classes are indicated in Fig. 7.

Comet type	Fly-by	Rendezvous	Landing	Sample return	Cryogenic SR
Centaur	M	M/L	L+		
Jupiter Family	<i>DI, DS1, EPOXI</i> etc.	<i>Rosetta</i>	<i>Philae</i>	<i>Stardust</i>	L++
Extinct JFC	F/M	M/L	L	L	
Returning OCC	<i>Giotto</i> etc.	L+	L++		
Dynamically new	<i>Comet Interceptor</i>				
Main Belt Comet	M	<i>ZhengHe-A</i>	L	M (<i>Stardust-like</i>) L (surface)	
Interstellar comet	M				

Fig. 7: Approximate mission classes for different mission and comet types, in increasing complexity from left to right, and covering different evolution stages of comets from the four possible reservoirs (Kuiper Belt, the Oort cloud, the Main Belt, and other planetary systems). Shading indicates approximate cost from yellow (F-class) through orange (M-class) to red (L-class or multi-agency flagship missions). Hatched boxes indicate that such a combination is not seen as feasible, mostly due to excessive Δv requirements. Past and planned missions are shown.

4.1 Lander

The delivery and deployment of a lander on a cometary nucleus is quite challenging due to the low gravity and cometary activity. The available mass for payloads, the generation of power, and the establishment of the communication link with the orbiter are further limitations to the science which could be achieved from an autonomous laboratory housed on a surface lander.

Philae was a complex device with sophisticated mechanisms, designed to cope with a wide range of possible cometary environment and conceptual redundancy (e.g., anchoring harpoons plus 'ice-screws' and a damping mechanism in the landing gear) (Biele and Ulamec, 2008; Ulamec and Biele, 2009). By taking advantage of technological developments of the last 25 years, a next generation lander could be improved in several aspects.

Miniaturization: Not only miniaturization in computers, but also in instrumentation, have advanced over the last years. This, however, does not necessarily apply for mechanisms (e.g., a drill designed to reach a certain depth can hardly be designed smaller). Nevertheless, sensors and electronics would require less mass and volume for a future lander (assuming similar performance). One could also consider a change in concept: while *Philae* was one large lander with ten instruments and sophisticated capabilities (like drilling or rotation), other systems (like the much smaller *MASCOT/Hayabusa2* lan-

der delivered to asteroid Ryugu (Ho et al., 2017)) are more modest in their technical requirements (and variety of science that can be performed), but also more flexible to use. Beyond *MASCOT* there is a trend to apply CubeSats for interplanetary missions, including landing (e.g., Juventas aboard the ESA *Hera* asteroid mission).

Enhanced landing system: While *Philae* descended passively after being ejected from the mother spacecraft, nowadays there exist highly developed systems with 3-axis stabilization, propulsion, and powerful Guidance, Navigation and Control (GNC) ready to be used for small spacecraft or landers. This could improve the flexibility in the landing scenario and increase reliability. Autonomous obstacle recognition and hazard avoidance would minimize landing risks due to surface roughness or boulders, and allow a very accurate touch-down with small landing uncertainties. This could possibly even allow targeting active spots or exposed ice patches that have been identified beforehand with orbiter instruments.

Mobility: In line with a 3-axis propulsion system (e.g., cold gas thrusters), mobility becomes also more easily achievable. Hopping by using the landing gear mechanism was considered for *Philae*, but given up in favor of safe anchoring (ironically, the anchoring did not work, and *Philae* made an un-controlled hop). *MASCOT* allowed some relocation, using an internal torquer (Jaumann et al., 2019). The concept worked fine, but the hops were very short in distance (~ 70 cm). For *MMX* (JAXA/Martian Moons eXplorer) it is foreseen to deliver a small (~ 25 kg) rover to Phobos, and drive several tens of meters in low gravity environment (Ulamec et al., 2019). A new lander design may well allow the investigation of several areas on a cometary surface via mobility, thus, giving insight into the heterogeneity of surface properties and enhancing possible (radar) sounding experiments.

Instrumentation: instrument development has improved since the 1990s and new types of instruments (e.g., Raman spectrometers, LIBS, tele-microscopes) now do have space applications. Other instrument types (like mass spectrometers) are available with better performance/resolution.

4.2 (Cryogenic) Sample return

Sample collection and return to Earth is the fourth step of robotic space exploration, after flybys, orbiter remote science, and lander in-situ science. Each step gets more challenging and usually builds up on the experiences of previous missions. So far, samples have been returned robotically from the Moon (*Apollo & Luna* missions), solar wind (*Genesis*), comet 81P/Wild 2 (*Stardust*), and asteroid Itokawa (*Hayabusa*). Sample return missions to other asteroids are underway or completed (*Hayabusa2* at Ryugu and *OSIRIS-REx* at Bennu), or in the phase of realization (Sect. 5). In the past, an advanced study of the Asteroid Redirect Robotic Mission (*ARRM*) was completed with the aim of grabbing a multi-ton boulder from the surface of a near-Earth object (Mazanek et al., 2015). Cometary sample return has been studied as the original con-

cept of *Rosetta* (Atzei et al., 1994), as well as for *CAESAR*/NASA mission returning to comet 67P (Glavin et al., 2019). In Europe, numerous industrial studies have been performed, in particular during phase A studies at ESA of the *MarcoPolo* and *MarcoPolo-R* asteroid sample return projects (Barucci et al., 2012) for the Cosmic Vision programme (2015-2025), including sampling tool technology and the re-entry capsule (e.g., heat shield material development, aerodynamic stability). The *Triple-F mission*, a Comet Nucleus Cryogenic Sample return, has also been proposed to the Cosmic Vision programme, in collaboration with the Russian space agency (Küppers et al., 2009).

For a future mission, considering an advance in technology, the requirements for sampling, storage, and return to Earth shall be beyond those expressed in the last NASA decadal survey (non-cryogenic, surface only), but should dare to attempt to go a step further. Table 2 lists different sampling strategies with increasing complexity.

Requirement	Possible technique	Studies with references
Coma dust	Aerogel capture	<i>Stardust</i> [1]
Surface dust only	Brush, air-blow system, etc	<i>OSIRIS-REx</i> [2], <i>CAESAR</i> [3]
Surface material, incl. consolidated material and volatiles	Corer, combined brush/rock-chipper	<i>CAESAR</i> [3] <i>CORSAIR-Sampler</i> [4]
Boulder capture	Grabber & manipulator arm	<i>ARRM</i> [5]
Subsurface material to a depth of few cm	Corer, digging system	<i>Philae</i> drill with retrieval system, <i>CORSAIR-Corer</i>
Sub-surface core, protecting the stratigraphy	Complex Corer	'original' <i>Rosetta</i> [6]
Subsurface material, kept at cryogenic temperatures, during sampling, storage, and return	Corer plus cryogenic chain	'original' <i>Rosetta</i> [6] NASA Tech. Study[7]

Table 2: Sample return mission requirements and collection techniques. References: [1] Brownlee et al. (2006); [2] Lauretta et al. (2017); [3] Glavin et al. (2019); [4] Völk et al. (2018); [5] Mazanek et al. (2015); [6] Atzei et al. (1994); [7] Veverka (2011).

The (Cryogenic) Sample Return option is the most ambitious scenario for cometary exploration, in which a sample of surface/subsurface material is collected and returned to Earth. A suitable protocol of cosmochemical measurements to be conducted in terrestrial laboratories to fully characterize the sample has been defined (Table 1). A further advantage of the sample return scenario is that the analysis techniques are not limited by the restricted resources available onboard the spacecraft and by the technology readiness available, but can make use of the full potentialities of terrestrial laboratories, for which future developments in analytical capabilities are possible.

In this scenario two possible options are feasible: 1) *Sample Return* will provide a surface/subsurface sample with the scope to characterize its mineral and organic material phases. The sample return canister will not be pressurized

nor stabilized at cryogenic temperatures, so the volatile fraction will not be preserved during the return voyage. 2) *Cryogenic Sample Return* is similar to the Sample Return option, but with the possibility to pressurize and thermally stabilize the sample at cryogenic temperatures during the entire return voyage to preserve the volatile fraction distribution. A major issue of this second option is the definition of the cryogenic storage temperature and overpressure of the sample after the collection, because these parameters will define the survivability of the volatile species.

From the *Rosetta* and *Philae* measurements, it appears that 67P's surface is characterized by a relatively hard sintered layer at a depth of a few centimetres, upon which sits a layer of loosely consolidated regolith (see Fig. 8). The observed presence of airfall implies that the thickness of this layer of 'sand' is likely to be highly variable but prevalent above the winter hemisphere at perihelion. This material is also subject to gas-driven transport leading to the formation of dune-like structures up to 3 meters in height (Hu et al., 2017; Keller et al., 2017). Hence, we need to be careful about what we actually mean by 'surface' because the regolith, being depleted of volatile species, may not be a representative sample, at least for some science goals. This is a dynamic environment comprising both the residues of particulate matter that have become concentrated as a consequence of the devolatilization operating in the surface layers, and the products of any alteration processes that arise from interactions of surface materials with insolation/radiation etc. It is inevitable that the chemical and physical properties of this surface layer will be different from those of the bulk comet. The nucleus's surface can be made of dehydrated dust (Capaccioni et al., 2015) or of more consolidated terrains (Sierks et al., 2015): due to periodic solar heating and the consequent outgassing of volatiles, the surface is mainly enriched by mineral and organic material but is almost depleted of ices, with the exception of very localized areas (detailed below). This is at the root of the difficulties in measuring the cometary ice-to-dust ratio, a fundamental parameter to link cometary formation conditions with interstellar grain models and observations. In order to respond to this question it is necessary to excavate the surface until ice-rich layers are reached.

The requirements on the cryogenic sample acquisition and storage during the return to Earth are the following: 1) a continuous cylindrical sample from the nucleus surface down to a desirable depth of about 3 m, but not less than 1 m, with a diameter of about 10 cm (corresponding to a volume of about 23.5 liters and mass of about 12 kg); 2) the sampled material should not be mechanically nor thermally altered by the extraction and storing process to maintain stratigraphy; the sample can be stored in a series of, e.g., 6 canisters corresponding to incremental depths of 0.5 m each; 3) since the volatile fraction can be expected preferentially in the lower part of the core sample the canisters housing the inner layers need to be sealed (to maintain controlled overpressure) and thermally stabilized at cryogenic temperature (≤ 90 K). Pressure and temperature of the canister need to be kept both under control to guarantee the survivability of the volatile species in the sample. Apart from very volatile species, like CO, for which saturation vapor pressure is very high (2.4 bar),




Fig. 8: Schematic diagram attempting to reconcile the information we currently have on the interior layers and surface of cometary nuclei. Adapted from Thomas et al. (2019).

an overpressure of about 1 bar is sufficient to maintain stable H_2O , CO_2 , and HCN ices at $T = 90$ K (see Table 6.3 of Veverka, 2011).

While the 3-meter-depth sample will offer the optimal approach to understand the internal structure and composition of a cometary nuclei surface, another possibility to collect volatile-rich cometary samples is offered by the presence of exposed and semi-buried boulders, some up to meter-size, directly accessible from the surface (Fig. 8). The interior of compact boulders may in fact still contain high amounts of water ice, as evidenced by high resolution images from *Rosetta*. Alternatively, several localized areas, up to a few tens of meters in size, indicate the presence of exposed water ice patches (Pommerol et al., 2015; Barucci et al., 2016; Filacchione et al., 2016a) or condensed water (De Sanctis et al., 2015) and carbon dioxide (Filacchione et al., 2016b) (Fig. 4) directly accessible on the surface, thus allowing access to condensed volatile species (albeit with not pristine relative abundances) without the necessity to excavate the interior layers. Grabbing and preserving at controlled temperature one of those boulders or volatile-rich patches would be an alternative approach for a Cryogenic Sample Return mission, although there remain questions as to whether these are representative of the volatile composition of the deep interior.

Any sampling strategy on a cometary body faces several technological challenges in addition to those well known for deep space missions, including the development of a navigation system in the proximity of the nucleus surface, the landing and anchoring system, the uncertainty of the nucleus' surface

properties (roughness, strength, compactness), the sampling/excavation system (stratigraphy preservation), and the uninterrupted cryogenic temperature and pressure controlling systems during the sampling process, handling, return to Earth, atmospheric re-entry, retrieval on Earth, and during curation.

4.3 M-class main belt comet/centaur rendezvous

In depth understanding of a single Jupiter-family comet that goes beyond what *Rosetta* achieved will require a large mission. However, there is also much to be learned by expanding our knowledge of different types of comets, or different evolutionary stages, with some populations still completely unexplored. A variety of different sized missions could potentially contribute to exploration of these populations (Fig. 7), from first reconnaissance flyby missions (e.g., the selected F-class *Comet Interceptor*) to more in depth in situ analysis with rendezvous missions. The different orbits of the various comet classes dictate the cost and complexity of these missions.

Extinct (or dormant) comets can be found within the near-Earth object population, so a first flyby to explore this end-state of comet evolution could be achieved at relatively low cost (e.g., F-class dedicated mission, or as part of a multi-comet/asteroid tour in the M-class category, similar to the *CASTAway* concept proposed to the M5 call (Bowles et al., 2018)). More advanced missions require a better understanding of which are truly extinct comets and how these are separated from D-type asteroids perturbed into higher eccentricity orbits. Also, since such comets are inactive, landing and (sub-)surface sampling would be required to make in situ composition measurements, implying a large mission. Extinct comets have elliptical orbits, meaning that rendezvous or landing missions require significant Δv (a measure of the impulse required to perform a maneuver).

At the other end of the short-period comet evolutionary path we have the Centaurs, objects with relatively unstable orbits in the giant planet region that are in a transition from Kuiper belt objects to JFCs (Peixinho et al., 2019). These have a wide range in orbits, from the nearly circular orbit of 29P/Schwassmann-Wachmann 1 just outside of Jupiter, to those that orbit beyond Uranus, and those with eccentric orbits that cross planet orbits. They also vary widely in size, including bodies much larger than any known comet nucleus, like Chariklo, a ~ 200 km diameter minor planet with its own ring system (Braga-Ribas et al., 2014). Only a fraction ($< 10\%$) of Centaurs are observed to show comet-like activity, triggered likely by the sublimation of hypervolatiles such as CO, as measured for 29P. The activity of Centaurs is often punctuated by large outbursts. For example, 29P shows almost constant activity and very regular large (many magnitudes increase in coma brightness) outbursts, the triggering mechanism of which being unknown. A mission to a Centaur would therefore be very interesting for comet science, to understand how comet activity works beyond the water ice snowline, and to see an earlier phase in comet evolution for comparison with results from JFC missions. The

challenges of such a mission are primarily due to the large distance from the Sun of the Centaurs, implying long cruise times and low power levels for a solar-powered spacecraft. Even a flyby mission would most likely be at least an M-class, and a rendezvous would certainly be at the upper end of M-class and into L-class. The scientific payload required would include remote sensing instruments to study the nucleus and in situ instrumentation to measure gas and dust composition, with the mission class and available budget dictating the choice between a minimal suite of a camera and mass spectrometer for a basic investigation, up to a *Rosetta*-like array of instruments for an in-depth study. The difficulty of such a mission depends greatly on how distant a target is chosen – 29P, just outside of Jupiter, is probably the most feasible, with its circular orbit lending itself well to electric propulsion. Finally, a first look at a Centaur could be provided by a flyby en-route to Uranus or Neptune, if an Ice Giants mission is selected by any agency, provided a suitable target is found.

The main belt comets are a relatively easily accessible population, as they have low eccentricity orbits within the asteroid belt. The *Castalia* MBC rendezvous mission was proposed to the M4 and M5 calls (Snodgrass et al., 2018), with the latter version passing programmatic and technical constraints and receiving positive feedback from the panel, even if it was not selected in the end. The goals of this mission were to make a first exploration of this new class of comets, confirm the presence of water ice in small main belt objects, understand how the activity of MBCs worked, test whether or not the water they contain is compatible with the main belt comets being the source of Earth’s water, and finally to use all of these results to better constrain models of Solar System formation and evolution. The proposed instrument payload built on *Rosetta* heritage to allow a direct comparison between MBCs and JFCs, and to enable the necessary very sensitive in situ measurements. The very low activity level of MBCs dictated a prolonged period at very close range to the comet during its few-month active period, but advances in spacecraft autonomy in navigation relative to *Rosetta*’s capabilities make such a proposal feasible as a medium sized mission. A (sub-)surface package (lander/penetrator) is not feasible for such a mission, but would certainly add further capabilities in a larger class mission, especially if it has the capability to reach volatile-rich layers, which are thought to be buried at some metres depth over most of the surface of MBCs. The recently approved *ZhengHe/AEM* (Asteroid Exploration Mission) Chinese mission plans to make the first visit to an MBC (Sect. 5), but does not expect to operate close to the comet during its active period, or sample the sub-surface, meaning that its ability to measure volatile composition will be very limited. Another proposal to the NASA Discovery call, *Proteus*, proposes a similar mission to *Castalia*, but, if not selected, an ESA MBC mission remains a compelling case for a future M-class comet mission.

Finally, the exploration of comets from more distant reservoirs has been very limited so far. *Giotto* and the other spacecraft in the ‘Halley armada’ flew by **1P/Halley**, an evolved returning **Oort cloud comet** that is now in a relatively short-period orbit, while the newly selected F-class mission *Comet Interceptor* is expected to make the first flyby of a new, pristine, Oort cloud

comet entering the inner Solar System for the first time. For such new comets, more advanced missions (rendezvous, etc) are unlikely to be feasible any time soon due to the short warning time between discovery and perihelion and the very large Δv that would be required to match the speed of the comet. A rendezvous with a returning comet, such as 1P on its next return in 2061, could be imagined, but would be a large mission, and the scientific gain versus a more capable mission at a short-period comet, including possible sample return, is not obvious.

Comets from other star systems, i.e., interstellar objects (ISOs), present a similar, but even more extreme problem, but would be very exciting mission targets (as the only feasible way to study extrasolar material in situ). Especially, if future examples are similar to the first discovered ‘Oumuamua and display little visible activity, they will only be discovered with very short warning time (Meech et al., 2017). It is expected that the Rubin Observatory (previously known as the Large Synodic Survey Telescope, LSST) will find more ISOs, perhaps 1 per year: we will have a better idea of the true population characteristics and arrival rate by the 2030s. A mission similar to *Comet Interceptor*, i.e., designed to wait in space for a suitable target to be found, and then make a fast flyby, could then be imagined to encounter an ISO. This would probably be a more expensive mission than *Comet Interceptor*, as it would likely require significantly more Δv and a rapid reaction operations scheme, but it could be proposed within the M-class budget.

5 International context of comet space and Earth-based exploration

Space exploration. Numerous space missions to small Solar System bodies have been proposed to ESA, NASA, JAXA, and CAST. Besides the selected ESA F-class *Comet Interceptor* mission, the only cometary mission in the landscape is the *ZhengHe/AEM* chinese mission to be launched in 2022. The current plan is a sample return from the small near-Earth asteroid 2016 HO3, followed by a rendezvous with the MBC 133P/Elst-Pizarro in 2030 (Zhang et al., 2019). The payload of the *ZhengHe/AEM* orbiter is expected to include wide/narrow angle cameras, visible/near-infrared imaging and thermal emission spectrometers, dust analysers, a mass spectrometer, γ -ray/neutron spectrometers, and a low frequency radar. *ZhengHe/AEM* also flies nano-landers for studies of surface and subsurface properties. The *ZhengHe/AEM* mission is technology driven and it remains unclear whether or not it will operate at 133P during its short active period, enabling comparable measurements to previous comet missions.

The Comet Astrobiology Exploration Sample Return (*CAESAR*) mission was pre-selected for a Phase A study in the New Frontiers NASA program, but did not pass the final selection. *CAESAR* was designed to acquire a minimum of 80 g of solid material from the surface of the comet, and to store sublimated volatiles after warming in a gas containment system (Lauretta et al., 2019).

CAESAR would have arrived in 2028 at 67P, and the capsule (provided by JAXA) would have been delivered in 2038.

Two sample return missions to asteroids are ongoing. After the success of the *Hayabusa* mission, which returned dust particles from S-type asteroid (25143) Itokawa, JAXA launched *Hayabusa2*, to study C-type asteroid (162173) Ryugu. *Hayabusa2*, whose payload includes a lander and three small rovers, returned surface material to Earth on 5 December 2020 (Watanabe et al., 2019). The NASA New Frontiers mission *OSIRIS-REx* reached B-type asteroid (101955) Bennu in December 2018, and will return the sample (> 60 g of regolith) in 2023 (Lauretta et al., 2017).

Earth-based observations of comets. Comets are mostly studied by telescopes. Observations are generally focusing on the physical and chemical properties of the coma and tails. Direct studies of nucleus properties, even size and color, are difficult due to their small size and the presence of the coma, though recent progress has been made thanks to the availability of large telescopes (e.g., Kokotanekova et al., 2017). Telescopic observations of comets are unavoidable to perform statistical studies and investigate differences and links between the various comet populations, including main belt comets and transition bodies such as centaurs. Advances in comet knowledge strongly benefitted from the development of new instrumentations combined with the legacy of several decades of dedicated surveys. Next generation telescopes will certainly provide breakthroughs in several aspects of comet science. For example, using the *James Webb Space Telescope (JWST)*, it will be possible to detect the main drivers of cometary activity, H₂O, CO₂, and CO, out to unprecedented heliocentric distances, and to conceivably test the cometary nature of main belt comets by the direct detection of water; spatially resolved infrared spectra will be used to study the properties of water ice particles, the nature and relative amounts of dust amorphous and crystalline silicates, and possibly detect nucleus surface signatures (Kelley et al., 2016). Giant telescopes in the optical and near-IR, such as the Extremely Large Telescope (ELT), also have an important potential for measuring, e.g., C, N, and H isotopic ratios in gas-phase species and detecting weakly abundant organic molecules. The Rubin Observatory is expected to vastly increase the number of known and characterized comet-like objects. In the submillimeter range, the high-resolution and sensitivity capabilities of ALMA make possible the detection of thermal emission from comet nuclei, from which insights on their size and surface thermal skin depth can be obtained; worth mentioning also is the study of gas-dust interrelations through the detailed mapping of the distributions of gases and dust particles in the coma with ALMA. Science objectives presented in this White Paper cannot be addressed by telescopic observations.

Laboratory developments. Laboratory experiments play a fundamental role in research programs designed to investigate the properties and evolution of comets. Experiments are diverse, and include the analyses of dust and ice analogues by complementary methods aimed at quantitative studies of mor-

phology, structure, chemistry, and optical behavior, and simulation of processes occurring at the early stages of Solar System formation, or at the surface and in the interior of cometary nuclei, cf, e.g., the KOSI experiment (Kochan et al., 1998). Such data aid in interpreting observations performed remotely or in situ, and provide strong constraints for theoretical models. Recently, the international network CoPhyLab (Comet Physics Laboratory), comprising scientists from TU Braunschweig, IWF Graz, University of Bern, MPS Göttingen, DLR, Berlin and the University of Stirling, was founded with the aim at experimentally and theoretically investigating the thermophysical behavior of dust-ice mixtures under cometary conditions, e.g., with various levels of insolation and with state-of-the-art dust and ice compositions and morphologies. Apart from this collaboration, the international network of laboratories is a growing community which includes many research groups involved in the characterization of physical and chemical properties of cometary materials.

6 Conclusion

This White Paper proposes that *AMBITION*, a Comet Nucleus Sample Return mission, be a cornerstone of ESA's Voyage 2050 programme. Rendezvous missions to main belt comets and centaurs are compelling cases for M-class missions, expanding our knowledge by exploring new classes of comets. *AMBITION* would engage a wide community, drawing expertise from a vast range of disciplines within planetary science and astrophysics. With *AMBITION*, Europe will continue its leadership in the exploration of the most primitive Solar System bodies.

References

- Ábrahám, P., Juhász, A., Dullemond, C. P., et al. 2009, *Nature*, 459, 224
- A'Hearn, M. F., Belton, M. J. S., Delamere, W. A., et al. 2005, *Science*, 310, 258
- A'Hearn, M. F., Belton, M. J. S., Delamere, W. A., et al. 2011, *Science*, 332, 1396
- Aikawa, Y., Herbst, E. 1999, *ApJ*, 526, 314
- Alexander, C. M. O. 'D ., Cody, G. D., De Gregorio, B. T., et al. 2017, *Chemie der Erde / Geochemistry*, 77, 227
- Altwegg, K., Balsiger, H., Bar-Nun, A., et al. 2015, *Science*, 347, 1261952
- Altwegg, K., Balsiger, H., Bar-Nun, A., et al. 2016, *Science Advances*, 2, e1600285
- Altwegg, K., Balsiger, H., Berthelier, J. J., et al. 2017a, *MNRAS*, 469, S130
- Altwegg, K., Balsiger, H., Berthelier, J. J., et al. 2017b, *Philosophical Transactions of the Royal Society of London Series A*, 375, 20160253
- Altwegg, K., Balsiger, H., Hänni, N., et al. 2020, *Nature Astronomy*, 4, 533
- André, P., Montmerle, T. 1994, *ApJ*, 420, 837
- Armitage, P. J. 2019, *Saas-Fee Advanced Course*, 45, 1
- Attree, N., Groussin, O., Jorda, L., et al. 2018, *A&A*, 611, A33
- Attree, N. et al., 2019, *A&A*, 630, A18
- Atzei, A., Hechler, M., Schwehm, G. and Mitchell, R., 1994, *Adv. Space Res.*, (12)197-205, 1994
- Bachiller, R. 1996, *ARAA*, 34, 111
- Bacmann, A., Taquet, V., Faure, A., et al. 2012, *A&A*, 541, L12
- Banerjee, R., Pudritz, R. E. 2006, *ApJ*, 641, 949
- Bardyn, A., Baklouti, D., Cottin, H., et al. 2017, *MNRAS*, 469, S712
- Barucci, M. A., Cheng, A. F., Michel, P., et al. 2012, *Experimental Astronomy*, 33, 645
- Barucci, M. A., Filacchione, G., Fornasier, S., et al. 2016, *A&A*, 565, A102
- Belton, M. J. S., Thomas, P., Veverka, J., et al. 2007, *Icarus*, 187, 332
- Bertaux, J.-L., Lallement, R. 2017, *MNRAS*, 469, S646
- Bianchi, E., Codella, C., Ceccarelli, C., et al. 2019, *MNRAS*, 483, 1850
- Biele, J., Ulamec, S. 2008, *Space Sci. Rev.*, 138, 275
- Bieler, A., Altwegg, K., Balsiger, H., et al. 2015, *Nature*, 526, 678
- Biver, N., Bockelée-Morvan, D., Moreno, R., et al. 2015, *Science Advances*, 1, 1500863
- Biver, N., Bockelée-Morvan, D., Hofstadter, M., et al. 2019, *A&A*, 630, A19
- Blum, J. 2018, *SSRev*, 214, 52
- Blum, J., Gundlach, B., Krause, M., et al. 2017, *MNRAS*, 469, S755
- Bockelée-Morvan, D., Gautier, D., Hersant, F., et al. 2002, *A&A*, 384, 1107
- Bockelée-Morvan, D., Calmonte, U., Charnley, S., et al. 2015, *Space Sci. Rev.*, 197, 47
- Bockelée-Morvan, D., Biver, N. 2017, *Philosophical Transactions of the Royal Society of London Series A*, 375, 20160252

- Boehnhardt, H., Bibring, J.-P., Apathy, I., et al. 2017, *Philosophical Transactions of the Royal Society of London Series A*, 375, 20160248
- Boss, A. P. 2008, *Earth and Planetary Science Letters*, 268, 102
- Bowles, N. E., Snodgrass, C., Gibbings, A., et al. 2018, *Advances in Space Research*, 62, 1998
- Bradley, J. P. 1994, *Science*, 265, 925
- Braga-Ribas, F., Sicardy, B., Ortiz, J. L., et al. 2014, *Nature*, 508, 72
- Brownlee, D., Tsou, P., Aléon, J., et al. 2006, *Science*, 314, 1711
- Brownlee, D. 2014, *Annual Review of Earth and Planetary Sciences*, 42, 179
- Burch, J. L., Gombosi, T. I., Clark, G., Mokashi, P., Goldstein, R. 2015, *GRL*, 42, 6575
- Busemann, H., Nguyen, A. N., Cody, G. D., et al. 2009, *Earth and Planetary Science Letters*, 288, 44
- Calmonte, U., Altwegg, K., Balsiger, H., et al. 2017, *MNRAS*, 469, S787
- Capaccioni, F., Coradini, A., Filacchione, G., et al., 2015, *Science*, 347, aaa0628
- Campins, H., Hargrove, K., Pinilla-Alonso, N., et al. 2010, *Nature*, 464, 1320
- Choukroun, M., Altwegg, K., Kührt, E., et al. 2020, *Space Sci. Rev.*, 216, 44
- Ciesla, F. J. 2009, *Meteoritics and Planetary Science*, 44, 1663
- Clayton, D. D., Nittler, L. R. 2004, *Ann. Rev. Astron. Astrophys.*, 42, 39
- Cochran, A. L., Cochran, W. D., Barker, E. S. 2000, *Icarus*, 146, 583
- Combe, J.-P., Raponi, A., Tosi, F., et al. 2019, *Icarus*, 318, 22
- Cottin, H., Kotler, J. M., Bartik, K., et al. 2017, *SSrev*, 209, 1
- Cravens, T. E. 1987, *Advances in Space Research*, 7, 147
- Cronin, J. R., Pizzarello, S. 1997, *Science*, 275, 951
- Dartois, E., Engrand, C., Duprat, J., et al. 2018, *A&A*, 609, A65
- Davidsson, B. J. R., Sierks, H., Güttler, C., et al. 2016, *A&A*, 592, A63
- DeMeo, F. E., Carry, B. 2014, *Nature*, 505, 629
- De Sanctis, M. C., Capaccioni, F., Ciarniello, M., et al. 2015, *Nature*, 525, 500
- Drozdovskaya, M. N., Walsh, C., van Dishoeck, E. F., et al. 2016, *MNRAS*, 462, 977
- Drozdovskaya, M. N., van Dishoeck, E. F., Rubin, M., et al. 2019, *MNRAS*, 490, 50
- Duprat, J., Dobrică, E., Engrand, C., et al. 2010, *Science*, 328, 742
- El-Maarry, M. R., Groussin, O., Thomas, N., et al. 2017, *Science*, 355, 1392
- El-Maarry, M. R., Groussin, O., Keller, H. U., et al. 2019, *Space Sci. Rev.*, 215, 36
- Engelhardt, I. A. D., Eriksson, A. I., Vigren, E., et al. 2018, *A&A*, 616, A51
- Engrand, C., Charon, E., Duprat, J., et al. 2018, *Lunar and Planetary Science Conference*, 2015
- Filacchione, G., De Sanctis, M. C., Capaccioni, F., et al. 2016a, *Nature*, 529, 368
- Filacchione, G., Raponi, A., Capaccioni, F., et al. 2016b, *Science*, 354, 1563
- Filacchione, G., Capaccioni, F., Ciarniello, M., et al. 2016c, *Icarus*, 274, 334
- Filacchione, G., Groussin, O., Herny, C., et al. 2019, *Space Sci. Rev.*, 215, 19
- Floss, C., Stadermann, F. J., Kearsley, A. T., et al. 2013, *ApJ*, 763, 140

- Fornasier, S., Mottola, S., Keller, H. U., et al. 2016, *Science*, 354, 1566
- Fornasier, S., Feller, C., Lee, J.-C., et al. 2017, *MNRAS*, 469, S93
- Fray, N., Bardyn, A., Cottin, H., et al. 2016, *Nature*, 538, 72
- Fray, N., Baklouti, D., Bardyn, A., et al. 2017a, EGU General Assembly Conference Abstracts, 12953
- Fray, N., Bardyn, A., Cottin, H., et al. 2017b, *MNRAS*, 469, S506
- Fulle, M., Altobelli, N., Buratti, B., et al. 2016a, *MNRAS*, 462, S2
- Fulle, M., Della Corte, V., Rotundi, A., et al. 2016b, *MNRAS*, 462, S132
- Fulle, M., Della Corte, V., Rotundi, A., et al. 2015, *ApJL*, 802, L12
- Fulle, M., Blum, J. 2017, *MNRAS*, 469, S39
- Fulle, M., Blum, J., Green, S. F., et al. 2019, *MNRAS*, 482, 3326
- Gasc, S., Altwegg, K., Balsiger, H., et al. 2017, *MNRAS*, 469, S108
- Glavin, D. P., Squyres, S. W., Chu, P. C., et al. 2019, *LPI*, 2132, id.2541
- Goetz, C., Volwerk, M., Richter, I., et al. 2017, *MNRAS*, 469, S268
- Goetz, C., et al., 2018, *A&A*, <https://doi.org/10.1051/0004-6361/201833544>
- Goesmann, F., Raulin, F., Bredehöft, J. H., et al. 2014, *P&SS*, 103, 318
- Gombosi, T. I., Burch, J. L., Horányi, M. 2015, *A&A*, 583, A23
- Gounelle, M., Morbidelli, A., Bland, P. A., et al. 2008, *The Solar System Beyond Neptune*, 525
- Gounelle, M., 2011, *Elements* 7, 29
- Gulkis, S., Allen, M., von Allmen, P., et al. 2015, *Science*, 347, aaa0709
- Gundlach, B., Schmidt, K. P., Kreuzig, C., et al. 2018, *MNRAS*, 479, 1273
- Güttler, C., Blum, J., Zsom, A., Ormel, C. W., Dullemond, C. P. 2010, *A&A*, 513, A56
- Hajra, R., Henri, P., Vallières, X., et al. 2017, *A&A*, 607, A34
- Hansen, K. C., Altwegg, K., Berthelier, J.-J., et al. 2016, *MNRAS*, 462, S491
- Hartogh, P., Lis, D. C., Bockelée-Morvan, D., et al. 2011, *Nature*, 478, 218
- Hässig, M., Altwegg, K., Balsiger, H., et al. 2015, *Science*, 347, aaa0276
- Hässig, M., Altwegg, K., Balsiger, H., et al. 2017, *A&A*, 605, A50
- Henri, P., Vallières, X., Hajra, R., et al. 2017, *MNRAS*, 469, S372
- Herique, A., Kofman, W., Beck, P., et al. 2016, *MNRAS*, 462, S516
- Herique, A., Kofman, W., Zine, S., et al. 2019, *A&A*, 630, A6
- Heritier, K. L., Altwegg, K., Balsiger, H., et al. 2017, *MNRAS*, 469, S427
- Heritier, K., et al., Reply to On the origin of molecular oxygen in cometary comae, 2018, *Nature Comm.*, 9, 2581
- Heritier, K. L., Galand, M., Henri, P., et al. 2018, *A&A*, 618, A77
- Ho, T.-M., Baturkin, V., Grimm, C., et al. 2017, *Space Sci. Rev.*, 208, 339
- Hoppe, P., Rubin, M., Altwegg, K. 2018, *Space Science Review*, 214, 106
- Hsieh, H. H., Jewitt, D. 2006, *Science*, 312, 561
- Hsieh, H. H., Denneau, L., Wainscoat, R. J., et al. 2015, *Icarus*, 248, 289
- Hu, X., Shi, X., Sierks, H., et al. 2017, *A&A*, 604, A114
- Ishii, H. A., Bradley, J. P., Dai, Z. R., et al. 2008, *Science*, 319, 447
- Isnard, R., Bardyn, A., Fray, N., et al. 2019, *A&A*, 630, A27
- Jaumann, R., Schmitz, N., Ho, T.M. et al. 2019, *Science*, 365, 817
- Jessberger, E. K., Christoforidis, A., Kissel, J. 1988, *Nature*, 332, 691
- Johansen, A., Oishi, J. S., Mac Low, M.-M., et al. 2007, *Nature*, 448, 1022

- Johansson, F. L., Odelstad, E., Paulsson, J. J. P., et al. 2017, *MNRAS*, 469, S626
- Jones, A. P. 2005, in *Chondrites and the protoplanetary disk*. ASP Conference Series, 341, 251
- Jutzi, M., Benz, W., Toliou, A., et al. 2017, *A&A*, 597, A61
- Kofman, W., Herique, A., Barbin, Y., et al. 2015, *Science*, 349,
- Kokotanekova, R., Snodgrass, C., Lacerda, P., et al. 2017, *MNRAS*, 471, 2974
- Keller, H. U., Arpigny, C., Barbieri, C., et al. 1986, *Nature*, 321, 320
- Keller, H. U., Mottola, S., Hviid, S. F., et al. 2017, *MNRAS*, 469, S357
- Kelley, M. S. P., Woodward, C. E., Bodewits, D., et al. 2016, *PASP*, 128, 018009
- Kissel, J., Brownlee, D. E., Buchler, K., et al. 1986, *Nature*, 321, 336
- Kissel, J., Krueger, F. R. 1987, *Nature*, 326, 755
- Kochan, H. W., Huebner, W. F., Sears, D. W. G. 1998, *Earth Moon and Planets*, 80, 369
- Kramer, T., Noack, M. 2016, *ApJL*, 823, L11
- Kramer, T., Läuter, M. 2019, *A&A*, 630, A4
- Krot, A. N., Petaev, M. I., Russell, S. S., et al. 2004, *Chemie der Erde / Geochemistry*, 64, 185
- Küppers, M., Keller, H. U., Kührt, E., et al. 2009, *Experimental Astronomy*, 23, 809
- Küppers, M., O'Rourke, L., Bockelée-Morvan, D., et al. 2014, *Nature*, 505, 525
- Lauretta, D. S., Balram-Knutson, S. S., Beshore, E., et al. 2017, *Space Sci. Rev.*, 212, 925
- Lauretta, D. S., Squyres, S. W., Bermúdez, L., et al. 2019, *Lunar and Planetary Science Conference*, 50, 2642
- Lavvas, P., Yelle, R.V., Koskinen, T., et al. 2013, *PNAS*, 110, 2729
- Lawler, M. E., Brownlee, D. E. 1992, *Nature*, 359, 810
- Levasseur-Regourd, A.-C., Agarwal, J., Cottin, H., et al. 2018, *Space Sci. Rev.*, 214, 64
- Lis, D. C., Bockelée-Morvan, D., Güsten, R., et al. 2019, *A&A*, 625, L5
- Lorek, S., Gundlach, B., Lacerda, P., Blum, J. 2016, *A&A*, 587, A128
- Lorek, S., Lacerda, P., Blum, J. 2018, *A&A*, 611, A18
- Luspay-Kuti, A., Hässig, M., Fuselier, S. A., et al. 2015, *A&A*, 583, A4
- Machida, M. N., Matsumoto, T. 2011, *MNRAS*, 413, 2767
- MacPherson, G.J., et al. 2005, in *Chondrites and the protoplanetary disk*. ASP Conference Series, 341, 225
- Mannel, T., Bentley, M. S., Schmied, R., et al. 2016, *MNRAS*, 462, S304
- Mannel, T., Bentley, M. S., Boakes, P. D., et al. 2019, *A&A*, 630, A26
- Marshall, D., Groussin, O., Vincent, J.-B., et al. 2018, *A&A*, 616, A122
- Marty, B., Avice, G., Sano, Y., et al. 2016, *Earth and Planetary Science Letters*, 441, 91
- Marty, B., Altwegg, K., Balsiger, H., et al. 2017, *Science*, 356, 1069
- Massironi, M., Simioni, E., Marzari, F., et al. 2015, *Nature*, 526, 402
- Matthews, L. S., Land, V., Hyde, T. W. 2012, *ApJ*, 744, 8
- Matzel, J. E. P., Ishii, H. A., Joswiak, D., et al. 2010, *Science*, 328, 483

- Mazanek, D. D., Merrill, R. G., Brophy, J. R., Mueller, R. P., 2015, *Acta Astronautica*, 117, 163
- Meech, K. J., Weryk, R., Micheli, M., et al. 2017, *Nature*, 552, 378
- Mendis, D. A., Horányi, M., *Rev Geophysics*, 2013, 51, 53
- Migliorini, A., Piccioni, G., Capaccioni, F., et al. 2016, *A&A*, 589, A45
- Modica, P., Meinert, C., de Marcellus, P., et al. 2014, *ApJ*, 788, 79
- Morbidelli, A., Rickman, H. 2015, *A&A*, 583, A43
- Morooka, M. W., Wahlund, J.-E., Eriksson, A. I., et al. 2011, *Journal of Geophysical Research (Space Physics)*, 116, A12221
- Mousis, O., Petit, J.-M., Wurm, G., et al. 2007, *A&A*, 466, L9
- Mumma, M. J., Charnley, S. B. 2011, *ARA&A*, 49, 471
- Nakashima, D., Ushikubo, T., Kita, N. T., et al. 2015, *Earth and Planetary Science Letters*, 410, 54
- Nilsson, H., Wieser, G. S., Behar, E., et al. 2017, *MNRAS*, 469, S252
- Nordheim, T. A., Jones, G. H., Halekas, J. S., Roussos, E., Coates, A. J. 2015, *P&SS*, 119, 24
- Ogliore, R. C., Huss, G. R., Nagashima, K., et al. 2012, *ApJL*, 745, L19
- Oró, J. 1961, *Nature*, 190, 389
- Oro, J., Cosmovici, C. B. 1997, *IAU Colloq. 161: Astronomical and Biochemical Origins and the Search for Life in the Universe*, 161, 97
- Pätzold, M., Andert, T. P., Hahn, M., et al. 2019, *MNRAS*, 483, 2337
- Peixinho, N., Thirouin, A., Tegler, S. C., et al. 2019, *arXiv:1905.08892*
- Pizzarello, S., Cooper, G. W., Flynn, G. J. 2006, *Meteorites and the Early Solar System II*, 625
- Poch, O., Istiqomah, I., Quirico, E., et al. 2020, *Science*, 367, aaw7462
- Podolak, M., Zucker, S. 2004, *Meteoritics and Planetary Science*, 39, 1859
- Pommerol, A., Thomas, N., El-Maarry, M. R., et al., 2015, *A&A*, 583, A25
- Poulet, F., Lucchetti, A., Bibring, J.-P., et al. 2016, *MNRAS*, 462, S23
- Quirico, E., Moroz, L. V., Schmitt, B., et al. 2016, *Icarus*, 272, 32
- Raponi, A., Ciarniello, M., Capaccioni, F., et al. 2020, *Nature Astronomy*, 4, 500
- Rotundi, A., Baratta, G. A., Borg, J., et al. 2008, *Meteoritics and Planetary Science*, 43, 367
- Rubin, M., Altwegg, K., Balsiger, H., et al. 2017, *A&A*, 601, A123
- Ruzicka, A., Floss, C., Hutson, M. 2012, *Geo. Chemica Acta*, 79, 79
- Schroeder, I. R. H. G., Altwegg, K., Balsiger, H., et al. 2019, *MNRAS*, 489, 4734
- Schwartz, S. R., Michel, P., Jutzi, M., et al. 2018, *Nature Astronomy*, 2, 379
- Shi, X., Hu, X., Sierks, H., et al. 2016, *A&A*, 586, A7
- Shu, F. H. 1977, *ApJ*, 214, 488
- Shu, F. H., Shang, H., Lee, T. 1996, *Science*, 271, 1545
- Sierks, H., Barbieri, C., Lamy, P. L., et al., 2015, *Science*, 347, aaa1044
- Simon Wedlund, C., Behar, E., Nilsson, H., et al. 2019, *A&A*, 630, A37
- Snodgrass C., et al. 2017, *A&ARv*, 25, 5
- Snodgrass, C., Jones, G. H., Boehnhardt, H., et al. 2018, *Advances in Space Research*, 62, 1947

- Soderblom, L. A., Becker, T. L., Bennett, G., et al. 2002, *Science*, 296, 1087
- Spohn, T., Knollenberg, J., Ball, A. J., et al. 2015, *Science*, 349,
- Taquet, V., van Dishoeck, E. F., Swayne, M., et al. 2018, *A&A*, 618, A11
- Taylor, M. G. G. T., Altobelli, N., Buratti, B. J., Choukroun, M. 2017, *Philosophical Transactions of the Royal Society of London Series A*, 375, 20160262
- Thomas, N., Sierks, H., Barbieri, C., et al. 2015, *Science*, 347, aaa0440
- Thomas, N., Ulamec, S., Kührt, E. et al., 2019, *Space Sci. Rev.*, 215, 47
- Tsiganis, K., Gomes, R., Morbidelli, A., Levison, H. F. 2005, *Nature*, 435, 459
- Ulamec, S., Biele, J. 2009, *Advances in Space Research*, 44, 847
- Ulamec, S., Michel, P., Grott, M., 20119, 70th International Astronautical Congress, IAC-19-A3.4.B8
- Veverka, J., 2011, Cryogenic Comet Nucleus Sample Return (CNSR) Mission Technology Study, in NASA Planetary Science Decadal Survey 2013-2022
- Vigren, E., Galand, M., Lavvas, P., Eriksson, A. I., Wahlund, J.-E. 2015, *ApJ*, 798, 130
- Vincent, J.-B., A’Hearn, M. F., Lin, Z.-Y., et al. 2016a, *MNRAS*, 462, S184
- Vincent, J.-B., Oklay, N., Pajola, M., et al. 2016b, *A&A*, 587, A14
- Völk, S., Ulamec, S., Biele, J., et al., 2018, *Acta Astron.*, 152, 218
- Walsh, K. J., Morbidelli, A., Raymond, S. N., O’Brien, D. P., & Mandell, A. M. 2011, *Nature*, 475, 206
- Walsh, C., Millar, T. J., Nomura, H., et al. 2014, *A&A*, 563, A33
- Watanabe, S., Hirabayashi, M., Hirata, N., et al. 2019, *Science*, 364, 268
- Weidenschilling, S. J. 1977, *MNRAS*, 180, 57
- Winske, D., et al., 2019, *Front. Astron. Space Sci.*, 5, 51
- Wooden, D. H., Butner, H. M., Harker, D. E., et al. 2000, *Icarus*, 143, 126
- Wurz, P., Rubin, M., Altwegg, K., et al. 2015, *A&A*, 583, A22
- Yao, Y., Giapis, K. P. 2017, *Nature Communications*, 8, 15298
- Zhang X., Huang J., Wang T., Huo Z., 2019, *LPI*, 50, 1045
- Zinner E. K., Holland H. D. and Turekian K. K. 2005, In *Meteorites, Comets and Planets: Treatise on Geochemistry* (eds. A. M. Davis, H. D. Holland and K. K. Turekian), pp. 17-40. Elsevier-Pergamon, Oxford.
- Zsom, A., Ormel, C. W., Güttler, C., Blum, J., Dullemond, C. P. 2010, *A&A*, 513, A57