



Publication Year	2020
Acceptance in OA	2025-03-03T09:49:37Z
Title	On the Metamorphosis of the Bailey Diagram for RR Lyrae Stars
Authors	Bono, G., BRAGA, Vittorio Francesco, Crestani, J., FABRIZIO, Michele, Sneden, C., MARCONI, Marcella, Preston, G. W., Mullen, J. P., Gilligan, C. K., FIORENTINO, Giuliana, PIETRINFERNI, Adriano, ALTAVILLA, Giuseppe, Buonanno, R., Chaboyer, B., OLIVEIRA DA SILVA, Ronaldo, DALL'ORA, Massimo, Degl'Innocenti, S., DI CARLO, Elisa, Ferraro, I., Grebel, E. K., Iannicola, G., INNO, LAURA, Kovtyukh, V., Kunder, A., Lemasle, B., Marengo, M., MARINONI, Silvia, MARRESE, Paola Maria, MARTINEZ VAZQUEZ, CLARA EUGENIA, Matsunaga, N., Monelli, M., Neeley, J., NONINO, Mario, Moroni, P. G. Prada, Prudil, Z., Stetson, P. B., Thévenin, F., Tognelli, E., Valenti, E., Walker, A. R.
Publisher's version (DOI)	10.3847/2041-8213/ab9538
Handle	http://hdl.handle.net/20.500.12386/36346
Journal	THE ASTROPHYSICAL JOURNAL LETTERS
Volume	896

On the Metamorphosis of the Bailey diagram for RR Lyrae stars

G. BONO,^{1,2} V. F. BRAGA,^{2,3} J. CRESTANI,^{1,2,4} M. FABRIZIO,^{2,3} C. SNEDEN,⁵ M. MARCONI,⁶
G. W. PRESTON,⁷ J. P. MULLEN,⁸ C. K. GILLIGAN,⁹ G. FIORENTINO,² A. PIETRINFERNI,¹⁰
G. ALTAVILLA,^{2,3} R. BUONANNO,¹⁰ B. CHABOYER,¹¹ R. DA SILVA,^{2,3} M. DALL'ORA,⁶
S. DEGL'INNOCENTI,^{12,13} E. DI CARLO,¹⁰ I. FERRARO,² E. GREBEL,¹⁴ G. IANNICOLA,² L. INNO,^{15,6}
V. KOVTYUKH,^{16,17} A. KUNDER,¹⁸ B. LEMASLE,¹⁴ M. MARENGO,⁸ S. MARINONI,^{2,3}
P. M. MARRESE,^{2,3} C. E. MARTÍNEZ-VÁZQUEZ,¹⁹ N. MATSUNAGA,²⁰ M. MONELLI,²¹ J. NEELEY,²²
M. NONINO,²³ P.G. PRADA MORONI,^{12,13} Z. PRUDIL,¹⁴ P. B. STETSON,²⁴ F. THÉVENIN,²⁵
E. TOGNETTI,¹³ E. VALENTI,²⁶ AND A. R. WALKER¹⁹

¹*Dipartimento di Fisica, Università di Roma Tor Vergata, via della Ricerca Scientifica 1, 00133 Roma, Italy*

²*INAF-Osservatorio Astronomico di Roma, via Frascati 33, 00078 Monte Porzio Catone, Italy*

³*Space Science Data Center, via del Politecnico snc, 00133 Roma, Italy*

⁴*Departamento de Astronomia, Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves 6500, Porto Alegre 91501-970, Brazil*

⁵*Department of Astronomy and McDonald Observatory, The University of Texas, Austin, TX 78712, USA*

⁶*INAF-Osservatorio Astronomico di Capodimonte, Salita Moiariello 16, 80131 Napoli, Italy*

⁷*The Observatories of the Carnegie Institution for Science, 813 Santa Barbara St., Pasadena, CA 91101, USA*

⁸*Department of Physics and Astronomy, Iowa State University, Ames, IA 50011, USA*

⁹*Department of Physics and Astronomy, Dartmouth College, Hanover, NBalmerH 03755, USA*

¹⁰*INAF-Osservatorio Astronomico d'Abruzzo, Via Mentore Maggini snc, Loc. Collurania, 64100 Teramo, Italy*

¹¹*Department of Physics and Astronomy, Dartmouth College, Hanover, NH 03755, USA*

¹²*INFN, Sezione di Pisa, Largo Pontecorvo 3, 56127, Pisa, Italy*

¹³*Dipartimento di Fisica "Enrico Fermi", Università di Pisa, Largo Pontecorvo 3, 56127, Pisa, Italy*

¹⁴*Astronomisches Rechen-Institut, Zentrum für Astronomie der Universität Heidelberg, Mönchhofstr. 12-14, D-69120 Heidelberg, Germany*

¹⁵*Università Parthenope di Napoli, Science and Technology Department, CDN IC4, 80143 Naples, Italy*

¹⁶*Astronomical Observatory, Odessa National University, Shevchenko Park, UA-65014 Odessa, Ukraine*

¹⁷*Newton Institute of Chile, Odessa branch, Shevchenko Park, UA-65014 Odessa, Ukraine*

¹⁸*Saint Martin's University, 5000 Abbey Way SE, Lacey, WA 98503, USA*

¹⁹*Cerro Tololo Inter-American Observatory, NSF's National Optical-Infrared Astronomy Research Laboratory, Casilla 603, La Serena, Chile*

²⁰*Department of Astronomy, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan*

²¹*Instituto de Astrofísica de Canarias, Calle Via Lactea s/n, E38205 La Laguna, Tenerife, Spain*

²²*Department of Physics, Florida Atlantic University, 777 Glades Rd, Boca Raton, FL 33431 USA*

²³*INAF-Osservatorio Astronomico di Trieste, Via G. B. Tiepolo 11, 34143 Trieste, Italy*

²⁴*Herzberg Astronomy and Astrophysics, National Research Council, 5071 West Saanich Road, Victoria, British Columbia V9E 2E7, Canada*

²⁵*Université de Nice Sophia-antipolis, CNRS, Observatoire de la Côte d'Azur, Laboratoire Lagrange, BP 4229, F-06304 Nice, France*

²⁶*European Southern Observatory, Karl-Schwarzschild-Str. 2, 85748 Garching bei Munchen, Germany*

ABSTRACT

We collected over 6000 high-resolution spectra of four dozen field RR Lyrae (RRL) variables pulsating either in the fundamental (39 RRab) or in the first overtone (9 RRC)

mode. We measured radial velocities (RVs) of four strong metallic and four Balmer lines along the entire pulsational cycle and derived RV amplitudes with accuracies better than $1\text{--}2\text{ km s}^{-1}$. The new amplitudes were combined with literature data for 23 RRab and 3 RRC stars (total sample 74 RRLs) which allowed us to investigate the variation of the Bailey diagram (photometric amplitude versus period) when moving from optical to mid-infrared bands and to re-cast the Bailey diagram in terms of RV amplitudes. We found that RV amplitudes for RRab are minimally affected by nonlinear phenomena (shocks) and multi-periodicity (Blazhko effect). The RV slope ($\log P\text{--}A(V_r)$) when compared with the visual slope ($\log P\text{--}A(V)$) is shallower and the dispersion, at fixed period, decreases by a factor of two. We constructed homogeneous sets of Horizontal Branch evolutionary models and nonlinear, convective pulsation models of RRLs to constrain the impact of evolutionary effects on their pulsation properties. Evolution causes, on the Bailey diagram based on RV amplitudes, a modest variation in pulsation period and a large dispersion in amplitude. The broad dispersion in period of the Bailey diagram is mainly caused by variation in RRL intrinsic parameters (stellar mass, chemical composition). Empirical evidence indicates that RV amplitudes are an optimal diagnostic for tracing the mean effective temperature across the RRab instability strip.

Keywords: stars: horizontal-branch — stars: oscillations — stars: variables: RR Lyrae — techniques: radial velocities

1. INTRODUCTION

The RR Lyrae (RRLs) are a very popular group of radial variables, because they can be easily identified thanks to the coupling between the pulsation periods ($0.25 \lesssim P \lesssim 1.0$ days) and the shapes of their light curves (Da Costa et al. 2010; Monelli et al. 2017).

RRLs pulsate in three different flavors: fundamental (RRab), first overtone (RRc) and mixed mode (pulsating simultaneously in the fundamental and in the first overtone). The Bailey diagram (Bailey et al. 1919)—luminosity amplitude versus logarithmic period—is a fundamental plane to identify the RRL pulsation mode. In this plane RRc variables cover the short period, low amplitude domain, and the shape of their amplitude-period relationship displays either a “bell” or a “hairpin” shape (Bono et al. 2011; Kunder et al. 2013; Fiorentino et al. 2017). The RRab variables cover the long-period, large amplitude area and display a well defined de-

crease in luminosity amplitude as period increases.

The Bailey diagram has also been the crossroad of a long-standing astrophysical problem, the so-called Oosterhoff dichotomy, identified over 80 years ago (Oosterhoff 1939). RRLs in Globular Clusters (GCs) can be split according to their mean periods in two groups: the Oosterhoff type I (OoI), with $\langle P_{ab} \rangle \sim 0.56$, $\langle P_c \rangle \sim 0.31$ days, and the Oosterhoff type II (OoII), with $\langle P_{ab} \rangle \sim 0.66$, $\langle P_c \rangle \sim 0.36$ days. Subsequent investigations identified a similar dichotomic distribution for the RRc, albeit with a difference of only ~ 0.05 days. Spectroscopic studies demonstrated that OoI GCs are more metal-rich and have a smaller ratio of RRc to total number of RRL than OoII GCs. More recently, Fabrizio et al. (2019) found a linear correlation between mean period and metallicity; thus, suggesting that this dichotomy is a natural consequence of the lack of Galactic, metal-intermediate GCs rich in RRLs.

RRL luminosity amplitudes steadily decrease when shifting to longer wavelengths. However, the amplitude attains an almost constant value at wavelengths longer than the K -band. This behavior is caused by the fact that luminosity amplitudes in the optical bands are mainly driven by temperature variations, while in the near-infrared (NIR)/mid-infrared (MIR), they are mainly driven by radius variations (Madore et al. 2013; Neeley et al. 2017).

The Bailey diagram relies on two observables that are independent of distance and reddening. The RRL distribution in the Bailey diagram depends on their intrinsic parameters (stellar mass, luminosity, effective temperature) and chemical composition, but it is also affected by degeneracies caused by three different phenomena. *i)*—Secondary modulations—Empirical evidence suggests that a fraction of RRL are affected by a secondary modulation (Blazhko effect, Blažko 1907) with a time scale ranging from tens of days to a few years. This fraction in large samples (in the Galactic Bulge) ranges from $\sim 6\%$ for RRc to $\sim 40\%$ for RRab (Netzel et al. 2018; Prudil & Skarka 2017). This means that the position of an RRL in the Bailey diagram can change by several tenths of a magnitude along the Blazhko cycle. The Blazhko effect partially explains why the RRL display, at fixed pulsation period and chemical composition, a well defined intrinsic spread in luminosity amplitude. *ii)*—Evolutionary phase—RRLs are low-mass helium burning stars and during this evolutionary phase they evolve off the Zero-Age-Horizontal-Branch (ZAHB). In their evolution they experience changes both in effective temperature and in surface gravity. These changes cause a variation both in pulsation period and in luminosity amplitude. We still lack empirical evidence concerning the variation of the position in the Bailey diagram directly caused by evolutionary effects. *iii)*—Nonlinear phenomena—Theory and observations indicate

that RRLs experience violent shocks approaching maximum compression (Preston et al. 2019). Shocks are strongly affected by the variation of the efficiency of convective motions (Bono & Stellingwerf 1994). This is the reason why RRab amplitudes display a steady decrease close to the red edge of the instability strip (IS, Bono et al. 1997; Braga et al. 2016).

The main aim of this investigation is to study the physical mechanisms affecting the distribution of RRLs in the Bailey diagram.

2. SPECTROSCOPIC AND PHOTOMETRIC DATASETS

The largest and most homogeneous subsample of high resolution (HR) spectra was collected with the Las Campanas Observatory du Pont echelle spectrograph. It includes 6,206 HR ($R \equiv \lambda/\delta\lambda = 35,000$) spectra of 192 RRLs. Their typical SNR, in the optical regime ($\lambda = 5,100 \text{ \AA}$), is ~ 40 .

These spectra were augmented with 272 HR ($R = 34,540\text{--}107,200$, $\text{SNR} \sim 20$; 68 RRLs) UVES@VLT spectra; 41 medium-resolution ($R = 4,300\text{--}18,000$, $\text{SNR} \sim 44$; 18 RRLs) X-Shooter@VLT spectra; 166 HR ($R = 115,000$, $\text{SNR} \sim 10$; 5 RRLs) HARPS@3.6m spectra and 55 HR ($R = 48,000$, $\text{SNR} \sim 4$; 3 RRLs) FEROS@2.2m (ESO, Chile) spectra.

Additionally we included HR spectra from SES@STELLA (15, $R = 55,000$, $\text{SNR} \sim 35$; 1 RRL), HRS@SALT (82, $R \sim 40,000$, $\text{SNR} \sim 50$; 70 RRLs), and HARPS-N@TNG (10, $R = 115,000$, $\text{SNR} \sim 40$; 4 RRLs).

In total we collected 6,572 spectra for 316 RRLs. Among them, 48 RRLs have a good/optimal coverage of the pulsation cycle (6,226 spectra). We label these variables as RV calibrators. These data were complemented with RV curves for 26 field and cluster RRLs available in the literature from Baade-Wesselink (BW) analyses (Storm et al. 1994; Bono et al. 2003). We ended up with a sample of 74 (62 RRab, 12 RRc) RV calibrators (see Table 1).

The periods from literature are usually precise enough to provide good phasing of the RV curves. However, we derived the pulsation period from RV measurements for 2 RRab and 1 RRC. We adopted Gaia G -band amplitudes (Gaia Collaboration et al. 2018; Marrese et al. 2019) transformed into the V band by using the equations provided by Evans et al. (2018). We also used V -band photometry from ASAS (Pojmanski 2002), ASAS-SN (Shappee et al. 2014; Jayasinghe et al. 2018) and from Catalina (Torrealba et al. 2015; Drake et al. 2013a,b).

The optical photometry was combined with MIR $W1$ -band photometry collected with NEOWISE (Mainzer et al. 2011). The MIR light curves have very good phase coverage (~ 120 phase points), and good photometric precision ($\sigma_{W1} \leq 0.05$ mag). Optical and MIR light curves were fitted with analytical functions (splines, PLOESS, Fourier parameters) following the same approach by Braga et al. (2018). The photometric amplitudes are based on the analytical fits.

3. RADIAL VELOCITY MEASUREMENTS

To measure RVs we chose three Fe I lines, from the multiplet 43, at $\lambda = 4045.81, 4063.59, 4071.74$ Å and the Sr II line at $\lambda = 4077.71$ Å. These metallic lines were strong enough to be easily identified in most spectra even at very low SNR or very high effective temperatures, enabling us to trace the whole pulsation cycle using individual spectra, i.e. without co-adding spectra. We also used four Balmer lines ($H_\alpha, H_\beta, H_\gamma, H_\delta, \lambda = 6562.80, 4861.36, 4340.46, 4101.74$ Å). The RVs were measured following the same approach employed by Fabrizio et al. (2011).

Figure 1 shows RV curves for the RRab, V Ind, and the RRC, MT Tel. The RV measurements plotted in this figure display the expected difference between metallic and Balmer lines (Preston 1959); the RV amplitudes for the latter ones are systematically larger (see also

Figure 9 of Chadid et al. (2017) for montages of RV and photometric phase plots). The difference is caused by the physical conditions at which the different lines form: smaller the optical depth the larger, the RV amplitude. The difference among H_α and metallic lines ranges from 50% for MT Tel to almost a factor of two for V Ind. In spite of the difference in amplitude, the secondary features (bumps, dips) appear to be largely co-phased.

We analytically fit the RV curves by using the PLOESS algorithm (Braga et al. 2018). The cyan lines in Figure 1 show these curves. Note that RVs for V Ind based on H_γ show a well defined “spike” around phases 0.70–0.76, near its point of minimum radius, as implied by the optical and MIR light curves of this star in the top panels of Figure 2.

This secondary spike feature was predicted by pulsation models (see Figs. 3, 16 in Bono & Stellingwerf 1994 and Fig. 12 in Marconi et al. 2015) suggesting that it is associated with the formation and propagation of shocks. Note that the duration of the spike is ~ 25 minutes and we succeeded in tracing it only because the du Pont spectra were collected with exposure times of 400 ± 200 seconds. Moreover, exposure times for V Ind are $\approx 30\%$ shorter than the mean, since it has a brighter mean visual magnitude ($V = 9.92 \pm 0.05$ mag). A similar feature is also present among other Balmer (mainly H_β, H_δ) and metallic (mainly Sr) lines, but it is less pronounced.

Note that the absolute minimum in RV for V Ind takes place, due to the phase lag (Castor 1971) at the same phases of the absolute maximum in the light curves. However, the absolute minimum in luminosity is not at the same phases of the absolute maximum in RV. Indeed, the latter takes place around the phases in which the optical light curves show a well defined bump. The difference appears to be caused by the interplay between the shock prop-




Figure 1. Panels a) and b): Radial velocity curves versus phase for the RRab V Ind based on metallic lines Fe I (λ : 4045.81, 4063.59, 4071.74 Å) and Sr II (λ : 4077.71 Å), and on Balmer (λ : 6562.80, 4861.36, 4340.46, 4101.74 Å) lines. Measurements based on different spectrographs are marked with different colors. The cyan lines show the analytical fits. The RV amplitude ($A(v_r)$ km s^{-1}) and the pulsation period (days) are also labeled. Panels c) and d): Same as the top, but for the RRC MT Tel.




Figure 2. Top: Optical (V , left) and MIR ($W1$, right) light curves for the RRab V Ind. The optical data were provided by ASAS-SN (Shappee et al. 2014; Jayasinghe et al. 2018), while the MIR ones by NEOWISE (Mainzer et al. 2011). The thin red lines show the analytical fits. Bottom: Same as the top, but for the RRC MT Tel.

agation and the increased efficiency of the convective transport in approaching the phases of minimum radius.

The RV curve of MT Tel is smoother than that of V Ind over the entire pulsation cycle. For this variable we have extensive measurements from both du Pont and HARPS spectrographs. Inspection of panels c) and d) of Figure 1 reveals that the RV amplitude from the HARPS dataset is $\sim 2\text{--}3 \text{ km s}^{-1}$ larger than that from du Pont. MT Tel is the only variable for which we found such a difference. We performed a number of tests that verified the reality of this offset. It cannot be caused by phasing problems, since the combined optical and NIR photometry allowed us very accurate phasing of the two datasets. The analytical fit displayed in Figure 1 is mainly following the du Pont measurements, because they have a more uniform sampling over the pulsation cycle. We suggest that the difference is intrinsic and possibly caused by the presence of a secondary

modulation (Blazhko effect), even though the light curves do not show yet a clear modulation at fixed phase. Note that HARPS spectra were collected in a single night seven years before the du Pont spectra (over five consecutive nights) and the periodogram lacks of relevant secondary peaks. This finding would suggest that accurate RV measurements might have a stronger sensitivity in detecting secondary modulations and in constraining the tomography of variable star atmospheres. In the next paper we will explore more aspects of these interesting issues.

4. RESULTS

We have investigated on a more quantitative basis the variation of the luminosity amplitudes when moving from optical to MIR, and to RV amplitudes and when moving from weak to strong lines. In Figure 3 we show the Bailey diagram for the RRLs for which we estimated both luminosity and velocity amplitudes. Panels a) and b) display quite clearly that the luminosity amplitudes in the $W1$ -band are roughly a factor of three smaller than in V -band. This trend is expected because the luminosity variation in the former band is mainly driven by temperature variations, while in the latter it is mainly driven by radius variations. Amplitude dispersion at fixed pulsation period also is smaller in the infrared. It is of the order of 20% for the V band ($\sigma_V \sim 0.19 \text{ mag}$), while for the $W1$ band it decreases to 10% ($\sigma_{W1} \sim 0.03 \text{ mag}$). This outcome applies to both RRab and RRC variables and suggests that the dispersion is driven by phenomena affecting the temperature variation along the pulsation cycle. Moreover, Blazhko RRLs (black crosses) in our sample display a variation in the visual band of $\sim 0.4 \text{ mag}$ and of $\approx 0.1 \text{ mag}$ in the $W1$ band (Jurcsik et al. 2018).

Note that the current sample is quite representative of the RRL pulsation properties indeed the analytical relations for OoI, Oosterhoff intermediate and OoII RRLs (Fabrizio et al. 2019) provide a plausible fit of optical amplitudes.




Figure 3. Bailey diagram, amplitude versus logarithmic period. Panels a) and b) display the luminosity amplitude in the visual and in the W1 band. The black lines display observed relations for OoI (solid), Oo-intermediate (dotted) and OoII (dashed) RRLs provided by Fabrizio et al. (2019). Panels c) and d) show RV amplitudes based either on metallic or on Balmer lines. The red symbols plotted in panel c) display BW RRLs. Hydrogen RV amplitudes are plotted with different colors (panel d). Blazhko RRLs are marked with crosses. The error bars plotted in the bottom left corner display uncertainties in amplitudes.

Figure 3 panels c) and d) display, for the first time, a Bailey diagram based on RV amplitudes. In panel c) together with the current velocity amplitudes we also included the velocity amplitudes for field and cluster RRLs (red symbols) for which a BW solution is available in the literature. Interestingly, the relative dispersion in RV amplitudes is smaller than for visual amplitudes. Indeed, the RV amplitudes are purely tracing the radius variation along the pulsation cycle. The result is quite clear for RRab for which we find that the dispersion is of the order of 10% ($\sigma_{Vr} \sim 6 \text{ km s}^{-1}$). Moreover, Blazhko RRLs show a variation in velocity amplitude significantly smaller ($\sim 20\%$) than in visual amplitude ($\sim 40\%$). Unfortunately, we cannot reach firm conclusions concerning RRLc, since the sample is too small.

Panel d) of Figure 3 also shows an interesting trend. The velocity amplitudes for RRab variables based on Balmer lines attain an almost constant value when moving from the hot (short-period) to the cool (long-period) edge of the fundamental IS. The mean RV amplitudes decrease from H_α to H_δ , but their values are constant over the entire period range. The dispersion is also modest and ranges from 5 to 11 km s^{-1} .

The RV amplitudes plotted in Figure 3 panels c) and d) clearly show that the dispersion, at fixed period, is significantly larger than intrinsic errors; thus, suggesting that other physical mechanisms can contribute to the observed dispersion. To investigate this intrinsic property quantitatively, we constructed four HB models covering a broad range in chemical compositions ($Z=0.0001, 0.001, 0.008, 0.02$; $Y=0.245, 0.245, 0.256, 0.27$). The stellar masses ($M=0.75, 0.65, 0.57, 0.55 M_\odot$) were selected to be centrally located inside the IS. The reason for this broad range in metallicity was driven by the relevant change in the evolutionary path inside the IS when moving from metal-poor to metal-




Figure 4. Top: Hertzsprung-Russell Diagram for HB models (Pietrinferni et al. 2006) constructed by assuming different chemical compositions and stellar masses (see labeled values). The symbols mark the position of the pulsation models we constructed by using evolutionary prescriptions (chemical composition, stellar mass, luminosity, effective temperature). The light blue numbers mark the first three and the last three models in the sequence of pulsation models. The symbols are large and bluish close to the ZAHB and become smaller and greenish-reddish along the off-ZAHB evolution. Middle: Bailey diagram for visual amplitudes. The symbols are the same as in panels a) and they are connected with solid lines following the same color coding. Bottom: Bailey diagram for RV amplitudes based on metallic lines. The red symbols display BW RRLs.

rich stellar structures. We also constructed, following [Marconi et al. 2015](#), a set of non-linear, convective hydrodynamical models of RRL stars along the HB evolutionary models to investigate the variation of the pulsation properties in their off-ZAHB evolution.

The top panels of Figure 4 show small Hertzsprung-Russell Diagram regions of the HB evolutionary models (black lines) together with the variable models that are pulsationally stable. The color and the size of the symbols change along the off-ZAHB evolution.

As expected, the evolution across the IS changes significantly when moving from the metal-poor to the metal-rich regime. Indeed, for the most metal-poor chemical composition the off-ZAHB evolution is redward, i.e. HB models become systematically brighter and cooler as a function of time. The evolutionary path becomes more complex at metal-intermediate regime, because the evolutionary path undergoes, in sequence, a redward, a blueward and eventually a redward evolution approaching its Asymptotic Giant Branch track. This means that we are dealing with triple points, i.e. evolutionary phases covering the same range in effective temperature, but slightly different luminosities ($\Delta \text{Log } L/L_{\odot} \sim 0.07$, $Z=0.0001$; ~ 0.09 , $Z=0.001$; ~ 0.11 , $Z=0.008$; ~ 0.05 , $Z=0.02$). This means a steady decrease in surface gravity, and in turn, a steady increase in pulsation period.

The change in the physical structure also implies variation in the pulsation properties. The middle and the bottom panels of Figure 4 display the comparison between observed and predicted amplitudes. This is the first time that evolutionary effects are investigated in detail along the off-ZAHB evolution. A glance at the data plotted in this figure shows that the models, when moving from metal-poor to metal-rich structures, attain, as expected, shorter periods and larger amplitudes. Interestingly enough,

the most metal-poor models display a steady decrease in amplitude when moving from hotter to cooler effective temperatures. Note that the smaller the symbols the faster is the evolutionary phase. This means that the lack of long-period small amplitude RRLs is a mix between the probability to find them and the sample size.

The comparison between theory and observations becomes even more interesting in the metal-intermediate and in the metal-rich regime, because pulsation models associated with blueward and redward evolutionary phases attain very similar values in the Bailey diagram. This means that RRLs in their off-ZAHB evolution move back and forth in the Bailey diagram covering a modest range in period, but a large range in amplitude. This indicates that a significant fraction of the dispersion in the Bailey diagram is caused by the evolutionary status of RRLs. Theory and observations agree quite well over the entire metallicity range. Note that predicted amplitudes are affected by uncertainties in the mixing length parameter adopted to treat time-dependent convective transport in hydrodynamical models of radial variables ([Bono & Stellingwerf 1994](#)). However, we are using predicted amplitudes in the relative sense, i.e. we are interested in the variation of the pulsation amplitudes when moving from the blue to the red edge of the IS and not in their absolute value.

The comparison in the Bailey diagram for RV amplitudes shows similar results. Note that to compare theory and observations we adopted a projection factor, the ratio between pulsation velocity and RV, $p=1.33$ ([Jones et al. 1992](#)). The comparison in this plane is even more compelling because RV amplitudes are marginally affected by nonlinear phenomena and by secondary modulations. Interestingly, predicted and observed velocity amplitudes display a quadratic variation when moving from

the blue to the red edge of the IS, while the luminosity amplitudes show a more linear trend.

5. CONCLUSIONS

We investigated the properties of the Bailey diagram based on luminosity (optical/MIR) and on RV amplitudes. We collected homogeneous RV measurements for four dozen of field RRLs. The current measurements double the sample of RRLs with a good coverage of the entire pulsation cycle. We investigated the dependence of luminosity amplitudes on nonlinear phenomena, secondary modulations and evolutionary effects. The impact of these effects is mitigated when moving from the visual to the MIR bands, because the variations along the pulsation cycle are mainly driven by radius than by temperature changes. This trend becomes even more compelling in dealing with RV amplitudes, indeed, we found that the dispersion, at fixed period, is systematically smaller when compared with luminosity amplitudes. Moreover, Blazhko RRLs display smaller variations in velocity amplitudes than in luminosity amplitudes.

We constructed four different HB evolutionary models covering a broad range in metal contents ($Z=0.0001, 0.001, 0.008, 0.02$; $Y=0.245, 0.245, 0.256, 0.27$) and a sizable sample

of nonlinear, convective hydrodynamical models of RRLs to constrain the impact of the off-ZAHB evolution on their pulsation properties. We found that evolutionary effects take account for the vertical structure (dispersion in amplitude) of the Bailey diagram based on RV amplitudes. Thus, we suggest that the dispersion in period is mainly caused by variation in the intrinsic parameters of RRLs (stellar mass, chemical composition).

The current findings are suggesting that RV amplitudes are not only independent of distance and reddening, but also less prone to amplitude variations caused by nonlinear phenomena and secondary modulations. This means that RV amplitudes are an optimal diagnostic for the variation of the mean effective temperature across the IS of RRab.

ACKNOWLEDGEMENTS

We thank an anonymous referee for his/her constructive suggestions on an early draft of this paper. We acknowledge financial support from INAF/ASI 2014-049-R.0 (SSDC), US NSF under Grants AST-1714534 (MM, JPM) and AST-1616040 (CS). Some of the observations were obtained with SALT. This publication uses NEOWISE data (JPL/Caltech, University of Arizona, NASA).

Table 1. Pulsation properties for the current RRLs.

Gaia DR2 ID	Classical ID	type ^a	Period days	A(V) mag	A(W1) mag	A(Vr) _{Fe} km s ⁻¹	flag ^b	A(Vr) _{Hα} km s ⁻¹	A(Vr) _{Hβ} km s ⁻¹	A(Vr) _{Hγ} km s ⁻¹	A(Vr) _{Hδ} km s ⁻¹
15489408711727488	X Ari	0	0.65117537	0.96	0.32	58.2±2.4	0	108.2±4.4	85.1±3.3	76.4±3.0	66.5±2.9
234108363683247616	AR Per	0	0.42556048	0.88	0.27	65.7±5.1	1	...	...	...	...
630421935431871232	RR Leo	0	0.45240000	1.25	0.32	63.3±6.8	1	...	...	...	...
886793515494085248	RR Gem	0	0.39730000	1.17	0.28	70.0±11.3	1	...	...	...	...
1009665142487836032	TT Lyn	0	0.59744301	0.67	0.25	50.5±3.1	1	...	...	...	...
1058066262817534336	SU Dra	0	0.66041178	0.98	0.28	57.6±3.5	1	...	...	...	...
119151003353849472	AN Ser	0	0.52207295	0.99	0.29	60.2±3.8	0	94.5±6.4	73.9±5.2	70.5±4.9	57.9±3.8
1286188056265485952	RS Boo	0	0.37736549	1.04	0.25	67.1±3.2	1	...	...	...	...
1360405567883886720	CJ* NGC 6341 SAW V1	0	0.70279828	0.87	0.26	61.2±4.9	1	...	...	...	...
1360408905076366208	CSS J171712.0+431225	0	0.63746100	0.76	0.26	61.4±3.9	1	...	...	...	...
1492230556717187456	TV Boo	1	0.31256000	0.63	0.12	35.0±5.6	1	...	...	...	...
1683444631037761024	SW Dra	0	0.56967009	0.94	0.28	62.6±4.5	1	...	...	...	...
1760981190300823808	DX Del	0	0.47261773	0.70	0.24	54.0±3.2	0	94.1±6.1	71.3±4.1	66.3±4.1	58.3±3.6
1793460115244988800	AV Peg	0	0.39038090	1.01	0.28	63.4±2.8	0	92.6±4.4	74.4±3.4	66.2±3.3	59.8±3.0
2381771781829913984	DN Aqr	0	0.63376712	0.74	0.24	54.6±2.1	0	104.3±4.0	79.6±3.4	79.6±3.5	63.8±3.0
2414817603803476864	UU Cet	0	0.60610163	0.60	0.23	53.0±3.8	1	...	...	...	...
2558296724402139392	RR Cet	0	0.55302505	0.90	0.28	60.1±3.1	0	106.2±5.5	84.1±4.3	81.2±4.3	68.7±3.6
2689556491246048896	SW Aqr	0	0.45930070	1.24	0.31	61.9±1.8	0	105.1±3.3	82.7±2.6	79.5±2.6	64.8±2.3
2720896455287475584	DH Peg	1	0.25551624	0.50	0.10	25.4±1.8	1	...	...	...	...
2857456207478683776	SW And	0	0.44230000	0.92	0.27	62.5±4.6	1	...	...	...	...
2981136563930816128	RX Eri	0	0.58725159	0.87	0.28	60.3±4.5	1	...	...	...	...
3479598373678136832	DT Hya	0	0.56798140	0.95	0.27	61.8±3.1	0	109.9±5.7	83.7±4.3	78.1±4.2	70.9±3.7
3535368455297644928	ASAS J110522-2641.0	1	0.29445920	0.33	...	14.3±1.7	0	16.4±2.2	...	...	...
3546458301374134528	W Crt	0	0.41201190	1.36	0.32	66.1±2.5	0	101.9±4.4	83.9±3.7	77.1±3.3	66.6±2.9
3604450388616968576	AM Vir	0	0.61510000	0.66	0.25	59.0±2.5	0	98.1±4.2	81.1±3.6	77.3±3.5	63.2±2.8
3626569264033312896	AS Vir	0	0.55341000	0.63	0.22	59.1±2.0	0	85.2±2.9	77.2±2.7	74.6±2.8	65.7±2.5
3652665558338018048	ST Vir	0	0.41080754	1.22	0.33	60.3±3.4	0	97.9±5.8	72.2±4.4	78.5±4.8	61.2±4.0
3698725337375614464	UU Vir	0	0.47560267	1.10	0.30	67.8±3.1	0	114.4±7.2	91.7±4.6	83.6±4.3	74.9±3.6
3797319369672686592	SS Leo	0	0.62632619	1.09	0.30	62.5±2.0	0	97.2±4.0	83.8±3.0	80.9±3.1	71.4±2.8
3846786226007324160	T Sex	1	0.32468493	0.42	0.08	27.0±2.3	1	...	...	...	...
3915998558830693888	ST Leo	0	0.47797595	1.23	0.31	64.0±4.3	0	112.4±7.6	82.9±5.4	85.0±6.1	65.3±4.9

Table 1 continued

Table 1 (*continued*)

Gaia DR2 ID	Classical ID	type ^a	Period days	A(V) mag	A(W1) mag	A(Vr) _{Fe} km s ⁻¹	flag ^b	A(Vr) _{Hα} km s ⁻¹	A(Vr) _{Hβ} km s ⁻¹	A(Vr) _{Hγ} km s ⁻¹	A(Vr) _{Hδ} km s ⁻¹
4022618712476736896	TU UMa	0	0.55690000	0.96	0.28	60.6 $\pm$ 4.3	1	...	...	...	...
4055098870077726976	V0494 Sco	0*	0.42727100	0.87	0.20	44.5 $\pm$ 1.7	0	68.6 $\pm$ 3.1	53.5 $\pm$ 2.5	53.7 $\pm$ 2.4	45.7 $\pm$ 2.4
4352084489819078784	V0445 Oph	0	0.39702600	0.77	0.26	57.4 $\pm$ 4.6	0	91.1 $\pm$ 7.1	68.6 $\pm$ 5.1	61.6 $\pm$ 5.1	58.8 $\pm$ 4.8
4417888542753226112	VY Ser	0	0.71410000	0.65	0.25	53.3 $\pm$ 4.3	0	90.6 $\pm$ 6.3	76.3 $\pm$ 11.2	74.2 $\pm$ 6.5	69.2 $\pm$ 6.0
4421571803630954752	CSS J151841.9+020232	0	0.54625080	0.65	0.26	51.8 $\pm$ 3.8	1	...	...	...	...
4421571803630958848	CJ* NGC 5904 SAW V28	0	0.54383658	0.71	...	60.9 $\pm$ 4.7	1	...	...	...	...
4433070255716036864	ASAS J162158+0244.5	1	0.32362580	0.41	0.09	20.2 $\pm$ 2.3	0	29.2 $\pm$ 3.2	29.0 $\pm$ 3.8	...	...
4454183799545435008	AT Ser	0	0.74655408	0.89	0.26	56.8 $\pm$ 4.1	0	99.8 $\pm$ 7.8	77.2 $\pm$ 5.6	77.0 $\pm$ 5.7	70.0 $\pm$ 5.4
4467433017735910912	VX Scl	0	0.45518030	1.28	0.33	60.6 $\pm$ 3.9	0	105.9 $\pm$ 7.1	81.0 $\pm$ 5.3	74.7 $\pm$ 5.0	64.9 $\pm$ 4.5
4596935593202765184	TW Her	0	0.39959577	1.33	0.33	68.3 $\pm$ 4.7	1	...	...	...	...
4689637956899105792	CJ* NGC 104 SAW V9	0	0.73720048	1.06	...	57.0 $\pm$ 5.3	1	...	...	...	...
4709830423483623808	W Tuc	0	0.64224028	1.14	0.31	63.7 $\pm$ 2.9	0	114.0 $\pm$ 5.8	79.9 $\pm$ 4.1	80.0 $\pm$ 4.0	66.2 $\pm$ 3.2
4860671839583430912	SX For	0	0.60534530	0.66	...	51.9 $\pm$ 2.9	0	93.5 $\pm$ 5.9	70.6 $\pm$ 4.3	66.6 $\pm$ 3.8	61.4 $\pm$ 3.7
4947090013255935616	CS Eri	1	0.31133020	0.52	...	28.9 $\pm$ 4.0	0	48.6 $\pm$ 3.9	34.7 $\pm$ 3.0	33.8 $\pm$ 2.9	28.2 $\pm$ 2.6
5022411786734718208	SV Scl	1	0.37735860	0.51	...	26.1 $\pm$ 2.1	0	42.2 $\pm$ 3.2	30.6 $\pm$ 2.6	29.4 $\pm$ 2.6	24.0 $\pm$ 2.6
5151789464548893184	RV Cet	0*	0.61802300	0.57	...	49.8 $\pm$ 2.3	0	89.3 $\pm$ 4.3	70.1 $\pm$ 3.3	65.6 $\pm$ 3.0	57.9 $\pm$ 2.7
5412243359495900928	CD Vel	0	0.57350788	0.86	0.27	52.5 $\pm$ 1.8	0	96.6 $\pm$ 4.1	74.0 $\pm$ 2.0	69.8 $\pm$ 2.8	60.7 $\pm$ 2.5
5461994297841116160	WY Ant	0	0.57434364	0.92	0.27	60.6 $\pm$ 2.9	0	106.5 $\pm$ 5.6	83.3 $\pm$ 4.4	77.1 $\pm$ 4.2	66.5 $\pm$ 3.7
5510293236607430656	HH Pup	0	0.39081190	1.22	0.32	67.9 $\pm$ 2.6	0	109.3 $\pm$ 4.3	81.4 $\pm$ 3.3	78.1 $\pm$ 3.4	68.9 $\pm$ 2.9
5707380936404638336	BB Pup	0	0.48055043	0.88	0.29	64.8 $\pm$ 4.8	1	...	...	...	...
5768557209320424320	UV Oct	0*	0.54257500	0.80	0.27	57.1 $\pm$ 1.1	0	90.3 $\pm$ 2.0	73.5 $\pm$ 1.6	73.0 $\pm$ 1.8	62.3 $\pm$ 1.7
5769986338215537280	RV Oct	0	0.57117800	1.05	0.30	65.1 $\pm$ 2.3	0	103.9 $\pm$ 3.6	82.7 $\pm$ 2.9	81.9 $\pm$ 3.0	70.1 $\pm$ 2.7
5773390391856998656	XZ Aps	0	0.58726739	1.15	0.29	63.0 $\pm$ 1.8	0	102.0 $\pm$ 3.2	81.7 $\pm$ 2.5	79.3 $\pm$ 2.7	65.1 $\pm$ 2.5
5806921716937210496	BS Aps	0	0.58256590	0.76	0.24	51.1 $\pm$ 1.7	0	85.2 $\pm$ 3.2	67.1 $\pm$ 2.5	62.0 $\pm$ 2.3	56.4 $\pm$ 2.3
5947570591534602240	S Ara	0*	0.45184650	0.96	0.29	60.6 $\pm$ 2.0	0	...	82.1 $\pm$ 3.0	77.9 $\pm$ 2.8	66.7 $\pm$ 2.5
6045464303643755008	CJ* NGC 6121 SAW V2	0	0.53568150	0.92	...	54.1 $\pm$ 6.5	1	...	...	...	...
6045485228725626752	CJ* NGC 6121 SAW V32	0	0.57909353	0.75	0.25	61.0 $\pm$ 6.8	1	...	...	...	...
6045491928874778624	CJ* NGC 6121 SAW V33	0	0.61484317	0.85	0.31	55.8 $\pm$ 8.3	1	...	...	...	...
6045502305516138368	CJ* NGC 6121 SAW V15	0	0.44379722	1.20	0.30	43.7 $\pm$ 6.0	1	...	...	...	...
6473212809741024256	ASAS J200431-5352.3	1	0.30022700	0.28	...	14.2 $\pm$ 1.9	0	16.1 $\pm$ 1.9	13.2 $\pm$ 2.2	15.4 $\pm$ 2.2	14.1 $\pm$ 2.3
6483680332235888896	V Ind	0	0.47959915	1.12	0.29	56.2 $\pm$ 1.2	0	101.7 $\pm$ 2.6	77.9 $\pm$ 2.0	73.8 $\pm$ 2.0	61.9 $\pm$ 1.7
6526559499016401408	RV Phe	0	0.59641862	0.67	0.25	54.6 $\pm$ 4.8	1	...	...	...	...
6541769554459131648	BO Gru	1	0.28272000	0.18	...	13.7 $\pm$ 1.5	0	16.0 $\pm$ 2.3	12.2 $\pm$ 2.3	15.6 $\pm$ 2.6	12.7 $\pm$ 2.6
6570585628216929408	TY Gru	0	0.57006515	0.91	0.27	59.2 $\pm$ 2.1	0	112.8 $\pm$ 4.7	82.5 $\pm$ 3.3	76.2 $\pm$ 3.2	64.6 $\pm$ 3.2

Table 1 *continued*

Table 1 (*continued*)

Gaia DR2 ID	Classical ID	type ^a	Period days	A(V) mag	A(W1) mag	A(Vr) _{Fe} km s ⁻¹	flag ^b	A(Vr) _{Hα} km s ⁻¹	A(Vr) _{Hβ} km s ⁻¹	A(Vr) _{Hγ} km s ⁻¹	A(Vr) _{Hδ} km s ⁻¹
6625215584995450624	AE PsA	0	0.54674000	1.13	0.32	65.1 $\pm$ 6.5	0	102.3 $\pm$ 8.7	76.6 $\pm$ 7.0	79.1 $\pm$ 7.6	64.5 $\pm$ 6.2
6662886605712648832	MT Tel	1	0.31689945	0.55	0.13	22.4 $\pm$ 1.4	0	35.6 $\pm$ 1.9	27.0 $\pm$ 1.6	26.0 $\pm$ 2.3	21.7 $\pm$ 1.9
6680420204104678272	V1645 Sgr	0	0.55292000	0.76	0.24	50.9 $\pm$ 1.9	0	74.0 $\pm$ 2.8	66.2 $\pm$ 2.5	65.8 $\pm$ 2.7	56.5 $\pm$ 2.4
6701821205809488384	ASAS J181215-5206.9	0	0.83753980	0.52	0.20	42.6 $\pm$ 5.6	0	...	72.0 $\pm$ 9.5	61.3 $\pm$ 8.2	54.2 $\pm$ 7.4
6771307454464848768	V0440 Sgr	0	0.47750000	0.77	0.30	62.7 $\pm$ 5.1	0	113.2 $\pm$ 8.9	84.0 $\pm$ 6.5	76.5 $\pm$ 6.4	67.1 $\pm$ 5.2
6787617919184986496	Z Mic	0	0.58692775	0.64	0.22	53.2 $\pm$ 1.8	0	92.1 $\pm$ 3.7	74.3 $\pm$ 3.0	69.3 $\pm$ 2.8	61.7 $\pm$ 2.6
6856027093125912064	ASAS J203145-2158.7	1	0.31071060	0.37	0.08	22.0 $\pm$ 2.0	0	29.7 $\pm$ 2.8	24.3 $\pm$ 2.6	21.1 $\pm$ 2.5	24.6 $\pm$ 4.2
6883653108749373568	RV Cap	0*	0.44774990	0.97	...	49.9 $\pm$ 2.0	0	90.8 $\pm$ 3.6	72.7 $\pm$ 3.1	62.4 $\pm$ 2.7	59.2 $\pm$ 2.7
6884361748289023488	YZ Cap	1	0.27345290	0.49	0.10	26.3 $\pm$ 2.1	0	35.8 $\pm$ 3.0	28.3 $\pm$ 2.8	29.1 $\pm$ 2.9	23.2 $\pm$ 2.6

^a Pulsation mode: 0=RRab; 1=RRc. Candidate Blazhko RRLs are marked with an asterisk.^b RV amplitudes: 0=current investigation; 1=BW sample.

REFERENCES

- Bailey, S. I., Leland, E. F., Woods, I. E., & Pickering, E. C. 1919, *Annals of Harvard College Observatory*, 78, 195
- Blažko, S. 1907, *Astronomische Nachrichten*, 175, 325
- Bono, G., Caputo, F., Castellani, V., & Marconi, M. 1997, *A&AS*, 121, 327
- Bono, G., Caputo, F., Castellani, V., et al. 2003, *MNRAS*, 344, 1097
- Bono, G., Dall’Ora, M., Caputo, F., et al. 2011, in *RR Lyrae Stars, Metal-Poor Stars, and the Galaxy*, ed. A. McWilliam, Vol. 5, 1
- Bono, G. & Stellingwerf, R. F. 1994, *ApJS*, 93, 233
- Braga, V. F., Stetson, P. B., Bono, G., et al. 2016, *AJ*, 152, 170
- Braga, V. F., Stetson, P. B., Bono, G., et al. 2018, *AJ*, 155, 137
- Castor, J. I. 1971, *ApJ*, 166, 109
- Chadid, M., Sneden, C., & Preston, G. W. 2017, *ApJ*, 835, 187
- Da Costa, G. S., Rejkuba, M., Jerjen, H., & Grebel, E. K. 2010, *ApJL*, 708, L121
- Drake, A. J., Catelan, M., Djorgovski, S. G., et al. 2013a, *ApJ*, 763, 32
- Drake, A. J., Catelan, M., Djorgovski, S. G., et al. 2013b, *ApJ*, 765, 154
- Evans, D. W., Riello, M., De Angeli, F., et al. 2018, *A&A*, 616, A4
- Fabrizio, M., Bono, G., Braga, V. F., et al. 2019, *ApJ*, 882, 169
- Fabrizio, M., Nonino, M., Bono, G., et al. 2011, *PASP*, 123, 384
- Fiorentino, G., Monelli, M., Stetson, P. B., et al. 2017, *A&A*, 599, A125
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *A&A*, 616, A1
- Jayasinghe, T., Kochanek, C. S., Stanek, K. Z., et al. 2018, *MNRAS*, 477, 3145
- Jones, R. V., Carney, B. W., Storm, J., & Latham, D. W. 1992, *ApJ*, 386, 646
- Jurcsik, J., Hajdu, G., Dékány, I., et al. 2018, *MNRAS*[1801.03436]
- Kunder, A., Stetson, P. B., Cassisi, S., et al. 2013, *AJ*, 146, 119
- Madore, B. F., Hoffman, D., Freedman, W. L., et al. 2013, *ApJ*, 776, 135
- Mainzer, A., Bauer, J., Grav, T., et al. 2011, *ApJ*, 731, 53
- Marconi, M., Coppola, G., Bono, G., et al. 2015, *ApJ*, 808, 50
- Marrese, P. M., Marinoni, S., Fabrizio, M., & Altavilla, G. 2019, *A&A*, 621, A144
- Monelli, M., Fiorentino, G., Bernard, E. J., et al. 2017, *ApJ*, 842, 60
- Neeley, J. R., Marengo, M., Bono, G., et al. 2017, *ApJ*, 841, 84
- Netzel, H., Smolec, R., Soszyński, I., & Udalski, A. 2018, *MNRAS*, 480, 1229
- Oosterhoff, P. T. 1939, *The Observatory*, 62, 104
- Pietrinferni, A., Cassisi, S., Salaris, M., & Castelli, F. 2006, *ApJ*, 642, 797
- Pojmanski, G. 2002, *AcA*, 52, 397
- Preston, G. W. 1959, *ApJ*, 130, 507
- Preston, G. W., Sneden, C., Chadid, M., Thompson, I. B., & Sheckman, S. A. 2019, *AJ*, 157, 153
- Prudil, Z. & Skarka, M. 2017, *MNRAS*, 466, 2602
- Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, *ApJ*, 788, 48
- Storm, J., Carney, B. W., & Latham, D. W. 1994, *A&A*, 290, 443
- Torrealba, G., Catelan, M., Drake, A. J., et al. 2015, *MNRAS*, 446, 2251