



Publication Year	2021
Acceptance in OA	2022-06-01T15:21:51Z
Title	Multiscale fractal dimension analysis of a reduced order model of coupled ocean-atmosphere dynamics
Authors	ALBERTI, TOMMASO, Donner, Reik V., Vannitsem, Stéphane
Publisher's version (DOI)	10.5194/esd-12-837-2021
Handle	http://hdl.handle.net/20.500.12386/32167
Journal	EARTH SYSTEM DYNAMICS
Volume	12



Multiscale fractal dimension analysis of a reduced order model of coupled ocean-atmosphere dynamics

Tommaso Alberti¹, Reik V. Donner^{2,3}, and Stéphane Vannitsem⁴

¹INAF-IAPS, via del Fosso del Cavaliere 100, 00133 Rome, Italy

²Department of Water, Environment, Construction and Safety, Magdeburg–Stendal University of Applied Sciences, Breitscheidstraße 2, 39114 Magdeburg, Germany

³Research Department IV – Complexity Science and Research Department I – Earth System Analysis, Potsdam Institute for Climate Impact Research (PIK) – Member of the Leibniz Association, Telegrafenberg A31, 14473 Potsdam, Germany

⁴Royal Meteorological Institute of Belgium, Bruxelles, Belgium

Correspondence: Tommaso Alberti (tommaso.alberti@inaf.it)

Abstract. Atmosphere and ocean dynamics display many complex features and are characterized by a wide variety of processes and couplings across different timescales. Here we demonstrate the application of Multivariate Empirical Mode Decomposition (MEMD) to investigate the multivariate and multiscale properties of a reduced order model of the ocean-atmosphere coupled dynamics. MEMD provides a decomposition of the original multivariate time series into a series of oscillating patterns with time-dependent amplitude and phase by exploiting the local features of the data and without any a priori assumptions on the decomposition basis. Moreover, each oscillating pattern, usually named Multivariate Intrinsic Mode Function (MIMF), represents a local source of information that can be used to explore the behavior of fractal features at different scales by defining a sort of multiscale/multivariate generalized fractal dimensions. With these two approaches, we show that the ocean-atmosphere dynamics presents a rich variety of features, with different multifractal properties for the ocean and the atmosphere at different timescales. For weak ocean–atmosphere coupling, the resulting dimensions of the two model components are very different, while for strong coupling for which coupled modes develop, the scaling properties are more similar especially at longer time scales. The latter result reflects the presence of a coherent coupled dynamics. Finally, we also compare our model results with those obtained from reanalysis data demonstrating that the latter exhibit a similar qualitative behavior in terms of multiscale dimensions and the existence of a scale-dependency of topological and geometric features for different regions, being related to the different drivers and processes occurring at different timescales in the coupled atmosphere-ocean system. Our approach can therefore be used to diagnose the strength of coupling in real applications.

1 Introduction

The atmosphere and the ocean form a complex system whose dynamical variability extends over a wide range of spatial and temporal scales (Liu, 2012; Xue et al., 2020). As an example, the tropical regions are markedly characterized by inter-/multi-



annual processes like the El Niño–Southern Oscillation (ENSO) (Neelin et al., 1994; Meehl et al., 2003), while the North Atlantic Oscillation (NAO) affects extra-tropical Northern regions at seasonal and decadal timescales (Ambaum et al., 2001). The sources of these processes have been widely investigated by means of multiple data analysis methods and various types of modelling (e.g., Philander, 1990; Czaja and Frankignoul, 2002; Van der Avoird et al., 2002; Mosedale et al., 2006; Kravtsov et al., 2007; Feliks et al., 2011; Liu, 2012; L’Hévéder et al., 2014; Farneti, 2017; Vannitsem and Ghil, 2017; Wang, 2019; Xue et al., 2020, and reference therein), highlighting how the atmospheric low-frequency variability (LFV) is strictly related to the ocean. The latter develops thanks to the interaction with the ocean mixed layer (OML) driven by a mixing process due to the development of an instability within the water column (Czaja and Frankignoul, 2002; D’Andrea et al., 2005; Wunsch and Ferrari, 2004; Gastineau et al., 2012) that also shows a strong seasonal variability. The relation between the OML and the LFV can be investigated from a dynamical systems point of view by developing suitable reduced order ocean-atmosphere models dealing with the modelling of the coupling between the atmosphere and the underlying surface layer of the ocean. Recently, by means of a 36-variable model displaying marked LFV Vannitsem et al. (2015) demonstrated that the LFV in the atmosphere could be a natural outcome of the ocean-atmosphere coupling.

The current work presents an investigation on how a recently introduced concept of multiscale generalized fractal dimensions can be used to analyze the topological and geometric properties of attractors in coupled ocean-atmosphere systems (Alberti et al., 2020a). This demonstration is done by means of the reduced order model developed in Vannitsem et al. (2015). Indeed, the dynamical properties of physical systems can be related to their support fractal dimension as well as its singularities by means of different established concepts like the box-counting dimension (Ott, 2002), generalized correlation integrals (Grassberger, 1983; Hentschel and Procaccia, 1983; Pawelzik and Schuster, 1987), the pointwise dimension method (Farmer et al., 1983; Donner et al., 2011), and related characteristics (Badii and Politi, 1984; Primavera and Florio, 2020). These methods are based on partitioning the phase-space into hypercubes of size ℓ to define a suitable invariant measure through the filling probability of the i -th hypercube by N_k points as $p_k = N_k/N$, with N being the total number of points. With $M(\ell)$ denoting the number of filled hypercubes, we can define some useful dynamical invariants such as the box-counting (or capacity or simply fractal) dimension

$$D_0 \doteq -\lim_{\ell \rightarrow 0} \lim_{N \rightarrow \infty} \frac{\log M(\ell)}{\log \ell}, \quad (1)$$

the information dimension

$$D_1 \doteq \lim_{\ell \rightarrow 0} \lim_{N \rightarrow \infty} \frac{\sum_{k=1}^{M(\ell)} p_k \log p_k}{\log \ell}, \quad (2)$$

and the correlation dimension

$$D_2 \doteq \lim_{\ell \rightarrow 0} \lim_{N \rightarrow \infty} \frac{\frac{1}{N^2} \sum_{i \neq j} \Theta(\ell - |x_i - x_j|)}{\log \ell}, \quad (3)$$

with $\Theta(\dots)$ being the Heaviside function. All these fractal dimension measures, as well as their higher order extensions D_q , have been used to characterize the global dynamical, topological, and geometric properties of a given system (Hentschel and Procaccia, 1983), however, without exploring how these properties evolve at different scales (Alberti et al., 2020a). More



recently, by means of a suitable combination between a state of the art time series decomposition method (the Empirical Mode Decomposition) and the concept of generalized fractal dimensions, Alberti et al. (2020a) introduced a multiscale approach to deal with the investigation of the evolution of topological properties in dynamical systems.

Here, we extend for the first time the concept of multiscale generalized fractal dimensions in a multivariate framework by means of the Multivariate Empirical Mode Decomposition (MEMD), allowing us to investigate the multiscale and multivariate properties of a reduced order model of the ocean-atmosphere coupled dynamics. By using the oscillating patterns forming the decomposition basis of the MEMD algorithm, usually named Multivariate Intrinsic Mode Function (MIMF), we define a sort of multiscale/multivariate generalized fractal dimensions. The MEMD results allow us to capture the essential dynamics of the phase-space trajectory that can be used for reconstructing the skeleton of the phase-space dynamics, while the evaluation of the fractal dimensions at different timescales provides a quantitative characterization of the intrinsic complexity of oscillating patterns that can be related to the attractor properties. Our results also allow for associating the geometric and topological properties to the dynamical regimes at different timescales of the coupled ocean-atmosphere system. Finally, our findings for the reduced order model well reconcile with corresponding results for reanalysis data, thus supporting and encouraging the use of reduced order models for investigating the essential aspects of the coupled ocean-atmosphere system in terms of topological and geometrical features.

2 The reduced order ocean-atmosphere model

Reduced order coupled ocean-atmosphere models are key tools in the hierarchy of climate models, allowing for an extensive analysis of the features of the coupled dynamics that would otherwise be impossible to evaluate (Lorenz, 1984; Nese and Dutton, 1993; Roebber, 1995; Jin, 1996; Timmermann et al., 2003; Van Veen, 2003; De Cruz et al., 2016; Vannitsem, 2017). These models allow for obtaining key insights into the role of coupling for the development of LFV in the atmosphere associated with the presence of the ocean.

Recently, dynamical analysis have been conducted by means of the development of a suitable reduced order model of the coupled ocean-atmosphere system. It has been developed starting from the quasi-geostrophic equations describing the interaction between a two-layer atmosphere and a one-layer ocean over an infinitely deep quiescent ocean layer (Vannitsem et al., 2015; Vannitsem, 2015; De Cruz et al., 2016; Vannitsem, 2017; De Cruz et al., 2018). The ocean flow passively advects the temperature within the ocean, while momentum, radiative, and heat transfer mechanisms realize the coupling between the atmosphere and the ocean. By expanding the solutions of these equations into Fourier series, by truncating them at low wavenumbers, and by projecting onto the Fourier modes retained, a set of ordinary differential equations is derived. The fields are defined over a rectangular domain with $0 \leq x \leq 2\pi L/n$ and $0 \leq y \leq \pi L$, being n the aspect ratio between the meridional and the zonal extents of the domain and L the characteristic spatial scale. Moreover, periodic boundaries along the zonal direction and free-slip along the meridional direction are chosen for the atmosphere, while a closed basin with no flux through the boundaries is imposed for the ocean.



85 In the reduced order coupled model version proposed in Vannitsem et al. (2015), a long-periodic attracting orbit combining atmospheric and oceanic variables emerges from a Hopf bifurcation for large values of the meridional gradient of radiative input and frictional coupling. Beyond a certain value of the meridional gradient for the radiative input, a chaotic behavior appears, which is still dominated by LFV on decadal and multi-decadal time-scales.

Here we used the original version of the model (Vannitsem et al., 2015) where the four relevant fields, i.e., the barotropic and baroclinic atmospheric streamfunctions, the ocean streamfunction and the ocean temperature, are given by $\psi_a = \sum_{i=1}^{10} \psi_{a,i} F_i$, $\theta_a = \sum_{i=1}^{10} \theta_{a,i} F_i$, $\Psi_o = \sum_{i=1}^8 \Psi_{o,i} \phi_i$ and $T_o = \sum_{i=1}^8 T_{o,i} \phi_i$, where F_i and ϕ_i are simplified notations for the sets of modes used, compatible with the boundary conditions of both the atmosphere and the ocean. The parameter values used are the ones given in Figs. 8 and 9 of Vannitsem (2017). Depending on the choice of the surface friction coefficient C , different solutions are found with a highly chaotic dynamics without marked LFV in the atmosphere for small values of C , but a more moderately chaotic dynamics with stronger LFV in both the ocean and the atmosphere (related to the development of a coupled mode) for larger values of C .

3 Methods

Traditional multivariate and/or spatiotemporal data analysis methods are commonly based on fixing an orthogonal decomposition basis, satisfying certain mathematical properties of completeness, convergence, linearity, and stationarity (Chatfield, 2016). However, these conditions are not usually met when real-world geophysical data are analyzed, which calls for more adaptive methods (Huang et al., 1998). Moreover, geophysical data are usually also characterized by scale-invariant features over a wide range of scales with different complexity and show a scale-dependent behavior due to several factors like forcings, coupling, intrinsic variability, and so on (Franzke et al., 2020). For the above reasons, in this work we put forward a novel approach based on combining two different data analysis methods for investigating the multiscale fractal behavior of the coupled ocean-atmosphere system: Multivariate Empirical Mode Decomposition (MEMD; Rehman and Mandic, 2010) and generalized fractal dimensions (Hentschel and Procaccia, 1983).

3.1 Multivariate Empirical Mode Decomposition (MEMD)

The Multivariate Empirical Mode Decomposition (MEMD) is the "natural" multivariate extension of the univariate Empirical Mode Decomposition (EMD) (Huang et al., 1998; Rehman and Mandic, 2010). MEMD directly works on the data domain, instead of defining a conjugate space as for Fourier or Wavelet transforms, with the aim of being as adaptive as possible to minimize mathematical assumptions and definitions (Huang et al., 1998) in extracting embedded structures in the form of so-called Multivariate Intrinsic Mode Functions (MIMFs) (Rehman and Mandic, 2010). Each MIMF is an oscillatory pattern of the multivariate coordinates having the same number (or differing at most by one) of local extremes and zero crossings, and whose upper and lower envelopes are symmetric (Huang et al., 1998; Rehman and Mandic, 2010). MIMFs are derived through the sifting process (Huang et al., 1998). This process is easily realized for univariate signals (Huang et al., 1998), while needs to be carefully implemented for multivariate processes (Rehman and Mandic, 2010), since it is based on the cubic spline



interpolation of local extremes that cannot easily be defined on a k -dimensional space (Rehman and Mandic, 2010). Rehman and Mandic (2010) proposed an alternative definition of local extremes for multivariate signals by considering the k -variate data as composed by k -dimensional signals projected onto appropriate directions in this k -dimensional space. This allows us to perform cubic spline interpolation in each direction, with the suitable directions chosen by means of a combination of a quasi-Monte Carlo-based low-discrepancy sequences and a uniform angular sampling method (Rehman and Mandic, 2010). These allow a uniform distribution of vectors and a proper estimate of the local mean estimates (see, e.g., Rehman and Mandic, 2010, for more details).

The main steps of the sifting process acting on a k -variate signal $\{\mathbf{s}(t)\}_{t \in T} = \{s_1(t), s_2(t), \dots, s_k(t)\}$ can be summarized as below:

1. identify local extremes (i.e., data points where abrupt changes in the local tendency of the series under study are observed);
2. interpolate local extremes separately by cubic splines (i.e., produce continuous functions with smaller error than other polynomial interpolation);
3. derive the upper $\{\mathbf{u}(t)\}_{t \in T}$ and the lower $\{\mathbf{l}(t)\}_{t \in T}$ envelopes;
4. derive the mean envelope $\{\mathbf{m}(t)\}_{t \in T}$ as $\{\mathbf{m}(t)\}_{t \in T} = \frac{\{\mathbf{u}(t)\}_{t \in T} + \{\mathbf{l}(t)\}_{t \in T}}{2}$;
5. evaluate the resulting candidate MIMF as $\{\mathbf{h}(t)\}_{t \in T}(t) = \{\mathbf{s}(t)\}_{t \in T} - \{\mathbf{m}(t)\}_{t \in T}$.

The previous steps are iteratively repeated until the obtained candidate MIMF $\{\mathbf{h}(t)\}_{t \in T}(t)$ can be identified as a Multivariate Intrinsic Mode Function (also called multivariate empirical mode) (Huang et al., 1998; Rehman and Mandic, 2010), while the full sifting process ends when no more MIMFs $\{\mathbf{c}_j(t)\}_{t \in T}$ can be filtered out from the data. Hence, we can write

$$\{\mathbf{s}(t)\}_{t \in T} = \sum_{j=1}^{N_j} \{\mathbf{c}_j(t)\}_{t \in T} + \{\mathbf{r}(t)\}_{t \in T}. \quad (4)$$

In this way a multivariate signal is decomposed into N_j k -dimensional functions, each containing the same frequency distribution, e.g., into a set of k -dimensional embedded oscillating patterns $\{\mathbf{c}_j(t)\}_{t \in T}$ which form the multivariate decomposition basis, plus a multivariate residue $\{\mathbf{r}(t)\}_{t \in T}$.

For each MIMF we can define a k^* -variate mean timescale as

$$\tau_{j,k^*} = \frac{1}{T} \int_0^T t' c_{j,k^*}(t') dt', \quad (5)$$

representing the typical oscillation scale of the j -th mode for the k^* -th univariate component c_{j,k^*} extracted from the multivariate signal $\{\mathbf{s}(t)\}_{t \in T} = \{s_{k^*}(t)\}$ for $k^* \in [1, k]$. Similarly, by ensemble averaging over the k -dimensional space we can introduce the concept of a multivariate mean timescale as

$$\tau_j = \frac{1}{T} \int_0^T t' \langle \{\mathbf{c}_j(t')\}_{t' \in T} \rangle_k dt', \quad (6)$$



with $\langle \dots \rangle_k$ denoting an ensemble average over the k -dimensional space. Thus, the k^* -variate timescale τ_{j,k^*} is evaluated for each mode and for each k^* -dimensional data, while the multivariate mean timescale τ_j is the mean over all $k^* \in [1, k]$. Moreover, as for univariate EMD (Huang et al., 1998), we can introduce the concepts of instantaneous amplitudes $\{\mathbf{a}_j(t)\}_{t \in T}$ and phases $\{\phi_j(t)\}_{t \in T}$ of each MEMD mode via the Hilbert Transform along the different directions of the k -dimensional space. The instantaneous energy content is then derived as $\{\mathbf{E}_j(t)\}_{t \in T} = \{\mathbf{a}_j(t)\}_{t \in T}^2$. Thereby, we can characterize the spectral content by introducing an alternative yet equivalent definition of the power spectral density (PSD) as

$$S(\tau) = \frac{1}{T^2} \int_0^T \langle \{\mathbf{E}_j(t')\}_{t' \in T} \rangle_k dt' \cdot \int_0^T t' \langle \{\mathbf{c}_j(t')\}_{t' \in T} \rangle_k dt' \doteq \sigma^2(\tau) \cdot \tau, \quad (7)$$

with $\sigma^2(\tau)$ being the k -variate variance of MIMFs and τ the mean timescale defined as in Eq. (6). Moreover, from the instantaneous energy content $\{\mathbf{E}_j(t)\}_{t \in T}$ the relative contribution e_j can be derived as

$$e_j = \frac{\frac{1}{T} \int_0^T \langle \{\mathbf{E}_j(t')\}_{t' \in T} \rangle_k dt'}{\sum_{j=1}^{N_j} \frac{1}{T} \int_0^T \langle \{\mathbf{E}_j(t')\}_{t' \in T} \rangle_k dt'}. \quad (8)$$

Finally, as for the univariate decomposition (Huang et al., 1998), also the MIMFs are empirically and locally orthogonal each other, the decomposition basis is a complete set (Rehman and Mandic, 2010) and partial sums of Eq. (4) can be obtained (Alberti, 2018; Alberti et al., 2020b).

3.2 Multivariate and multiscale generalized fractal dimensions

The behavior of complex systems usually consists of a collection of scales whose dynamical features determine their collective behavior. Nevertheless, vast efforts have been made to determine collective properties of systems (e.g., Hentschel and Procaccia, 1983), instead of considering to measure scale-dependent features. Recently, Alberti et al. (2020a) introduced a new formalism allowing measuring information at different scales by combining a data-adaptive decomposition method and the classical concept of generalized fractal dimensions. The starting point is that a multivariate signal manifesting a multiscale behavior can be written as

$$\{\mathbf{s}(t)\}_{t \in T} = \langle \{\mathbf{s}\}_{t \in T} \rangle + \sum_{\tau} \delta \{\mathbf{s}_{\tau}(t)\}_{t \in T} = \{\mathbf{s}_0\} + \{\mathbf{s}_1(t)\}_{t \in T}, \quad (9)$$

with $\langle \dots \rangle$ representing a steady-state average operation and δ indicating a fluctuation at scale τ . For any given τ we can introduce a local natural probability measure $d\mu_{\tau}$ such that the probability p_i of visiting the i -th hypercube $B_{\mathbf{s}^*, \tau}(\ell)$ of size ℓ centered at the point $\{\mathbf{s}^*\}$ on the considered (d -dimensional) phase-space of $\{\mathbf{s}_1(t)\}_{t \in T}$ can be defined as

$$p_i \doteq \int_{\mathbf{s}_1 \in B_{\mathbf{s}^*, \tau}(\ell)} d\mu_{\tau}. \quad (10)$$

By defining a q -th order partition function

$$\Gamma_q(\mu_{\tau}, B_{\mathbf{s}^*, \tau}(\ell)) = \sum_i p_i^q = \int d\mu_{\tau}(s) \mu_{\tau}(B_{\mathbf{s}^*, \tau}(\ell))^q \quad (11)$$



and taking the limit $\ell \rightarrow 0$, the multiscale generalized fractal dimensions are derived as

$$D_{q,\tau} = \frac{1}{q-1} \lim_{\ell \rightarrow 0} \frac{\log \Gamma_q(\mu_\tau, B_{s^*,\tau}(\ell))}{\log \ell}. \quad (12)$$

175 Here we identify the intrinsic oscillations by using the MEMD and then we investigate the phase-space properties at different scales by deriving the generalized dimensions (Alberti et al., 2020a). Summarizing:

1. we extract multiscale components from $\{s(t)\}_{t \in T}$ by using the MEMD;
2. we evaluate the intrinsic scale τ_j of each MIMF;
3. we evaluate reconstructions of modes by means of Eq. (4)

$$180 \quad \sum_{\tau} \delta\{s_{\tau}(t)\}_{t \in T} \rightarrow F_{j^*}(t) = \sum_{j=1}^{j^*} \{c_j(t)\}_{t \in T} \quad (13)$$

with $j^* = 1, \dots, N_j$ (by construction, MIMFs are ordered from short to long scales, i.e., $\tau_j < \tau_{j'}$ if $j < j'$);

4. we evaluate the generalized dimensions $D_{q,\tau}$ from $F_{j^*}(t)$ for each j^* (i.e., for each scale τ_{j^*});
5. we evaluate the singularities and singularity spectrum

$$\alpha_{\tau} = \frac{d}{dq} [(q-1)D_{q,\tau}] \quad (14)$$

$$185 \quad f_{\tau} = f(\alpha_{\tau}) = q\alpha_{\tau} - [(q-1)D_{q,\tau}]. \quad (15)$$

From Eq. (13) we can inspect local properties of fluctuations in terms of the geometry of the phase-space, thus providing a characterization of dynamical features of different regimes and disentangling the different dynamical components of (possibly) different origin. Finally, it is expected (for ensuring convergence) that when $j^* \rightarrow N_j$ then $D_{q,\tau} \rightarrow D_q$, with D_q being the standard generalized fractal dimensions proposed by Hentschel and Procaccia (1983).

190 4 Results

4.1 Multivariate Empirical Mode Decomposition

Figure 1 reports the 3-D projection of the full system attractor in the subspace $(T_{o,2}, \Psi_{o,2}, \psi_{a,1})$ for two representative values of the friction coefficient C (0.008 and 0.015 $\text{kg m}^{-2} \text{s}^{-1}$ as indicated by red and black points, respectively). In the following, we will omit the physical units of this parameter for the sake of brevity. The considered subspace characterizes the dynamics of the system as represented by the dominant mode of the meridional temperature gradient in the ocean ($T_{o,2}$), by the double-gyre transport within the ocean ($\Psi_{o,2}$), and by the vertically averaged zonal flow within the atmosphere ($\psi_{a,1}$), respectively.

The behavior of the system is clearly dependent on the friction coefficient, with both the location and the topology of the attractor changing as C is increased from 0.008 (red points in Fig. 1) to 0.015 (black points in Fig. 1). This behavior has also




Figure 1. 3-D projection of the full system attractor in the subspace $(T_{o,2}, \Psi_{o,2}, \psi_{a,1})$ for $C = 0.008$ (red) and $C = 0.015$ (black), respectively.

been previously reported by Vannitsem et al. (2015) and Vannitsem (2015), indicating a drastic qualitative change of the nature of the dynamics at about $C = 0.011$ above which substantial LFV emerges (Vannitsem et al., 2015; Vannitsem, 2015, 2017). However, all model components are clearly characterized by multiscale variability, spanning a wide range of timescales that can contribute to the dynamics in different ways, depending on the values of the friction coefficient and the intrinsic variability of the coupled ocean-atmosphere system.

Figure 2 displays the behavior of the spectral energy content $S(\tau)$ of the different MIMFs as a function of their mean timescales τ as in Eq. (7) for the full system (atmosphere+ocean) and for the two subsystems separately (i.e., the atmosphere and the ocean, respectively). First of all, it is important to underline that a different number of MIMFs has been identified for the two different cases: $N_j = 17$ for $C = 0.008$ and $N_j = 22$ for $C = 0.015$. This underlines that the respective dynamical behavior of the system is different, being characterized by different sets of empirical modes and consequently by a different number of relevant timescales. Moreover, by keeping in mind that for pure noise the expected number of MIMFs is $\log_2 N$ with N being the number of data points, both situations cannot be related with a purely stochastic dynamics. Indeed, in both cases we have used $N = 10^5$ data points, thus the expected number of MIMFs is $N_j^{\text{noise}} = 16$ (Flandrin et al., 2004). However, an interesting feature is that for the lower C value a number of MIMFs closer to that expected for noisy data is found, possibly related to the more irregular dynamics in this low friction coefficient case. Conversely, a marked departure from $N_j = 16$ is found for the higher C case, corresponding to a more regular dynamics characterized by significant LFV.

Furthermore, from Fig. 2 it is easy to note that the behavior of $S(\tau)$ depends on both the friction coefficient C and the different components of the model. For the full system (i.e., atmosphere+ocean) $S(\tau)$ decreases as τ increases for both values of C , while it is characterized by increasing spectral energy content at larger scales (i.e., at lower frequencies). By discriminating




Figure 2. Spectral energy content $S(\tau)$ of the different MIMFs as a function of their mean timescales τ as in Eq. (7) for the full system (atmosphere+ocean, blue circles), only for the atmosphere (orange asterisks), and only for the ocean (yellow diamonds). Left and right panels refer to the two values of the friction coefficient, $C = 0.008$ and $C = 0.015$, respectively.

between the atmospheric and the oceanic contribution we are able to see that (as expected), the short-term variability of the full system can be attributed to the atmosphere, while the long-term one is a reflection of the ocean dynamics. Moreover, when C increases we note an increase of the spectral energy content at all timescales, together with a flattening of the atmospheric spectral behavior, while the ocean dynamics seems to preserve its spectral features. These behaviors can be related to the existence of multiscale variability of the full system that can be linked to the different components operating at different timescales and to the different dynamics of the system as the friction coefficient C is changed.




Figure 3. Relative contribution (in percentage) $E_{\chi,\tau}$ of each variable $\chi = \{\psi_{a,i}, \theta_{a,i}, \Psi_{o,i}, T_{o,i}\}$ in dependence on the mean timescale τ . Left and right panels refer to the two values of the friction coefficient $C = 0.008$ and $C = 0.015$, respectively. The white line separates the atmospheric variables from the oceanic ones.



To further clarify the latter aspect, we evaluate the relative contribution (in percentage) $E_{\chi,\tau}$ of the different MIMFs (i.e., at
 225 different timescales τ) for each variable $\chi = \{\psi_{a,i}, \theta_{a,i}, \Psi_{o,i}, T_{o,i}\}$ as reported in Fig. 3. It can be clearly noted that the oceanic
 variability mainly contributes to the low-frequency dynamics ($E_{\chi,\tau} > 95\%$ for $\chi = \{\Psi_{o,i}, T_{o,i}\}$ and $\tau \gtrsim 10^4$ days), while the
 atmosphere is mainly characterized by short-term variability for $C = 0.008$ ($E_{\chi,\tau} > 95\%$ for $\chi = \{\psi_{a,i}, \theta_{a,i}\}$ and $\tau \lesssim 10$ days)
 and by both short- and long-term dynamics for $C = 0.015$. This points towards the C -dependent behavior of the atmospheric
 dynamics, with the ocean multiscale variability being less affected by changes in the values of the friction coefficient, and to
 230 the role of the ocean in developing LFV in the atmosphere as C increases.

Thanks to the completeness property of the MEMD we can explore the dynamics of the system as reproduced by the most
 energetic empirical modes via partial sums of Eq. (4). By using the information coming from the energy percentage distribution
 across the different timescales for each variable χ we can provide MIMF reconstructions accounting for a certain percentage of
 energy with respect to the total spectral energy content. By ordering the empirical modes with decreasing relative contribution
 235 e_j and summing up those contributing at least 95% of the total spectral content, we are able to investigate the 3-D projection
 of the full system attractor onto the subspace $(T_{o,2}, \Psi_{o,2}, \psi_{a,1})$ and compare it with the projection obtained by considering all
 timescales (as in Figure 1). Thus, for each variable $\chi = \{\psi_{a,i}, \theta_{a,i}, \Psi_{o,i}, T_{o,i}\}$ we can define a reconstruction based on empirical
 modes, $R_{\chi,95\%}$, as

$$R_{\chi,95\%}(t) \doteq \sum_{j'|e_{j'} \geq 95\%} \{\mathbf{c}_{\chi,j'}(t)\}_{t \in T} \quad (16)$$

240 with $\{\mathbf{c}_{\chi,j'}(t)\}_{t \in T}$ being the j' -th multivariate empirical mode extracted via the MEMD of the variables χ . The 3-D pro-
 jection onto the subspace $(T_{o,2}, \Psi_{o,2}, \psi_{a,1})$ of $R_{\chi,95\%}$ is shown in Fig. 4, while Tab. 1 summarizes the mode indices j' and
 corresponding k^* -variate timescales τ_{j',k^*} (see Eq. (5)) used for the reconstruction.

Table 1. Mode indices j' and corresponding k^* -variate timescales τ_{j',k^*} (see Eq. (5)) used for the reconstruction based on empirical modes
 $R_{\chi,95\%}$.

C	χ	j'	τ_{j',k^*} [days]
0.008	$\psi_{a,1}$	1, 2	3, 5
	$\Psi_{o,2}$	14, 15, 16	631, 1333, 2086
	$T_{o,2}$	14, 15, 16	599, 1132, 1913
0.015	$\psi_{a,1}$	21	2690
	$\Psi_{o,2}$	19, 20, 21	829, 1469, 2449
	$T_{o,2}$	19, 20, 21	735, 1506, 2598

By comparing Figs. 1 and 4 it can be easily noted that the underlying structure of the 3-D projection of the full attractor
 is essentially the same, thus suggesting that the subspace topological and geometric information can be recovered by a subset
 245 of multivariate empirical modes. This underlines that the dynamics of the full system can be reproduced by only few relevant




Figure 4. 3-D projection of the full system attractor in the subspace $(T_{o,2}, \Psi_{o,2}, \psi_{a,1})$ for $C = 0.008$ (red) and $C = 0.015$ (black), respectively, as obtained from reconstructions based on the multivariate empirical modes $R_{\chi,95\%}(t)$ accounting for 95% of the total variance of the model dynamics.

timescales without too much loss of information, thus reducing the complexity of the low order model itself. These results appear relevant if put into the wider context of coupled ocean-atmosphere dynamics, allowing us to recover the main features by only considering the most relevant (in terms of energy) timescale dynamical components.

4.2 Multiscale generalized fractal dimensions

250 Under general conditions, the complexity of a dynamical system can be conveniently investigated by means of the nonlinear properties of its phase-space trajectory (e.g., its attractor or repeller in case of dissipative dynamics) (Ott, 2002). One of the most common ways to characterize the topology of an attractor is to compute its spectrum of generalized fractal dimensions, allowing us to statistically characterize important properties of the dynamics as reflected by its phase-space geometry, including its information content, complexity, and underlying fractal structure (Grassberger, 1983; Hentschel and Procaccia, 1983; Donner
 255 et al., 2011). However, classical approaches can only provide global information on the phase-space topology (Hentschel and Procaccia, 1983; Ott, 2002), while multiscale dynamical systems can be characterized by topological properties changing as different scales are considered (Alberti et al., 2020a). For this purpose, we investigate the topological properties of the attractor of the coupled ocean-atmosphere model by evaluating the multiscale generalized fractal dimensions described in Section 3.2. Figures 5 and 6 report the behavior of the correlation dimension D_2 for both values of the friction coefficient and for three
 260 different cases: (a) for each MIMF individually (D_2^j), (b) for reconstructions of MIMFs summing up from $j = 1$ to N_j ($D_2^{\sum_j}$), and (c) for reconstructions of MIMFs performed separately for each variable $\chi = \{\psi_{a,i}, \theta_{a,i}, \Psi_{o,i}, T_{o,i}\}$.




Figure 5. Multiscale correlation dimension $D_{2,\tau}$ for $C = 0.008$ at different timescales τ_j for different cases: (a) for each MIMF individually (D_2^j), (b) for reconstructions of MIMFs summing up from $j = 1$ to N_j ($D_2^{\sum_j}$), and (c) for reconstructions of MIMFs separately for each variable (barotropic modes - blue circles, baroclinic modes - orange asterisks, transport modes - yellow diamonds, and temperature modes - violet symbol). Each panel also shows the 95% confidence intervals as error bars.

As expected, the multiscale correlation dimension for each MIMF decreases with increasing timescale, being representative of a more regular, less stochastic/chaotic, behavior of large-scale MIMFs as compared with the short-term ones (Alberti et al., 2020a). Particularly, when approaching the largest timescales, $D_{2,\tau} \rightarrow 1$ suggesting the existence of fixed-scale MIMFs, i.e.,
 265 with the instantaneous frequencies being almost constant (as expected, e.g., Rehman and Mandic, 2010). Conversely, when the multiscale correlation dimensions are evaluated by summing up the different MIMFs, starting from the shortest up to the largest scale, a clearly scale-independent behavior of $D_{2,\tau}$ is highlighted for both values of the friction coefficient C . However, the role of C clearly emerges in determining the values of $D_{2,\tau}$, being lower for the larger C value. Indeed, $D_{2,\tau} \sim 8$ for $C = 0.008$, while $D_{2,\tau} \sim 1.5$ for $C = 0.015$. This reflects the different topological properties of the attractor of the full system associated
 270 with a different dynamical behavior of the model variables (Faranda et al., 2019). However, the most interesting features emerge when the different variables of both atmosphere and ocean are separately investigated by means of the multiscale generalized fractal dimensions.

The described findings are not only valid for the multiscale correlation dimension $D_{2,\tau}$ but are also observed for both the multiscale capacity dimension $D_{0,\tau}$ and the multiscale information dimension $D_{1,\tau}$ as reported in Figs. 7 and 8, together with
 275 the multiscale correlation dimension $D_{2,\tau}$, for both values of C . Our formalism reveals the expected property that for $q < q'$,




Figure 6. Same as in Fig. 5, but for $C = 0.015$.

$D_{q,\tau} > D_{q',\tau} \forall \tau$ (Alberti et al., 2020a; Hentschel and Procaccia, 1983). Moreover, when evaluating the multiscale generalized fractal dimensions for each MIMF separately (e.g., Figs. 7(a) and 8(a)) a decreasing value for D_q^j is found as τ increases, with all D_q^j converging towards the same value of 1 at large timescales. Conversely, when the $D_{q,\tau}$ are evaluated for reconstructions based on MIMFs a completely different behavior emerges between the oceanic and the atmospheric variables. In this case, the atmospheric variables are characterized by scale-independent $D_{q,\tau}$, being representative of a high-dimensional system whose prime dynamics occurs at short timescales and with little effects of large-scale processes on the collective dynamics of the atmosphere. By contrast, a clearly scale-dependent behavior is found for the oceanic variables, with the multiscale generalized dimensions decreasing at larger timescales, reflecting the effects of large-scale dynamics dominating with respect to the short-term one for the ocean variability.

By estimating the Lyapunov spectra (not shown) separately for the ocean and the atmosphere we obtained that for $C = 0.008$ the instability is large for the atmosphere with a Lyapunov dimension $D_L \sim 10$, while for $C = 0.015$ the instability is weaker for the atmosphere, and the Lyapunov dimension is a bit larger than 4. Following the Kaplan-Yorke conjecture (Kaplan and Yorke, 1979), the Lyapunov dimension can be used as a proxy of the Hausdorff and, hence, capacity dimension. Hence, our results are clearly consistent with the dimension estimates for the atmosphere. For the ocean, however, there seems to be a less good agreement, with $D_L \approx 2$ while we found that $D_{0,\tau} \approx 4$. This quantitative disagreement could be related to the fact that the ocean can be viewed as a relatively stable system perturbed by high-frequency "noise" provided the atmosphere. Deeper investigations will be devoted to clarify this point in future research.




Figure 7. Multiscale capacity dimension $D_{0,\tau}$, multiscale information dimension $D_{1,\tau}$, and multiscale correlation dimension $D_{2,\tau}$ for $C = 0.008$ at different timescales τ_j for different cases: (a) for each MIMF individually (D_q^j), (b) for reconstructions of MIMFs summing up from $j = 1$ to N_j ($D_q^{\sum_j}$), and (c)-(f) for reconstructions of MIMFs separately for each variable (barotropic modes - (c), baroclinic modes - (d), transport modes - (e), and temperature modes - (f)).




Figure 8. Same as in Fig. 7, but for $C = 0.015$.

As a further step, we evaluate the full spectrum of generalized fractal dimensions for each MIMF by considering all moments $q \in [-20, 20]$, thus providing an estimate of the asymptotic values $D_{\pm\infty,\tau}$. This analysis allows characterizing how the



295 (multi)fractal properties of the system evolve with the timescale τ . Indeed, there are ongoing discussions on the fractal structure of both, the atmosphere and the ocean, especially dealing with the short-term variability and in terms of scaling-law behavior and statistics of increments (e.g., Franzke et al., 2020).




Figure 9. $D_{q,\tau}$ spectra for the coupled ocean-atmosphere dynamics at different timescales τ_j (indicated by different line colors) for reconstructions of MIMFs summing up from $j = 1$ to N_j ($D_q^{\sum_j}$) for (a) $C = 0.008$ and (b) $C = 0.015$.

The $D_{q,\tau}$ spectrum for $q \in [-20, 20]$ is reported in Fig. 9, where colored lines correspond to different timescales. It can be observed that for both values of the friction coefficient C , the full system exhibits signatures of multifractality at all timescales, especially at very short and very long timescales. By defining the multifractal width as $\Delta_\infty \doteq D_{-\infty,\tau} - D_{+\infty,\tau}$ we observe (see
 300 Fig. 10(a,b), black circles) that $\Delta_\infty = 3$ for $\tau \in [\tau_S, \tau_L]$ days, while $\Delta_\infty > 3$ for both $\tau < \tau_S$ and $\tau > \tau_L$, with $\tau_S \sim 20$ days and $\tau_L \sim 1$ year. This behavior could be the reflection of processes operating at different timescales for both the atmosphere (at short timescales) and the ocean (at long timescales). In order to further disentangle those processes, we also evaluated the full spectra of the generalized multifractal dimensions by considering all orders $q \in [-20, 20]$ for each subsystem (i.e., atmosphere
 305 and ocean) individually. For both values of C , the corresponding results are shown in Fig. 11.

We clearly see that for the atmosphere, there is a scale-independent behavior of $D_{q,\tau}$ for all q , rendering the different curves almost invariant with respect to the scale. By contrast, a scale-dependent behavior emerges for the ocean for the lower value of C . Indeed, it is evident that as the timescale increases the multiscale generalized dimensions tend to decrease for all values of q , moving from $D_{q,\tau_1} \in [5, 8]$ to $D_{q,\tau_{17}} \in [1, 3]$ for $C = 0.008$. Conversely, although there is an overall reduction in the
 310 $D_{q,\tau}$ values for $C = 0.0015$ with respect to those evaluated for $C = 0.008$, the decrease with the timescale is less evident for this higher C value, although it is still present for $\tau > 1$ year (see orange and red curves in comparison with the blue ones in Fig. 11(d)). This clearly suggests that the presence of strong multifractality in the full system can be essentially attributed to the atmosphere, with only a marginal role of the ocean variability in determining the fractal structure of the full system. By evaluating the difference between $D_{-\infty,\tau}$ and $D_{+\infty,\tau}$ we can clearly see that larger values, of the order of 4, are found for the
 315 atmosphere, at almost all timescales (and especially at shorter timescales), for both values of C . Conversely, larger values are found at shorter timescales for both values of C for the ocean. As the timescale increases, this difference tends to be reduced




Figure 10. Multifractal width Δ_∞ at different timescales τ_j for reconstructions of MIMFs summing up from $j = 1$ to N_j ($D_q^{\sum_j}$) for (a) $C = 0.008$ and (b) $C = 0.015$. The different colors refer to the full system (atmosphere+ocean, black circles), only the atmosphere (red circles), and only the ocean (green circles), respectively.




Figure 11. $D_{q,\tau}$ spectra for the dynamics of atmosphere and ocean individually at different timescales τ_j (indicated by different line colors) for reconstructions of MIMFs summing up from $j = 1$ to N_j ($D_q^{\sum_j}$) for (a,c) $C = 0.008$ and (b,d) $C = 0.015$. Panels (a,b) refer to the atmosphere, (c,d) to the ocean.



to values close to 2, suggesting a reduced multifractality of the ocean with respect to the atmosphere, especially for the lower value of C at larger timescales when the role of the ocean becomes dominant as compared to the atmosphere (see Fig. 2).

4.3 Comparison with regional averages from reanalysis data

320 As a final step we compare our previous results for the reduced order coupled ocean-atmosphere model with those obtained from reanalysis data (Poli, 2015). More specifically, we use three different sets of regional time series based on the European Centre for Medium-range Weather Forecasts (ECMWF) ORA-20C project (De Boissésou and Balmaseda, 2016; De Boissésou et al., 2017) that is a 10-member ensemble of ocean reanalyses covering the complete 20th century using atmospheric forcing from the ERA-20C reanalysis (<https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era-20c>). Here, we focus on
325 data from January 1958 to December 2009 at monthly resolution in terms of different monthly-averaged time series, the set of data also used previously in Vannitsem and Ekelmans (2018). This period has been chosen in the latter study because of the ocean reanalysis dataset showing here smaller uncertainties than during the first half of the 20th century (De Boissésou and Balmaseda, 2016).

Three different representative regions are chosen: the North Atlantic region, corresponding to the domain defined by $\lambda \in$
330 $[55^\circ\text{W}, 15^\circ\text{W}]$ and $\phi \in [25^\circ\text{N}, 60^\circ\text{N}]$, the North Pacific region, i.e., a spherical-rectangle domain with $\lambda \in [165^\circ\text{E}, 225^\circ\text{E}]$ and $\phi \in [25^\circ\text{N}, 60^\circ\text{N}]$, and the Tropical Pacific region, corresponding to $\lambda \in [165^\circ\text{E}, 225^\circ\text{E}]$ and $\phi \in [25^\circ\text{S}, 25^\circ\text{N}]$ (Vannitsem and Ekelmans, 2018). The individual series for the two extratropical regions have been derived by projecting the reanalysis fields on two dominant Fourier modes: (i) $F_1 = \sqrt{2} \cos(\pi y/L_y)$, and (ii) $\phi_2 = 2 \sin(\pi x/L_x) \sin(2\pi y/L_y)$ (Vannitsem and Ekelmans, 2018). For the Tropical Pacific region, the series are formed by spatial averages. In this way, we obtain two sets of three time
335 series each for both the North Atlantic and the North Pacific (i.e., one for the atmosphere and two for the ocean), and a third set of three time series for the Tropical Pacific (two for the atmosphere at two different pressure levels and one for the ocean). This allows us to build up a 3-D projection of the local atmosphere-ocean coupled dynamics for each region (see Vannitsem and Ekelmans, 2018, for more details).

By using the MEMD analysis to investigate the multivariate patterns of reanalysis data we found the same number of $N_j = 9$
340 MIMFs for each region, whose mean timescales range from ~ 2 months up to ~ 20 years, suggesting the existence of multiscale variability over a wide range of scales. As for the reduced order model, we first investigate the behavior of the spectral energy content $S(\tau)$ of the different MIMFs as a function of their mean timescales τ as in Eq. (7) for the three different regions as shown in Fig. 12. We clearly observe an increase of the spectral energy content up to a timescale $\tau \sim 1$ year for all regions, then declining for both the North Atlantic and the North Pacific. Conversely, the Tropical Pacific is characterized by larger
345 spectral content also for timescales larger than 1 year, up to $\tau \sim 5$ years, which coincide with the typical timescales of the El Niño–Southern Oscillation (ENSO). Furthermore, for all regions a decreasing spectral energy content is found at the largest timescales (i.e., $\tau > 5$ years).

To further compare our above model results with those obtained for the reanalysis data, we evaluate the multiscale generalized fractal dimensions for the three different regions. For each region, we derive both the multifractal width $\Delta_\infty \doteq$
350 $D_{-\infty, \tau} - D_{+\infty, \tau}$ and the full multiscale multifractal spectrum at different timescales τ_j for reconstructions of MIMFs sum-




Figure 12. Spectral energy content $S(\tau)$ of the different MIMFs as a function of their mean timescales τ as in Eq. (7) for the North Atlantic (blue circles), the North Pacific (orange asterisks), and the Tropical Pacific (yellow diamonds).

ming up from $j = 1$ to N_j ($D_q^{\sum_j}$). Figure 13 shows the corresponding results for the North Atlantic region, the North Pacific region, and the Tropical Pacific region, respectively.

First of all, it is important to underline that the multiscale generalized fractal dimensions are clearly different with respect to those obtained from the ocean-atmosphere model. This directly follows from the different numbers of variables (time series) in the model, being a 36-dimensional dynamical system, with respect to the reanalysis data, being a 3-dimensional projection of the regional ocean-atmosphere dynamics. Nevertheless, although different in terms of absolute values, both the model and the reanalysis data show a similar qualitative behavior with varying scale τ , although some differences are found between the different regions.

On the one hand, both the North Atlantic and the North Pacific regions (see Fig. 13(d,e)) are characterized by a scale-dependent behavior, with decreasing $D_{q,\tau}$ as τ increases. Moreover, by looking at the multifractal width as a function of the scale (Fig. 13(a,b)) we find evidence for a decreasing Δ_∞ as τ increases, being representative of a transition from a short-term multifractal nature to long-term monofractal one. These features can be interpreted in terms of the different multiscale dynamical processes affecting the atmosphere on short scales and the ocean on larger scales.

On the other hand, by looking at the Tropical Pacific region we clearly see an enhancement of Δ_∞ , i.e., the emergence of multifractal features (see Fig. 13(c)), at annual/multi-annual timescales (i.e., $\tau \sim 1 - 8$ years), being also characterized by the largest values of the multiscale generalized fractal dimensions (see Fig. 13(f)). This could be related to the role of the El Niño–Southern Oscillation (ENSO) cycle manifesting at these timescales (between 2 and 7 years), which is likely responsible for the different scale-dependent behavior of $D_{q,\tau}$ as compared to the two other extratropical regions.




Figure 13. (a)-(c) Multifractal width Δ_∞ and (d)-(f) $D_{q,\tau}$ spectra at different timescales τ_j for reconstructions of MIMFs summing up from $j = 1$ to N_j ($D_q^{\sum_j}$) for (a,d) the North Atlantic, (b,e) the North Pacific, and (c,f) the Tropical Pacific, respectively.

In summary, by means of the reanalysis data, we have been able to demonstrate that i) the reduced order coupled ocean-
 370 atmosphere model and the reanalysis data show some qualitatively similar behavior of the multiscale generalized fractal di-
 mensions, although characterized by different absolute values due to the different numbers of variables considered in the
 model and the projections on a few modes of the reanalysis data, and that ii) interesting features emerge when looking at the
 scale-dependency of topological and geometric features for different regions, being the reflection of different driving mecha-
 nisms and processes operating at different timescales in the coupled ocean-atmosphere system. However, further investigations
 375 are needed to characterize the role of the different processes as well as their intrinsic dimensionality, occurrence, and spatial
 dependency in more detail. Such an in-depth investigation is outlined as a part of our future work.

5 Conclusions

We have provided a first time systematic investigation of the multiscale dynamics of a reduced order coupled ocean-atmosphere
 model (Vannitsem et al., 2015) as described by means of the topological and geometric features (Alberti et al., 2020a).

380 First, by means of the Multivariate Empirical Mode Decomposition (MEMD) we have been able to detect oscillating patterns
 with time-dependent amplitude and phase that are directly linked to a rich variety of features of the coupled ocean-atmosphere



system. We have found that the underlying structure of the 3-D projection of the full attractor is essentially reproduced by a subset of Multivariate Intrinsic Mode Functions (MIMFs) corresponding to the most relevant timescales without too much loss of information, thus further reducing the complexity of the reduced order model itself. These results appear relevant if put into
385 the wider context of coupled ocean-atmosphere dynamics, allowing us to recover the main features by only considering the most relevant (in terms of energy) timescale dynamical components.

Second, by exploiting the novel concept of multiscale/multivariate generalized fractal dimensions we have investigated the different multifractal properties for the ocean and the atmosphere at different timescales. We have demonstrated that for weak ocean-atmosphere coupling (i.e., for low values of the friction coefficient C), the resulting dimensions of the two model
390 components are very different, while for strong coupling (larger C) at which coupled modes develop at low frequencies, the scaling properties are more similar especially at longer time scales. These results suggest that as C increases, we observe the development of a coherent coupled dynamics, primarily at large timescales. In terms of the underlying fractal structure, we have found that for both considered values of the friction coefficient C , the full system exhibits signatures of multifractality at all timescales, especially pronounced at short and long as compared to intermediate timescales. By means of the full spectrum
395 of generalized fractal dimensions, we have clearly evidenced that for the atmosphere, there is a scale-independent behavior of $D_{q,\tau}$ for all q , rendering the multifractal spectra almost invariant with respect to the timescale. By contrast, a scale-dependent behavior emerges for the ocean for the lower value of C . This clearly suggests that the presence of strong multifractality in the full system can be attributed to the atmosphere, with only a marginal role of the ocean variability in determining the fractal structure of the full system.

400 Finally, we have compared our results for the reduced order coupled ocean-atmosphere model with those derived from reanalysis data (Poli, 2015) by using three sets of different regional time series from the ORA-20C project (De Boissésou and Balmaseda, 2016; De Boissésou et al., 2017). Although the resulting multiscale generalized fractal dimensions clearly differ quantitatively from those obtained from the ocean-atmosphere model – which can be easily understood by considering the different dimensions of the model (a 36-dimensional dynamical system) and the reanalysis data (3-dimensional projections of
405 the local ocean-atmosphere dynamics) – we observed a similar qualitative behavior with the scale τ . Interestingly, the multiscale multifractal features of different regions show different scale-dependent behaviors. Specifically, both the North Atlantic and the North Pacific regions are characterized by a scale-dependent behavior, with decreasing $D_{q,\tau}$ as τ increases, with a transition from a short-term multifractal nature to long-term monofractal one. These features can be interpreted in terms of the different multiscale dynamical processes affecting the atmosphere at short timescales and the ocean at longer timescales. Conversely,
410 the Tropical Pacific region is characterized by the emergence of multifractal features at annual/multi-annual timescales (i.e., $\tau \sim 1 - 8$ years), being also characterized by the largest values of the multiscale generalized fractal dimensions. This behavior can be seen as a manifestation of the El Niño–Southern Oscillation (ENSO) cycle that typically acts at these timescales and can be considered the key driving factor of a different scale-dependent behavior of $D_{q,\tau}$ as compared to the two other extratropical regions.

415 Our findings for both the model and the reanalysis data suggest that our approach can be used to diagnose the strength of coupling in the ocean-atmosphere system and to investigate the topological features of the system. We have demonstrated that the



420 model and the reanalysis data show qualitatively similar behavior of the multiscale generalized fractal dimensions. However, the different scale-dependency of topological and geometric features for different regions can contribute to a better understanding on the different driving mechanisms and processes operating at different timescales in the coupled ocean-atmosphere system. These observations suggest that further investigations are needed to better characterize the role of the different processes as well as their intrinsic dimensionality, occurrence, and spatial dependency, which shall be further addressed in our future work.

Code availability. All codes used for the analysis and generating the figures can be obtained from the authors upon request.

425 *Data availability.* The time series of the model used in the present manuscript are available from the authors upon request. The reanalysis dataset is available on Zenodo, <https://doi.org/10.5281/zenodo.1135134>.

Author contributions. TA, RVD, and SV designed the study. TA conducted the analysis and drafted the manuscript. RVD and SV contributed to the interpretation of the results. All authors contributed to the writing of the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

430 *Acknowledgements.* This work has been partially supported by the German Federal Ministry for Education and Research (BMBF, grant no. 01LP2002B) and the Belgian Science Policy Office (Belspo) through the JPI Climate/JPI Oceans project ROADMAP.



References

- Alberti, T.: Multivariate empirical mode decomposition analysis of Swarm data, *Il Nuovo Cimento C*, 41, 1-10, 2018.
- Alberti, T., Consolini, G., Ditlevsen, P. D., Donner, R. V., and Quattrociochi, V.: Multiscale measures of phase-space trajectories, *Chaos*, 30, 123116, doi:10.1063/5.0008916, 2020.
- 435 Alberti, T., Giannattasio, F., De Michelis, P., and Consolini, G.: Linear Versus Nonlinear Methods for Detecting Magnetospheric and Ionospheric Current Systems Patterns, *Earth Space Sci.*, 7, e2019EA000559, 2020.
- Ambaum, M.H.P., Hoskins, B. J., and Stephenson, D. B.: Arctic Oscillation or North Atlantic Oscillation?, *J. Clim.*, 14,3495–3507, 2001.
- Badii, R., and Politi, A.: Hausdorff dimension and uniformity factor of strange attractors, *Phys. Rev. Lett.*, 52, 1661–1664, 1984.
- Chatfield, C.: *The analysis of time series – an introduction*, Chapman and Hall/CRC, 6th edition, London, 2016.
- 440 Czaja, A., and Frankignoul, C.: Observed impact of Atlantic SST anomalies on the North Atlantic Oscillations, *J. Clim.*, 15, 606-623, 2002.
- D’Andrea, F., Czaja, A., and Marshall, J.: Impact of anomalous ocean heat transport on the North Atlantic Oscillation, *J. Clim.*, 18, 4955-4969, 2005.
- De Boissésou, E. and Balmaseda, M.: An ensemble of 20th century ocean reanalyses for providing ocean initial conditions for CERA-20C coupled streams, ERA report series, 24, European Centre for Medium-range Weather Forecasts, Reading, UK, 2016.
- 445 De Boissésou, E., Balmaseda, M., and Mayer, M.: Ocean heat content variability in an ensemble of twentieth century ocean reanalyses, *Clim. Dyn.*, 50, 3783-3798, doi:10.1007/s00382-017-3845-0, 2018.
- De Cruz, L., Demaeyer, J., and Vannitsem, S.: The Modular Arbitrary-Order Ocean-Atmosphere Model: MAOOAM v1.0, *Geoscient. Mod. Dev.*, 9, 2793, 2016.
- De Cruz, L., Schubert, S., Demaeyer, J., Lucarini, V., and Vannitsem, S.: Exploring the Lyapunov instability properties of high-dimensional atmospheric and climate models, *Nonlin. Proc. Geophys.*, 25, 387–412, <https://doi.org/10.5194/npg-25-387-2018>, 2018.
- 450 Donner, R.V., Heitzig, J., Donges, J.F., Zou, Y., Marwan, N., and Kurths, J.: The geometry of chaotic dynamics – a complex network perspective, *Eur. Phys. J. B*, 84, 653–672, 2011.
- Faranda, D., Messori, G., and Vannitsem, S.: Attractor dimension of time-averaged climate observables: insights from a low-order ocean-atmosphere model, *Tellus A: Dynamic Meteorology and Oceanography*, 71:1, 10.1080/16000870.2018.1554413, 2019.
- 455 Farmer, J.D., Ott, E., and Yorke, J.A.: The dimension of chaotic attractors, *Physica D*, 7, 153-180, 1983.
- Farneti, R.: Modelling interdecadal climate variability and the role of the ocean, *WIREs Clim. Change*, 8, e441, 2017.
- Feliks, Y., Ghil, M., and Robertson, A. W.: The atmospheric circulation over the North Atlantic as induced by the SST field, *J. Clim.*, 24(2), 522–542, doi: 10.1175/2010JCLI3859.1, 2011.
- Flandrin, P., Rilling, G., and Goncalves, P.: Empirical Mode Decomposition as a Filter Bank, *IEEE Sign. Proc. Lett.*, 11, 112, doi:10.1109/LSP.2003.821662, 2004.
- 460 Franzke, C.L.E., Barbosa, S., Blender, R., Fredriksen, H.-B., Laepple, T., Lambert, F., et al.: The Structure of Climate Variability Across Scales, *Rev. Geophys.*, 58, e00657. doi:10.1029/2019RG000657, 2020.
- Gastineau, G., D’Andrea, F., and Frankignoul, C.: Atmospheric response to the North Atlantic Ocean variability on seasonal to decadal time scales, *Clim. Dyn.*, 40, 2311–2330, 10.1007/s00382-012-1333-0, 2013.
- 465 Grassberger, P.: Generalized dimensions of strange attractors, *Phys. Lett. A*, 97, 227–230, 1983.
- Hentschel, H. G. E. and Procaccia, I.: The infinite number of generalized dimensions of fractals and strange attractors, *Physica D*, 8, 435-444, 1983.



- Huang, N. E., Shen, Z., Long, S. R., et al.: The Empirical Mode Decomposition and the Hilbert Spectrum for Nonlinear and Nonstationary Time Series Analysis, *Proc. R. Soc. London A*, 454, 903, 1998.
- 470 Huang, N. E., and Wu, Z.: A review on Hilbert-Huang transform: Method and its applications to geophysical studies, *Rev. Geophys.*, 46, RG2006, 2008.
- Jin, F.-F.: Tropical Ocean-Atmosphere interaction, the Pacific cold tongue, and the El-Niño-Southern Oscillation, *Science*, 274, 76-78, 1996.
- Kaplan, J. L., and Yorke, J. A.: Chaotic behavior of multidimensional difference equations, in: Peitgen, H.-O., and Walther, H.-O. (eds) *Functional Differential Equations and Approximation of Fixed Points*, Lecture Notes in Mathematics, vol 730, pp. 204-227, Springer, 475 Berlin, Heidelberg, doi: 10.1007/BFb0064319, 1979.
- Kravtsov, S., Dewar, W. K., Berloff, P., Ghil, M. and McWilliams, J. C.: A highly nonlinear coupled mode of decadal variability in a mid-latitude ocean-atmosphere model, *Dyn. Atmos. Oceans*, 43, 123-150, 2007.
- L'Hévéder, B., Codron, F., and Ghil, M.: Impact of anomalous northward oceanic heat transport on global climate in a slab-ocean setting, *J. Clim.*, 28, 2650-2664, 2014.
- 480 Liu, Z.: Dynamics of interdecadal climate variability: A historical perspective, *J. Clim.*, 25, 1963-1995, 2012.
- Lorenz, E. N., Formulation of a low-order model of a moist general circulation, *J. Atmos. Sci.*, 41, 1933-1945, 1984.
- Meehl, G. A., Arblaster, J. M., and Loschnigg, J.: Coupled Ocean-Atmosphere Dynamical Processes in the Tropical Indian and Pacific Oceans and the TBO, *J. Clim.*, 16(13), 2138-2158, 2003.
- Mosedale, T. J., Stephenson, D. B., Collins, M., and Mills, T. C.: Granger Causality of Coupled Climate Processes: Ocean Feedback on the 485 North Atlantic Oscillation, *J. Clim.*, 19(7), 1182-1194, 2006.
- Neelin, J. D., Latif, M., and Jin, F.-F.: Dynamics of Coupled Ocean-Atmosphere Models: The Tropical Problem, *Ann. Rev. Fluid Mech.*, 26:1, 617-659, 1994.
- Nese, J.M., and Dutton, J.A.: Quantifying predictability variations in a low-order ocean-atmosphere model: A dynamical system approach. *J. Clim.*, 6, 185-203, 1993.
- 490 Ott, E.: *Chaos in Dynamical Systems - 2nd Edition*, Cambridge University Press, pp. 490, doi:10.2277/0521811961, 2002.
- Pawelzik, K., and Schuster, H.G.: Generalized dimensions and entropies from a measured time series, *Phys. Rev. A*, 35, 481-484, 1987.
- Philander, S. G. H.: *El Niño and the Southern Oscillation*, Academic Press, New York, 1990.
- Poli, P., Hersbach, H., Tan, D. G. H., Dee, D. P., Thépaut, J.-N., Simmons, A., Peubey, C., Laloyaux, P., Komori, T., Berrisford, P., Dragani, R., Trémolet, Y., Hólm, E. V., Bonavita, M., Isaksen, L., and Fisher, M.: The data assimilation system and initial performance evaluation 495 of the ECMWF pilot reanalysis of the 20th-century assimilating surface observations only, ERA report series, 14, European Centre for Medium-range Weather Forecasts, Reading, UK, 2015.
- Primavera, L., and Florio, E.: Parallel Algorithms for Multifractal Analysis of River Networks, in: Sergeyev Y., Kvasov D. (eds) *Numerical Computations: Theory and Algorithms*, Lecture Notes in Computer Science, 11973, Springer, Cham, 2020.
- Rehman, N. and Mandic, D. P.: Multivariate empirical mode decomposition, *Proceedings of the Royal Society A*, 466, 1291-1302, 2010.
- 500 Roebber, P. J.: Climate variability in a low-order coupled atmosphere-ocean model, *Tellus A*, 47, 473-494, 1995.
- Timmermann, A., Jin, F.-F., and Abshagen, J.: A nonlinear theory for El Niño bursting. *J. Atmos. Sci.*, 60, 152-165, 2003.
- Van der Avoird, E., Dijkstra, H.A., Nauw, J. J., and Schuurmans, C. J. E.: Nonlinearly induced low-frequency variability in a midlatitude coupled ocean-atmosphere model of intermediate complexity, *Clim. Dyn.*, 19, 303-320, 2002.
- Van Veen, L.: Overturning and wind driven circulation in a low-order ocean-atmosphere model, *Dynam. Atmos. Oceans*, 37, 197-221, 2003.



- 505 Vannitsem, S.: The role of the ocean mixed layer on the development of the North Atlantic Oscillation: A dynamical system's perspective, *Geophys. Res. Lett.*, 42, 8615, 2015.
- Vannitsem, S.: Predictability of large-scale atmospheric motions: Lyapunov exponents and error dynamics, *Chaos*, 27, 032101, 2017.
- Vannitsem, S. and Ekkelmans, P.: Causal dependences between the coupled ocean–atmosphere dynamics over the tropical Pacific, the North Pacific and the North Atlantic, *Earth Syst. Dynam.*, 9, 1063–1083, <https://doi.org/10.5194/esd-9-1063-2018>, 2018.
- 510 Vannitsem, S., and Ghil, M.: Evidence of coupling in ocean-atmosphere dynamics over the North Atlantic, *Geophys. Res. Lett.*, 44, 2016–2026, doi:10.1002/2016GL072229, 2017.
- Vannitsem, S., Demaeyer, J., de Cruz, L., and Ghil, M.: Low-frequency variability and heat transport in a low-order nonlinear coupled ocean-atmosphere model, *Physica D*, 309, 71–85, 2015.
- Wang, C.: Three-ocean interactions and climate variability: a review and perspective, *Clim. Dyn.* 53, 5119–5136, 2019).
- 515 Wunsch, C., and Ferrari, R.: Vertical mixing, energy, and the general circulation of the ocean, *Annu. Rev. Fluid Mech.*, 36, 281–314, 2004.
- Xue, P., Malanotte-Rizzoli, P., Wei, J., and Eltahir, E. A. B.: Coupled ocean-atmosphere modeling over the Maritime Continent: A review, *J. Geophys. Res. Oceans*, 125, e2019JC014978, <https://doi.org/10.1029/2019JC014978>, 2020.