



<b>Publication Year</b>	2015
<b>Acceptance in OA</b>	2020-12-30T12:12:51Z
<b>Title</b>	The evolution of the disc variability along the hard state of the black hole transient GX 339-4
<b>Authors</b>	De Marco, B., PONTI, GABRIELE, Muñoz-Darias, T., Nandra, K.
<b>Publisher's version (DOI)</b>	10.1093/mnras/stv1990
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/29368">http://hdl.handle.net/20.500.12386/29368</a>
<b>Journal</b>	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
<b>Volume</b>	454

# The evolution of the disc variability along the hard state of the black hole transient GX 339-4

B. De Marco,<sup>1★</sup> G. Ponti,<sup>1</sup> T. Muñoz-Darias<sup>2,3,4</sup> and K. Nandra<sup>1</sup>

<sup>1</sup>Max-Planck-Institut für Extraterrestrische Physik, Giessenbachstrasse 1, D-85748 Garching, Germany

<sup>2</sup>Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain

<sup>3</sup>Departamento de astrofísica, Univ. de La Laguna, E-38206 La Laguna, Tenerife, Spain

<sup>4</sup>Department of Physics, Astrophysics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK

Accepted 2015 August 25. Received 2015 August 5; in original form 2015 August 4

## ABSTRACT

We report on the analysis of hard-state power spectral density function (PSD) of GX 339-4 down to the *soft* X-ray band, where the disc significantly contributes to the total emission. At any luminosity probed, the disc in the hard state is intrinsically more variable than in the soft state. However, the fast decrease of disc variability as a function of luminosity, combined with the increase of disc intensity, causes a net drop of fractional variability at high luminosities and low energies, which reminds the well-known behaviour of disc-dominated energy bands in the soft state. The peak frequency of the high-frequency Lorentzian (likely corresponding to the high-frequency break seen in active galactic nuclei, AGN) scales with luminosity, but we do not find evidence for a linear scaling. In addition, we observe that this characteristic frequency is energy dependent. We find that the normalization of the PSD at the peak of the high-frequency Lorentzian decreases with luminosity at all energies, though in the *soft* band this trend is steeper. Together with the frequency shift, this yields quasi-constant high-frequency (5–20 Hz) fractional rms at high energies, with less than 10 per cent scatter. This reinforces previous claims suggesting that the high-frequency PSD solely scales with black hole mass. On the other hand, this constancy breaks down in the *soft* band (where the scatter increases to  $\sim 30$  per cent). This is a consequence of the additional contribution from the disc component, and resembles the behaviour of optical variability in AGN.

**Key words:** accretion, accretion discs – X-rays: binaries – X-rays: individual: GX 339-4.

## 1 INTRODUCTION

Black hole X-ray binaries (BHXRB) spend most of their time in a quiescent state, which is occasionally interrupted by outbursts of activity. As the outburst evolves the source passes through different accretion states and both the variability and spectral properties change (e.g. Miyamoto et al. 1992; Belloni et al. 2005; Remillard & McClintock 2006). The variability properties of these sources have been extensively studied in the past, revealing the existence of trends as a function of accretion state and/or luminosity (e.g. Zdziarski et al. 2004; Belloni, Motta & Muñoz-Darias 2011; Muñoz-Darias, Motta & Belloni 2011). However, due to the sensitivity limits of the detectors employed (particularly the Proportional Counter Array, PCA, on the *Rossi X-ray Timing Explorer*, *RXTE*), these studies have been mostly limited to energies  $> 3$  keV. At these energies the broad-frequency-band (e.g. in the typical frequency range  $\sim 0.1$ –60 Hz) aperiodic variability progressively decreases as the outburst

evolves, with the fractional root-mean-square (rms) variability amplitude (e.g. Nandra et al. 1997; Vaughan et al. 2003; Ponti et al. 2004) dropping from values of tens of percent in the canonical hard state, down to a few percent in the canonical soft state (Belloni et al. 2005; Muñoz-Darias, Motta & Belloni 2011). During the hard state the fractional rms shows an ‘inverted spectrum’ (decreasing with energy), which switches to a ‘hard spectrum’ (increasing with energy, at least above  $\sim 5$  keV) during intermediate and soft states (e.g. Zdziarski et al. 2004; Belloni et al. 2011, as illustrated in the inset of Fig. 4). These behaviours are usually ascribed to variations of the relative contribution of the main X-ray spectral components (e.g. Gierliński & Zdziarski 2005). Indeed, over the past years, the general idea was that the strong broad-frequency-band variability characterizing the hard state was associated with the Comptonized hard X-ray emission (power law) from the corona (or the base of a radio-emitting jet; e.g. Zdziarski et al. 1998; Markoff, Nowak & Wilms 2005; Droulans et al. 2010), since this component dominates the X-ray spectrum during this state. On the other hand, the disc-thermal component, known to be variable on the time-scales of the outburst, was thought to be intrinsically constant on shorter

\* E-mail: [bdemarco@mpe.mpg.de](mailto:bdemarco@mpe.mpg.de)

**Table 1.** Log of the analysed *XMM-Newton* and simultaneous *RXTE* observations: (1) observation ID; (2) date; (3) detector used for the analysis and corresponding observing mode (for the *RXTE* observations the Proportional Counter Units – PCU – co-added in the analysis are also reported); (4) effective exposure (the *XMM-Newton* effective exposures refer to the duration of the observation after removal of high soft proton background, the percentage of this exposure affected by telemetry dropouts is reported within parentheses) (5) mean count rates in the energy bands used for the analysis (i.e. bands 0.5–1.5 keV – soft – and 2–9 keV – hard – for *XMM-Newton*, and band 10–30 keV – very hard – for *RXTE*).

<i>XMM-Newton</i>				
(1) Obs ID	(2) Date	(3) Instrument/Mode	(4) Exp (drop outs) (s)	(5) Mean count rate (soft/hard) (cts s <sup>-1</sup> )
0204730201	2004-03-16	Epic-PN/Timing	104 200 (20.7 per cent)	90/108
0204730301	2004-03-18	Epic-PN/Timing	71 000 (21 per cent)	98/117
0605610201	2009-03-26	Epic-PN/Timing	32 220 (0.2 per cent)	43/61
0654130401	2010-03-28	Epic-PN/Timing	33 630 (23.7 per cent)	357/434
<i>RXTE</i>				
Obs ID	Date	Instrument/Mode	Exp (s)	Mean count rate (very hard) (cts s <sup>-1</sup> PCU <sup>-1</sup> )
90118-01-05-00	2004-03-16	PCU 0,2,3/Good Xenon	2324	55
90118-01-06-00	2004-03-17	PCU 0,2/Good Xenon	1660	58
90118-01-07-00	2004-03-18	PCU 0,2/Good Xenon	6505	60
94405-01-03-00	2009-03-26	PCU 0,2/Good Xenon	3055	38
94405-01-03-01	2009-03-26	PCU 2/Good Xenon	3305	37
94405-01-03-02	2009-03-26	PCU 1,2/Good Xenon	1807	39
95409-01-12-01	2010-03-28	PCU 2/Generic Event	19 195	180

(than about hours–minutes) time-scales (e.g. Gierliński & Zdziarski 2005). Given that the disc dominates the spectrum up to energies of  $\sim 10$  keV during the soft state, it would then be responsible for the drop of broad-frequency-band variability amplitude, and for the change of shape of the fractional rms spectrum.

The way these variability properties extrapolate to soft X-ray energies ( $< 3$  keV) during the hard state is currently poorly known. Since, during the hard state the disc component gives non-negligible contribution in the soft X-ray energy band ( $\lesssim 1$  keV; Tomsick et al. 2008; Kolehmainen, Done & Díaz-Trigo 2014), to some extent we would expect to observe behaviours similar to those characterizing the disc during the soft state. However, the recent use of data from the EPIC-pn detector onboard *XMM-Newton* (sensitive down to  $\sim 0.3$  keV), for the spectral-timing analysis of BHXRBs, unveiled the existence of high levels of intrinsic (on time-scales of  $\gtrsim 1$  s) disc variability in the hard state (Wilkinson & Uttley 2009). Though this result has been initially obtained studying a single data set for each of the two sources GX 339-4 and Swift J1753.5–0127, it has been recently confirmed by the analysis of Swift data (Kalamkar et al. 2013, 2015), suggesting this might represent a general property of BHXRBs. Thus, extending timing studies down to soft X-ray energies is crucial in order to gain a complete understanding of the disc–corona variability processes.

Another important property of X-ray variability in BHXRBs is that the characteristic time-scales vary during the outburst. During the hard state all the variability components (commonly modelled with Lorentzians; Belloni, Psaltis & van der Klis 2002) in the power spectral density function (PSD) shift towards higher frequencies as the luminosity increases, while keeping tight correlations between each other (e.g. Belloni et al. 2005; Done & Gierliński 2005). This indicates strong dependence of X-ray variability on the luminosity/mass-accretion rate of the source. The shift of the characteristic frequencies as a function of mass-accretion rate is a general property of accreting systems. Indeed, as in BHXRBs, a scaling (consistent with being linear; McHardy et al. 2006; Koerding et al. 2007) also links the ‘high-frequency break’ (Edelson & Nandra 1999) and the mass-accretion rate in active

galactic nuclei (AGN). However, despite all these variations in both characteristic frequencies and broad-frequency-band variability amplitude, the high-frequency PSD (i.e. above the peak frequency of the high-frequency Lorentzian) remains remarkably constant during the entire outburst (Gierliński, Niłojajuk & Czerny 2008), indicating little or no dependence on the luminosity and the accretion state of the source. In analogy with BHXRBs, in AGN the high-frequency PSD (above the high-frequency break) does not show any clear dependence on the luminosity/mass-accretion rate (e.g. O’Neill et al. 2005; Zhou et al. 2010; Ponti et al. 2012).

In this paper we aim at investigating the evolution of the PSD as a function of luminosity/mass-accretion rate, extending the analysis down to the soft X-ray energy band. For this reason, we study the hard-state PSD of GX 339-4, the best monitored BHXRB. We used data collected with *XMM-Newton*, and, in order to obtain a broad-band coverage, we considered also simultaneous *RXTE* observations. The data selection and reduction are presented in Section 2, while the PSD estimate and analysis procedures are described in Section 3. Given the different, frequency-dependent behaviours outlined before, which characterize the PSD at energies  $\gtrsim 3$  keV, we will first study the broad-frequency-band variability properties (Section 4.1–4.2), and then more specifically focus on the high-frequency part (Section 4.3). Results are discussed in Section 5.

## 2 DATA REDUCTION

GX 339-4 has been observed 18 times by *XMM-Newton* (as of May 2014). In this paper we focus on all the archived observations with exposures  $> 10$  ks, which caught the source in a hard state, i.e. four observations in total. In order to further increase the spectral band coverage we considered also all the available simultaneous *RXTE* observations. Details about the analysed data sets are reported in Table 1.

Among the four selected *XMM-Newton* observations, two (2004 March 16–18) were carried out during consecutive satellite revolutions. Hereafter, we will show results obtained by combining these two observations together, and we will refer to them as one single

observation. The remaining two observations belong to the same outburst of the source, but they have been carried out one year apart (in 2009 and 2010).


We used EPIC-pn (Strüder et al. 2001) and PCA (Jahoda et al. 2006) data, which ensure high effective area over a broad spectral range.

The *XMM-Newton* data reduction was done using the *XMM* Science Analysis System (SAS v13.5), and the latest current calibration files (CCF, as of 2014 May). The analysed data sets are all in Timing mode. We followed standard reduction procedures, and applied Rate Dependent PHA (RDPHA) and X-ray loading (XRL) corrections (Guainazzi & Smith 2013; Guainazzi 2014, and references therein). We selected time intervals free from background proton flares (GTI, the final effective exposures are listed in Table 1) and considered only events with PATTERN  $\leq 4$ . The source counts were extracted from the region which includes 90 per cent of the total collected counts. For all the observations this region falls between RAWX columns 31 and 45.

Being all the observations in Timing mode, pile-up is expected to be an issue only when the count rate exceeds  $\sim 800$  cts  $s^{-1}$  (in the 0.7–10 keV range; Guainazzi et al. 2014). This threshold is well above the average count rate registered during the 2004 and 2009 observations considered here. However, the 2010 observation exceeds this threshold. In addition, variability may cause pile-up during short time intervals even when the average count rate is below the threshold. Therefore we verified the pile-up level during each observation using the SAS tool ‘epatplot’. In the 2010 and 2004 observations, the observed *single* and *double* events fractions slightly deviate from the expected trend for data not affected by pile-up (respectively by  $\sim 1.2$  and 12 per cent in the 2–10 keV band for both observations), but this deviation is systematic at all energies. As noticed in Done & Díaz Trigo (2010), these systematic deviations might be either due to the uncertainties related to the modelling of the *singles* and *doubles* fraction for the Timing mode, or might indicate that the data are actually slightly affected by pile-up. Thus, we tested our results against pile-up by repeating the analysis (described in Section 3) of the 2010 observation (which has the highest EPIC-pn count rate) on the event file obtained by excising the central three pixels (i.e. RAWX=37–39), which reduces the excess of *double* events due to pile-up by a factor  $\sim 5$  in the 2–10 keV band. We found results perfectly consistent with those presented in Section 4, the only difference being the expected reduction of variability power signal-to-noise (S/N) ratio in the PSD.

For the *RXTE* data analysis we followed standard procedures<sup>1</sup> using the HEASOFT software v6.13. For the timing analysis, we used either Good Xenon or Generic Event configuration modes, depending on the availability (see Table 1). To maximize the S/N ratio we extracted data from all the layers of the Proportional Counter Units (PCU) that were simultaneously and continuously switched on during each observation. On the other hand, the luminosities are computed from data in Standard2 configuration, considering only the top layer of PCU2.

The position of the analysed observations within the hardness–intensity diagram (HID) is displayed in Fig. 1. This plot includes all the *RXTE* data relative to the 2004 (light grey dots) and 2009 (dark grey triangles) outbursts. Fluxes are computed in the 3–30 keV energy range, assuming a simple power law plus a Gaussian at the energy of the Fe K  $\alpha$  line. The hardness ratio is computed as



**Figure 1.** The hardness–intensity diagram of GX 339-4 as obtained from all the archived *RXTE* observations relative to the 2004 (light grey dots) and 2009 (dark grey triangles) outbursts. The luminosity is computed in the range 3–30 keV, the hardness is computed as the ratio between the 13–30 and 3–13 keV flux. Coloured points mark the position of the analysed *XMM-Newton* and simultaneous *RXTE* observations in the diagram. The labels refer to the nomenclature used in the paper to indicate the 2009 (low luminosity), 2004 (medium luminosity), and 2010 (high luminosity) observations. The error bar on the left-hand side of the plot reports the uncertainty on the HID normalization due to the uncertainties on the distance and the BH mass of GX 339-4.

the ratio between the 3–13 and 13–30 keV fluxes. The fluxes are converted into Eddington-scaled luminosities. Eddington luminosities are obtained by considering the currently available estimates and uncertainties on the distance and the BH mass of the source (i.e.  $\sim 6$ –10 kpc, and  $\sim 6$ –10  $M_{\odot}$ , respectively; Hynes et al. 2003, 2004; Muñoz-Darias, Casares & Martínez-Pais 2008) and assuming a reasonable mean value, i.e. 8 kpc and 8  $M_{\odot}$ . The error bar displayed on the left-hand side of the plot highlights the uncertainty on the plot normalization, as due to the uncertainties on the distance and the BH mass. Note that, when considering the relative increase of luminosity, the relevant uncertainty is the error on the flux measurements, which however is very small (i.e. of the order of  $\sim 0.1$ –0.3 per cent), thus it has not been reported in the plot. The analysed observations show luminosities  $L_{3-30\text{keV}}/L_{\text{Edd}} \sim 0.007$  (2009 observations),  $\sim 0.02$  (2004 observations), and  $\sim 0.07$  (2010 observations). In the following we will refer to these observations respectively as ‘low’, ‘medium’, and ‘high luminosity’.

### 3 DATA ANALYSIS

For each observation we extracted light curves with time bins of 1 ms (we verified that this choice of the time bin does not cause any spurious feature within the analysed frequency window, which might affect the analysis and results presented in this paper) in the three energy bands: 0.5–1.5 keV, 2–9 keV (using *XMM-Newton* data), and 10–30 keV (using *RXTE* data). Hereafter we will refer to these bands respectively as soft, hard, and very hard. According to previous analyses of the same data sets (Kolehmainen et al. 2014; Plant et al. 2015), the chosen soft band is where the accretion disc gives the most significant contribution. In the other two energy bands the dominant component is the primary hard X-ray power law (this is shown also in Section 4.1.1).

To study the evolution of the variability characteristics of GX 339-4 at different energies, we estimated the PSD within each of

<sup>1</sup> The reduction procedures are described in the *RXTE* data reduction cookbook: [http://heasarc.nasa.gov/docs/xte/recipes/cook\\_book.html](http://heasarc.nasa.gov/docs/xte/recipes/cook_book.html).

these energy bands. To this aim, the light curves were sampled over intervals shorter than the total exposure and of equal length within each observation. The length of the segments for the *XMM-Newton* light curves was chosen so as to exclude the gaps due to telemetry dropouts (which occur mostly when the target source is very bright, i.e. during the high- and medium-luminosity observations), while ensuring a sufficiently broad frequency coverage. For consistency, the *RXTE* light curves have been sampled according to the corresponding simultaneous *XMM-Newton* light curves (i.e. 21, 59, and 500 s for the high-, medium- and low-luminosity observations, respectively). We estimated the PSD (e.g. Nowak et al. 1999a) of each light-curve segment, using routines implemented with IDL v6.4, and averaged them to obtain an estimate of the intrinsic PSD for every observation and energy band. The level of variability power due to Poisson noise was determined fitting the high-frequency part (above 100 Hz) of the PSD with a constant. Then, the average PSDs have been Poisson noise-subtracted, and renormalized adopting the squared fractional rms normalization (Miyamoto et al. 1991). We corrected the *RXTE* data for instrumental dead-time (which produces an overestimate of the intrinsic Poisson noise level) using the dead-time model of Zhang et al. (1995, 1996). Since the variability power S/N ratio drops at high frequencies, we limited the analysis to frequencies  $\lesssim 20$  Hz.

The hard-state PSDs of BHXRBs are usually very complex. They contain several structures which can be modelled with multiple Lorentzian components (e.g. Nowak 2000, Belloni et al. 2002, Pottschmidt et al. 2003). We followed this same approach for the modelling of the PSDs of GX 339-4.


#### 4 RESULTS

The hard-state PSDs of GX 339-4 are shown in Figs 2 and 3. In Fig. 2 we show for each observation, the PSDs at different energies, while in Fig. 3 we show for each energy band, the PSDs at different luminosities.

All the PSDs show a similar shape, with two broad humps, whose intensity and characteristic frequencies clearly change as a function of both luminosity and energy. These two broad humps are a characteristic feature of BHXRBs in the hard state (e.g. Nowak, Wilms & Dove 1999b; Wilkinson & Uttley 2009; Böck et al. 2011; Cassatella, Uttley & Maccarone 2012; Grinberg et al. 2014). At frequencies lower than those covered here the variability power per unit logarithmic interval ( $\nu P_\nu$ ) is expected to drop in the hard state (*band-limited* noise). The chosen length of light-curve segments allows us to obtain a good sampling of the broad-band noise of GX 339-4, down to the onset of this drop. We verified the absence of additional low-frequency broad structures by inspecting the PSDs obtained using longer segments of *RXTE* data. This confirmed that the analysed PSDs are band-limited, at least in the hard and very hard bands. We note that the soft band PSD shows the same structures as those seen at higher energies, therefore we assume that also the soft band PSD is band-limited.


We used two Lorentzian components to model the two humps and more rigorously quantify their variations. A third Lorentzian is required in all the fits to account for residual structures.<sup>2</sup> The best-fitting centroid frequency and full width at half-maximum have

<sup>2</sup>The third Lorentzian fits the narrow peak of the low-frequency broad-band noise component of the high-luminosity observation. In the other two observations, this Lorentzian fits an additional broad component at intermediate frequencies (see Table 2).



**Figure 2.** The PSDs of GX 339-4 in the three energy bands 0.5–1.5 keV (soft), 2–9 keV (hard), and 10–30 keV (very hard), for each observation at different luminosity. The arrows mark the best-fitting peak frequency of the low-frequency ( $\nu_l$ ) and the high-frequency ( $\nu_h$ ) Lorentzians for each energy band.

been used to estimate the peak frequency (i.e. the frequency of the maximum power per logarithmic interval; Belloni et al. 2002) of each Lorentzian (see Table 2) using the formula reported in Section 4.2. The quality of the data is high enough, that our simple model, though providing a good description of each PSD, does not always return sufficiently small values of the reduced  $\chi^2$  (see Table 2). We verified that this is mainly due to the presence of small residuals which are left unmodelled. These residuals are usually accounted for by increasing the number of Lorentzian components (e.g. Nowak 2000, Belloni et al. 2005). However, comparing our results with those found in the literature, both for GX 339-4 (e.g. Nowak et al. 1999b; Nowak 2000) and other BHXRBs (e.g. Cassatella et al. 2012; Grinberg et al. 2014) we conclude that three variability components are sufficient to obtain a good description of the hard-state broad-band noise of GX 339-4 within the analysed frequency window (i.e. up to frequencies of 20 Hz). Our best-fitting models are plotted in Fig. 2 (solid lines) together with the



**Figure 3.** Overplotted are the PSDs of GX 339-4 at different luminosities, for each of the energy bands soft (upper panel), hard (middle panel), and very hard (lower panel).

single components (dotted curves). To avoid confusion, only the components associated with the very hard band PSD are shown in the plots. These best-fitting PSD models were first used to study the variations of the broad-frequency-band (between 0.002 and 20 Hz) fractional rms variability amplitude (Section 4.1). Then we used them to constrain the characteristic frequencies and study their variations as a function of energy and luminosity (Section 4.2). Finally, we focused on the variability properties of the high-frequency (between 5–20 Hz) portion of the PSD (Section 4.3).

#### 4.1 Trends of broad-frequency-band fractional rms as a function of energy and luminosity

We integrated the best-fitting PSD models for each luminosity and energy band, over the frequency interval 0.002–20 Hz (broad-frequency-band fractional rms), respectively for the high-, medium-, and low-luminosity observations. Note that there is no evidence of additional low-frequency broad noise structures in the *RXTE* data

**Table 2.** Results obtained from the fit of the PSDs of GX 339-4 with a model comprising three broad Lorentzians. Best-fitting values of the peak-frequencies (in units of Hz) of the lowest, intermediate, and highest frequency Lorentzian (respectively,  $\nu_1$ ,  $\nu_i$ , and  $\nu_h$ ) are reported, in the soft (0.5–1.5 keV, S), hard (2–9 keV, H), and very hard (10–30 keV, VH) band, and at different luminosities. Errors are reported at 90 per cent confidence level.

Band	Low $L$	Med $L$	High $L$
		$\nu_1$	
S	$0.0096 \pm 0.0014$	$0.042 \pm 0.002$	$0.146 \pm 0.013$
H	$0.0092 \pm 0.0014$	$0.043 \pm 0.003$	$0.172 \pm 0.026$
VH	$0.0080 \pm 0.0035$	$0.043 \pm 0.004$	$0.179 \pm 0.016$
		$\nu_i$	
S	$0.18 \pm 0.04$	$0.56 \pm 0.05$	$0.166 \pm 0.008$
H	$0.22 \pm 0.02$	$0.76 \pm 0.05$	$0.161 \pm 0.008$
VH	$0.17 \pm 0.06$	$0.55 \pm 0.08$	$0.161 \pm 0.012$
		$\nu_h$	
S	$1.15 \pm 0.15$	$1.18 \pm 0.10$	$1.85 \pm 0.15$
H	$1.88 \pm 0.15$	$2.79 \pm 0.29$	$2.89 \pm 0.11$
VH	$1.90 \pm 0.42$	$2.58 \pm 0.44$	$3.05 \pm 0.28$
		$\chi^2/\text{dof}$	
S	80.6/42	161.9/66	21.4/25
H	95.9/42	87.1/66	60.3/25
VH	44.4/42	89.4/30	36.8/25

(Section 4), and therefore these best-fitting models describe the data well down to 0.002 Hz, at least in the hard and very hard bands. Given that the soft band PSD shows similar structures as observed at higher energies, we assume that the extrapolation of our best-fitting model describes the data well down to very low frequencies also in this band. We also estimated the fractional rms associated with the low- and high-frequency Lorentzian (obtained integrating the single components of the best-fitting PSD models within the same broad-frequency-band interval). Results are reported in Table 3. The broad-frequency-band fractional rms, independently of luminosity and energy band, is always high ( $\gtrsim 30$  per cent), covering values typical of the hard state (Muñoz-Darias et al. 2011). In Fig. 4 we plot the broad-frequency-band fractional rms as a function of energy (within the three energy bands adopted throughout this paper). Note that the fractional rms of each Lorentzian shows the same energy-dependence as the broad-frequency-band fractional rms. For comparison we also report (inset of Fig. 4) the spectral shape of the 0.1–32-Hz fractional rms of GX 339-4 during a (high luminosity) hard and the soft state, at the energies covered by the *RXTE* PCA (adapted from Belloni et al. 2011).


Our main results can be summarized as follows.

(i) Above  $\sim 2$  keV, where the power-law component dominates the source emission, we always observe a decrease of broad-frequency-band fractional rms as a function of energy. Therefore such a decrease of variability with energy appears to be an intrinsic property of the power-law component. This might be produced by variations of the spectral index, with a high-energy pivot point. Extrapolating this trend on to the soft energy band, we see that the broad-frequency-band fractional rms in this energy band is lower than expected. This is consistent with the presence of a disc-component, less variable than the power law.

(ii) Comparing the decrease of broad-frequency-band fractional rms within each energy band as a function of luminosity, we find that the largest drop is observed in the soft energy band (a factor  $\sim 2.1$ , as compared to  $\sim 1.6$  and  $1.4$  in the hard and very hard energy bands). In Section 4.1.1, we show that, though the intrinsic fractional rms of

**Table 3.** The fractional rms (e.g. Vaughan et al. 2003) as obtained integrating the PSD best-fitting models of each observation at different luminosities, and in each of the three bands 0.5–1.5 keV (soft), 2–9 keV (hard), 10–30 keV (very hard). The broad-frequency-band fractional rms and the fractional rms associated with each Lorentzian are obtained integrating over the frequency interval 0.002–20 Hz. Finally we report the high-frequency fractional rms obtained integrating over the interval 5–20 Hz. Errors are computed at the 90 per cent confidence level.

	Soft	Hard	Very hard
Broad-frequency-band			
High $L$	$0.318 \pm 0.022$	$0.376 \pm 0.024$	$0.318 \pm 0.020$
Med $L$	$0.492 \pm 0.034$	$0.496 \pm 0.031$	$0.374 \pm 0.024$
Low $L$	$0.670 \pm 0.036$	$0.600 \pm 0.036$	$0.433 \pm 0.024$
Low-frequency Lorentzian			
High $L$	$0.231 \pm 0.024$	$0.251 \pm 0.026$	$0.203 \pm 0.022$
Med $L$	$0.363 \pm 0.037$	$0.349 \pm 0.035$	$0.261 \pm 0.027$
Low $L$	$0.426 \pm 0.046$	$0.384 \pm 0.041$	$0.281 \pm 0.023$
High-frequency Lorentzian			
High $L$	$0.187 \pm 0.020$	$0.223 \pm 0.022$	$0.189 \pm 0.019$
Med $L$	$0.312 \pm 0.032$	$0.240 \pm 0.028$	$0.223 \pm 0.021$
Low $L$	$0.386 \pm 0.030$	$0.335 \pm 0.031$	$0.245 \pm 0.026$
5–20 Hz			
High $L$	$0.075 \pm 0.007$	$0.118 \pm 0.011$	$0.103 \pm 0.010$
Med $L$	$0.109 \pm 0.010$	$0.138 \pm 0.011$	$0.112 \pm 0.011$
Low $L$	$0.132 \pm 0.010$	$0.147 \pm 0.012$	$0.108 \pm 0.010$



**Figure 4.** Variations of the broad-frequency-band fractional rms spectrum with luminosity. Different symbols refer to the energy bands: soft (black dots), hard (red triangles), and very hard (blue squares). The inset shows the typical spectral shape of the 0.1–32-Hz fractional rms of GX 339-4 during a (high luminosity) hard and soft state observation in the *RXTE* bandpass (adapted from Belloni et al. 2011).

the disc-component is high in the hard state, the fraction of variable disc flux in the soft band decreases fast (faster than the power-law component), thus causing the observed drop of soft band variability as the luminosity increases.


(iii) In both the medium- and low-luminosity observations, the broad-frequency-band fractional rms spectrum has an overall ‘inverted’ shape (decreasing with energy), typical of the hard state (e.g. Zdziarski et al. 2004; Belloni et al. 2011; inset of Fig. 4). However, during the high-luminosity observation, the large drop of broad-frequency-band fractional rms observed in the soft band causes a shift of the peak of variability power towards intermediate energies (i.e. in the hard band). This appears to be the onset of a transition

towards a ‘hard spectrum’, typical of intermediate and soft states (inset of Fig. 4).

In order to characterize the nature of the drop of broad-frequency-band fractional rms observed in the soft energy band at high luminosity, and investigate how this can be associated with an increasing contribution of disc-thermal emission in this band, we carried out fits of the *XMM-Newton* energy and covariance spectra (e.g. Wilkinson & Uttley 2009; Uttley et al. 2011, 2014).

#### 4.1.1 Is the disc variable or constant during the hard state?

Previous analyses of the same (or part of the) data sets analysed in this paper (Wilkinson & Uttley 2009; Kolehmainen et al. 2014) revealed the presence of a significant contribution from the accretion disc-thermal emission in the soft band. Thus we used the *XMM-Newton* data to measure the disc-thermal emission-to-total flux ratio in the soft band and verify whether the drop of fractional rms is related to spectral variations in this band. We fit the *XMM-Newton* spectra between 0.5–10 keV using a simple model for the continuum (i.e.  $tbabs*(diskbb + nthcomp)$ , with the Comptonization high-energy cut-off fixed at 100 keV and the seed photon temperature tied to the disc temperature), and excluding the range of energies 4–7 keV, dominated by the Fe K  $\alpha$  line. We also excluded the energies around the edges of the response matrix, since residuals at these energies might be an artefact of uncorrected XRL effects and/or charge transfer inefficiency (Kolehmainen et al. 2014). The contribution from the disc emission strongly depends on the cold absorption column density parameter,  $N_H$ . To be more conservative, we let this parameter free. Hence, all the uncertainties stated below account for the uncertainty on  $N_H$ . Nevertheless, the derived best-fitting values range between  $N_H \sim 5\text{--}6 \times 10^{21} \text{ cm}^{-2}$ , in agreement with the expected values (Dickey & Lockman 1990; Kong et al. 2000). In Fig. 5 we plot the broad-frequency-band fractional rms as a function of disc thermal emission-to-total (absorbed) flux ratio in the soft band, for the different luminosities. A net increase of



**Figure 5.** The broad-frequency-band fractional rms as a function of disc-to-total (absorbed) flux ratio in the soft band.

the disc relative contribution to the total flux as the luminosity increases is observed. This increase coincides with the net decrease of fractional rms, and the relation between the two is consistent with being linear (the best-fitting slope being  $-1.26 \pm 0.24$ ). Thus the observed drop of broad-frequency-band fractional rms at soft X-ray energies is strongly linked to the progressive increase of a disc component, in analogy with what would be expected from a comparison with disc-dominated states.


In order to produce a drop of the fractional rms, the disc component should be less variable than the power law. Thus we estimated the fractional rms intrinsic to the disc and the power-law components. To this aim we computed the broad-frequency-band covariance spectra (with respect to the 0.5–10-keV reference band) of the *XMM-Newton* data (see Wilkinson & Uttley 2009; Uttley et al. 2011, 2014), which allow us to determine the fraction of variable disc emission. Note that, strictly speaking, the covariance spectrum gives the spectral shape of the components whose variability is linearly correlated with the variability in the reference band. However, for intrinsic coherence close to unity (as in this case, see also De Marco et al. 2015), the covariance spectrum is equivalent, though with smaller error bars (Wilkinson & Uttley 2009; Uttley et al. 2014), to the rms spectrum, which gives the spectral shape of all the variable components. We fit both the covariance and the energy spectra with the same model used above, and with the  $N_{\text{H}}$  parameter

tied to the value obtained from the fit of the energy spectra. Fig. 6 shows the ratios to the best-fitting power law, once the normalization of the disc-component is set to zero. All the covariance spectra show an excess at soft X-ray energies. This demonstrates the presence of a variable disc component at all the luminosities spanned by the analysed hard-state observations. The intensity of the excess in the covariance spectra is lower than in the energy spectra in the medium- and high-luminosity observations, indicating that some fraction of the total disc emission is constant or that the disc is less variable. We used the best-fitting models to derive estimates of the intrinsic fractional rms of the disc and the power-law components (for each of the two components this is computed as the ratio between the absorbed flux in the covariance and in the energy spectrum). These are reported in Table 4. In addition, we report the fractions of variable/constant power-law/disc flux contributing to the total soft band flux.

We observe that the disc is intrinsically variable at all luminosities. Its fractional rms is comparable with that intrinsic to the power-law component (at low luminosities it is consistent with being as variable as the power law), and much higher than typically observed in soft states (a few percents; Muñoz-Darias et al. 2011). However, the disc fractional rms decreases with luminosity faster than the power-law fractional rms. Combined with the variation of the relative contribution from the disc and the power law to the total soft band flux (Fig. 5), the result is a net increase (by a factor of  $\sim 4$ ) of constant disc emission in the soft band ( $D_{\text{const}}$ ) as a function of luminosity. In comparison, the constant power-law emission in the soft band ( $P_{\text{const}}$ ) increases only by a factor  $\sim 1.6$ . Thus we conclude that the drop of fractional rms in the soft band is driven by the net increase of constant disc flux in this band. It is worth stressing that all these considerations refer to the broad-frequency-band, while a more detailed treatment of the high-frequency (i.e. above the peak frequency of the high-frequency Lorentzian) PSD properties is deferred to Section 4.3.

#### 4.2 Testing the correlation between the characteristic frequencies of the PSD in the soft band


Fig. 7 reports the best-fitting peak frequency of the low-frequency Lorentzian,  $\nu_1$ , plotted against the high-frequency one,  $\nu_h$  (these two frequencies are marked by arrows in the plots of Fig. 2). The peak frequency of each Lorentzian is computed using the formula (e.g.



**Figure 6.** Data-to-model ratios of the energy (red squares) and covariance spectra of the low- (left-hand panel), medium- (middle panel), and high-luminosity (right-hand panel) observations. The model used to fit the data is *tbabs\*(diskbb + nthcomp)*, and the normalization of the *diskbb* component has been set to zero to make the plots. All the covariance spectra have been computed using the 0.5–10 keV range as the reference band. The broad-frequency-band covariance spectra (black circles) are obtained integrating over the entire frequency range covered by each data set up to frequencies of 20 Hz (i.e. 0.002–20, 0.017–20, and 0.04–20 Hz, respectively, for the low-, medium-, and high-luminosity observations). The high-frequency covariance spectra (light grey triangles) are obtained integrating over the frequency range 5–20 Hz.

**Table 4.** The fractional rms of the disc (in the soft band) and the power-law (in both the soft and the hard band) components, and the fractions of the total soft band flux due to variable/constant disc/power-law emission ( $D_{\text{var}}$ ,  $P_{\text{var}}$  and  $D_{\text{const}}$ ,  $P_{\text{const}}$ ). These estimates have been obtained by fitting the broad-frequency-band (0.002–20, 0.017–20, and 0.04–20 Hz, respectively, for the low-, medium-, and high-luminosity observations) covariance spectra of GX 339-4 and its *XMM-Newton* energy spectra with the model described in Section 4.1.1. All these estimates have been derived considering the absorbed fluxes. Errors are reported at 90 per cent confidence level.

	Low $L$	Med $L$	High $L$
Disc (0.5–1.5 keV)			
Fractional rms	$0.65 \pm 0.20$	$0.40 \pm 0.03$	$0.20 \pm 0.03$
$D_{\text{var}}$	$0.17 \pm 0.03$	$0.15 \pm 0.01$	$0.09 \pm 0.01$
$D_{\text{const}}$	$0.09 \pm 0.08$	$0.24 \pm 0.01$	$0.36 \pm 0.02$
Power law (0.5–1.5 keV)			
Fractional rms	$0.68 \pm 0.02$	$0.47 \pm 0.01$	$0.34 \pm 0.01$
$P_{\text{var}}$	$0.51 \pm 0.04$	$0.29 \pm 0.01$	$0.19 \pm 0.01$
$P_{\text{const}}$	$0.23 \pm 0.03$	$0.32 \pm 0.01$	$0.37 \pm 0.01$
Power law (2–9 keV)			
Fractional rms	$0.59 \pm 0.02$	$0.42 \pm 0.01$	$0.33 \pm 0.01$




**Figure 7.** The best-fitting peak frequencies of the high- and low-frequency Lorentzians. The colours refer to the energy bands: soft (black), hard (red), and very hard (blue). The grey points are the frequencies of the high- and low-frequency Lorentzian components as obtained by Belloni et al. (2005) from the analysis of all the hard-state *RXTE* observations of GX 339-4 during the 2002/2003 outburst. The dashed curve is a linear best fit to the Belloni et al. (2005) data points, while the dotted curve is the same linear model (with only the intercept left free to vary) fit to the soft band data points derived in this paper.

Belloni et al. 2002)

$$\nu = \sqrt{\nu_0^2 + \Delta^2},$$

where  $\nu_0$  is the centroid frequency and  $\Delta$  is the half width at half-maximum of the Lorentzian. Note that flat-top, broad-band noise components are best fit by Lorentzians with  $\nu_0$  consistent with zero. This is the case for most of our fits. However, in some cases, the high-frequency Lorentzian fit required a more peaked profile, with  $\nu_0$  different from zero. The error on the peak frequency is computed from the 90 per cent confidence contours of the  $\nu_0$  and  $\Delta$  parameters. Both  $\nu_l$  and  $\nu_h$  show the characteristic shift towards high frequencies as a function of luminosity. While  $\nu_l$  shifts by a factor of  $\sim 20$ ,  $\nu_h$  shifts by a factor of  $\sim 1.6$ , for a variation in luminosity of a factor



**Figure 8.** The scaling as a function of  $L/L_{\text{Edd}}$  (where  $L$  is the 3–30 keV luminosity) of (a) the peak frequency of the high-frequency Lorentzian,  $\nu_h$ ; (b) the normalization of the PSD at the peak frequency of the high-frequency Lorentzian,  $N_{\nu_h}$ ; (c) the high-frequency (5–20 Hz) fractional rms. The dashed lines are the best-fitting models. In the plots, the colours refer to the energy bands: soft (black), hard (red), very hard (blue).

$\sim 10$ . The black dots show that this same shift is observed also in the soft energy band. In addition we observe that  $\nu_h$  is systematically and significantly (at  $>3\sigma$  confidence level) offset towards lower frequencies in the soft band. Note that this offset is driven by a variation of  $\Delta$ , indicating that the high-frequency Lorentzian is systematically narrower in the soft band, than at higher energies. It is also worth noting that, if the high-frequency PSDs (e.g. 1.5 times above  $\nu_h$ ) are fitted with a simple power law, a significant steepening of the slope is observed in the soft band of the high-luminosity observation. This resembles the behaviour observed in AGN in the disc-dominated optical band (e.g. Mushotzky et al. 2011). On the other hand,  $\nu_l$  is consistent with remaining constant among the different energy bands of the same observation. Indeed, for each luminosity, the low-frequency Lorentzians in the three energy bands are co-aligned in the plots of Fig. 2, while an offset of  $\nu_h$  in the soft band is clearly observed (see also Fig. 8a and Section 4.3

for a description of this figure). Note that the peak frequency of the additional third Lorentzian does not show any significant energy-dependence (see also Table 2). These results suggest that the high-frequency PSD behaves in a different manner in the soft band than at higher energies. This aspect will be explored in more detail in Section 4.3.

In order to compare our results with those reported in the literature, we overplot in Fig. 7 (grey points) the characteristic frequencies associated with the high- and low-frequency Lorentzians (in the energy band  $E = 3.8\text{--}15.3$  keV) as obtained by Belloni et al. (2005) from the analysis of all the *RXTE* observations of the 2002/2003 outburst of GX 339-4 (only the values relative to the hard state of the source are shown in the plot). The two main Lorentzians of the hard- and very hard-band PSDs analysed in this paper (red and blue points in Fig. 7) follow the same trend and span the same range of characteristic frequencies as those of the 2002/2003 *RXTE* data. This trend seems to characterize also the Lorentzians in the soft-band PSD, despite the significant offset. To show that the correlation between  $\nu_l$  and  $\nu_h$  is preserved in the soft band, we fit the Belloni et al. (2005) data points with a linear model (in the log-linear plot of Fig. 7 the model is represented by the dashed curve). Then we fixed the slope to the best-fitting value, and letting the intercept free to vary, we fit the soft band data points (dotted curve in Fig. 7). Though the number of data points is small, we obtain a very good description of the  $\nu_l$  versus  $\nu_h$  trend in the soft band, thus in agreement with the idea that the two Lorentzians are linked by the same linear relation as observed at high energies. According to these fits, the offset of  $\nu_h$  as measured in the soft band and at higher energies is of a factor  $\sim 2$ .

### 4.3 The high-frequency PSD of GX 339-4

We now focus on the high-frequency PSD of GX 339-4 (between 5 and 20 Hz, i.e. above the peak of the high-frequency Lorentzian). In the analysed data sets we observe that the peak of the high-frequency Lorentzian varies with luminosity by a factor of  $\sim 1.6$  (Section 4.2), as also shown in Fig. 8(a), where we plot  $\nu_h$  as a function of  $L/L_{\text{Edd}}$  (note that  $L$  is computed in the 3–30 keV range). A fit with a linear model in the log–log space returns a best-fitting slope of  $0.26 \pm 0.11$ ,  $0.17 \pm 0.06$ , and  $0.19 \pm 0.16$ , respectively, in the soft, hard, and very hard band. We note that these slopes are consistent with being the same, but consistently and significantly different from a slope of 1. However the soft band has a significantly lower intercept (differing by a factor  $\sim 0.7 \pm 0.2$ ), this being a consequence of the energy dependence of  $\nu_h$  (Section 4.2).

We also looked at the variations of the normalization of the PSD at the peak frequency of the high-frequency Lorentzian ( $N_{\nu_h}$ , in units of  $\text{Hz}^{-1}$ ) as a function of luminosity. We derived  $N_{\nu_h}$  and its related uncertainties (at the 90 per cent confidence level) from the best-fitting PSD models. Results are shown in Fig. 8(b). A fit with a linear model in the log–log space gives best-fitting slopes of  $-0.81 \pm 0.13$ ,  $-0.52 \pm 0.08$ , and  $-0.42 \pm 0.20$ , respectively, in the soft, hard, and very hard band (dashed lines in Fig. 8b). These results point to the existence of an inverse-power scaling relation between  $N_{\nu_h}$  and  $L/L_{\text{Edd}}$ . Once again we note a consistent behaviour between the hard, and very hard bands, and a significant deviation in the soft band. Indeed, in the soft band this trend appears steeper than in the other bands.

Finally, we computed the fractional rms in the 5–20 Hz frequency range. Results are reported in Table 3 and plotted in Fig. 8(c). Despite the significant variations of the high-frequency Lorentzian as

a function of luminosity, the high-frequency fractional rms (above the peak frequency of the high-frequency Lorentzian) in the hard and very hard bands is consistent with being remarkably constant. This is due to the combined effect of the frequency shift and the decrease of normalization of the high-frequency PSD with luminosity. These combined variations produce a constant high-frequency fractional rms. Indeed, fitting the data with a constant we measure a scatter around this best-fitting value of  $\sim 10$  and  $\sim 4$  per cent (respectively in the hard and very hard band), i.e. of the order of the corresponding average uncertainty (Fig. 3, middle and lower panel). On the other hand, the constancy of the high-frequency fractional rms with luminosity breaks down in the soft energy band (Fig. 3, upper panel). If fitted with a constant model, the scatter increases to  $\sim 30$  per cent. Instead, a fit with a linear model in the log–log space yields a best-fitting slope of  $-0.24 \pm 0.08$ , and the scatter around the model is of  $\sim 3$  per cent. This is due to the fact that the normalization of the high-frequency PSD decreases with luminosity faster in the soft band than at higher energies.

## 5 DISCUSSION

In this paper we investigated the luminosity dependence of the characteristic frequencies and variability power of GX 339-4 in the hard state. These properties have been extensively studied in the past, through the use of *RXTE* data, which cover energies  $> 3$  keV. Our work aims at extending this analysis down to the soft X-ray energy band ( $E \geq 0.5$  keV), where the contribution of the disc thermal emission becomes non-negligible. For this reason we used *XMM-Newton* and simultaneous *RXTE* observations, covering the energy band 0.5–30 keV. The selected observations belong to two different outbursts of the source. None the less, the variability properties of BHXRBs are extremely similar from outburst to outburst (Muñoz-Darias et al. 2011, Grinberg et al. 2014), thus justifying our approach of considering observations taken during different activity periods.

### 5.1 Disc variability and the shape of the broad-frequency-band fractional rms spectra

The drop of fractional rms that characterizes the transition from the hard to the soft state, is ascribed to the emergence of a disc component, which, in the soft state, is more intense and less variable (with typical fractional rms values of few per cent) than the power law. The fractional rms spectrum is observed to change accordingly, from an ‘inverted’ (in the hard state) to a ‘hard’ (in the soft state) spectral shape.

Extending our analysis of the hard-state variability properties of GX 339-4 down to the soft X-ray band, we observe a drop of fractional rms that correlates with the progressive increase of disc flux relative to the total flux in that band. This resembles the behaviour seen in soft states, although to a much lower extent (indeed the disc fractional rms is much higher and the drop less intense than in soft states, see inset of Fig. 4), and is expected if the disc has lower fractional rms than the power law. This raises the question as to whether or not the disc in the hard state has similar timing properties as in the soft state. Analysing the energy and covariance spectra of GX 339-4, we observe intrinsic broad-frequency-band disc variability at all the luminosities spanned by the hard-state observations. The presence of a variable disc in a hard state observation (corresponding to our medium luminosity observation) was first reported by Wilkinson & Uttley (2009). We detect high

levels of disc variability ( $\sim 20$ – $65$  per cent) in the three observations, comparable to the variability intrinsic to the power-law component, and significantly higher than observed in soft states. However, the disc fractional rms decreases with luminosity faster than the power-law fractional rms. At the same time, the fraction of soft band flux due to the disc (power law) component increases (decreases) with luminosity (Fig. 5). The net effect is a strong increase (a factor of  $\sim 4$ ) of constant disc-flux fraction in the soft band, which causes the observed drop of fractional rms. It is worth noting that, without the contribution from the disc component, the fractional rms spectra always show an ‘inverted’ shape (as typically seen in the *RXTE* data; e.g. Belloni et al. 2011), with variability power decreasing as a function of energy. We conclude that a relatively high fraction of disc flux is variable during the hard state of GX 339-4. However, the increasing fraction of constant disc flux with luminosity in the soft band is responsible for the net drop of fractional rms as compared to power-law-dominated energy bands. We argue that this trend continues throughout the outburst, thus leading to the disc being mostly constant during the soft state, and significantly different from the disc seen in hard states.

## 5.2 The scaling of the high-frequency PSD with luminosity in BHXRBs and AGN

A number of correlations link the high-frequency variability properties of BHXRBs and AGN, suggesting that the dominant variability process at these frequencies is the same in both the classes of objects. In particular, considering a sample of 10 AGN and 2 BHXRBs in the soft state, McHardy et al. (2006) found a linear scaling of the high-frequency break (a characteristic frequency of AGN PSDs, Edelson & Nandra 1999, commonly associated with  $\nu_h$  in BHXRBs) with the BH mass ( $M_{\text{BH}}$ ) and the mass-accretion rate ( $\dot{m}_{\text{Edd}}$ , i.e. normalized by the mass-accretion rate required to reach the Eddington luminosity), which defines the so-called ‘variability plane’. Koerding et al. (2007) found a similar scaling relation also for sources in the hard state (but offset with respect to the corresponding relation valid for soft-state sources).

As a consequence, we would expect to observe  $\nu_h$  scaling linearly with the mass-accretion rate also in the hard-state observations of GX 339-4 analysed in this paper. However, we see  $\nu_h$  scaling with  $L/L_{\text{Edd}}$  (with  $L = L_{3-30\text{keV}}$ ), with an average slope of  $\sim 0.21 \pm 0.11$ , therefore not consistent with the results by McHardy et al. (2006) and Koerding et al. (2007).<sup>3</sup> We note, however, that our result is in agreement with the recent estimate obtained from the PSD analysis of a sample of 104 nearby AGN observed with *XMM-Newton* (Gonzalez-Martín & Vaughan 2012), which reports a best-fitting slope of  $0.24 \pm 0.28$ . This therefore suggests that, indeed, the same engine is at work in both AGN and BHXRBs, but that the dependence of the high-frequency variability on luminosity/mass-accretion is significantly shallower than previously thought (McHardy et al. 2006; Koerding et al. 2007). We also investigated the dependence of the normalization of the PSD at the break,  $N_{\nu_h}$ , on the luminosity. We find that  $N_{\nu_h}$  decreases as a function of  $L/L_{\text{Edd}}$ . Interestingly, this trend can counterbalance the  $\nu_h$  versus  $L/L_{\text{Edd}}$  scaling, producing the same high-frequency variability power at different luminosities.

<sup>3</sup> Here we assume  $L_{\text{bol}} \propto L$  and  $L_{\text{bol}}/L_{\text{Edd}} \propto \dot{m}_{\text{Edd}}^\beta$ , with  $\beta = 1$ – $2$ , the exact value depending on the accretion efficiency of the flow (Fender & Muñoz-Darias 2015).

Previous studies have pointed out that the high-frequency PSD does not show a clear scaling with the luminosity/mass-accretion rate both in BHXRBs (Gierliński et al. 2008) and AGN (O’Neill et al. 2005; Zhou et al. 2010; Ponti et al. 2012). We observe that the high-frequency fractional rms of GX 339-4 is constant in the hard and very hard band (dominated by the hard X-ray Comptonization continuum) at any luminosity probed. In particular we find that this quantity has a small scatter (4–10 per cent, in agreement with results by Muñoz-Darias et al. 2011 and by Heil, Vaughan & Uttley 2012) in these two energy bands. The remarkable constancy of the high-frequency PSD provides us with a tool to measure the BH mass of accreting objects (Nikołajuk, Papadakis & Czerny 2004; Gierliński et al. 2008; Kelly, Sobolewska & Siemiginowska 2011; Ponti et al. 2012; La Franca et al. 2014). Indeed the high-frequency fractional rms (or equivalently the ‘excess variance’, e.g. Nikołajuk et al. 2004; Ponti et al. 2012), while being independent of the mass-accretion rate, scales inversely with the BH mass, with a very small scatter (which mostly depends on the uncertainty on  $M_{\text{BH}}$ ), of the order of less than 0.2–0.4 dex (e.g. O’Neill et al. 2005; Zhou et al. 2010; Kelly et al. 2011; Ponti et al. 2012).

As observed in Section 4.3, the dependences of  $\nu_h$  and  $N_{\nu_h}$  are significantly different in the soft band. In particular, the constancy of the high-frequency fractional rms as a function of luminosity breaks down at low energies, with a scatter around the best-fitting constant model increasing to  $\sim 30$  per cent. We ascribe this to the presence of the additional contribution from the disc component. Interestingly the same behaviour is observed in AGN. Indeed, it is observed that the scatter in the relation between the short-time-scale variability and the BH mass is larger in the optical band (dominated by the disc emission) than in the X-ray band (dominated by the power law), and the optical variability of AGN is significantly anticorrelated with the optical luminosity (Kelly et al. 2013).

## 5.3 A possible interpretation for the offset of the high-frequency Lorentzian in the soft band

The characteristic frequencies in the PSD of accreting systems (i.e. the peak frequency of each Lorentzian) are likely related to fundamental time-scales of the accretion flow at specific radii (e.g. Psaltis, Belloni & van der Klis 1999; Churazov, Gilfanov & Revnivtsev 2001; Motta et al. 2014). Though not yet conclusive, the general picture that emerged over the years is that the low-frequency Lorentzian in the hard state of BHXRBs is associated with a characteristic time-scale (most probably the viscous time-scale) at the inner edge of the accretion disc (e.g. Churazov et al. 2001; Done, Gierliński & Kubota 2007). According to recent studies, variations intrinsic to the seed-photons flux produce the long-time-scale variability (i.e. the low-frequency Lorentzian) observed in the soft energy band (Section 4.1.1; Wilkinson & Uttley 2009; Uttley et al. 2014). These variations are due to perturbations in the accretion rate, which travel inwards and modulate the hard X-ray emission from the inner corona, thus producing the long-time-scale variability observed also in harder energy bands (e.g. Lyubarskii 1997; Kotov, Churazov & Gilfanov 2001; Arévalo & Uttley 2006; Ingram & van der Klis 2013). The existence of additional variability components at frequencies higher than those corresponding to the inner edge of the disc, can be explained provided the corona extends down to the innermost stable circular orbit (Motta et al. 2014) and is intrinsically variable on short time-scales (e.g. Churazov et al. 2001). This inner hot flow/corona is thus responsible for producing the high-frequency Lorentzian observed in hard and very hard energy bands.

Such a high-frequency component is observed also in the soft band, and is linearly correlated with that in harder energy bands (as seen from covariance spectra, light grey triangles in Fig. 6; see also Uttley et al. 2011; De Marco et al. 2015). We showed that the high-frequency Lorentzian is significantly offset, by a factor of  $\sim 2$ , towards lower frequencies in the soft band with respect to higher-energy bands. The covariance spectra (light grey triangles in Fig. 6) show that the high-frequency PSD contains significant contribution from variable disc emission, so the offset, soft band high-frequency Lorentzian is likely linked to variability coming from the disc component. However, within the scenario depicted above, intrinsic disc variability due to accretion rate perturbations occurring on time-scales shorter than the viscous time-scale at the inner edge of the disc is expected to be highly suppressed. Thus, a possible explanation is that this high-frequency variability is not intrinsic to the disc, but results from reprocessing of the short-time-scale variable corona emission (e.g. thermal reprocessing; Wilkinson & Uttley 2009; Uttley et al. 2011; De Marco et al. 2015). In standard disc-corona geometries, the hard X-rays would irradiate a relatively *extended* region of the disc. Thus, the reprocessing region would act as a low-pass filter, damping the high-frequency variability power in the soft band (e.g. Arévalo & Uttley 2006; Uttley et al. 2014). As a consequence the soft band high-frequency Lorentzian would be narrower, and its peak frequency would result slightly shifted towards low frequencies with respect to the corresponding peak in harder energy bands. This resembles the behaviour observed in the data of GX 339-4. According to this interpretation, the relative offset of the peak frequency of the high-frequency Lorentzian in the different bands should trace the variations of the geometry of the accretion disc/corona during the outburst.

## ACKNOWLEDGEMENTS

This work is based on observations obtained with *XMM-Newton*, an ESA science mission with instruments and contributions directly funded by ESA Member States and NASA. GP acknowledges support via the Bundesministerium für Wirtschaft und Technologie/Deutsches Zentrum für Luft und Raumfahrt (BMWi/DLR, FKZ 50 OR 1408) and the Max Planck Society. This project was funded in part by European Research Council Advanced Grant 267697 4- $\pi$ -sky: Extreme Astrophysics with Revolutionary Radio Telescopes. TMD acknowledges support by the Spanish Ministerio de Economía y competitividad (MINECO) under grant AYA2013-42627. The authors thank the anonymous referee for helpful comments which significantly improved the paper.

## REFERENCES

Arévalo P., Uttley P., 2006, *MNRAS*, 367, 801  
 Belloni T., Psaltis D., van der Klis M., 2002, *ApJ*, 572, 392  
 Belloni T., Homan J., Casella P., van der Klis M., Nespoli E., Lewin W. H. G., Miller J. M., Méndez M., 2005, *A&A*, 440, 207  
 Belloni T. M., Motta S. E., Muñoz-Darias T., 2011, *Bull. Astron. Soc. India*, 39, 409  
 Böck M. et al., 2011, *A&A*, 533, 8  
 Cassatella P., Uttley P., Maccarone T. J., 2012, *MNRAS*, 427, 2985  
 Churazov E., Gilfanov M., Revnivtsev M., 2001, *MNRAS*, 321, 759  
 De Marco B., Ponti G., Muñoz-Darias T., Nandra K., 2015, *ApJ*, submitted  
 Dickey J. M., Lockman F. J., 1990, *ARA&A*, 28, 215  
 Done C., Díaz Trigo M., 2010, *MNRAS*, 407, 2287  
 Done C., Gierliński M., 2005, *MNRAS*, 364, 208  
 Done C., Gierliński M., Kubota A., 2007, *A&AR*, 15, 1  
 Droulans R., Belmont R., Malzac J., Jourdain E., 2010, *ApJ*, 717, 1022

Edelson R., Nandra K., 1999, *ApJ*, 514, 682  
 Fender R., Muñoz-Darias T., 2015, in Haardt et al., eds, *Lecture Notes in Physics, Astrophysical Black Holes*. Springer, Berlin, preprint (arXiv:1505.03526)  
 Gierliński M., Zdziarski A. A., 2005, *MNRAS*, 363, 1349  
 Gierliński M., Nikolajuk M., Czerny B., 2008, *MNRAS*, 383, 741  
 Gonzalez-Martín O., Vaughan S., 2012, *A&A*, 544, 80  
 Grinberg V. et al., 2014, *A&A*, 565, 1  
 Guainazzi M., 2014, *XMM-CCF-REL-312*, available at: <http://xmm2.esac.esa.int/docs/documents/CAL-SRN-0312-1-4.pdf>  
 Guainazzi M., Smith M., 2013, *XMM-CCF-REL-302*, available at: <http://xmm2.esac.esa.int/docs/documents/CAL-SRN-0302-1-5.pdf>  
 Guainazzi M., Kirsch M., Haberl F., de Juan Ovelar M., Rodríguez-Pascual P., Smith M., Carpano S., 2014, *XMM-SOC-CAL-TN-0083*, available at: <http://xmm.vilspa.esa.es/docs/documents/CAL-TN-0083.pdf>  
 Heil L. M., Vaughan S., Uttley P., 2012, *MNRAS*, 422, 2620  
 Hynes R. I., Steeghs D., Casares J., Charles P. A., O'Brien K., 2003, *ApJ*, 583, 95  
 Hynes R. I., Steeghs D., Casares J., Charles P. A., O'Brien K., 2004, *ApJ*, 609, 317  
 Ingram A., van der Klis M., 2013, *MNRAS*, 434, 1476  
 Jahoda K., Markwardt C. B., Radeva Y., Rots A. H., Stark M. J., Swank J. H., Strohmayer T. E., Zhang W., 2006, *ApJS*, 163, 401  
 Kalamkar M., van der Klis M., Heil L., Homan J., 2015, *ApJ*, 808, 144  
 Kalamkar M., van der Klis M., Uttley P., Altamirano D., Wijnands R., 2013, *ApJ*, 766, 89  
 Kelly B. C., Sobolewska M., Siemiginowska A., 2011, *ApJ*, 730, 52  
 Kelly B. C., Treu T., Malkan M., Pancoast A., Woo J.-H., 2013, *ApJ*, 779, 187  
 Koerding E. G., Migliari S., Fender R., Belloni T., Knigge C., McHardy I., 2007, *MNRAS*, 380, 301  
 Kolehmainen M., Done C., Díaz Trigo M., 2014, *MNRAS*, 437, 316  
 Kong A. K. H., Kuulkers E., Charles P. A., Homer L., 2000, *MNRAS*, 312, 49  
 Kotov O., Churazov E., Gilfanov M., 2001, *MNRAS*, 327, 799  
 La Franca F., Bianchi S., Ponti G., Branchini E., Matt G., 2014, *ApJ*, 787, 12  
 Lyubarskii Y. E., 1997, *MNRAS*, 285, 604  
 McHardy I. M., Koerding E., Knigge C., Uttley P., Fender R. P., 2006, *Nature*, 444, 730  
 Markoff S., Nowak M. A., Wilms J., 2005, *ApJ*, 635, 1203  
 Miyamoto S., Kimura K., Kitamoto S., Dotani T., Ebisawa K., 1991, *ApJ*, 383, 784  
 Miyamoto S., Kitamoto S., Iga S., Negoro H., Terada K., 1992, *ApJ*, 391, 21  
 Motta S., Belloni T. M., Stella L., Muñoz-Darias T., Fender R., 2014, *MNRAS*, 437, 2554  
 Muñoz-Darias T., Casares J., Martínez-Pais I. G., 2008, *MNRAS*, 385, 2205  
 Muñoz-Darias T., Motta S., Belloni T. M., 2011, *MNRAS*, 410, 679  
 Mushotzky R. F., Edelson R., Baumgartner W., Gandhi P., 2011, *ApJ*, 743, 12  
 Nandra K., George I. M., Mushotzky R. F., Turner T. J., Yaqoob T., 1997, *ApJ*, 476, 70  
 Nikolajuk M., Papadakis I. E., Czerny B., 2004, *MNRAS*, 350, 26  
 Nowak M. A., 2000, *MNRAS*, 318, 361  
 Nowak M. A., Vaughan B. A., Wilms J., Dove J. B., Begelman M. C., 1999a, *ApJ*, 510, 874  
 Nowak M. A., Wilms J., Dove J. B., 1999b, *ApJ*, 517, 355  
 O'Neill P. M., Nandra K., Papadakis I. E., Turner T. J., 2005, *MNRAS*, 358, 1405  
 Plant D. S., Fender R. P., Ponti G., Muñoz-Darias T., Coriat M., 2015, *A&A*, 573, 120  
 Ponti G., Cappi M., Dadina M., Malaguti G., 2004, *A&A*, 417, 451  
 Ponti G., Papadakis I., Bianchi S., Guainazzi M., Matt G., Uttley P., Bonilla N. F., 2012, *A&A*, 542, 83  
 Pottschmidt K. et al., 2003, *A&A*, 407, 1039  
 Psaltis D., Belloni T., van der Klis D., 1999, *ApJ*, 520, 262  
 Remillard R. A., McClintock J. E., 2006, *ARA&A*, 44, 49

- Strüder L. et al., 2001, *A&A*, 365, 18  
Tomsick J. A. et al., 2008, *ApJ*, 680, 593  
Uttley P., Wilkinson T., Cassatella P., Wilms J., Pottschmidt K., Hanke M., Böck M., 2011, *MNRAS*, 414, 60  
Uttley P., Cackett E. M., Fabian A. C., Kara E., Wilkins D. R., 2014, *A&AR*, 22, 72  
Vaughan S., Edelson R., Warwick R. S., Uttley P., 2003, *MNRAS*, 345, 1271  
Wilkinson T., Uttley P., 2009, *MNRAS*, 397, 666  
Zdziarski A. A., Poutanen J., Mikołajewska J., Gierliński M., Ebisawa K., Johnson W. N., 1998, *MNRAS*, 301, 435  
Zdziarski A. A., Gierliński M., Mikołajewska J., Wardziński G., Smith D. M., Harmon B. A., Kitamoto S., 2004, *MNRAS*, 351, 791  
Zhang W., Jahoda K., Swank J. H., Morgan E. H., Giles A. B., 1995, *ApJ*, 449, 930  
Zhang W., Morgan E. H., Jahoda K., Swank J. H., Strohmayer T. E., Jernigan G., Klein R. I., 1996, *ApJ*, 469, 29  
Zhou X.-L., Zhang S.-N., Wang D.-X., Zhu L., 2010, *ApJ*, 710, 16

This paper has been typeset from a  $\text{\TeX/L\AA\TeX}$  file prepared by the author.