



Publication Year	2019
Acceptance in OA	2020-12-29T11:20:35Z
Title	Unveiling the Galaxy Cluster - Cosmic Web Connection with X-ray observations in the Next Decade
Authors	Walker, Stephen, Nagai, Daisuke, Simionescu, A., Markevitch, M., Akamatsu, H., Arnaud, M., Avestruz, C., Bautz, M., BIFFI, Veronica, BORGANI, STEFANO, Bulbul, E., Churazov, E., Dolag, K., Eckert, D., ETTORI, STEFANO, Fujita, Y., GASPARI, MASSIMO, Ghirardini, V., Kraft, R., Lau, E. T., Mantz, A., Matsushita, K., McDonald, M., Miller, E., Mroczkowski, T., Nulsen, P., Okabe, N., Ota, N., Pointecouteau, E., Pratt, G., Sato, K., Shi, X., Tremblay, G., Tremmel, M., VAZZA, FRANCO, Zhuravleva, I., Zinger, E., ZuHone, J.
Handle	http://hdl.handle.net/20.500.12386/29271
Journal	BULLETIN OF THE AMERICAN ASTRONOMICAL SOCIETY
Volume	51

Astro2020 Science White Paper

Unveiling the Galaxy Cluster – Cosmic Web Connection with X-ray observations in the Next Decade

- Thematic Areas:**
- Planetary Systems
 - Star and Planet Formation
 - Formation and Evolution of Compact Objects
 - Cosmology and Fundamental Physics
 - Stars and Stellar Evolution
 - Resolved Stellar Populations and their Environments
 - Galaxy Evolution
 - Multi-Messenger Astronomy and Astrophysics

Principal Authors:

Name: Stephen A. Walker¹, Daisuke Nagai²
Institution: 1) NASA GSFC, 2) Yale University,
Email: stephen.a.walker@nasa.gov; daisuke.nagai@yale.edu
Phone: +1 (301) 286-9882; +1 (203) 432-5370

Co-authors: A. Simionescu (SRON), M. Markevitch (NASA GSFC), H. Akamatsu (SRON), M. Arnaud (CEA), C. Avestruz (U.Chicago), M. Bautz (MIT), V. Biffi (CfA), S. Borgani (UniTS/INAF), E. Bulbul (CfA), E. Churazov (MPA), K. Dolag (USM/MPA), D. Eckert (MPE), S. Etori (INAF), Y. Fujita (Osaka), M. Gaspari (Princeton), V. Ghirardini (CfA), R. Kraft (CfA), E. T. Lau (Miami), A. Mantz (Stanford), K. Matsushita (TUS), M. McDonald (MIT), E. Miller (MIT), T. Mroczkowski (ESO), P. Nulsen (CfA), N. Okabe (Hiroshima), N. Ota (Nara), E. Pointecouteau (IRAP), G. Pratt (CEA), K. Sato (Saitama), X. Shi (SWIFAR), G. Tremblay (CfA), M. Tremmel (Yale), F. Vazza (Bologna), I. Zhuravleva (U.Chicago), E. Zinger (Heidelberg), J. ZuHone (CfA)

Abstract: In recent years, the outskirts of galaxy clusters have emerged as one of the new frontiers and unique laboratories for studying the growth of large scale structure in the universe. Modern cosmological hydrodynamical simulations make firm and testable predictions of the thermodynamic and chemical evolution of the X-ray emitting intracluster medium. However, recent X-ray and Sunyaev-Zeldovich effect observations have revealed enigmatic disagreements with theoretical predictions, which have motivated deeper investigations of a plethora of astrophysical processes operating in the virialization region in the cluster outskirts. Much of the physics of cluster outskirts is fundamentally different from that of cluster cores, which has been the main focus of X-ray cluster science over the past several decades. A next-generation X-ray telescope, equipped with sub-arcsecond spatial resolution over a large field of view along with a low and stable instrumental background, is required in order to reveal the full story of the growth of galaxy clusters and the cosmic web and their applications for cosmology.

1 Introduction

As the largest gravitationally bound structures in the universe, galaxy clusters continue to grow and accrete matter in their outskirts (see [1] for a review). The majority of the gas lies in the outskirts beyond the cluster’s virial radius (denoted by r_{200} ¹) and in the intergalactic medium (IGM) within filaments that connect clusters to the cosmic web. However, the low gas density means that these regions are extremely faint in X-rays, and at present beyond the sensitivity limits of current X-ray telescopes (Fig.1, left panel). As a result, the full story of large-scale structure formation is hidden from us. Cosmological simulations of galaxy cluster formation predict the outskirts to be a hive of activity with gas continuing to accrete, and ongoing mergers with small sub-clusters and clumps of gas (Fig.1, right panels). A plethora of unexplored structure formation physics is believed to be operating in the outskirts, and these physical processes are fundamentally different from the physics in the cores of clusters that has been the focus of X-ray cluster science over the past several decades.

The outskirts of galaxy clusters is a new territory for addressing the following outstanding questions at the crossroads of cosmology and astrophysics: **How do galaxy clusters grow? How do they connect to the cosmic web? What is the chemical composition in and around the most massive structures in the Universe?** A great leap forward in the sensitivity of X-ray telescopes is required to answer these questions.

2 How do galaxy clusters grow and connect to the cosmic web?

Our current understanding of the gas in cluster outskirts has been achieved through recent pioneering measurements of the intracluster medium (ICM) in X-rays and microwaves via the Sunyaev-Zeldovich (SZ) effect that map the hot gas distribution near the virial radius of nearby galaxy clusters. The *Suzaku* X-ray Observatory, with its low and stable background afforded by its low Earth orbit, was able to obtain temperature and density measurements out to the virial radius [e.g., 2, 3]. Combining density measurements from deep *XMM-Newton* observations of nearby bright clusters with *Planck* SZ pressure profile data has also allowed statistical constraints on the thermodynamic profiles of the outskirts ICM to be made [4, 5]. Initial results from these observations have been puzzling, as they disagree with theoretical predictions [e.g., 2–5, see also Fig.2, left panel]. In particular, the gas entropy just outside the virial radii has been observed to lie well below the predictions from purely gravitational collapse (shown as black line in the left panel of Fig.2). Future high angular resolution and large field of view (FoV) X-ray and SZ observations with high sensitivity are required to address the apparent disagreement, and to capture the rich structures in cluster outskirts predicted by modern cosmological hydrodynamical simulations (see Fig.1, right panels). This will allow us to investigate the physics of galaxy and cluster formation, especially in the unexplored territory where matter from the cosmic web accrete into galaxy clusters. Below, we highlight several transformative cluster outskirts sciences for the coming decade.

ICM Clumping: Both *Suzaku* and *XMM-Newton/Planck* measurements of the ICM outskirts have found evidence for increasing levels of gas clumping, as predicted by cosmological simulations [10–16]. Gas clumping is believed to be due to infalling gas substructures. Simulations indicate that the level of gas clumping should increase dramatically in the $(1-2)r_{200}$ region of clusters, where the infalling clumps have not yet been entirely ram-pressure stripped by the ICM. If not resolved,

¹The radius within which the enclosed total mean overdensity is 200 times the critical density of the Universe.




Figure 1: *Left*: Hydrodynamical cosmological simulation of large-scale structure formation, showing that galaxy clusters reside at the nodes of IGM filaments [6]. With *Chandra* and *XMM-Newton*, only the central regions of clusters can be explored in detail (inner white circle). To begin to explore large-scale structure in its entirety (outer white contours) requires at least a factor of 50 improvement in sensitivity to low surface brightness extended emission. *Right*: Simulated X-ray mosaic of the RomulusC cluster [7], a low mass ($10^{14} M_{\odot}$) cluster at $z = 0.05$, comparing 100ks deep coverage from *Chandra* (top) and *AXIS/Lynx* (bottom). The mock *Chandra* simulation includes background point sources from the cosmic X-ray background. Due to the high background and small collecting areas of current X-ray telescopes, we are only able to see the brightest X-ray emission in the cluster cores. The bulk of the ICM in the outskirts beyond the virial radius (dashed white circle), and the IGM filaments that connect clusters together, remain hidden from view.

clumping will bias gas profile measurements by overestimating the gas density and gas mass fraction, and underestimating the gas temperature, leading to biases in hydrostatic mass estimates. Correcting for clumping will help explain the lower observed entropy compared to predictions, the excess baryon fraction beyond the cosmic value, and help correct biases in hydrostatic mass. Detecting and resolving gas clumps, that are associated with infalling galaxies in the outskirts of clusters, is also crucial for gaining insights into the stripping of their circumgalactic medium (CGM) through their interactions with the ICM, quenching of cluster galaxies, and chemical enrichment process in the outskirts of galaxy clusters (see §3). Current X-ray telescopes are unable to resolve faint gas clumps that contribute to the bulk of clumping (Fig. 1, right), and their physical properties are almost completely unexplored. Future X-ray observations with high sensitivity and high spatial resolution, such as *AXIS/Lynx*, will allow routine detection of clumps and open up a new window for studying the physical processes of galaxy and cluster formation and evolution.

Bulk & Turbulent Gas Motions: Gas motions in cluster outskirts are generated from mergers and accretion during cluster formation. Thus the amount of these gas motions is a direct probe of the cluster’s dynamical state. Cosmological hydrodynamical simulations predict that such non-thermal




Figure 2: *Left panel*: Disagreement between the observed entropy profiles in X-ray from Suzaku (coloured points) and the entropy profile predicted by gravitational collapse (solid black line, [8]) just beyond the virial radius (highlighted by the red oval) [1]. *Right panel*: Simulated entropy map of a galaxy cluster, illustrating filamentary streams of high-density, low-entropy gas penetrating deep into the cluster interior [9].

pressure becomes more significant with radius in the density stratified ICM, reaching $\sim 50\%$ of the thermal gas pressure at $r \approx 1.5r_{200}$ [17, 18]. Such non-thermal pressure is believed to be responsible for biasing hydrostatic cluster masses at a level of 10% – 30% [19–24] by providing extra support against gravity. Upcoming high-resolution spectral X-ray observations with *Athena* and *Lynx* will enable us to directly measure the level of bulk and turbulent gas motions in cluster outskirts through Doppler shifting and broadening of the X-ray emission lines from the ICM [25, 26, see also the white paper by Bulbul et al.], allowing us to fully account for the hydrostatic mass bias when combined with gravitational lensing masses from optical surveys such as *WFIRST* and *Euclid* [27]. High-resolution X-ray and SZ spectral imaging observations will also provide complementary constraints on the level of non-thermal pressure support [28] and ICM microphysics [29] in the outskirts of galaxy clusters.

Filamentary gas streams: Cosmological simulations predict that high density, low entropy gas in the cosmic web filaments can penetrate deep into the cluster interior [9, see Fig.2]. These filamentary streams transport the warm-hot gas from the outskirts to the cluster core regions, stirring up the gas, generating bulk and turbulent gas motions, and producing shocks and contact discontinuities generated by the interaction of the cold penetrating stream with the surrounding hot ICM [30]. These streams can deposit energy, mass, and enrich the ICM as they break up and mix with the surrounding gas via fluid instabilities. Direct observations of these streams are not possible at present as these low entropy, low temperature streams are predicted to be very X-ray faint. High spatial resolution X-ray observations with *AXIS/Lynx*, with large azimuthal and radial coverage (from the cluster centers to large radii) will enable us to identify the transformation from cosmic web gas filaments in the outskirts into penetrating gas streams in the cluster interior, providing direct evidence for cosmic web filaments feeding the growth of galaxy clusters.

Thermodynamics of accretion shocks: The outer boundary of the ICM is defined by the accretion shock. This is an important location where cool-warm gas accreting from the cosmic web is shock

heated to very high temperature, generating entropy via high Mach number ($M \approx 10 - 100$) accretion shocks [31–33]. Detecting accretion shocks around clusters and measuring their strength will provide the first direct evidence of the primary physical process that defines the fate of the hot, diffuse baryons that make up more than half of the normal matter in the local Universe.

Non-equilibrium electrons: In the low-density region in the outskirts of galaxy clusters, the collision rate of electrons and protons becomes longer than the age of the universe, potentially causing the electron temperature to be lower than the ion temperature, especially in more massive and less relaxed systems [34, 35].² X-ray and SZ measurements of galaxy cluster outskirts, which probe the electron densities and temperatures of the ICM, therefore hold promise for shedding new light on the thermalization process operating during the cluster formation, and probing for the first time a regime where the physics of non-equilibrium plasmas beyond the simple hydrodynamics approximation becomes important in the ICM.

3 How do metals spread in galaxy clusters & the cosmic web?

Galaxy clusters, with their deep potential wells, are ‘closed-box’ systems ideal for studying galaxy formation physics. In particular, metallicity in the outskirts of galaxy clusters is a powerful probe of feedback physics [36–38], as the metal distribution in the ICM is strongly dependent on the chemical enrichment histories. Feedback from active galactic nuclei (AGN) at early times ($z \gtrsim 2$) is found to be effective at removing pre-enriched gas from galaxies, spreading metals uniformly throughout the cluster outskirts (see the solid black curve in Fig.3, left). On the other hand, late-time enrichment leads to inhomogeneous distributions of ICM metals that rapidly decline with cluster-centric radius (red curve in Fig.3, left). The relative composition of various elements of the ICM originating from different types of supernova explosions (core-collapse versus Ia) also place constraints on the star formation histories and the chemical evolution of the universe [39].

Measurements of metal abundance in the outskirts are extremely challenging, and are currently limited to within half the virial radius for *Chandra* and *XMM-Newton*. With *Suzaku* only very few clusters have measurements reaching the virial radius (blue crosses in Fig.3 from [40]) which also come with large uncertainties. With the high angular resolution and high sensitivity observations of *AXIS* and *Lynx*, we will be able to accurately map the metal abundance out to $2r_{200}$ for the first time, heralding a new era in our understanding of the metal enrichment of the ICM and IGM (Fig.3, left).

4 Key advances in observation capability required

Over the next decade, and into the 2030s, transformative advances in the spectral resolution (*XRISM* [41] and *Athena*) and the grasp (*eROSITA*[42, 43]) of X-ray observatories will be made, allowing us to continue the rapid advancement in the field of cluster outskirts [1, 44]. However all of these missions are fundamentally limited by their large PSFs. An enormous gap in our science capability will remain due to the lack of high spatial resolution, high sensitivity X-ray imaging. Such capability is also vital to allow us to fully exploit improvements in the resolution and sensitivity of SZ observations [45–47, for a review and white papers]. In order to probe the diffuse

²When an electron-ion plasma passes through the accretion shock, most of the kinetic energy goes into heating heavier ions, causing $T_i \gg T_e$. After the shock, electrons and ions equilibrate via Coulomb interactions over an electron-ion equilibration timescale, t_{ei} .




Figure 3: *Left*: Simulated metallicity profiles beyond the virial radius with stellar and AGN feedback (black solid curve) and only stellar feedback without AGN (red dashed curve), highlighting AGN feedback plays an important role in elevating the metallicity level in clusters outskirts [37]. Suzaku X-ray measurements are shown in blue. Simulated data from *AXIS/Lynx* are shown as the black crosses for high and low metallicity scenarios, showing that highly accurate metallicity measurements can be made out to $1.8r_{200}$. *Right*: The ratio of sky signal to detector background, which determines the sensitivity to low surface brightness emission, for major X-ray imaging instruments (blue and red are future and proposed, respectively). A is effective area at 1 keV, B is detector background (assuming 1 at low-Earth orbit and 4 at high-Earth orbit for simplicity) and F is the focal length. *Lynx* and *AXIS* are >50 times more sensitive than *Chandra*.

gas in the outskirts of galaxy clusters and the cosmic web, an X-ray telescope is required which satisfies 3 key criteria:

- An effective area to background ratio at least $50\times$ better than *Chandra's* must be combined with a low and stable particle background level. The orbits of *Chandra* and *XMM-Newton* are ill suited to this. A low-Earth orbit, placing the telescope within the protection of the Earth's magnetic field, provides the lowest and most stable particle background. An orbit around L1 or L2 also promises to provide a stable particle background.
- Subarcsecond spatial resolution must be achieved across a large FoV with low levels of vignetting. This is required to ensure background AGN in the cosmic X-ray background (CXB) can be resolved and removed throughout the FoV. As shown in the top-right panel of Fig.1, our ability to detect faint diffuse X-ray emission is limited by our ability to resolve out AGN point sources. Furthermore, gas clumps associated with infalling galaxies and groups are either too faint to be resolved directly, or impossible to distinguish from background AGN due to low photon counts. A telescope is needed which can resolve these faint gas clumps as extended objects to separate them from background AGN.
- A large FoV with low vignetting is required to allow the large areas subtended by clusters to be explored within a feasible exposure time. For a massive ($M_{200} = 10^{15}M_{\odot}$) cluster at an intermediate redshift of 0.1, the virial radius is around 20 arcmins. To allow the cluster outskirts to be revealed in their entirety with reasonable exposure times, the size of the imaging detector therefore needs to be at least 20×20 arcmins, with low vignetting.

The imaging detectors on *AXIS/Lynx* fulfill all the criteria listed above (see also Fig. 3, right). With these new instruments, we can measure the thermodynamical properties and chemical composition of the bulk of the ICM up to $2r_{200}$ (8 times more volume than in the best current studies) and into the filamentary cosmic web.

References

- [1] Stephen Walker, Aurora Simionescu, Daisuke Nagai, Nobuhiro Okabe, Dominique Eckert, Tony Mroczkowski, et al. The Physics of Galaxy Cluster Outskirts. *Space Science Reviews*, 215:7, Jan 2019. [arXiv:1810.00890](#), [doi:10.1007/s11214-018-0572-8](#).
- [2] A. Simionescu, S. W. Allen, A. Mantz, N. Werner, Y. Takei, R. G. Morris, et al. Baryons at the Edge of the X-ray-Brightest Galaxy Cluster. *Science*, 331:1576, March 2011. [arXiv:1102.2429](#), [doi:10.1126/science.1200331](#).
- [3] S. A. Walker, A. C. Fabian, J. S. Sanders, A. Simionescu, and Y. Tawara. X-ray exploration of the outskirts of the nearby Centaurus cluster using Suzaku and Chandra. *MNRAS*, 432:554–569, June 2013. [arXiv:1303.4240](#), [doi:10.1093/mnras/stt497](#).
- [4] V. Ghirardini, D. Eckert, S. Ettori, E. Pointecouteau, S. Molendi, M. Gaspari, et al. The universal thermodynamic properties of the intracluster medium over two decades in radius in the X-COP sample. *ArXiv e-prints*, April 2018. [arXiv:1805.00042](#).
- [5] D. Eckert, V. Ghirardini, S. Ettori, E. Rasia, V. Biffi, E. Pointecouteau, et al. Non-thermal pressure support in X-COP galaxy clusters. *ArXiv e-prints*, April 2018. [arXiv:1805.00034](#).
- [6] K. Dolag, M. Meneghetti, L. Moscardini, E. Rasia, and A. Bonaldi. Simulating the physical properties of dark matter and gas inside the cosmic web. *MNRAS*, 370:656–672, Aug 2006. [arXiv:astro-ph/0511357](#), [doi:10.1111/j.1365-2966.2006.10511.x](#).
- [7] M. Tremmel, T. R. Quinn, A. Ricarte, A. Babul, U. Chadayammuri, P. Natarajan, et al. Introducing ROMULUSC: a cosmological simulation of a galaxy cluster with an unprecedented resolution. *MNRAS*, 483:3336–3362, Mar 2019. [arXiv:1806.01282](#), [doi:10.1093/mnras/sty3336](#).
- [8] G. M. Voit, S. T. Kay, and G. L. Bryan. The baseline intracluster entropy profile from gravitational structure formation. *MNRAS*, 364:909–916, December 2005. [arXiv:astro-ph/0511252](#).
- [9] E. Zinger, A. Dekel, Y. Birnboim, A. Kravtsov, and D. Nagai. The role of penetrating gas streams in setting the dynamical state of galaxy clusters. *MNRAS*, 461:412–432, Sep 2016. [arXiv:1510.05388](#), [doi:10.1093/mnras/stw1283](#).
- [10] D. Nagai and E. T. Lau. Gas Clumping in the Outskirts of Λ CDM Clusters. *APJ Letters*, 731:L10, April 2011. [arXiv:1103.0280](#), [doi:10.1088/2041-8205/731/1/L10](#).
- [11] I. Zhuravleva, E. Churazov, A. Kravtsov, E. T. Lau, D. Nagai, and R. Sunyaev. Quantifying properties of ICM inhomogeneities. *MNRAS*, 428:3274–3287, February 2013. [arXiv:1210.6706](#), [doi:10.1093/mnras/sts275](#).
- [12] M. Roncarelli, S. Ettori, S. Borgani, K. Dolag, D. Fabjan, and L. Moscardini. Large-scale inhomogeneities of the intracluster medium: improving mass estimates using the observed azimuthal scatter. *MNRAS*, 432:3030–3046, July 2013. [arXiv:1303.6506](#), [doi:10.1093/mnras/stt654](#).
- [13] F. Vazza, D. Eckert, A. Simionescu, M. Brüggén, and S. Ettori. Properties of gas clumps and gas clumping factor in the intra-cluster medium. *MNRAS*, 429:799–814, February 2013. [arXiv:1211.1695](#), [doi:10.1093/mnras/sts375](#).
- [14] E. Rasia, E. T. Lau, S. Borgani, D. Nagai, K. Dolag, C. Avestruz, et al. Temperature Structure of the Intracluster Medium from Smoothed-particle Hydrodynamics and Adaptive-mesh Refinement Simulations. *APJ*, 791:96, August 2014. [arXiv:1406.4410](#), [doi:10.1088/0004-637X/791/2/96](#).

- [15] N. Battaglia, J. R. Bond, C. Pfrommer, and J. L. Sievers. On the Cluster Physics of Sunyaev-Zel'dovich and X-Ray Surveys. IV. Characterizing Density and Pressure Clumping due to Infalling Substructures. *APJ*, 806:43, June 2015. [arXiv:1405.3346](#), [doi:10.1088/0004-637X/806/1/43](#).
- [16] S. Planelles, D. Fabjan, S. Borgani, G. Murante, E. Rasia, V. Biffi, et al. Pressure of the hot gas in simulations of galaxy clusters. *MNRAS*, 467:3827–3847, June 2017. [arXiv:1612.07260](#), [doi:10.1093/mnras/stx318](#).
- [17] K. Nelson, E. T. Lau, D. Nagai, D. H. Rudd, and L. Yu. Weighing Galaxy Clusters with Gas. II. On the Origin of Hydrostatic Mass Bias in Λ CDM Galaxy Clusters. *APJ*, 782:107, February 2014. [arXiv:1308.6589](#), [doi:10.1088/0004-637X/782/2/107](#).
- [18] X. Shi, E. Komatsu, K. Nelson, and D. Nagai. Analytical model for non-thermal pressure in galaxy clusters - II. Comparison with cosmological hydrodynamics simulation. *MNRAS*, 448:1020–1029, March 2015. [arXiv:1408.3832](#), [doi:10.1093/mnras/stv036](#).
- [19] E. T. Lau, A. V. Kravtsov, and D. Nagai. Residual Gas Motions in the Intracluster Medium and Bias in Hydrostatic Measurements of Mass Profiles of Clusters. *APJ*, 705:1129–1138, November 2009. [arXiv:0903.4895](#), [doi:10.1088/0004-637X/705/2/1129](#).
- [20] E. T. Lau, D. Nagai, and K. Nelson. Weighing Galaxy Clusters with Gas. I. On the Methods of Computing Hydrostatic Mass Bias. *APJ*, 777:151, November 2013. [arXiv:1306.3993](#), [doi:10.1088/0004-637X/777/2/151](#).
- [21] V. Biffi, S. Borgani, G. Murante, E. Rasia, S. Planelles, G. L. Granato, et al. On the Nature of Hydrostatic Equilibrium in Galaxy Clusters. *APJ*, 827:112, August 2016. [arXiv:1606.02293](#), [doi:10.3847/0004-637X/827/2/112](#).
- [22] X. Shi, E. Komatsu, D. Nagai, and E. T. Lau. Analytical model for non-thermal pressure in galaxy clusters - III. Removing the hydrostatic mass bias. *MNRAS*, 455:2936–2944, January 2016. [arXiv:1507.04338](#), [doi:10.1093/mnras/stv2504](#).
- [23] X. Shi, D. Nagai, and E. T. Lau. Multiscale analysis of turbulence evolution in the density-stratified intracluster medium. *MNRAS*, 481:1075–1082, November 2018. [arXiv:1806.05056](#), [doi:10.1093/mnras/sty2340](#).
- [24] S. Ettori, V. Ghirardini, D. Eckert, E. Pointecouteau, F. Gastaldello, M. Sereno, et al. Hydrostatic mass profiles in X-COP galaxy clusters. *Astron. Astrophys.*, 621:A39, January 2019. [arXiv:1805.00035](#), [doi:10.1051/0004-6361/201833323](#).
- [25] N. Ota, D. Nagai, and E. T. Lau. Constraining hydrostatic mass bias of galaxy clusters with high-resolution X-ray spectroscopy. *PASJ*, 70:51, June 2018. [arXiv:1507.02730](#), [doi:10.1093/pasj/psy040](#).
- [26] M. Roncarelli, M. Gaspari, S. Ettori, V. Biffi, F. Brighenti, E. Bulbul, et al. Measuring turbulence and gas motions in galaxy clusters via synthetic Athena X-IFU observations. *Astron. Astrophys.*, 618:A39, October 2018. [arXiv:1805.02577](#), [doi:10.1051/0004-6361/201833371](#).
- [27] G. W. Pratt, M. Arnaud, A. Biviano, D. Eckert, S. Ettori, D. Nagai, et al. The galaxy cluster mass scale and its impact on cosmological constraints from the cluster population. *arXiv e-prints*, 215:25, February 2019. [arXiv:1902.10837](#).
- [28] R. Khatri and M. Gaspari. Thermal SZ fluctuations in the ICM: probing turbulence and thermodynamics in Coma cluster with Planck. *MNRAS*, 463:655–669, November 2016. [arXiv:1604.03106](#), [doi:10.1093/mnras/stw2027](#).

- [29] M. Gaspari and E. Churazov. Constraining turbulence and conduction in the hot ICM through density perturbations. *Astron. Astrophys.*, 559:A78, November 2013. [arXiv:1307.4397](#), [doi:10.1051/0004-6361/201322295](#).
- [30] E. Zinger, A. Dekel, Y. Birnboim, D. Nagai, E. Lau, and A. V. Kravtsov. Cold fronts and shocks formed by gas streams in galaxy clusters. *MNRAS*, 476:56–70, May 2018. [arXiv:1609.05308](#), [doi:10.1093/mnras/sty136](#).
- [31] S. M. Molnar, N. Hearn, Z. Haiman, G. Bryan, A. E. Evrard, and G. Lake. Accretion Shocks in Clusters of Galaxies and Their SZ Signature from Cosmological Simulations. *APJ*, 696:1640–1656, May 2009. [arXiv:0902.3323](#), [doi:10.1088/0004-637X/696/2/1640](#).
- [32] F. Vazza, G. Brunetti, A. Kritsuk, R. Wagner, C. Gheller, and M. Norman. Turbulent motions and shocks waves in galaxy clusters simulated with adaptive mesh refinement. *Astron. Astrophys.*, 504:33–43, September 2009. [doi:10.1051/0004-6361/200912535](#).
- [33] E. T. Lau, D. Nagai, C. Avestruz, K. Nelson, and A. Vikhlinin. Mass Accretion and its Effects on the Self-similarity of Gas Profiles in the Outskirts of Galaxy Clusters. *APJ*, 806:68, June 2015. [arXiv:1411.5361](#), [doi:10.1088/0004-637X/806/1/68](#).
- [34] D. H. Rudd and D. Nagai. Non-equilibrium Electrons and the Sunyaev-Zel’dovich Effect of Galaxy Clusters. *accepted to the ApJL*, June 2009. [arXiv:0907.1287](#).
- [35] C. Avestruz, D. Nagai, E. T. Lau, and K. Nelson. Non-equilibrium Electrons in the Outskirts of Galaxy Clusters. *APJ*, 808:176, August 2015. [arXiv:1410.8142](#), [doi:10.1088/0004-637X/808/2/176](#).
- [36] Norbert Werner, Ondrej Urban, Aurora Simionescu, and Steven W. Allen. A uniform metal distribution in the intergalactic medium of the Perseus cluster of galaxies. , 502:656–658, Oct 2013. [arXiv:1310.7948](#), [doi:10.1038/nature12646](#).
- [37] V. Biffi, S. Planelles, S. Borgani, E. Rasia, G. Murante, D. Fabjan, et al. The origin of ICM enrichment in the outskirts of present-day galaxy clusters from cosmological hydrodynamical simulations. *MNRAS*, 476:2689–2703, May 2018. [arXiv:1801.05425](#), [doi:10.1093/mnras/sty363](#).
- [38] F. Mernier, V. Biffi, H. Yamaguchi, P. Medvedev, A. Simionescu, S. Ettori, et al. Enrichment of the Hot Intracluster Medium: Observations. *Space Science Reviews*, 214:129, December 2018. [arXiv:1811.01967](#), [doi:10.1007/s11214-018-0565-7](#).
- [39] A. Simionescu, N. Werner, O. Urban, S. W. Allen, Y. Ichinohe, and I. Zhuravleva. A Uniform Contribution of Core-collapse and Type Ia Supernovae to the Chemical Enrichment Pattern in the Outskirts of the Virgo Cluster. *APJ*, 811:L25, Oct 2015. [arXiv:1506.06164](#), [doi:10.1088/2041-8205/811/2/L25](#).
- [40] O. Urban, N. Werner, S. W. Allen, A. Simionescu, and A. Mantz. A uniform metallicity in the outskirts of massive, nearby galaxy clusters. *MNRAS*, 470:4583–4599, Oct 2017. [arXiv:1706.01567](#), [doi:10.1093/mnras/stx1542](#).
- [41] M. Tashiro, H. Maejima, K. Toda, R. Kelley, L. Reichenthal, J. Lobell, et al. Concept of the X-ray Astronomy Recovery Mission. In *Space Telescopes and Instrumentation 2018: Ultraviolet to Gamma Ray*, volume 10699 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, page 1069922, July 2018. [doi:10.1117/12.2309455](#).
- [42] P. Predehl, R. Andritschke, H. Böhringer, W. Bornemann, H. Bräuninger, H. Brunner, et al.

- eROSITA on SRG. In *Space Telescopes and Instrumentation 2010: Ultraviolet to Gamma Ray*, volume 7732 of , page 77320U, July 2010. [arXiv:1001.2502](#), [doi:10.1117/12.856577](#).
- [43] A. Merloni, P. Predehl, W. Becker, H. Böhringer, T. Boller, H. Brunner, et al. eROSITA Science Book: Mapping the Structure of the Energetic Universe. *arXiv e-prints*, September 2012. [arXiv:1209.3114](#).
- [44] S. Etori, G. W. Pratt, J. de Plaa, D. Eckert, J. Nevalainen, E. S. Battistelli, et al. The Hot and Energetic Universe: The astrophysics of galaxy groups and clusters. *ArXiv e-prints*, June 2013. [arXiv:1306.2322](#).
- [45] T. Mroczkowski, D. Nagai, K. Basu, J. Chluba, J. Sayers, R. Adam, et al. Astrophysics with the Spatially and Spectrally Resolved Sunyaev-Zeldovich Effects. A Millimetre/Submillimetre Probe of the Warm and Hot Universe. *Space Science Reviews*, 215:17, February 2019. [arXiv:1811.02310](#), [doi:10.1007/s11214-019-0581-2](#).
- [46] T. Mroczkowski, D. Nagai, P. Andreani, M. Arnaud, J. Bartlett, N. Battaglia, et al. A High-resolution SZ View of the Warm-Hot Universe. *arXiv e-prints*, March 2019. [arXiv:1903.02595](#).
- [47] Sehgal et al. Science from an Ultra-Deep, High-Resolution Millimeter-Wave Survey. *arXiv e-prints*, March 2019. [arXiv:1903.03263](#).