



| | |
|----------------------------------|--|
| Publication Year | 2016 |
| Acceptance in OA | 2020-05-12T13:55:33Z |
| Title | ERRATUM: "The variable spin-down rate of the transient magnetar XTE J1810-197" |
| Authors | Pintore, Fabio, BERNARDINI, Federico, MEREGHETTI, Sandro, ESPOSITO, PAOLO, Turolla, Roberto, Rea, Nanda, Coti Zelati, Francesco, ISRAEL, Gian Luca, TIENGO, ANDREA, Zane, Silvia |
| Publisher's version (DOI) | 10.1093/mnras/stw928 |
| Handle | http://hdl.handle.net/20.500.12386/24743 |
| Journal | MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY |
| Volume | 459 |

ERRATUM: ‘The variable spin-down rate of the transient magnetar XTE J1810-197’

by Fabio Pintore,¹★ Federico Bernardini,^{2,3} Sandro Mereghetti,¹ Paolo Esposito,¹ Roberto Turolla,^{4,5} Nanda Rea,^{6,7} Francesco Coti Zelati,^{6,8,9} Gian Luca Israel,¹⁰ Andrea Tiengo^{1,11,12} and Silvia Zane⁵

¹INAF – IASF Milano, Via E. Bassini 15, I-20133 Milano, Italy

²New York University Abu Dhabi, Saadiyat Island, Abu Dhabi, 129188, United Arab Emirates

³INAF – Osservatorio Astronomico di Capodimonte, Salita Moiariello 16, I-80131 Napoli, Italy

⁴Dipartimento di Fisica e Astronomia, Università di Padova, via F. Marzolo 8, I-35131 Padova, Italy

⁵MSSL, University College London, Holmbury St. Mary, Dorking Surrey RH5 6NT, UK

⁶Anton Pannekoek Institute for Astronomy, University of Amsterdam, Postbus 94249, NL-1090-GE Amsterdam, the Netherlands

⁷Instituto de Ciencias de l’Espacio (ICE, CSIC-IEEC), Carrer de Can Magrans, S/N, E-08193 Barcelona, Spain

⁸Università dell’Insubria, via Valleggio 11, I-22100 Como, Italy

⁹INAF – Osservatorio Astronomico di Brera, Via Bianchi 46, I-23807 Merate (LC), Italy

¹⁰INAF – Osservatorio Astronomico di Roma, via Frascati 33, I-00040 Monteporzio Catone, Roma, Italy

¹¹IUSS – Istituto Universitario di Studi Superiori, piazza della Vittoria 15, I-27100 Pavia, Italy

¹²Istituto Nazionale di Fisica Nucleare, Sezione di Pavia, via A. Bassi 6, I-27100 Pavia, Italy

Key words: errata, addenda – magnetic fields – stars: magnetars – stars: neutron – pulsars: individual: XTE J1810-197 – X-rays: stars.

Prompted by the recent paper by Camilo et al. (2016), we re-examined our phase-connected timing solution for XTE J1810-197 (Pintore et al. 2016), and we found a flaw in the procedure to compute the errors during some steps of our analysis. Due to this mistake, the phase-connected solution on 3000 d of X-ray data (reported in table 2 and fig. 2 of Pintore et al. 2016) is wrong.

With the new analysis of the data, we can phase-connect 13 observations with a good fit [χ^2_v (dof) = 0.9(9); solution 1 in Table 1 and fig. 1-left], from MJD 55079 to MJD 55814 (observations from 18 to 30 of Pintore et al. 2016). The inclusion of also the two observations at MJD 55976 and MJD 56071 (observations 31 and 32) yields best-fitting parameters (solution 2 in Table 1 and Fig. 1-right) consistent with those obtained by Camilo et al. (2016) for the same set of observations, but with a higher χ^2_v with respect to solution 1.

The table and figures reported here supersede table 2 and fig. 2 of Pintore et al. (2016). We note that these changes do not affect the main conclusions of that paper.

Table 1. Best-fitting timing solutions for 13 (solution 1) and 15 (solution 2) *XMM-Newton* and *Chandra* observations. Errors are at 1σ .

| Parameter | Solution 1 | Solution 2 |
|------------------------------------|-------------------------------|-------------------------------|
| Time range (MJD) | 55079–55814 | 55079–56071 MJD TDB |
| T_0^a (MJD TDB) | 55444.0 | 55444.0 |
| ν_0 (Hz) | 0.180 481 213 35(44) | 0.180 481 215 99(27) |
| $\dot{\nu}$ (Hz s ⁻¹) | $-9.2059(16) \times 10^{-14}$ | $-9.2085(16) \times 10^{-14}$ |
| $\ddot{\nu}$ (Hz s ⁻²) | $5.7(3) \times 10^{-23}$ | $3.80(13) \times 10^{-23}$ |
| P (s) | 5.540 742 892(14) | 5.540 742 811(8) |
| \dot{P} (s s ⁻¹) | $2.8262(5) \times 10^{-12}$ | $2.8270(5) \times 10^{-12}$ |
| \ddot{P} (s s ⁻²) | $-1.75(9) \times 10^{-21}$ | $-1.16(4) \times 10^{-21}$ |
| χ^2_v (dof) | 0.9 (9) | 5.0 (11) |

Note. ^aReference epoch.

REFERENCES

- Camilo F. et al., 2016, ApJ, 820, 110
Pintore F. et al., 2016, MNRAS, 458, 2088

* E-mail: pintore@iasf-milano.inaf.it




Figure 1. Phase-connection of ~ 800 (left) and ~ 1000 (right) days of *XMM-Newton* and *Chandra* data using a third-order polynomial function. Top panels: the red points are the measured phases, one for each observation, and the solid line is the best-fitting model; bottom panels: residuals with respect to the best-fitting model.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.