



Publication Year	2007
Acceptance in OA	2022-12-27T14:33:09Z
Title	VLBI observations of H ₂ O and CH ₃ OH masers in two high-mass YSOs
Authors	Goddi, C., MOSCADELLI, Luca, SANNA, ALBERTO, CESARONI, Riccardo, Minier, V.
Publisher's version (DOI)	10.1017/S1743921307012768
Handle	http://hdl.handle.net/20.500.12386/32804
Journal	PROCEEDINGS OF THE INTERNATIONAL ASTRONOMICAL UNION
Volume	242

VLBI observations of H₂O and CH₃OH masers in two high-mass YSOs

C. Goddi¹, L. Moscadelli², A. Sanna³, R. Cesaroni² and V. Minier⁴

¹Harvard-Smithsonian Center for Astrophysics, 60 Garden St. Cambridge, MA 02138, USA

²INAF, Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy

³Dipartimento di Fisica, Università degli Studi di Cagliari, S.P. Monserrato-Sestu Km 0.7, 09042 Cagliari, Italy

⁴Service d'Astrophysique, DAPNIA/DSM/CEA Saclay, 91191 Gif-sur-Yvette, France

Abstract. We have conducted phase-reference VLBI observations of H₂O and CH₃OH masers toward two high-mass star forming regions, Sh 2-255 IR and AFGL 5142. In Sh 2-255 infrared water masers are aligned along a direction close to the orientation of a large scale H₂ jet, tracing possibly shocked material in a precessing jet, or, alternatively, the disk-wind emerging from the disk atmosphere. In AFGL 5142 water masers trace expansion at the base of a protostellar jet, whilst methanol masers are more probably tracing infalling than outflowing gas. Our results suggest that water and methanol masers trace different kinematic structures in the circumstellar gas.

1. Introduction

Most previous high-angular ($\leq 0''.1$) resolution studies of molecular masers in high-mass star forming regions (SFRs) have concentrated mainly on either H₂O or CH₃OH masers. While Very Long Baseline Interferometry (VLBI) multi-epoch observations have clarified that water masers originate from shocks associated with protostellar jets (e.g. Goddi *et al.* 2005), different environments have been proposed in several sources to explain the origin of CH₃OH masers (Keplerian disks, jets, hot molecular cores, and HII regions). Comparing the spatio-kinematic distribution of both maser species in the same YSO at high-angular resolution better constrains the interpretation of the kinematic structures traced by the masers. So far only a few studies have been performed in both maser types with sufficient angular resolution ($< 0''.1$) to investigate the nature of the CH₃OH maser birthplace in SFRs and the association between the H₂O and CH₃OH maser emission in the same YSO. Here, we present the results of a multi-epoch VLBI campaign conducted on two high-mass SFRs, Sh 2-255 IR and AFGL 5142, where both the 6.7 GHz methanol and 22.2 GHz water masers are observed.

2. Observational results

We performed Very Long Baseline Array (VLBA) multi-epoch observations of the 22.2 GHz water masers and European VLBI Network (EVN) single-epoch observations of the 6.7 GHz methanol masers toward two high-mass YSOs: Sh 2-255 IR and AFGL 5142.

In Sh 2-255 IR, water maser emission consists of three main clusters of features aligned along a direction close to the orientation of a H₂ jet (Fig. 1, left panel) observed on angular scales of 1–10 arcsec (Howard *et al.* 1997). This might suggest that the water maser features are associated with the inner part of the jet. However, all the measured proper motions form large angles with the outflow axis. The observed spatial and velocity distribution of water masers might be explained considering either jet precession or rotation




Figure 1. Positions and velocities of VLBA H₂O (filled circles) and EVN CH₃OH (filled triangles) masers, for Sh 2-255 IR (left panel) and AFGL 5142 (right panel), respectively. Different grayscale tones denote the maser line-of-sight velocities and the arrows indicate the measured absolute proper motions of the H₂O masers. The dotted lines indicate the axes of the H₂ jet (Howard *et al.* 1997) and the CO outflow (Zhang *et al.* 2007) observed in Sh 2-255 IR and AFGL 5142, respectively. The cross indicates the position uncertainty of the VLA 15 GHz continuum emission peak detected in Sh 2-255 IR (Rengarajan *et al.* 1996), whereas the contour map shows the 22 GHz VLA continuum emission in AFGL 5142 (Goddi & Moscadelli 2006).

around the jet axis. The data in hand do not allow us to discriminate unambiguously between the two scenarios. The 6.7 GHz methanol masers were observed with the EVN by Minier *et al.* (2000). The present data are too scarce even to attempt a qualitative interpretation of their birthplace. New multi-epoch EVN observations of the 6.7 GHz methanol masers are scheduled in order to determine their absolute positions and proper motions.

In AFGL 5142, the water masers show a bipolar spatial and line-of-sight (LOS) velocity distribution; their proper motions clearly indicate that H₂O masers trace expansion from the YSO along a NW-SE direction (Fig. 1, right panel). Recent SMA observations (Zhang *et al.* 2007) revealed a CO outflow oriented along a similar direction, suggesting that water masers might trace the innermost portion of the CO outflow. The methanol masers, which on the plane of the sky lie closer to the flow center than H₂O masers, show higher LOS velocities, in disagreement with a Hubble flow (as observed in the CO emission). Since the 6.7 GHz masers are seen in projection against the HII region, which is optically thick at 6.7 GHz (Goddi & Moscadelli 2006), the 6.7 GHz masers must be located in the foreground of the HII region. The fact that most of the spots present strongly red-shifted l.o.s. velocities suggests that they are tracing gas infalling toward the YSO.

3. Conclusions

Our VLBI data reveal a *true* association of both H₂O 22.2 GHz and CH₃OH 6.7 GHz masers with the same massive YSO in AFGL 5142. Notwithstanding that, the two species appear to trace different kinematic structures, in particular expansion at the base of a molecular outflow and infall in a molecular envelope, respectively.

References

- Goddi, C., Moscadelli, L., Alef, W., Tarchi, A., & Pani, M. 2005, *A&A* 432, 161
 Goddi, C., & Moscadelli, L. 2006, *A&A* 447, 577
 Howard, E. M., Pipher, J. L., & Forrest, W. J. 1997, *ApJ* 481, 327
 Minier, V., Booth, R. S., & Conway, J. E. 2000, *A&A* 362, 1093
 Rengarajan, T. N., & Ho, P. T. P. 1996, *ApJ* 465, 363
 Zhang, Q., *et al.* 2006, *astroph-12027*