



Publication Year	2023
Acceptance in OA	2025-01-22T11:49:51Z
Title	A Spatially Resolved X-Ray Polarization Map of the Vela Pulsar Wind Nebula
Authors	Liu, Kuan, XIE, FEI, Liu, Yi-han, Ng, Chi-Yung, BUCCIANTINI, Niccolo', Romani, Roger W., Weisskopf, Martin C., COSTA, ENRICO, DI MARCO, Alessandro, LA MONACA, Fabio, MULERI, FABIO, SOFFITTA, PAOLO, Deng, Wei, Meng, Yu, Liang, En-wei
Publisher's version (DOI)	10.3847/2041-8213/ad0bfc
Handle	http://hdl.handle.net/20.500.12386/35709
Journal	THE ASTROPHYSICAL JOURNAL LETTERS
Volume	959



A Spatially Resolved X-Ray Polarization Map of the Vela Pulsar Wind Nebula

Kuan Liu¹, Fei Xie^{1,2}, Yi-han Liu^{3,4}, Chi-Yung Ng³, Niccolò Bucciantini^{5,6,7}, Roger W. Romani⁸, Martin C. Weisskopf⁹, Enrico Costa², Alessandro Di Marco², Fabio La Monaca², Fabio Muleri², Paolo Soffitta², Wei Deng¹, Yu Meng¹, and En-wei Liang¹

¹ Guangxi Key Laboratory for Relativistic Astrophysics, School of Physical Science and Technology, Guangxi University, Nanning 530004, People's Republic of China; xief@gxu.edu.cn

² INAF Istituto di Astrofisica e Planetologia Spaziali, Via del Fosso del Cavaliere 100, I-00133 Roma, Italy

³ Department of Physics, The University of Hong Kong, Pokfulam, Hong Kong

⁴ School of Physics and Astronomy, Sun Yat-Sen University, Guangzhou 510275, People's Republic of China

⁵ INAF Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, I-50125 Firenze, Italy

⁶ Dipartimento di Fisica e Astronomia, Università degli Studi di Firenze, Via Sansone 1, I-50019 Sesto Fiorentino (FI), Italy

⁷ Istituto Nazionale di Fisica Nucleare, Sezione di Firenze, Via Sansone 1, I-50019 Sesto Fiorentino (FI), Italy

⁸ Department of Physics and Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, Stanford, CA 94305, USA

⁹ NASA Marshall Space Flight Center, Huntsville, AL 35812, USA

Received 2023 October 25; revised 2023 November 12; accepted 2023 November 13; published 2023 December 6

Abstract

In this paper, we present a full spatially resolved polarization map for the Vela pulsar wind nebula (PWN) observed by IXPE. By employing effective background discrimination techniques, our results show a remarkably high degree of local polarization in the outskirt region, exceeding 60% (55%) with a probability of 95% (99%), which approaches the upper limit predicted by the synchrotron emission mechanism. The high degree of polarization suggests that the turbulent magnetic energy is at most 33% of the ordered one. In addition, the X-ray polarization map exhibits a toroidal magnetic field pattern that is most consistent with the field revealed by radio observations across the entire nebula. This consistency reveals that the observed X-ray and radio emissions are radiated by electrons from the same magnetic field. Different from the Crab PWN, the consistency observed in the Vela PWN may be attributed to the interaction between the reverse shock of the supernova blast wave and the PWN, which leads to a displacement between the synchrotron-cooled nebula and the fresh nebula close to the pulsar. These findings deepen our understanding of the structure and evolution of the Vela PWN and the magnetohydrodynamic interaction in PWNe.

Unified Astronomy Thesaurus concepts: Pulsar wind nebulae (2215); Magnetic fields (994); Polarimetry (1278)

1. Introduction

A pulsar wind nebula (PWN) is an intriguing astrophysical phenomenon that provides a unique environment for studying the interaction processes between the magnetized pulsar wind and the surrounding materials. The Vela PWN, located at a distance of 290 pc (Dodson et al. 2003a), is one of the best objects for investigating the interplay between the PWN and the reverse shock from the supernova explosion. The nebula is powered by the Vela pulsar (also known as PSR B0833-45) with an age of 11.4 kyr (Caraveo et al. 2001), and resides in the north of the extended radio structure Vela X. The Vela X is considered as a PWN relic, and the offset between the pulsar and the Vela X's center is thought to result from the interaction between the nebula and the asymmetric reverse shock from supernova blast-wave (Blondin et al. 2001; Slane et al. 2018).

Multiwavelength observations have been employed to comprehensively study and characterize the Vela PWN. The Chandra X-ray Observatory has provided detailed information into the innermost features (see the left panel of Figure 1), revealing the presence of two prominent arcs and a jet toward the northwestern direction within a radius of $\sim 1'$ centered on the Vela pulsar (Helfand et al. 2001). A deeper observation shows a soft emission shell encompassing the arc-jet structure,

and a faint, diffuse emission structure that extends toward the southwestern region with a scale of several arc minutes (Kargaltsev & Pavlov 2004). Observations in the radio band, at 5 GHz, display an extended image with two distinctive and asymmetric lobes (see the middle panel of Figure 1). These lobes are located in the northeast and southwest of the center pulsar, with a diameter exceeding $10'$ (Dodson et al. 2003b). However, the radio nebula does not show significant features resembling the X-ray arc-jet structure observed in the central region. Instead, it primarily overlaps with the dim diffuse X-ray emission observed in the southwest region. Despite the significant differences in multiwavelength images, the southwest region exhibits notably nebular emissions in both X-ray and radio bands. Thus, the presence of the southwestern nebula could provide some insights into the multiwavelength observations of Vela PWN.

The magnetic field plays a crucial role in the evolution of PWNe, which could be investigated through a polarization study. Dodson et al. (2003b) showed the polarization map of the radio nebula, and a high polarization degree (PD) of 60% was found. The observation revealed a large-scale toroidal magnetic field with a symmetry axis aligned with the direction of the jet axis and the pulsar proper motion. Recently, Xie et al. (2022) reported a remarkably high X-ray PD in the central region (the white square region in the right panel of Figure 1), using the Imaging X-ray Polarimeter Explorer (IXPE; Soffitta et al. 2021; Weisskopf et al. 2022). It is found that the X-ray PD approached the upper limit permitted by the synchrotron




Figure 1. Chandra (left), ATCA (middle, 6 cm), and IXPE (right) images of the Vela PWN. In the ATCA image, the contour from the Chandra observation is overlaid. The white box in the right panel is the region analyzed in Xie et al. (2022), and the white circle marks the boundary of a radius of $300''$ centered at the pulsar. The red panda region and three black squared boxes identify the background and the source regions in the analysis in Section 3, respectively. Each black box is $90'' \times 90''$, and the background region extends from an inner radius of $140''$ up to an outer radius of $300''$ centered on the pulsar, with an open angle of 160° . The white (black) lines in the middle (right) panel show the magnetic field direction revealed by the radio (X-ray) polarization.

emission mechanism, suggesting a highly ordered magnetic field near the particle acceleration site. In addition, the polarization angle (PA) of the X-ray shows a good agreement with that of the radio in the central region, despite their distinct shape.

The X-ray polarization study provides new information for understanding the formation and evolution of the Vela PWN. In this paper, we present a full X-ray polarization study for the entire Vela PWN. We extend the findings presented in Xie et al. (2022) by analyzing the polarization characteristics observed in the southwestern diffuse emission region, estimating the upper limit of the magnetic field strength ratio between the random and the ordered magnetic field, and conducting a comprehensive comparison between the X-ray and radio polarization pattern.

2. Observation and Data Reduction

2.1. X-Ray Polarization Data

IXPE, a Small EXplorers Mission funded by NASA and the Agenzia Spaziale Italiana (ASI), was launched on 2021 December 9th (Weisskopf et al. 2022). It is the first spatially resolved observatory fully dedicated to X-ray polarimetry in the energy band 2–8 keV. IXPE consists of three identical telescopes, each one having a polarimeter, the Gas Pixel Detector (Costa et al. 2001; Baldini et al. 2021), sensitive to linear X-ray polarization and placed at the focus of the coaligned Wolter-1 mirror module assemblies (MMAs). The IXPE MMAs have an angular resolution in half power diameter (HPD) of $\sim 30''$ and a field of view of $12'.9 \times 12'.9$.

The Vela PWN was observed on 2022 April for a total exposure of ~ 860 ks. Data are publically available on NASA'S HEASARC archive, and analysis is performed with IXPEOBS-SIM V30.5.0 (Baldini et al. 2022), which is developed by the IXPE collaboration following the formalism in Kislat et al. (2015). Details on data extraction are reported in Xie et al. (2022).

The right panel of Figure 1 shows the images of the Vela PWN observed by IXPE, as well as the $150'' \times 150''$ squared region analyzed in Xie et al. (2022), and three $90'' \times 90''$ black boxes in the diffuse emission region. All regions, including source and background, are chosen within a radius of $300''$ centered on the pulsar (the white circle) to avoid the

geometrical edge effects, as discussed in Di Marco et al. (2023). The energy range was narrowed down to 25 keV for further increment of the signal-to-noise ratio (S/N, hereafter). In addition, we applied the data selections provided in Di Marco et al. (2023) and Xie et al. (2021), which successfully remove approximately 40% of the instrumental background, reducing the background contamination significantly.

2.2. Radio Polarization Data

We also produced radio polarization images of Vela PWN at 6, 13, and 21 cm for a comparison with the X-ray results. All of the radio data were taken from the Australia Telescope Compact Array (ATCA), and Table 1 shows detailed information about selected observations in every band. There are five observations in the 6 cm band with integration times of 53.7 hr, covering a $u-v$ range from 0.35 to 115 k λ ; four 13 cm mosaic observations give 0.8 hr integration times and a $u-v$ coverage from 0.25 to 40.5 k λ ; five other observations have 5.6 hr integrated observation time at 21 cm, which covers a $u-v$ coverage from 0.1 to 29.6 k λ .

The MIRIAD package helps to reduce all the radio data in this study (Sault et al. 1995). According to the standard procedures, we flagged bad data affected by radio frequency interference, obtained calibrated observation solutions (e.g., flux, bandpass, and gains) from primary and secondary calibrators, and applied these solutions to the target source for every observation. We also eliminated contamination from the strong pulsar flux by excluding emission in the on-pulse phase with the help of ATCA pulsar binning mode (see Table 1 the last column). As is shown in Table 1, every observation in an individual band was combined together to produce Stokes I , Q , and U radio images of the Vela PWN. The rms noises of images are around 0.03, 0.04, and 0.08 mJy beam $^{-1}$ at 6, 13, and 21 cm, respectively. We weighted data inversely proportional to the noise level and the Briggs's robust parameter of 0.5 (Briggs 1995) and plotted images in every band with beam sizes of $30''$. Then, we deconvolved the initial images and combined final Stokes I , Q , and U images to produce the PA images of polarized emission from Vela PWN in each band. Finally, we linearly fit the PA of every pixel in each band to measure the Faraday effect and derive the corrected PA map of the magnetic field in the radio PWN.

Table 1
Radio Observations of Vela PWN Used for the Polarization Analysis

Telescope	Obs. Date	Array Config.	Center Freq. (MHz)	Usable Bandwidth (MHz)	No. of Channels	Integration Time (hr)	Pulsar Binning Mode
6 cm							
ATCA	2018 Jan 09	6C	5997.5	2048	513	9.5	Y
ATCA	2001 Feb 24	375	4800, 5696	104	13	11.5	Y
ATCA	2001 Mar 17	1.5D	4800, 5696	104	13	10.4	Y
ATCA	2001 Mar 30	6E	4800, 5696	104	13	10.7	Y
ATCA	2001 Apr 18	750D	4800, 5696	104	13	10.6	Y
13 cm							
ATCA	1996 Jan 09	750C	2368	104	13	0.2	Y
ATCA	1996 Jan 24	750B	2368	104	13	0.2	Y
ATCA	1996 May 21	750D	2368	104	13	0.2	Y
ATCA	1996 Nov 22	750A	2368	104	13	0.2	Y
21 cm							
ATCA	1996 Jan 09	750C	1344	104	13	0.2	Y
ATCA	1996 Jan 24	750B	1344	104	13	0.2	Y
ATCA	1996 May 21	750D	1344	104	13	0.2	Y
ATCA	1996 Jul 31	6C	1344, 1432	104	13	4.8	Y
ATCA	1996 Nov 22	750A	1344	104	13	0.2	Y

3. Polarization Result

We performed a spatially resolved polarization analysis of Vela PWN in X-ray and radio bands. The top panel of Figure 2 displays the X-ray PD map using $45'' \times 45''$ grids, convoluted with a Gaussian kernel function with a kernel size of $67''.5$ and overlaid with Chandra contours in the 2–5 keV band. The green box marks the region analyzed in Xie et al. (2022), and the three blue boxes are the outskirts regions we focus on (same as the 1, 2, and 3 labeled square regions in Figure 1). The direction of the black lines represents the projected magnetic field, perpendicular to the X-ray PA (defined from north to east), with its length proportional to the PD. The white lines indicate the Faraday-rotation-corrected magnetic field derived from radio data (Liu 2023). Only the grids with X-ray PD larger than MDP_{99} (i.e., the minimum detectable polarization at 99% confidence level) have polarization vectors overlaid, and the corresponding polarization protractor plots are presented in the bottom panel. In these protractor plots, the confidence levels of 50%, 90%, 99%, 99.9%, and 99.99% are enclosed by the red, yellow, green, purple, and blue contours, and the MDP_{99} are filled with pink shadow. The radio polarization is marked by the black star, and the black dashed line highlights the direction of the radio PA. Owing to the fact that the error of the radio PA is at the level of 1° , which is much smaller than that of the X-ray, it is neglected in our discussion.

Xie et al. (2022) have shown that the X-ray polarization is well consistent with the radio in the central region. In this paper, we extend this consistency to the entire nebula. The maximum discrepancy in PA between the X-ray and the radio is below 20° , and the average PA difference is $\sim 17^\circ$ with a mean error of 5° . In the areas exhibiting the most difference, such as pixels (5, 2) and (5, 1), the X-ray PA errors are notably large, implying that the discrepancies may be heavily affected by a statistic effect. In addition, in pixels like (3, 4), the radio polarization deviates from the X-ray polarization by more than 4σ (i.e., beyond the contour of 99.99% confidence level). This

region is close to the pulsar, thus it may be affected by the pulsar’s polarization. Furthermore, the geometry structure near the pulsar seen in the Chandra image is highly curved; therefore, the underlying PA may vary rapidly, leading to a discrepancy in polarization between the radio and X-ray. Future X-ray polarimetry with a smaller point-spread function (PSF) will offer more insight into this matter.

The outskirts of the Vela PWN is highly polarized, as shown in Figure 2, although it is much dimmer than the central region. To obtain statistically significant polarization results, we rebin the diffuse emission into three $90'' \times 90''$ squared regions, labeled 1, 2, and 3, and each of them contains four pixels. All of these three boxes display a high level of significance exceeding 7σ (calculated as the PD divided by its 1σ error), which indicates a strong polarization in the diffuse emission region. We also consider the possible systematic effect on the diffuse emission, including the PSF and polarization leakage (Bucciantini et al. 2023a), using the simulation tools in *ixpeobssim* (Baldini et al. 2022) and GEANT4 (Agostinelli et al. 2003). It is found that these effects cause negligible impact on the X-ray polarization results within 1σ uncertainty, and thus are neglected in our analysis.

Moreover, for the area with low S/N, such as the diffuse emission region, background subtraction could be crucial. We consider a local background sampled from the northeast side of the Vela PWN (the red panda region in the right panel of Figure 1). This background is unpolarized, with an upper limit of 7.6%. Its normalized Stokes parameters are subtracted from the source data, with results presented in Table 2. In general, the PD substantially increases to almost twice its original value after background subtraction, ranging from $\sim 50\%$ to $\sim 80\%$, with 1σ error of $\sim 10\%$ and a significance level $\sim 7\sigma$. To enhance the significance, we combine regions 2 and 3, which are the two most polarized areas in the diffuse emission. The combined region has a PD of $76.2\% \pm 7.2\%$ with a significance level larger than 10σ .




Figure 2. (Top) The PD map of the Vela PWN in the 2–5 keV energy range with a pixel size of 45'' overlaid with the Chandra contour. Pixels with X-ray PD > MDP₉₉ have polarization vectors on top. The length of the black (white) lines indicates the PD measure in X-ray (radio), and orientation indicates the projected magnetic field. (Bottom) The polarization protractor plot for pixels with the polarization vector overlaid on the top panel, covering the same region, with radius and polar angle indicating the PD and PA, respectively.

4. Discussions

We detect a high PD in the outskirt region of the Vela PWN. For the most polarized region, at a confidence level of 90% (95%), the PD is larger than ~60% (~55%) after background subtraction. The maximum PD allowed by the synchrotron

mechanism for a power-law photon spectrum, with a photon index of Γ , is given by $\Pi = 3\Gamma / (3\Gamma + 2)$ (Rybicki & Lightman 2004). For the diffuse emission, with a spectral index of 1.3 ~ 1.4 (i.e., Γ is 0.3 ~ 0.4; Kargaltsev & Pavlov 2004; Kargaltsev et al. 2008), the maximum PD is 66% ~ 67%. Thus the observed PD of the outer diffuse emission is close to the

Table 2
The Normalised Stokes Parameters Q/I , U/I , and the Significance Level of the Diffuse Emission and Background Are Reported, along with the Background Subtracted Polarization

Region	Q/I	U/I	Sig(σ)	PD(%)	PA($^\circ$)
1	-0.35 ± 0.05	0.02 ± 0.05	7.00	35.0 ± 5.0	88.1 ± 4.1
2	-0.37 ± 0.06	-0.20 ± 0.06	7.33	41.6 ± 5.7	104.1 ± 3.9
3	-0.24 ± 0.06	-0.38 ± 0.06	7.82	44.5 ± 5.7	118.8 ± 3.7
Background	0.06 ± 0.02	-0.03 ± 0.02	2.72	<7.6	...
Background Subtracted Polarization					
1	-0.56 ± 0.08	0.05 ± 0.08	7.21	56.4 ± 7.8	87.5 ± 4.0
2	-0.69 ± 0.10	-0.33 ± 0.10	7.36	76.7 ± 10.4	102.6 ± 3.9
3	-0.48 ± 0.11	-0.65 ± 0.11	7.60	80.7 ± 10.6	116.9 ± 3.8
2 + 3	-0.59 ± 0.07	-0.49 ± 0.07	10.49	76.2 ± 7.2	108.8 ± 2.8

Note. Additionally, the PD, PA, and their uncertainties are calculated at 68.3% confidence level. For PD < MDP₉₉, the upper limit in terms of MDP₉₉ is reported.

maximum. The diffuse emission is located $2' \sim 4'$ away from the pulsar, which is almost 10 times further than the arc structure in the center PWN (Ng & Romani 2004). The high PD of the diffuse emission reveals that the magnetic field in the outflow remains undisturbed as the magnetized plasma travels from the acceleration site to the outer region.

Following Bandiera & Petruk (2016) and Bucciantini et al. (2017), we estimate the upper limit of the energy ratio between the random and ordered magnetic fields. The observed PD for a local region can be defined as:

$$\Pi = \frac{\alpha + 1}{\alpha + 5/3} \frac{3 + \alpha \sin^2 \theta_B}{4} \frac{{}_1F_1((1 - \alpha)/2, 3; -\sin^2 \theta_B/2\sigma^2)}{2\sigma^2} \frac{{}_1F_1(-1 + \alpha)/2, 1; -\sin^2 \theta_B/2\sigma^2)}{{}_1F_1(-1 + \alpha)/2, 1; -\sin^2 \theta_B/2\sigma^2)}. \quad (1)$$

Here, α is the spectral index, ${}_1F_1(a, b; x)$ is the Kummer confluent hypergeometric function, with a and b representing the upper and lower parameters, and x being the argument of the function, θ_B is the angle between the magnetic field direction and line of sight in the comoving coordinate; $\sigma = \sqrt{E_r/3E_o} = B_r/\sqrt{3}B_o$ is related to the energy ratio between the random magnetic field and the ordered magnetic field, assuming a three-dimensional isotropic Gaussian random field with a variance of $(B\sigma)^2$ in each direction; the B_r , B_o , and B are the strength of the random, ordered, and total magnetic field in the comoving coordinate, respectively. The spectral index α of Vela diffuse emission is measured as 1.4 (Kargaltsev & Pavlov 2004; Kargaltsev et al. 2008). The relation between the θ'_B and the upper limit of B_r/B_o is shown in Figure 3, in cases of given PD of 55%, 60%, and 65%. With a PD of 55% and under an extreme condition $\theta_B \sim 90^\circ$, the maximum of B_r/B_o is 57%, corresponding to a magnetic energy ratio E_r/E_o of 33%. Considering a more possible case with higher PD and smaller θ_B (for example, PD $\sim 60\%$ and $\theta_B \sim 50^\circ$), the upper limit of B_r/B_o (E_r/E_o) would be $\sim 33\%$ ($\sim 10\%$).

Besides the strong PD in X-ray, we also found a strong correlation between the polarization of the X-ray and the radio throughout the entire nebula, as shown in Figure 2. Both X-ray and radio observations exhibit regions of high polarization, approaching the upper limit allowed by the synchrotron mechanism. This supports that the X-ray and radio emissions




Figure 3. The relationship between the upper limit of the ratio between the random and ordered magnetic field strength and the angle between the light of sight and the magnetic field direction, with given PD of 55% (blue), 60% (orange), and 65% (green).

are produced by the electrons in the same magnetic field (Kargaltsev & Pavlov 2004).

It is worth noting that such a correlation is not observed in the Crab PWN (Aumont et al. 2010; Bucciantini et al. 2023b). This disparity might be attributed to their different evolutionary stages. The Crab is a relatively young nebula at an age of ~ 1 kyr, while the Vela is ~ 11.4 kyr, whose reverse shock of the supernova blast wave is interacting with the nebula. The outer nebular bubble undergoes compression, detaching from the pulsar, and forming the PWN relic, Vela X. In the absence of the outer nebula, we can directly observe the X-ray and radio nebula near the termination site. The predicted X-ray and radio polarization should be consistent, according to the magnetic hydrodynamic numerical simulations (Lyubarsky 2002; Del Zanna et al. 2004; Komissarov & Lyubarsky 2004; Bucciantini et al. 2005). A young nebula, such as the Crab, is expanding in its supernova ejecta and has not yet been interrupted by the reverse shock (Gaensler & Slane 2006). The fresh X-ray-emitting nebula is surrounded by the outer cool nebula, which predominantly radiates in the radio band, leading to the observed disparity between the radio and X-ray polarization.

Furthermore, the high PD observed in the Vela PWN implies limited turbulence in the magnetic field, indicating some

suppression of instability. A strong magnetic field could play a crucial role in suppressing the growth of instability (Bucciantini et al. 2004). The magnetic field in the Vela could be strengthened due to the compression of the reverse shock, resulting in the observed high PD.

The boomerang PWN G106.6+2.96 is similar to Vela PWN (Kothes et al. 2006), where the original nebula has been crushed by the reverse shock. Its radio observations reveal a thick toroidal polarization pattern, akin to that observed in the Vela. If our understanding holds true, its X-ray polarization should also align closely with the radio polarization. Future observations will validate our conjecture.

5. Conclusion

X-ray polarization is important for understanding the magnetic fields of PWNe. Using data from IXPE we made the first full map of the X-ray polarization for Vela PWN as a function of position. In addition to the polarization detected in the inner regions closest to the pulsar, we find after careful background subtraction, statistically significant high PD is detected in the extended X-ray emission, which is close to the upper limit allowed by the synchrotron emission mechanism. The result suggests that the magnetic field in these regions remains highly uniform, and the turbulent magnetic field strength is estimated to be up to around half of the ordered magnetic field strength in the most polarized region.

We also found a strong correlation between the X-ray and radio polarization, which implies that the X-ray and radio are emitted by the electrons in the same magnetic field. Such consistency observed in the Vela PWN could be attributed to the interaction between the nebula and the reverse shock of the supernova blast wave, which leads to a displacement between the synchrotron-cooled nebula and the fresh nebula close to the pulsar. The compression by the reverse shock amplifies the magnetic field, suppresses the growth of instability, and leads to an ordered magnetic field, thus the high degree of polarization. Future X-ray polarization observations of other similar nebular systems, such as boomerang PWN G106.6+2.9, will gain a deeper understanding of them.








Acknowledgments

We thank the anonymous referee of this work for useful comments and suggestions that improved the paper. This work is supported by National Key R&D Program of China (grant No. 2023YFE0117200) and National Natural Science Foundation of China (grant No. 12373041 and grant No. 12133003). N.B. was supported by the INAF MiniGrant “PWNnumpol—Numerical Studies of Pulsar Wind Nebulae in The Light of IXPE.” C.-Y.N. is supported by a GRF grant of the Hong Kong Government under HKU 17305419.

The Imaging X-ray Polarimetry Explorer (IXPE) is a joint US and Italian mission. The US contribution is supported by the National Aeronautics and Space Administration (NASA) and led and managed by its Marshall Space Flight Center (MSFC), with industry partner Ball Aerospace (contract NNM15AA18C). The Italian contribution is supported by the Italian Space Agency (Agenzia Spaziale Italiana, ASI) through contract ASI-OHBI2017-12-I.0, agreements ASI-INAF-2017-12-H0 and ASI-INFN2017.13-H0, and its Space Science Data Center (SSDC), and by the Istituto Nazionale di Astrofisica (INAF) and the Istituto Nazionale di Fisica Nucleare (INFN) in

Italy. This research used data products provided by the IXPE Team (MSFC, SSFC, INAF, and INFN) and distributed with additional software tools by the High-Energy Astrophysics Science Archive Research Center (HEASARC), at NASA Goddard Space Flight Center (GSFC). The X-ray data for the Chandra observations were downloaded from the public access site, the Chandra Data Archive (CDA). This is part of the Chandra X-ray Observatory Science Centre (CXC), which is operated for NASA by the Smithsonian Astrophysical Observatory. The Australia Telescope Compact Array is part of the Australia Telescope, funded by the Commonwealth of Australia for operation as a National Facility, and managed by CSIRO.

ORCID iDs

Kuan Liu  <https://orcid.org/0009-0007-8686-9012>
 Fei Xie  <https://orcid.org/0000-0002-0105-5826>
 Chi-Yung Ng  <https://orcid.org/0000-0002-5847-2612>
 Niccolò Bucciantini  <https://orcid.org/0000-0002-8848-1392>
 Roger W. Romani  <https://orcid.org/0000-0001-6711-3286>
 Martin C. Weisskopf  <https://orcid.org/0000-0002-5270-4240>
 Enrico Costa  <https://orcid.org/0000-0003-4925-8523>
 Alessandro Di Marco  <https://orcid.org/0000-0003-0331-3259>
 Fabio La Monaca  <https://orcid.org/0000-0001-8916-4156>
 Fabio Muleri  <https://orcid.org/0000-0003-3331-3794>
 Paolo Soffitta  <https://orcid.org/0000-0002-7781-4104>
 Wei Deng  <https://orcid.org/0000-0002-9370-4079>
 En-wei Liang  <https://orcid.org/0000-0002-7044-733X>

References

- Agostinelli, S., Allison, J., Amako, K., et al. 2003, *NIMPA*, 506, 250
 Aumont, J., Conversi, L., Thum, C., et al. 2010, *A&A*, 514, A70
 Baldini, L., Barbanera, M., Bellazzini, R., et al. 2021, *APh*, 133, 102628
 Baldini, L., Bucciantini, N., Lalla, N. D., et al. 2022, *SoftX*, 19, 101194
 Bandiera, R., & Petruk, O. 2016, *MNRAS*, 459, 178
 Blondin, J. M., Chevalier, R. A., & Frierson, D. M. 2001, *ApJ*, 563, 806
 Briggs, D. 1995, *BAAS*, 27, 1444
 Bucciantini, N., Amato, E., Bandiera, R., Blondin, J. M., & Del Zanna, L. 2004, *A&A*, 423, 253
 Bucciantini, N., Bandiera, R., Olmi, B., & Del Zanna, L. 2017, *MNRAS*, 470, 4066
 Bucciantini, N., Del Zanna, L., Amato, E., & Volpi, D. 2005, *A&A*, 443, 519
 Bucciantini, N., Di Lalla, N., Romani, R. W. R., et al. 2023a, *A&A*, 672, A66
 Bucciantini, N., Ferrazzoli, R., Bachetti, M., et al. 2023b, *NatAs*, 7, 602
 Caraveo, P. A., Luca, A. D., Mignani, R. P., & Bignami, G. F. 2001, *ApJ*, 561, 930
 Costa, E., Soffitta, P., Bellazzini, R., et al. 2001, *Natur*, 411, 662
 Del Zanna, L., Amato, E., & Bucciantini, N. 2004, *A&A*, 421, 1063
 Di Marco, A., Soffitta, P., Costa, E., et al. 2023, *AJ*, 165, 143
 Dodson, R., Legge, D., Reynolds, J. E., & McCulloch, P. M. 2003a, *ApJ*, 596, 1137
 Dodson, R., Lewis, D., McConnell, D., & Deshpande, A. A. 2003b, *MNRAS*, 343, 116
 Gaensler, B. M., & Slane, P. O. 2006, *ARA&A*, 44, 17
 Helfand, D. J., Gotthelf, E. V., & Halpern, J. P. 2001, *ApJ*, 556, 380
 Kargaltsev, O., & Pavlov, G. 2004, in *IAU Symp.* 218, *Spatially Resolved Spectrum of the Vela PWN*, 218 (Cambridge: Cambridge Univ. Press), 195
 Kargaltsev, O., Pavlov, G. G., Bassa, C., et al. 2008, in *AIP Conf. Proc.* 983, *40 YEARS OF PULSARS: Millisecond Pulsars, Magnetars and More*, ed. C. Bassa (Melville, NY: AIP), 171
 Kislak, F., Clark, B., Beilicke, M., & Krawczynski, H. 2015, *APh*, 68, 45
 Komisarov, S. S., & Lyubarsky, Y. E. 2004, *MNRAS*, 349, 779
 Kothes, R., Reich, W., & Uyaniker, B. 2006, *ApJ*, 638, 225
 Liu, Y. 2023, PhD thesis, HKU Theses Online (HKUTO)
 Lyubarsky, Y. E. 2002, *MNRAS*, 329, L34
 Ng, C.-Y., & Romani, R. W. 2004, *ApJ*, 601, 479

Rybicki, G. B., & Lightman, A. P. 2004, *Radiative Processes in Astrophysics*, Physics Textbook (Weinheim: Wiley)

Sault, R., Teuben, P., Wright, M., et al. 1995, in *ASP Conf. Ser. 77, Astronomical Data Analysis Software and Systems IV*, ed. A. Shaw, H. E. Payne, & J. J. E. Hayes (San Francisco, CA: ASP), 433

Slane, P., Lovchinsky, I., Kolb, C., et al. 2018, *ApJ*, 865, 86

Soffitta, P., Baldini, L., Bellazzini, R., et al. 2021, *AJ*, 162, 208

Weisskopf, M. C., Soffitta, P., Baldini, L., et al. 2022, *JATIS*, 8, 026002

Xie, F., Di Marco, A., La Monaca, F., et al. 2022, *Natur*, 612, 658

Xie, F., Ferrazzoli, R., Soffitta, P., et al. 2021, *APh*, 128, 102566