









<b>Publication Year</b>	2023
<b>Acceptance in OA</b>	2024-03-08T10:57:28Z
<b>Title</b>	Characterising TOI-732 b and c: New insights into the M-dwarf radius and density valley
<b>Authors</b>	Bonfanti, A., Brady, M., Wilson, T.G., Venturini, J., Egger, J.A., Brandeker, A., Sousa, S.G., Lendl, M., Simon, A.E., Queloz, D., Olofsson, G., Adibekyan, V., Alibert, Y., Fossati, L., Hooton, M.J., Kubyshkina, D., Luque, R., Murgas, F., Mustill, A.J., Santos, N.C., Van Grootel, V., Alonso, R., Asquier, J., Bandy, T., Bárczy, T., Barrado Navascues, D., Barros, S.C., Baumjohann, W., Bean, J., Beck, M., Beck, T., Benz, W., BERGOMI, Maria, Billot, N., BORSATO, LUCA, Broeg, C., Collier Cameron, A., Csizmadia, Sz., Cubillos, P.E., Davies, M.B., Deleuil, M., Deline, A., Delrez, L., Demangeon, O.D.S., Demory, B.-O., Ehrenreich, D., Erikson, A., Fortier, A., Fridlund, M., Gandolfi, D., Gillon, M., Güdel, M., Günther, M.N., Heitzmann, A., Helling, Ch., Hoyer, S., Isaak, K.G., Kasper, D., Kiss, L., Lam, K.W.F., Laskar, J., Lecavelier des Etangs, A., MAGRIN, DEMETRIO, Maxted, P.F.L., Mordasini, C., NASCIMBENI, VALERIO, Ottensamer, R., PAGANO, Isabella, Pallé, E., Peter, G., Piotto, G., Pollacco, D., RAGAZZONI, Roberto, Rando, N., Rauer, H., Ribas, I., SCANDARIATO, GAETANO, Ségransan, D., Seifahrt, A., Smith, A.M.S., Stalport, M., Stefánsson, G., Steinberger, M., Stürmer, J., Szabó, M. Gy., Thomas, N., Udry, S., Villaver, E., Walton, N.A., Westerdorff, K., Zingales, T.
<b>Publisher's version (DOI)</b>	10.1051/0004-6361/202348180
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/34949">http://hdl.handle.net/20.500.12386/34949</a>
<b>Journal</b>	ASTRONOMY & ASTROPHYSICS

# Characterising TOI-732 b and c: New insights into the M-dwarf radius and density valley<sup>★</sup>

A. Bonfanti<sup>1</sup> , M. Brady<sup>2</sup> , T. G. Wilson<sup>3</sup> , J. Venturini<sup>4</sup> , J. A. Egger<sup>5</sup> , A. Brandeker<sup>6</sup> , S. G. Sousa<sup>7</sup> , M. Lendl<sup>4</sup> , A. E. Simon<sup>5,8</sup> , D. Queloz<sup>9,10</sup> , G. Olofsson<sup>6</sup> , V. Adibekyan<sup>7</sup> , Y. Alibert<sup>8,5</sup> , L. Fossati<sup>1</sup> , M. J. Hooton<sup>10</sup> , D. Kubyskhina<sup>1</sup> , R. Luque<sup>2</sup> , F. Murgas<sup>11,12</sup> , A. J. Mustill<sup>13</sup> , N. C. Santos<sup>7,14</sup> , V. Van Grootel<sup>15</sup> , R. Alonso<sup>16,17</sup> , J. Asquier<sup>18</sup> , T. Bandy<sup>5</sup> , T. B arczy<sup>19</sup> , D. Barrado Navascues<sup>20</sup> , S. C. C. Barros<sup>7,14</sup> , W. Baumjohann<sup>1</sup> , J. Bean<sup>2</sup> , M. Beck<sup>4</sup> , T. Beck<sup>5</sup> , W. Benz<sup>5,8</sup> , M. Bergomi<sup>21</sup> , N. Billot<sup>4</sup> , L. Borsato<sup>21</sup> , C. Broeg<sup>5,8</sup> , A. Collier Cameron<sup>22</sup> , Sz. Csizmadia<sup>23</sup> , P. E. Cubillos<sup>24,1</sup> , M. B. Davies<sup>25</sup> , M. Deleuil<sup>26</sup> , A. Deline<sup>4</sup> , L. Delrez<sup>27,15</sup> , O. D. S. Demangeon<sup>7,14</sup> , B.-O. Demory<sup>8,5</sup> , D. Ehrenreich<sup>4,28</sup> , A. Erikson<sup>23</sup> , A. Fortier<sup>5,8</sup> , M. Fridlund<sup>29,30</sup> , D. Gandolfi<sup>31</sup> , M. Gillon<sup>27</sup> , M. G udel<sup>32</sup> , M. N. G unther<sup>18</sup> , A. Heitzmann<sup>4</sup> , Ch. Helling<sup>1,33</sup> , S. Hoyer<sup>26</sup> , K. G. Isaak<sup>18</sup> , D. Kasper<sup>2</sup> , L. L. Kiss<sup>34,35</sup> , K. W. F. Lam<sup>23</sup> , J. Laskar<sup>36</sup> , A. Lecavelier des Etangs<sup>37</sup> , D. Magrin<sup>21</sup> , P. F. L. Maxted<sup>38</sup> , C. Mordasini<sup>5,8</sup> , V. Nascimbeni<sup>21</sup> , R. Ottensamer<sup>32</sup> , I. Pagano<sup>39</sup> , E. Pall e<sup>16,17</sup> , G. Peter<sup>40</sup> , G. Piotto<sup>21,41</sup> , D. Pollacco<sup>3</sup> , R. Ragazzoni<sup>21,41</sup> , N. Rando<sup>18</sup> , H. Rauer<sup>23,42,43</sup> , I. Ribas<sup>44,45</sup> , G. Scandariato<sup>39</sup> , D. S egransan<sup>4</sup> , A. Seifahrt<sup>2</sup> , A. M. S. Smith<sup>23</sup> , M. Stalport<sup>15,27</sup> , G. Stef ansson<sup>46\*\*</sup> , M. Steinberger<sup>1</sup> , J. St urmer<sup>47</sup> , Gy. M. Szab o<sup>48,49</sup> , N. Thomas<sup>5</sup> , S. Udry<sup>4</sup> , E. Villaver<sup>16,17</sup> , N. A. Walton<sup>50</sup> , K. Westerdorff<sup>40</sup> , and T. Zingales<sup>41,21</sup> 

(Affiliations can be found after the references)

## ABSTRACT

**Context.** TOI-732 is an M dwarf hosting two transiting planets that are located on the two opposite sides of the radius valley. Inferring a reliable demographics for this type of systems is key to understanding their formation and evolution mechanisms.

**Aims.** By doubling the number of available space-based observations and increasing the number of radial velocity (RV) measurements, we aim at refining the parameters of TOI-732 b and c. We also use the results to study the slope of the radius valley and the density valley for a well-characterised sample of M-dwarf exoplanets.

**Methods.** We performed a global Markov chain Monte Carlo analysis by jointly modelling ground-based light curves and CHEOPS and TESS observations, along with RV time series both taken from the literature and obtained with the MAROON-X spectrograph. The slopes of the M-dwarf valleys were quantified via a support vector machine (SVM) procedure.

**Results.** TOI-732 b is an ultrashort-period planet ( $P = 0.76837931^{+0.00000039}_{-0.00000042}$  d) with a radius  $R_b = 1.325^{+0.057}_{-0.058} R_\oplus$ , a mass  $M_b = 2.46 \pm 0.19 M_\oplus$ , and thus a mean density  $\rho_b = 5.8^{+1.0}_{-0.8} \text{ g cm}^{-3}$ , while the outer planet at  $P = 12.252284 \pm 0.000013$  d has  $R_c = 2.39^{+0.10}_{-0.11} R_\oplus$ ,  $M_c = 8.04^{+0.50}_{-0.48} M_\oplus$ , and thus  $\rho_c = 3.24^{+0.55}_{-0.43} \text{ g cm}^{-3}$ . Even with respect to the most recently reported values, this work yields uncertainties on the transit depths and on the RV semi-amplitudes that are smaller up to a factor of  $\sim 1.6$  and  $\sim 2.4$  for TOI-732 b and c, respectively. Our calculations for the interior structure and the location of the planets in the mass-radius diagram lead us to classify TOI-732 b as a super-Earth and TOI-732 c as a mini-Neptune. Following the SVM approach, we quantified  $d \log R_{p,\text{valley}} / d \log P = -0.065^{+0.024}_{-0.013}$ , which is flatter than for Sun-like stars. In line with former analyses, we note that the radius valley for M-dwarf planets is more densely populated, and we further quantify the slope of the density valley as  $d \log \hat{\rho}_{\text{valley}} / d \log P = -0.02^{+0.12}_{-0.04}$ .

**Conclusions.** Compared to FGK stars, the weaker dependence of the position of the radius valley on the orbital period might indicate that the formation shapes the radius valley around M dwarfs more strongly than the evolution mechanisms.

**Key words.** planets and satellites: fundamental parameters – stars: fundamental parameters – techniques: photometric – techniques: radial velocities

## 1. Introduction

M dwarfs are the most common stars in the Universe (e.g. Bastian et al. 2010). Because of their low mass and small radius, they are quite attractive in the domain of exoplanetology: It is easier to detect low-mass planets with the transit method (e.g.

Winn 2010) and the radial velocity (RV) technique (e.g. Hatzes 2016). In addition, the habitable zone (HZ; Kasting et al. 1993; Kopparapu et al. 2013) around M dwarfs is located closer to the host star than in stars of earlier spectral type, and therefore, it is more likely that planets in the habitable zone (HZ) of M dwarfs are found because both the transit method and the RV technique preferentially detect close-in planets.

<sup>★</sup> This article uses data from CHEOPS programme CH\_PR100031

<sup>\*\*</sup> NASA Sagan Fellow

A few mechanisms may act against the possibility that M dwarfs harbour life, such as their strong magnetic activity (e.g. Saar & Linsky 1985; Reiners et al. 2009; Shulyak et al. 2019), accompanied by flares and high-energy emission that induce atmospheric escape (e.g. Luger & Barnes 2015; Tilley et al. 2019), or the likely scenario of tidally locked close-in exoplanets that lead to extreme surface temperature gradients (e.g. Barnes 2017). However, different works highlighted ways in which planets might be able to become and remain habitable despite the unfavourable stellar environment (Kay et al. 2016; Sergeev et al. 2020; Childs et al. 2022; Ojha et al. 2022; Lobo et al. 2023).

The first exoplanet discovered around an M dwarf is GJ 876 b. This Jovian-mass planet was independently detected via the RV technique by Delfosse et al. (1998) and Marcy et al. (1998). Subsequent studies have also revealed smaller exoplanets (e.g. GJ 436 b; Butler et al. 2004; Gillon et al. 2007) or multiplanet systems containing Neptune-sized planets and super-Earths, such as GJ 581 (Bonfils et al. 2005; Udry et al. 2007). Since then, the search for exoplanets orbiting M dwarfs has been rather prolific; the community has never lost interest in M-dwarf exoplanets, as proven by several recent discoveries, such as TOI-244 b (Castro-González et al. 2023), TOI-715 b (Dransfield et al. 2023), K2-416 b and K2-417 b (Incha et al. 2023), TOI-3785 b (Powers et al. 2023), TOI-3984 A b and TOI-5293 A b (Cañas et al. 2023), TOI-1680 b (Ghachoui et al. 2023), and TOI-2084 b and TOI-4184 b (Barkaoui et al. 2023).

In this work, we characterise TOI-732. This M dwarf is orbited by an ultrashort-period planet (TOI-732 b,  $P \sim 0.77$  d) and an outer planet (TOI-732 c,  $P \sim 12.25$  d). The system has been studied by Cloutier et al. (2020 hereafter C20), Nowak et al. (2020 hereafter N20), and Luque & Pallé (2022 hereafter L22). We add a new TESS (Transiting Exoplanet Survey Satellite; Ricker et al. 2015) sector, 25 novel space-based light curves (LCs) observed by CHEOPS (Characterising Exoplanet Satellite; Benz et al. 2021), and 39 RV data points taken with the high-precision echelle spectrograph MAROON-X (Seifahrt et al. 2018, 2022) to the already published data. Even with respect to the most recent analysis by L22, we almost doubled the number of space-based transit events, and we significantly increased the number of RV data points, which allowed us to significantly reduce the uncertainties in the planetary parameters.

From the point of view of planet formation and evolution, TOI-732 is an interesting system because the planets are located on the two opposite sides of the radius valley (Fulton et al. 2017). The paucity of exoplanets with orbital periods  $P < 100$  d in the  $R_p$  1.5–2.0  $R_\oplus$  radius range determines a bimodal  $R_p$  distribution that peaks at  $\sim 1.3$  and  $\sim 2.4 R_\oplus$  (e.g. Fulton et al. 2017; Fulton & Petigura 2018; Van Eylen et al. 2018). One interpretation of this distribution suggests that sub-Neptunes likely form with rocky cores with sizes 1.5  $R_\oplus$  or smaller, surrounded by an envelope with a low mean molecular weight that is subject to photo-evaporation (e.g. Lammer et al. 2003; Lopez & Fortney 2013; Owen & Wu 2013; Chen & Rogers 2016). However, atmospheric erosion may also be driven by cooling of rocky cores (core-powered mass loss; e.g. Ginzburg et al. 2018; Gupta & Schlichting 2019) and by energy release following cohesive giant impacts during proto-planet formation (impact erosion; e.g. Kegerreis et al. 2020). Finally, the smallest sub-Neptunes might also be the result of late formation within gas-depleted discs (gas-poor formation; e.g. Lee et al. 2014; Lee & Connors 2021; Lee et al. 2022).

These mechanisms that account for the radius valley assume that super-Earths and sub-Neptunes assembled from the same underlying population of dry rocky cores, which might or might

not retain a H-He atmosphere. However, global planet formation models show that migration is a key mechanism delivering water-rich sub-Neptunes at short orbital periods (e.g. Alibert et al. 2013; Venturini et al. 2020; Emsenhuber et al. 2021), especially for low-mass planets around M dwarfs (Alibert 2017; Miguel et al. 2020; Burn et al. 2021). In particular, Venturini et al. (2020) showed that the radius valley emerges from a combination of formation and evolution processes that separate small rocky from larger water-rich planets that formed beyond the ice line. Observational support for this scenario was recently found by L22, who studied a sub-sample of M-dwarf exoplanets and reported a clear density gap that separated super-Earths (identified as rocky planets) from mini-Neptunes (identified as water-ice-rich worlds and not as rocky cores surrounded by H-He). They also concluded that the radius dispersion, especially among puffy exoplanets, may be the consequence of the different accretion histories of H-He envelopes and not of the atmospheric mass loss.

Obtaining observational data is key to investigating the relative importance of the different formation and evolution scenarios. So far, most of the studies have investigated the nature of the radius valley by focusing on FGK stars (e.g. Van Eylen et al. 2018; MacDonald 2019; Martinez et al. 2019; Ho & Van Eylen 2023), while only a few works specifically drew attention to low-mass stars (Cloutier & Menou 2020; Van Eylen et al. 2021; Luque & Pallé 2022). The discoveries of M-dwarf systems in which planets straddle the radius gap have steadily increased. They comprise, for example, TOI-776 (Luque et al. 2021; Fridlund et al. 2023), TOI-1634 (Cloutier et al. 2021b; Hirano et al. 2021), TOI-270 (Van Eylen et al. 2021), TOI-1468 (Chaturvedi et al. 2022), K2-3 (Diamond-Lowe et al. 2022), TOI-2096 (Pozuelos et al. 2023), and LHS 1903 (Wilson et al. 2023). More generally, the parameters of planets orbiting low-mass stars are progressively known with increasingly better precision. This work therefore also aims at describing the characteristics of the radius valley better for planets orbiting M dwarfs.

This paper is organised as follows. Sect. 2 presents the stellar properties, Sects. 3 and 4 describe the photometric and RV data, that were used to characterise the system as explained in Sect. 5. Sect. 6 investigates the radius and density valleys of M-dwarf exoplanets from a quantitative perspective by using the most precise collection of planetary parameters available so far. Finally, Sect. 7 gathers our conclusions.

## 2. Host star properties

TOI-732 is an M4 V star (Scholz et al. 2005) located  $\sim 22$  pc away from us (Gaia Collaboration et al. 2023), with magnitudes  $V = 13.14 \pm 0.04$  (Zacharias et al. 2012) and  $K = 8.204 \pm 0.021$  (Cutri et al. 2003). It is part of a visual binary system, and its companion is known as LP 729-55 and is located at an angular separation  $\theta = 15.81 \pm 0.15''$ , which implies a projected orbital distance of  $348 \pm 3$  AU (N20). LP 729-55 is fainter by  $\sim 2$   $K$ -band mag than TOI-732, and its spectral type has been estimated by N20 as M5.0 V.

To estimate the stellar effective temperature  $T_{\text{eff}}$  and metallicity [Fe/H] of TOI-732, we used the ODUSSEAS<sup>1</sup> code (Antoniadis-Karnavas et al. 2020), and we input spectroscopic data taken from the ESO archive. Although we were able to combine data from ESPRESSO (Pepe et al. 2021) and HARPS (Mayor et al. 2003), we used the latter because it provided a higher signal-to-noise ratio (S/N) of the combined spectrum.

<sup>1</sup> <https://github.com/AlexandrosAntoniadis/ODUSSEAS>

ODUSSEAS uses the ridge regression (Hoerl & Kennard 1970) implemented via the machine-learning PYTHON package *scikit-learn* (Pedregosa et al. 2011), which is trained to measure the pseudo-equivalent widths of more than 4000 stellar absorption lines. Using a library of HARPS spectra for several M stars with well-defined reference parameters from interferometric calibrations (Antoniadis-Karnavas et al. in prep.), ODUSSEAS derived  $T_{\text{eff}} = 3358 \pm 92$  K and  $[\text{Fe}/\text{H}] = 0.06 \pm 0.11$  dex. The trigonometric surface gravity was estimated using  $T_{\text{eff}}$  and  $[\text{Fe}/\text{H}]$  in combination with the Gaia parallax (Gaia Collaboration et al. 2023) and photometry, following the same procedure as described in Sousa et al. (2021), which yields  $\log g = 4.85 \pm 0.11$ .

Because of the heavy line blending, the determination of the individual elemental abundances of M dwarfs from visible spectra is challenging (e.g. Maldonado et al. 2020). In this work, we estimated the abundance of Mg and Si following the procedure presented in Demangeon et al. (2021a). In brief, we used the systemic radial velocity ( $RV_{\text{sys}}$ ), parallax ( $\pi$ ), right ascension ( $\alpha$ ), declination ( $\delta$ ), and proper motions ( $\mu_{\alpha}$  and  $\mu_{\delta}$ ) from Gaia DR3 (Gaia Collaboration et al. 2023) to derive the Galactic space velocity UVW of TOI-732 via the `GalVel_Pop.py` routine<sup>2</sup>. We obtained  $U = 4.0 \pm 0.1$  km s<sup>-1</sup>,  $V = -10.3 \pm 0.3$  km s<sup>-1</sup>, and  $W = -27.5 \pm 0.2$  km s<sup>-1</sup> with respect to the local standard of rest (LSR), adopting the solar peculiar motion from Schönrich et al. (2010). Based on these velocities, adopting the characteristic parameters of Galactic stellar populations from Reddy et al. (2006), and following Adibekyan et al. (2012), we estimated that the star belongs to the Galactic thin disc with a 97% probability. Then, from the APOGEE DR17 (Abdurro'uf et al. 2022), we selected cool stars with metallicities similar to that of TOI-732 that belong to the chemically defined Galactic thin disc. We obtained a sample of several thousand stars, for which we calculated the mean abundance of Mg and Si, and their standard deviation (star-to-star scatter). After taking the stellar metallicity into account, we obtained  $[\text{Mg}/\text{H}] = 0.04 \pm 0.20$  dex and  $[\text{Si}/\text{H}] = 0.02 \pm 0.21$  dex.

We computed the infrared flux method (IRFM) (Blackwell & Shallis 1977) radius of TOI-732 using a modified Markov chain Monte Carlo (MCMC) approach (Schanche et al. 2020). We constructed spectral energy distributions (SEDs) by constraining stellar atmospheric models from three catalogues (Kurucz 1993; Castelli & Kurucz 2003; Allard 2014) with the results of our spectral analysis. From these, we calculated the stellar bolometric flux via comparison of synthetic and observed broadband photometry in the following bandpasses: Gaia  $G$ ,  $G_{\text{BP}}$ , and  $G_{\text{RP}}$ , 2MASS  $J$ ,  $H$ , and  $K$ , and WISE  $W1$  and  $W2$  (Gaia Collaboration et al. 2023; Skrutskie et al. 2006; Wright et al. 2010). The bolometric flux was first converted into effective temperature and angular diameter and then into stellar radius using the offset-corrected Gaia parallax (Lindgren et al. 2021). The stellar atmospheric modelling uncertainties were accounted for by using a Bayesian modelling that averaged the radius posterior distributions. The complex spectral features of M-dwarfs can cause degeneracies in the strengths of molecular lines and thus in the bolometric flux computation within the MCMC when using different atmospheric models. This propagates to large errors on M-dwarf IRFM radii compared to using empirical relations (see C20 and N20). The consistency between our estimate and the outcomes in both C20 and N20 is well below  $1\sigma$  and therefore, we attributed the typical uncertainty to  $R_{\star}$  as derived from empirical relations. We obtained  $R_{\star} = 0.380 \pm 0.012 R_{\odot}$ .

We used  $T_{\text{eff}}$ ,  $[\text{Fe}/\text{H}]$ , and  $R_{\star}$  along with their error bars to then derive the stellar mass  $M_{\star}$  from two different evolutionary models. In detail, we applied the isochrone placement algorithm (Bonfanti et al. 2015, 2016), which is designed to interpolate the set of input parameters within pre-computed grids of PARSEC<sup>3</sup> v1.2S (Marigo et al. 2017) isochrones and tracks, and we obtained a first estimate for the mass. A second estimate was instead obtained via the Code Liègeois d'Évolution Stellaire (CLEM; Scufflaire et al. 2008), which builds the best-fit evolutionary tracks on the fly following the Levenberg-Marquadt minimisation scheme (Salmon et al. 2021). As outlined in Bonfanti et al. (2021), the consistency of the two results is checked through a  $\chi^2$ -based criterion, after which the mass distributions inferred from the two different evolutionary models are merged together. We finally obtained  $M_{\star} = 0.381^{+0.024}_{-0.034} M_{\odot}$ .

Both C20 and N20 have derived the stellar mass and obtained  $M_{\star, \text{C20}} = 0.401 \pm 0.012 M_{\odot}$  and  $M_{\star, \text{N20}} = 0.379 \pm 0.016 M_{\odot}$ . These estimates are consistent with ours, but are more precise by a factor of  $\sim 2$ , but the uncertainties appear to be underestimated. In detail, C20 used the mass-luminosity relation from Benedict et al. (2016). Even considering the K-band luminosity, which yields the most satisfactory fit, the average root mean square (rms) of the residuals is  $0.014 M_{\odot}$ , which is larger than the reported estimate. Furthermore, the mass residuals in the neighbourhood of the TOI-732 absolute stellar magnitude (i.e.  $M_K = 6.494 \pm 0.021$  mag) as displayed in Benedict et al. (2016 Fig. 23, right panel) are higher than the average value by about a factor of two. Instead, N20 used the mass-radius relation from Schweitzer et al. (2019), whose rms inherent to the fit is  $0.02 M_{\odot}$ . When the fit-related source of errors is accounted for, the uncertainties on  $M_{\star}$  from both C20 and N20 become similar to ours. Therefore, our mass uncertainty is probably genuine and robust, also considering that it comes from evolutionary models employing different physical ingredients and was inferred using different derivation algorithms (see Bonfanti et al. 2021 for further details).

As is well known, M dwarfs evolve very slowly. Any age inference via isochrone fitting is therefore inconclusive. However, due to stellar interactions that manifest themselves as kinematic disturbances over the lifetimes of stars, we can estimate the stellar age based on kinematics alone (Wielen 1977; Nordström et al. 2004; Casagrande et al. 2011; Maciel et al. 2011). We used the method of Almeida-Fernandes & Rocha-Pinto (2018), which allows for age estimates based on kinematic-age probability distributions that were formalised and bench-marked using a sample of 9000 stars in the Geneva-Copenhagen Survey whose isochronal ages are known. For this study, we computed the age of TOI-732 using the Galactic  $U$ ,  $V$ , and  $W$  velocities and Gaia DR3 Galactic reference coordinates (Gaia Collaboration et al. 2023), and we obtained an age of  $3.10^{+6.20}_{-0.98}$  Gyr. All the relevant stellar parameters are reported in Table 1.

We further investigated the evolutionary stage of TOI-732 by computing the equivalent width (EW) of the H $\alpha$  emission component, which has been related to the age of M dwarfs by Kiman et al. (2021). To this end, we used the HARPS and ESPRESSO combined spectra. Following the procedure described in Schmidt et al. (2015) and West et al. (2011), we calculated a H $\alpha$  EW of 0.64 Å from the HARPS spectra and 0.52 Å from the ESPRESSO spectra.

Kiman et al. (2021) defined a boundary that separates active from inactive M dwarfs, which latter have an H $\alpha$  EW below a

<sup>2</sup> [https://github.com/vadibekyan/GalVel\\_Pop/blob/main/GalVel\\_Pop.py](https://github.com/vadibekyan/GalVel_Pop/blob/main/GalVel_Pop.py)

<sup>3</sup> PAдова and TRIeste Stellar Evolutionary Code: <http://stev.oapd.inaf.it/cgi-bin/cmd>

Table 1: Stellar properties

TOI-732		
TIC 36724087		
Star names	LTT 3780	
	LP 729-54	
Gaia DR2 3767281845873242112		
Parameter	Value	Source
$\alpha$ [°]	154.64485	Gaia DR3
$\delta$ [°]	-11.71784	Gaia DR3
$\mu_\alpha$ [mas yr <sup>-1</sup> ]	-341.537 ± 0.032	Gaia DR3
$\mu_\delta$ [mas yr <sup>-1</sup> ]	-247.747 ± 0.032	Gaia DR3
$RV_{\text{sys}}$ [km s <sup>-1</sup> ]	+0.27 ± 0.34	Gaia DR3
$\pi$ [mas]	45.382 ± 0.030	Gaia DR3 <sup>(a)</sup>
$T_{\text{eff}}$ [K]	3358 ± 92	Spectroscopy
log $g$	4.85 ± 0.11	Trigonometric
[Fe/H]	0.06 ± 0.11	Spectroscopy
[Mg/H]	0.04 ± 0.20	Thin disc
[Si/H]	0.02 ± 0.21	Thin disc
$R_\star$ [ $R_\odot$ ]	0.380 ± 0.012	IRFM
$M_\star$ [ $M_\odot$ ]	0.381 <sup>+0.024</sup> <sub>-0.034</sub>	Isochrones
$t_\star$ [Gyr]	3.10 <sup>+6.20</sup> <sub>-0.98</sub>	Kinematics
$L_\star$ [ $L_\odot$ ]	0.0165 ± 0.0021	$R_\star$ & $T_{\text{eff}}$
$\rho_\star$ [ $\rho_\odot$ ]	6.94 ± 0.84	$R_\star$ & $M_\star$

**Notes.** <sup>(a)</sup> Correction from Lindegren et al. (2021) applied.

colour-dependent threshold value. Given the  $G - G_{\text{RP}} = 1.197$  colour (Gaia Collaboration et al. 2023) of TOI-732, the corresponding activity boundary is  $H\alpha\text{-EW}_{\text{bound}} = 0.85 \text{ \AA}$ . Because both our  $H\alpha\text{-EW}$  estimates derived from HARPS and ESPRESSO are below the threshold value, TOI-732 can be categorised as inactive. Kiman et al. (2021) pointed out that inactive stars can be found at different evolutionary stages, and their age therefore cannot be well constrained in this way. However, they noted an increasing number of stars with low  $H\alpha$  EW as age increases. In particular, mid-M-type stars show a strong  $H\alpha$  decline after 1 Gyr, and TOI-732 is therefore likely to be older than one billion years, which is consistent with our kinematic age estimate.

A further indication for the evolutionary stage of TOI-732 may come from M-dwarf gyrochronology. Pass et al. (2022) found that M dwarfs usually start spinning down at about 2-3 Gyr. Given the  $H\alpha$ -based inactivity of TOI-732, it is likely that the star is older than the turning-point age of 2-3 Gyr, which again agrees with the stellar evolutionary stage we inferred from kinematics.

### 3. Photometric data

Both C20 and N20 have performed a photometric analysis of the system based on one TESS sector and several observations taken with ground-based facilities (see Table 2). L22 used exclusively space-based observations instead, but added a second TESS sector for the photometric characterisation. In addition to using all the ground- and space-based data that were published in the literature, we added a significant number of space-based data as we

Table 2: Photometric observations from ground-based facilities.

Telescope	Planet	Start Date [UTC]	Duration [h]	Filter
CTIO	b	2019-06-09	2.4	z'
CTIO	b	2019-06-16	2.7	z'
SAAO	b	2019-06-17	3.1	g' z'
SSO	c	2020-01-04	3.8	B
Trappist-N	c	2019-11-12	4.4	z'
OSN	c	2019-11-12	3.5	V R
OAA	b	2020-01-31	6.5	I
MEarth	c	2020-02-10	3.8	RG715
MuSCAT2	b	2019-12-29	2.7	g' r' i' z'
MuSCAT2	b	2020-01-25	2.7	g' r' i' z'
MuSCAT2	b	2020-01-28	3.1	g' r' i' z'
MuSCAT2	b	2020-01-31	1.9	g' r' i' z'
MuSCAT2	c	2019-12-11	3.4	g' r' i' z'
MuSCAT2	c	2020-01-29	6.0	g' r' i' z'

benefited from a further TESS sector and collected 25 CHEOPS 287 visits, 17 of which contain transit events, while the remaining 288 8 are short observations that were not time constrained (fillers) 289 with the aim of monitoring stellar activity (see Table 3). There- 290 fore, our photometric analysis is based on a total of  $\sim 140$  transit 291 events (spread over 132 different LCs), which enabled us to con- 292 siderably improve the photometric properties of the system. All 293 details of the available LCs are given below. 294

#### 3.1. TESS observations

The Transiting Exoplanet Survey Satellite (TESS, Ricker et al. 296 2015) observed the system in cycle 1 (Sector 8; from 28 Febru- 297 ary to 26 March 2019) in cycle 3 (Sector 35; from 9 February to 298 7 March 2021), and in cycle 5 (Sector 62; from 12 February to 299 10 March 2023). The data were downloaded from the Mikulski 300 Archive for Space Telescopes (MAST)<sup>4</sup>, and we used the pre- 301 search data conditioned simple aperture photometry (PDCSAP) 302 LCs, as processed by the Science Processing Operation Center 303 (SPOC; Jenkins et al. 2016). 304

After rejecting data with a poor-quality flag and performing a 305 five median-absolute-deviation (MAD) clipping on the flux val- 306 ues to discard the outliers, we extracted the temporal windows 307 that were centred on each transit event containing  $\sim 4$  hours of 308 out-of-transit data both before and after the transit for detrend- 309 ing purposes. Following this procedure, we obtained 81 TESS LCs, 310 5 of which contain the transits of both planets because their tran- 311 sits are very close in time. Each LC lists the epoch of observation 312 ( $t$ ), the normalised PDCSAP flux with its uncertainty, and other 313 parameters that are available from the TESS data products, such 314 as  $\text{MOM\_CENTR1}$ ,  $\text{MOM\_CENTR2}$  (hereafter denoted with  $x$  and  $y$ , 315 respectively), and  $\text{POS\_CORR1}$  and  $\text{POS\_CORR2}$  (hereafter denoted 316 with  $dx$  and  $dy$ , respectively)<sup>5</sup>. 317

#### 3.2. CHEOPS observations

The Characterising Exoplanet Satellite (CHEOPS, Benz et al. 318 2021) collected 25 LCs of TOI-732 from 8 January to 10 April 319 2022. Because nearby bright background stars strongly contam- 320 inate the observations, we used a mask of 321

<sup>4</sup> <https://mast.stsci.edu/portal/Mashup/Clients/Mast/Portal.html>

<sup>5</sup> See <https://tasoc.dk/docs/EXP-TESS-ARC-ICD-TM-0014-Rev-F.pdf> for further details about the TESS Science Data Products.

Table 3: Details of the CHEOPS visits.

Counter	File key	Planet	Start date [UTC]	Duration [h]
CH 1	CH_PR100031_TG047301_V0200	b c	2022-01-08T14:21:12.4	6.79
CH 2	CH_PR100031_TG047201_V0200	b	2022-01-18T12:45:11.5	7.10
CH 3	CH_PR100031_TG047601_V0200	b c	2022-01-20T20:47:11.6	6.92
CH 4	CH_PR100031_TG047202_V0200	b	2022-01-24T15:56:12.3	6.50
CH 5	CH_PR100031_TG047203_V0200	b	2022-01-25T10:22:12.3	6.50
CH 6	CH_PR100031_TG048301_V0200	0	2022-01-26T15:15:11.5	2.57
CH 7	CH_PR100031_TG047204_V0200	b	2022-01-26T23:29:11.5	5.85
CH 8	CH_PR100031_TG047205_V0200	b	2022-01-30T01:40:11.5	5.89
CH 9	CH_PR100031_TG048501_V0200	b c	2022-02-01T23:28:12.4	12.71
CH 10	CH_PR100031_TG048302_V0200	0	2022-02-02T14:28:12.6	3.20
CH 11	CH_PR100031_TG048401_V0200	b	2022-02-03T12:49:11.6	10.05
CH 12	CH_PR100031_TG047206_V0200	b	2022-02-04T10:14:11.6	6.05
CH 13	CH_PR100031_TG048303_V0200	0	2022-02-26T11:40:12.5	3.20
CH 14	CH_PR100031_TG049701_V0200	b	2022-03-04T18:42:11.4	9.64
CH 15	CH_PR100031_TG048304_V0200	0	2022-03-05T04:32:12.5	3.20
CH 16	CH_PR100031_TG049501_V0200	b c	2022-03-10T20:47:12.6	11.66
CH 17	CH_PR100031_TG049702_V0200	b	2022-03-12T09:46:12.6	9.79
CH 18	CH_PR100031_TG049703_V0200	b	2022-03-16T07:34:12.6	11.26
CH 19	CH_PR100031_TG049704_V0200	b	2022-03-19T09:33:12.5	10.14
CH 20	CH_PR100031_TG048305_V0200	0	2022-03-22T04:34:12.5	3.20
CH 21	CH_PR100031_TG049705_V0200	b	2022-03-22T11:22:12.5	10.17
CH 22	CH_PR100031_TG048306_V0200	0	2022-03-26T22:23:12.5	3.20
CH 23	CH_PR100031_TG048307_V0200	b	2022-03-29T11:42:12.4	3.20
CH 24	CH_PR100031_TG048308_V0200	0	2022-04-01T05:19:13.0	3.20
CH 25	CH_PR100031_TG048309_V0200	0	2022-04-10T15:48:12.5	3.08

**Notes.** Within the visits targeting TOI-732 c, also transits of the ultra-short period planet TOI-732 b are present. The “0” flag in the third column indicates a filler visit.

322 inate the aperture photometry, we opted to extract point-spread  
 323 function (PSF) photometry using the PIPE package<sup>6</sup> (Morris  
 324 et al. 2021; Brandeker et al. 2022). The raw CHEOPS LCs are  
 325 shown in Appendix A.

326 In addition to the parameters given by PIPE (i.e. the stellar  
 327 flux and the  $x$ - and  $y$ -location of the target PSF centroid on the  
 328 detector), we added a few more vectors to the information com-  
 329 prising the CHEOPS LCs that were to be used for the following  
 330 data detrending. In detail, these vectors are produced by the de-  
 331 fault data reduction pipeline (DRP, Hoyer et al. 2020) v.13.1, and  
 332 they are the spacecraft roll angle (`roll`), the flux due to contami-  
 333 nating background stars (`cont a`), the smearing effect that is seen  
 334 as trails on the CCD (`smear`), and the background flux (`bg`) due  
 335 to zodiacal light, for example.

336 Among these data products, the stellar flux measured by  
 337 CHEOPS usually exhibits a highly variable pattern against the  
 338 roll angle (see e.g. Bonfanti et al. 2021 for a broader discussion  
 339 about this topic). As our global LC+RV modelling (see Sec. 5.1)  
 340 only accounts for polynomials when the time series is decorre-  
 341 lated, it would be hard to model the flux versus `roll` pattern.  
 342 Therefore, after masking out the in-transit data points, we pre-  
 343 liminary detrended the PIPE flux against roll angle via Gaussian  
 344 processes (GPs; Rasmussen & Williams 2005) using a Matérn  
 345  $3/2$  kernel (Foreman-Mackey et al. 2017). We duly increased the  
 346 error bars of the flux by adding the standard deviation of the GP  
 347 model in quadrature.

### 3.3. Ground-based observations

348 Several LCs taken with ground-based facilities from 2019 up  
 349 to 2020 are available on the ExoFOP webpage<sup>7</sup>. In particular,  
 350 we downloaded the data obtained with the following one-meter-  
 351 class telescopes that are part of the Las Cumbres Observatory  
 352 Global Telescope (LCOGT) network (Brown et al. 2013), which  
 353 are located at the Cerro Tololo Inter-American Observatory  
 354 (CTIO), the South Africa Astronomical Observatory (SAAO),  
 355 and the Siding Spring Observatory (SSO). In addition, we down-  
 356 loaded LCs acquired with: (i) the 60 cm Trappist-North telescope  
 357 (Jehin et al. 2011; Barkaoui et al. 2019) at Oukaimeden Obser-  
 358 vatory in Morocco; (ii) the 150 cm (T150) telescope at the Ob-  
 359 servatorio de Sierra Nevada (OSN)<sup>8</sup> in Granada, Spain; (iii) the  
 360 40 cm telescope at the Observatori Astronòmic Albanya<sup>9</sup>  
 361 in Catalonia, Spain; (iv) the 40 cm telescope array at the Fred  
 362 Lawrence Whipple Observatory (FLWO) in Arizona (MEarth  
 363 project; Charbonneau et al. 2008).  
 364

365 We also retrieved four transit LCs of TOI-732 b and two tran-  
 366 sit LCs of TOI-732 c each observed in four different filters with  
 367 the MuSCAT2 multi-colour imager (Narita et al. 2019) installed  
 368 on the 1.5 m Telescopio Carlos Sánchez (TCS) at the Teide Ob-  
 369 servatory in Tenerife, Spain. MuSCAT2 is equipped with four  
 370 CCDs, each of which has  $1024 \times 1024$  pixels with a field of view  
 371 of  $7.4 \times 7.4$  square arcmin. The instrument is capable of obtain-  
 372 ing simultaneous images in the  $g'$ ,  $r'$ ,  $i'$ , and  $z_s$  bandpasses. The  
 373 basic data reduction (i.e. dark and flat-field calibrations) was per-

<sup>7</sup> <https://exofop.ipac.caltech.edu/tess/target.php?id=36724087>

<sup>8</sup> <https://www.osn.iaa.csic.es/en/>

<sup>9</sup> <https://www.observatorialbanya.com/en/albanya-astronomical-observatory>

<sup>6</sup> <https://github.com/alphapsa/PIPE>

Table 4: RV data employed in the combined analysis.

Instrument	Start date [UTC]	Time span [d]	Data points [#]
HARPS	2019-06-21	247	33
IRD	2019-12-10	1	4
HARPS-N	2019-12-14	92	30
CARMENES	2019-12-27	54	52
iSHELL	2020-01-25	37	8
MAROON-X	2021-02-22	102	38

formed by the MuSCAT2 pipeline (Parviainen et al. 2019). This pipeline is also capable of fitting a transit model including instrumental systematics and a photometric aperture optimised to reduce the light-curve scatter. For all the transits of planets b and c observed by MuSCAT2, we found an optimal aperture for the target star of 13.92 arcsec.

Finally, all LCs that were taken with a time cadence shorter than one minute were downsampled by binning the data to a one-minute cadence. Further details about the telescope properties and the observational setups can be found in C20 and N20.

## 4. Radial velocity data

Both C20 and N20 combined photometric and RV data. In particular, C20 analysed the RV time series obtained with HARPS and HARPS-N (Cosentino et al. 2012), while N20 separately analysed the RV time series obtained with CARMENES (Quirrenbach et al. 2014, 2018), IRD (Kotani et al. 2018), and iSHELL (Rayner et al. 2016, 2022). L22 performed an RV analysis using the data points coming from these five spectrographs together, while in our case, we further added MAROON-X observations as detailed below for a total of 165 data points. A summary of the RV data employed in our global analysis is given in Table 4.

### 4.1. Literature radial velocity data

The RV measurements available in the literature were retrieved directly from C20 and N20, who also provided a detailed description of the RV data reduction. We briefly recall here that C20 obtained 33 spectra with the HARPS echelle spectrograph mounted at the ESO 3.6 m telescope at the La Silla Observatory in Chile and 30 spectra with the HARPS-N echelle spectrograph mounted at Telescopio Nazionale Galileo (TNG; Cosentino et al. 2000; Oliva 2006) in the Canary Islands, Spain. The corresponding RV measurements along with their error bars were extracted using the TERRA reduction pipeline (Anglada-Escudé & Butler 2012).

Instead, N20 obtained 52 spectra with the CARMENES spectrograph mounted on the 3.5 m Calar Alto Observatory in Almería, Spain. They obtained the RV measurements using the `serval` code (Zechmeister et al. 2018) and applied the necessary corrections following Trifonov et al. (2018) and Kaminski et al. (2018). N20 also took five spectra with the IRD instrument mounted on the Subaru 8.2 m telescope in Mauna Kea, Hawaii. After discarding one low-quality observation, they reduced the spectra using `iraf` (Tody 1986, 1993) and extracted the RV measurements through the Subaru/IRD dedicated pipeline (Hirano et al. 2020). Using the iSHELL spectrometer mounted on the NASA Infrared Facility (IRTF) in Mauna Kea, Hawaii, N20 further collected eight RV measurements by applying the spectral reduction method presented in Cale et al. (2019).

Each of the five instruments is characterised by its own offset and is affected by a different jitter term. We therefore organised these RV time series as five independent data sets. These five RV time series contain the vectors of epochs, RV measurements, and RV error bars as found in the literature.

### 4.2. MAROON-X

We observed TOI-732 with MAROON-X, which is a high-precision echelle spectrograph installed on the 8.1 m telescope Gemini-North (Seifahrt et al. 2018, 2022), 19 times between February and June 2021. The MAROON-X data were reduced with a python3 pipeline based on the pipeline originally used for the CRIRES instrument (Bean et al. 2010), and the RVs were calculated with a version of `serval` (Zechmeister et al. 2020) modified to work on MAROON-X data. `serval` calculates RVs by least-squares fitting each individual spectrum to a template created by co-adding all spectra together. The `serval` routine also extracts the chromatic index (`crx`), the differential line width (`dlw`), and the  $H\alpha$  index, which may be useful for data detrending. The wavelength calibration is accomplished by simultaneously observing the science target with an etalon spectra, and the etalons themselves are calibrated using a ThAr lamp.

MAROON-X has two separate CCDs, each with slightly different wavelength coverages, which are exposed simultaneously. The blue channel (500—670 nm) and the red channel (650—920 nm) were treated as two separate instruments for the purposes of this analysis because they have a different wavelength coverage and thus capture different stellar signals. We achieved a median S/N of 200 in the red channel and 77 in the blue channel, which corresponded to median RV uncertainties of  $0.5 \text{ m s}^{-1}$  in the red channel and  $1 \text{ m s}^{-1}$  in the blue channel. The higher signal in the red channel is expected for the late stellar spectral type.

MAROON-X is a visitor instrument on Gemini-North, and it is thus connected and disconnected multiple times over the course of a semester. It organises its data into discretised runs. In particular, the TOI-732 data were collected over the course of three runs in 2021 (one in February, one in April, and one in May). Combined with the roughly  $2.5 \text{ cm s}^{-1} \text{ d}^{-1}$  RV drift of the etalon calibrations, this results in small offsets between the RVs of MAROON-X data taken in separate runs. We therefore treated each run of the MAROON-X data as an independent RV time series and further distinguished the data taken via the red and blue channel. That is, we fit six independent MAROON-X RV time series. Accounting for the five RV time series described in Sec. 4.1, we analysed a total of 11 RV time series.

## 5. Methods and results

### 5.1. Global light-curve and radial-velocity modelling

We jointly analysed the 132 LCs and 11 RV time series using the MCMCI code (Bonfanti & Gillon 2020), where we switched off the interaction with stellar evolutionary models to avoid a dramatic increase in computational time due to the large data sets. In short, the code fit the LCs against the photometric model of Mandel & Agol (2002) and the RV data against a Keplerian model using an MCMC approach.

On the stellar side, we adopted  $T_{\text{eff}}$ ,  $[\text{Fe}/\text{H}]$ ,  $M_{\star}$ , and  $R_{\star}$  as jump parameters that were subject to Gaussian priors based on the values reported in Table 1. The reason for this choice is twofold. On the one hand, both  $M_{\star}$  and  $R_{\star}$  induce a prior on the mean stellar density  $\rho_{\star}$ , which better constrains the transit fitting. On the other hand, stellar parameters are the start-

ing point for interpolating within ATLAS9<sup>10</sup> grids of quadratic limb-darkening (LD) coefficients ( $u_1, u_2$ ), which were set up for each of the 11 photometric filters using the code by Espinoza & Jordán (2015). Gaussian priors were then imposed on the 11 interpolated pairs ( $u_1, u_2$ ) as summarised in Table C.1, but the actual LD-related jump parameters were derived from a linear combination of ( $u_1, u_2$ ) following Holman et al. (2006) to reduce their mutual correlation.

For each planet, the jump parameters were the transit depth  $dF \equiv \left(\frac{R_p}{R_*}\right)^2$ , the impact parameter  $b$ , the mid-transit time  $T_0$ , the orbital period  $P$ , and the RV semi-amplitude  $K$ . We assumed a circular orbit for TOI-732 b as its ultrashort orbital period implies a tide-induced circularisation timescale of  $\sim 15$  Myr (Matsumura et al. 2008). We instead fit for the eccentricity of TOI-732 c using the parametrisation ( $\sqrt{e} \cos \omega$ ,  $\sqrt{e} \sin \omega$ ), where  $e$  is the eccentricity and  $\omega$  is the argument of periastron. In the case of TESS observations, N20 noted that a close-in star, namely TIC 36724077, was located within the aperture mask. We therefore fitted for a dilution factor following their estimate. All planetary jump parameters were subject to uniform unbounded priors (except for the physical limits). For details about the adopted parametrisations, we refer to Bonfanti & Gillon (2020 § 2.1.2) and references therein.

The MCMCI tool is able to detrend data against time and the ancillary vectors of the LC and RV time series along the MCMC process via polynomial interpolation. To find the best polynomial order for each detrending parameter of each LC and RV time series, we launched several preliminary MCMC runs and changed the polynomial order of one detrending parameter at a time. We finally selected the best detrending polynomial baseline (see Table C.2) according to the Bayesian information criterion (BIC; Schwarz 1978).

We then launched a first MCMC run of 200 000 steps (burn-in: 40 000 steps) to evaluate the impact of the white and red noise as detailed in Pont et al. (2006) and Bonfanti & Gillon (2020) to properly rescale the photometric errors and provide reliable uncertainties on the fitted parameters. After this, we performed the final MCMCI analysis made of two independent runs (each comprising 200 000 steps with a burn-in of 40 000 steps) to check the posterior distribution convergence through the Gelman-Rubin (GR) test (Gelman & Rubin 1992).

The chains converged according to the GR statistic ( $\hat{R} \lesssim 1.006$  for all the jump parameters), and we obtained planetary radii of  $R_b = 1.325^{+0.057}_{-0.058} R_\oplus$  and  $R_c = 2.39^{+0.10}_{-0.11} R_\oplus$ , masses of  $M_b = 2.46 \pm 0.19 M_\oplus$  and  $M_c = 8.04^{+0.50}_{-0.48} M_\oplus$ , and thus densities of  $\rho_b = 5.8^{+1.0}_{-0.8} \text{ g cm}^{-3}$  and  $\rho_c = 3.24^{+0.55}_{-0.43} \text{ g cm}^{-3}$ . All relevant system parameters as derived from our MCMC global analysis are listed in Tables 5, C.1, and C.3. The phase-folded and detrended LCs of both TOI-732 b and c, as observed by both TESS and CHEOPS, are shown in Fig. 1, while the LCs taken by ground-based facilities are shown in Appendix B. Finally, the phase-folded and detrended RV time-series of both TOI-732 b and c are displayed in Fig. 2.

The bulk densities obtained for both planets are at the  $\sim 15\%$  precision level, and only  $\sim 20\%$  of all known planets orbiting M dwarfs have been characterised to a similar or better precision<sup>11</sup>. This is a consequence of the precision we reached on both the transit depths of TOI-732 b and TOI-732 c (4.4% and 4.0%, respectively) and the radial velocity semi-amplitudes (6.2% and

Table 5: Parameters of the TOI-732 system.

Parameter	TOI-732 b	TOI-732 c
$P$ [d]	$0.76837931^{+0.00000039}_{-0.00000042}$	$12.252284 \pm 0.000013$
$T_0^{(a)}$ [BJD]	$9606.58098^{+0.00032}_{-0.00040}$	$9600.54227^{+0.00066}_{-0.00065}$
$b$	$0.462^{+0.063}_{-0.094}$	$0.794^{+0.023}_{-0.027}$
$dF$ [ppm]	$1032^{+44}_{-45}$	$3355^{+140}_{-130}$
$\frac{R_p}{R_*}$	$0.03212^{+0.00068}_{-0.00072}$	$0.0579^{+0.0012}_{-0.0011}$
$W$ [min]	$47.90 \pm 0.73$	$92.5^{+1.7}_{-1.6}$
$i$ [°]	$86.10^{+0.92}_{-0.68}$	$88.958^{+0.074}_{-0.068}$
$a$ [AU]	$0.01195^{+0.00028}_{-0.00029}$	$0.0757 \pm 0.0018$
$\frac{a}{R_*}$	$6.79^{+0.29}_{-0.25}$	$43.0^{+1.8}_{-1.6}$
$K$ [m s <sup>-1</sup> ]	$3.24 \pm 0.20$	$4.22 \pm 0.16$
$e$	0 (fixed)	$0.024^{+0.032}_{-0.017}$
$\omega$ [°]	90 (fixed)	$-66^{+110}_{-50}$
$T_{\text{eq}}^{(b)}$ [K]	$903 \pm 26$	$359 \pm 10$
$S$ [ $S_\oplus$ ]	$111^{+13}_{-12}$	$2.76^{+0.33}_{-0.31}$
$R_p$ [ $R_\oplus$ ]	$1.325^{+0.057}_{-0.058}$	$2.39^{+0.10}_{-0.11}$
$M_p$ [ $M_\oplus$ ]	$2.46 \pm 0.19$	$8.04^{+0.50}_{-0.48}$
$\rho_p$ [g cm <sup>-3</sup> ]	$5.8^{+1.0}_{-0.8}$	$3.24^{+0.55}_{-0.43}$

**Notes.** Uncertainties are defined as the 68.3% credible intervals of the posterior distributions. All fitted parameters, that is  $P$ ,  $T_0$ ,  $b$ ,  $dF$ ,  $K$ ,  $e$ , and  $\omega$ , were subject to uniform unbounded priors (except for physical limits) following the parameterisations detailed in Bonfanti & Gillon (2020).

<sup>(a)</sup> Shifted by  $-2\,450\,0000$ . <sup>(b)</sup> Assuming zero albedo.

Table 6: Comparison between literature uncertainties and those derived in this work on the orbital periods  $P$ , the transit depths  $dF$ , and the RV semi-amplitudes  $K$  of the planets.

Planet	Uncertainty	C20	N20	L22	This work
TOI-732 b	$\Delta P$ [s]	4.7	0.12	0.045	0.035
	$\frac{\Delta dF}{dF}$ [%]	9.3	6.5	6.9	4.3
	$\frac{\Delta K}{K}$ [%]	18	10	8.5	6.2
TOI-732 c	$\Delta P$ [s]	251	5.9	3.0	1.1
	$\frac{\Delta dF}{dF}$ [%]	12	5.5	9.6	4.0
	$\frac{\Delta K}{K}$ [%]	17	10	8.0	3.8

3.8%, respectively), which marks a significant improvement over what was reported so far in the literature, as summarised in Table 6.

Based on the large amount of available data and the broad temporal baseline spanning four years, we were able to reduce the uncertainties on the orbital periods of both planets by more than two orders of magnitude with respect to what was reported by C20. Even comparing our results with those of L22, who derived the most precise ephemerides so far, we improved the uncertainty on the planetary orbital periods by a factor of  $\sim 1.3$  and  $\sim 2.7$  for planets b and c, respectively (see Table 6). By propagating our ephemerides, we computed that the  $1\sigma$  uncertainties on the transit timings of the two planets are comparable to the respective transit durations after  $\sim 170$  years from now.

<sup>10</sup> <http://kurucz.harvard.edu/grids.html>
<sup>11</sup> Source: Nasa Exoplanet Archive, <https://exoplanetarchive.ipac.caltech.edu/>




Fig. 1: Phase-folded and detrended LCs showing the transit of TOI-732 b (first column) and TOI-732 c (second column) as observed by CHEOPS (first row) and TESS (second row). The original data points are shown in blue, the binned data points are shown in black (binning of 10 min), and the transit model is displayed in red.

## 5.2. Internal structure of the planets

We modelled the internal structure of both TOI-732 b and c using a neural-network-based Bayesian inference scheme following the method that was described in detail in Leleu et al. (2021) and is based on Dorn et al. (2017). As input parameters, we used transit depths, periods, and the mass relative to that of the star for both planets, as well as some of the stellar parameters, namely mass, radius, age, effective temperature, [Si/H], [Mg/H], and [Fe/H]. We modelled both planets simultaneously, assuming that they consist of four fully distinct layers that we modelled according to the equations of state of Hakim et al. (2018) (an inner iron core with up to 19% sulphur), Sotin et al. (2007) (a silicate mantle consisting of Si, Mg, and Fe) and Haldemann et al. (2020) (a condensed water layer), with a H-He envelope modelled following (Lopez & Fortney 2014) on top. Furthermore, we assumed that the Si, Mg, and Fe ratios of both planets match those of the star (Thiabaud et al. 2015), even if we note that despite an expected trend between stellar and planetary composition, the correlation might not necessarily be strict (Adibekyan et al. 2021).

As the problem of determining the internal structure of a planet is highly degenerate, the results of our analysis depend

on our choice of prior. For the mass fractions of the inner iron core (i.e. the mantle layer and the water layer), all calculated with respect to the inner part of the planet without the H-He layer, we sampled from a prior that is uniform on the simplex on which they all add up to 1. Furthermore, we implemented an upper limit for the water-mass fraction of 0.5, in accordance with Thiabaud et al. (2014) and Marboeuf et al. (2014). We also used a prior that is log-uniform for the mass of the H-He envelope.

The results of our analysis are summarised in Figures 3 and 4. The derived posteriors of the internal structure parameters show us that TOI-732 b is unlikely to host a H-He layer given its density. Meanwhile, the presence of a water layer is possible, but not necessary, as the derived mass and radius values also agree with a purely rocky structure. For TOI-732 c, the posterior distribution of the gas mass is instead quite well constrained, with a median of  $M_{\text{gas},c} = 0.02^{+0.05}_{-0.02} M_{\oplus}$ , which corresponds to a thickness of  $R_{\text{gas},c} = 0.40^{+0.24}_{-0.27} R_{\oplus}$  (errors are the 5th and 95th percentile of the distribution). However, the presence of a water layer is completely unconstrained.

Figure 5 locates TOI-732 b and TOI-732 c on the mass-radius (MR) diagram along with M-dwarf planets with  $R_p < 4 R_{\oplus}$  and  $M_p < 30 M_{\oplus}$  whose precision on the radius and mass are better than 8% and 25%, respectively. When TOI-732 b and c are in-




Fig. 2: Phase-folded and detrended RV time series of TOI-732 b (Top) and TOI-732 c (Bottom), obtained after subtracting the signal of the other planet. The corresponding Keplerian model is superimposed in red. The different colours mark different instruments, namely HARPS (black), IRD (light green), HARPS-N (blue), CARMENES (magenta), iSHELL (cyan), MAROON-X (yellow, orange, and deep green for observations taken in February, April, and May 2021, respectively). As MAROON-X has two different channels, full and empty symbols represent data acquired using the blue and red channel, respectively (see text for further details).

598 cluded, this exoplanet sample (hereafter denoted as  $M_{\text{sample}}^{12}$ )  
599 is made of 45 well-characterised planets (a mean planetary  
600 bulk density above the  $3\sigma$  level). The main parameters of the  
601  $M_{\text{sample}}$  are listed in Tab. C.4. Along with the planets belonging  
602 to the  $M_{\text{sample}}$ , Fig. 5 also displays two sets of theoretical models  
603 for a planet composition that correspond to  $T_{\text{eq}} = T_{\text{eq},b} = 900$   
604 K (solid lines) and  $T_{\text{eq}} = T_{\text{eq},c} = 360$  K (dashed lines) using the  
605 BICEPS model (Haldemann et al. 2023). In addition, we further  
606 collected the MR model as computed by Aguichine et al.  
607 (2021) for steam worlds made of 50% water + 50% rocks with  
608  $T_{\text{eq}} = 400$  K  $\approx T_{\text{eq},c}$  (the dashed cyan line). Theoretical models  
609 of rocky and/or iron worlds do not depend upon  $T_{\text{eq}}$ , but dif-

<sup>12</sup> Planetary data of M dwarfs (that is stars with  $T_{\text{eff}} < 4000$  K) were properly filtered and downloaded from the NASA Exoplanet Archive as of 27 July 2023




Fig. 3: Corner plot showing the posteriors of the main parameters of our internal structure analysis for TOI-732 b. The titles of each column correspond to the median of the distribution, with the 5<sup>th</sup> and 95<sup>th</sup> percentiles as the uncertainty values. From left to right, the depicted internal structure parameters are the mass fractions of the inner iron core and of the water layer (both calculated with respect to the condensed part of the planet without the H-He layer), the molar fractions of Si and Mg in the mantle, the molar fraction of Fe in the inner core, and the total mass of H-He in Earth masses on a logarithmic scale. The mass fractions of the inner core and the water layer add up to one, with the mass fraction of the mantle layer (not shown) by construction.




Fig. 4: Same as Figure 3, but for TOI-732 c.




Fig. 5: Mass-radius diagram of M-dwarf planets with  $R_p < 4 R_\oplus$  and  $M_p < 30 M_\oplus$  whose  $R_p$  and  $M_p$  precision is better than 8% and 25%, respectively. All planets are colour-coded with respect to their equilibrium temperature ( $T_{\text{eq}}$ ) and in particular, TOI-732 b and TOI-732 c are marked by a star. Following the colour scheme given in the legend, two different sets of planet composition models generated with BICEPS (Haldemann et al. 2023) are displayed. The solid and dashed lines are obtained for  $T_{\text{eq}} = T_{\text{eq,b}} = 900$  K and  $T_{\text{eq}} = T_{\text{eq,c}} = 360$  K, respectively. The 50% steam + 50% Earth-like line corresponds to the model of Aguichine et al. (2021) for  $T_{\text{eq}} = 400$  K. An Earth-like composition implies a mixture of 32.5% iron and 67.5% silicates. The dotted black lines correspond to the loci of constant density, that is, 0.5, 1, 3, 5, and 10  $\text{g cm}^{-3}$  (from top to bottom).

602 ferences become noticeable when water and/or H-He envelopes  
 603 are added to the planet structure. The MR diagram confirms that  
 604 TOI-732 b is likely rocky with a possible iron core, while TOI-  
 605 732 c is likely rich in volatiles. As shown above, inferring the  
 606 internal planet structure from observables is a degenerate prob-  
 607 lem and other mixtures of silicates, gas, and water (which is in-  
 608 deed unconstrained according to our modelling of TOI-732 c)  
 609 may produce a  $(M_p, R_p)$  pair consistent with the observations.  
 610 For example, we note that the MR location of TOI-732 c is com-  
 611 patible with either a rocky planet surrounded by a H-He envelope  
 612 (1% by mass) or a steam world consisting of water and rocks in  
 613 the same proportion by mass.

## 622 6. Radius valley of M-dwarf planets

623 According to their radii, TOI-732 b and TOI-732 c are located on  
 624 the two opposite sides of the radius valley. Although some degen-  
 625 eracy is expected when modelling the internal structures of  
 626 planets, we concluded in Sect. 5.2 that TOI-732 b is not likely  
 627 to hold any gaseous envelope, while TOI-732 c cannot be just  
 628 purely rocky. When we also consider the mean planetary den-  
 629 sities,  $\rho_b > \rho_c$ , which can lead to a classification of the inner  
 630 planet as a super-Earth and of the outer one as a mini-Neptune,  
 631 the TOI-732 system has a quite common architecture (e.g. Ciardi  
 632 et al. 2013; Weiss et al. 2018; Mishra et al. 2023).

## 633 6.1. Radius valley dependence on orbital period

634 Taking a step further, we studied the radius valley  $R_{p,\text{valley}}$  for  
 635 M dwarfs as a function of planet orbital period  $P$  by using our  
 636 Msample. Several theoretical studies (e.g. Owen & Wu 2017;  
 637 Lopez & Rice 2018; Gupta & Schlichting 2019; Wyatt et al.  
 638 2020; Lee & Connors 2021; Rogers et al. 2021; Affolter et al.  
 639 2023) have quantified different  $d \log R_{p,\text{valley}} / d \log P$  slopes char-  
 640 acterising the radius valley depending on the specific formation  
 641 and evolution mechanisms causing it (e.g. impact erosion, photo-  
 642 evaporation, core-powered mass loss, or late planet formation in  
 643 either gas-poor or even gas-empty discs). It is worth emphasising  
 644 that planets formed in a gas-poor environment may also be sub-  
 645 ject to thermally driven mechanisms (i.e. photo-evaporation and  
 646 core-powered mass loss). Hereafter, the discussion of thermally  
 647 driven mechanisms is intended to involve planets that have not  
 648 formed in a gas-poor environment, unless stated otherwise.

649 As summarised in Tab. 7, a negative slope is theoretically  
 650 expected for both impact erosion and thermally driven mass-  
 651 loss mechanisms, with the slope becoming milder when passing  
 652 from the former to the latter. Furthermore, in the case of late-  
 653 time planet formation within a gas-poor environment, the slope  
 654 is even shallower (but still negative) when photo-evaporation is  
 655 considered to be at play afterwards. As emphasised by Lee &  
 656 Connors (2021), a positive  $d \log R_{p,\text{valley}} / d \log P$  is sometimes  
 657 incorrectly associated to late-time planet formation, according to  
 658 the work by Lopez & Rice (2018). However, Lopez & Rice  
 659 (2018) computed the expected scaling between  $R_p$  and  $P$  as-  
 660 suming a gas-empty scenario, and the positive slope line they  
 661 derived therefore just corresponds to the maximum radius that  
 662 can be reached by a purely rocky planet. Therefore, this loc-  
 663 us of points does not trace the radius valley dividing rocky  
 664 planets from sub-Neptune simply because no sub-Neptunes may  
 665 form in a gas-empty environment. Nonetheless, we kept the  
 666  $d \log R_{p,\text{valley}} / d \log P_{L18} = +0.11$  in Table 7 because it sets the  
 667 upper limit of the radius valley slope for a sample of heteroge-  
 668 neous exoplanets in the  $R_p$ - $P$  plane. On the one hand, the purely  
 669 rocky exoplanets that are born in a gas-empty disc would be dis-  
 670 tributed following a positive trend, whose upper limit is given  
 671 by  $d \log R_{p,\text{valley}} / d \log P_{L18}$ . On the other hand, from a disc with  
 672 gas, both super-Earth and sub-Neptunes would be generated, and  
 673 they would appear on the two opposite sides of a descending radi-  
 674 us valley. The full picture that we would see a posteriori in the  
 675  $R_p$ - $P$  plane would be the overlap of these two groups of exo-  
 676 planets, which would show a radius valley with an intermediate  
 677 slope, possibly even positive, depending on the weights of the  
 678 formation mechanisms at play.

679 To study the dependence of the radius valley on planetary  
 680 orbital period, we followed the same approach as in Van Eylen  
 681 et al. (2018) and Ho & Van Eylen (2023), but focused on M-  
 682 dwarf hosts ( $M_\star \lesssim 0.6 M_\odot$ ). This complements the stellar mass  
 683 range spanned by the F, G, and K type stars investigated by Ho  
 684 & Van Eylen (2023). In detail, we first clustered our M-dwarf  
 685 exoplanets into two different groups, according to their location  
 686 with respect to the radius valley (above or below), by perform-  
 687 ing a Gaussian mixture model selection (e.g. Huang et al. 2017;  
 688 Fruhwirth-Schnatter et al. 2018). To this end, we employed the  
 689 PYTHON *sklearn* GridSearchCV class, which allows specifying  
 690 four different covariance types to define the clustering. After  
 691 rescaling the period  $P$  by a factor of five to avoid misclassifica-  
 692 tion (Ho & Van Eylen 2023), we fit the selection model within  
 693 the  $\log R_p$ - $\log P$  plane, and we finally selected the model inferred  
 694 from the spherical covariance type, which has the lowest associ-  
 695 ated BIC.

Table 7: Radius valley slopes  $m \equiv d \log R_{p,\text{valley}}/d \log P$  as predicted from theory for different scenarios.

Model	$m$	Reference
Impact erosion	-0.33	Wyatt et al. (2020)
Photo-evaporation	[-0.25, -0.16]	Owen & Wu (2017)
Thermally-driven mass loss	-0.10	Affolter et al. (2023)
Photo-evaporation in gas-poor discs	[-0.15, -0.08]	Lee & Connors (2021)
Gas-empty formation	+0.11	Lopez & Rice (2018)

**Notes.** The photo-evaporation model has been developed using the energy-limited formula (e.g. Watson et al. 1981; Erkaev et al. 2007) and accounting for different efficiency values of stellar high-energy photons in the atmospheric mass removal. Instead, the thermally-driven mechanisms have been modelled via hydrodynamic simulations that couple photo-evaporation and core-powered mass loss. [a, b] denotes a range of values from a to b.

696 After this, we followed a support vector machine (SVM) procedure (e.g. Cortes & Vapnik 1995; Ben-Hur et al. 2002) implemented via the *sklearn* SVC() class. After we set a linear kernel and a penalty parameter  $C = 10$  (see Van Eylen et al. 2018), the *fit* method of SVC() was able to compute the best-fit line separating the two groups of exoplanets in any desired space of covariates. In particular, we obtained

$$\log R_{p,\text{valley}} = -0.065_{-0.013}^{+0.024} \log P + 0.344_{-0.018}^{+0.008}, \quad (1)$$

703 where the uncertainties (at the  $1 \sigma$  level) were computed by bootstrapping the *Msample* 10 000 times and repeating the algorithm outlined above.

706 When compared to the outcome obtained by Ho & Van Eylen (2023) ( $d \log R_{p,\text{valley}}/d \log P_{\text{H}23} = -0.11 \pm 0.02$ ), the slope value we obtained differs by almost a factor of two (tension at the  $2 \sigma$  level), which may suggest that formation and evolution mechanisms enter with different weights in the case of exoplanets orbiting M dwarfs or FGK stars. Instead, when we performed a homogeneous comparison with other works targeting the  $R_{p,\text{valley}}$  slope of exoplanets around low-mass stars, our  $d \log R_{p,\text{valley}}/d \log P$  value is consistent within  $1 \sigma$  with the estimate from L22 ( $d \log R_{p,\text{valley}}/d \log P_{\text{L}22} = -0.02 \pm 0.05$ ), it is milder than the slope found by Van Eylen et al. (2021) ( $d \log R_{p,\text{valley}}/d \log P_{\text{V}21} = -0.11_{-0.04}^{+0.05}$ ), but still consistent at the  $\sim 1 \sigma$  level, and it differs from the outcome of Cloutier & Menou (2020) ( $d \log R_{p,\text{valley}}/d \log P_{\text{C}20} = +0.058 \pm 0.022$ ). The sample of Cloutier & Menou (2020) also comprises planets orbiting K dwarfs (with a spectral type later than K3.5V, i.e.  $M_{\star} \lesssim 0.8 M_{\odot}$ ), and the reason for the difference in slope may be that the  $R_p$  precision for half of the planets they analysed is lower than our 8% threshold (the 99<sup>th</sup>-quantile of their  $R_p$  relative uncertainties is  $\sim 26\%$ ). Instead, both Van Eylen et al. (2021) and L22 focused on planets orbiting M dwarfs alone, and the difference with our  $d \log R_{p,\text{valley}}/d \log P$  value decreases as the selection threshold for the sample is set to a lower  $R_p$  uncertainty (below 20% for Van Eylen et al. (2018) and below 8% for L22). Only the sample by L22 reaches the same precision level as our *Msample* (because we adopted the same selection criteria), but our sample contains 30% more planets (45 versus 34 planets).

734 A visual synthesis of our results is given in Fig. 6, where the best-fit line marking the radius valley (solid grey line) is compared with the theoretical slopes expected from a thermally driven mass-loss model (solid red line; Affolter et al. 2023) and a gas-empty formation model (dashed red line; Lopez &




Fig. 6:  $R_p$  vs  $P$  distribution representing the planets in our *Msample*. Planets classified above and below the radius valley are shown in blue and green, respectively, while the red star-shaped markers are for TOI-732 b and c. The radius valley inferred via the SVM-based method is marked by the solid grey line with the shaded region highlighting the  $1 \sigma$  limits of the best-fit line. The two parallel dashed grey lines are the median boundaries passing through the supporting vectors that determine the location of the solid line. Finally, the red lines represent the theoretically expected  $R_{p,\text{valley}}$  boundary in case of a thermally-driven mass-loss scenario (solid line as computed from Affolter et al. (2023); negative slope) and the  $R_p$  upper limit of planets borned rocky in a gas-empty disc (dashed line as computed from Lopez & Rice (2018); positive slope).

738 Rice 2018). When compared with the theoretical slope expected from a mixed scenario, where both photo-evaporation and core-powered mass loss are at play (that is  $-0.10$ ; Affolter et al. 2023), the negative  $d \log R_{p,\text{valley}}/d \log P$  slope we computed (i.e.  $-0.065_{-0.013}^{+0.024}$ ) is shallower by a factor of  $\sim 1.5$  (tension at the  $\sim 3$  sigma level). Slopes milder than  $-0.10$  possibly tending towards positive values indicate a stronger impact of gas-poor formation according to Lopez & Rice (2018); Lee & Connors (2021). Therefore, we may conclude that although thermally driven mechanisms appear to be statistically prevalent, the currently observed properties of some of the planets orbiting M dwarfs may be caused by late formation in gas-depleted discs. This scenario has indeed been proposed for a few M-dwarf systems, such as TOI-1634 (Cloutier et al. 2021b), where the composition of the close-in USP TOI-1634 b is inconsistent with that of the Earth, or LHS 1903 (Wilson et al. 2023), where the outermost planet at  $P \sim 29.3$  d lacks any gaseous envelope, in contrast to some of the inner planets. An alternative scenario explaining our  $d \log R_{p,\text{valley}}/d \log P$  findings is investigated in Sect. 6.3.

757 As the strength of core-powered mass-loss experienced by a planet scales proportionally to  $R_p T_{\text{eq}}^4$  (Gupta & Schlichting 2019), the colour-coding in Fig. C.1 is an attempt of investigating the impact of core-powered mass loss in shaping the radius valley. However, the colour gradient from the bottom right to top left just reflects the increase in  $R_p T_{\text{eq}}^4$  at greater radii and lower orbital period (hotter planets). The radii of billion-year-old planets dominating the *Msample* are thought to have significantly shrunk during their evolution due to planetary cooling and evaporation, which effect is correlated to the strength of the atmo-

spheric escape (Lopez & Fortney 2014; Chen & Rogers 2016; Kubyskhina & Fossati 2022; Affolter et al. 2023). Thus, the present-day radii cannot unambiguously define the strength of core-powered mass loss because they are not indicative of the escape rates during the early evolution phases, when core-powered mass loss can dominate. In addition, the equilibrium temperature strongly correlates with the (poorly constrained) amount of XUV radiation received by the planet because both  $T_{\text{eq}}$  and XUV radiation scale with the planet distance. Thus, the  $T_{\text{eq}}$  dependence does not allow us to distinguish the inputs from the core-powered and XUV-driven escape mechanisms sufficiently well.

On the other hand, we know from hydrodynamic modeling that core-powered mass loss dominates the atmospheric escape completely if the atmospheric density is sufficiently high in the upper atmospheric layers to prevent the penetration of XUV radiation inside the planetary Roche lobe (Kubyskhina et al. 2018; Kubyskhina 2023). This situation occurs most likely for planets with low masses and small Roche radii (comprising a few  $R_p$  at young ages); of these two parameters, the Roche radius carries more information than the planetary mass alone. Along this line, the two panels of Fig. 7 still represent the planets of our Msample in the  $R_p$ - $P$  plane, but with a specific focus on the role of the core-powered mass-loss mechanism by tracing the size of the planetary Roche radius. We computed the Roche-lobe radius (Eggleton 1983) of each planet

$$R_{\text{Roche}} = \frac{0.49q^{\frac{2}{3}}}{0.6q^{\frac{2}{3}} + \ln(1 + q^{\frac{1}{3}})} a, \quad \text{being } q \equiv \frac{M_p}{M_\star}, \quad (2)$$

as a measure of the region within which a possible atmospheric envelope is bounded to the planet. The larger  $R_{\text{Roche}}$ , the less effective the core-powered atmospheric escape. After normalisation to  $R_p$  (top panel),  $R_{\text{Roche}}$  still maintains the linear dependence upon the semi-major axis  $a$ , and indeed,  $R_{\text{Roche}}/R_p$  increases with the orbital period. On the one hand, while this trend is expected, this panel emphasises on the other hand, that planets on long-period orbits are less subject to core-powered mass loss. Therefore, a rocky planet (i.e. without a low mean-molecular weight envelope) farther away from its host is more likely to be born in a gas-depleted environment.

The bottom panel of Fig. 7 is similar to the top panel, but this time, the colour-coding follows  $R_{\text{Roche}}$  normalised to  $a$ . In this way, we removed the linear dependence of the Roche radius on  $a$ , which means that  $R_{\text{Roche}}$  depends solely on the  $M_p/M_\star$  ratio. Now,  $R_{\text{Roche}}/a$  increases as  $R_p$  increases, with the highest  $R_{\text{Roche}}/a$  values clustering above the radius valley. The larger Roche radius of these planets enabled them to keep their atmospheric envelope, and they therefore appear to be more puffy than the planets below the radius valley.

## 6.2. Dependence of the radius valley on stellar mass

Considering the increasing interest in exploring the trend between the radius valley and the spectral type of the host star (e.g. Wu 2019; Gupta & Schlichting 2020; Rogers et al. 2021; Ho & Van Eylen 2023; Berger et al. 2023), we repeated the SVM analysis described above, but assuming the covariate pair  $(R_p, M_\star)$ , and we derived

$$\log R_{p,\text{valley}} = +0.054_{-0.034}^{+0.049} \log M_\star + 0.319_{-0.016}^{+0.022} \quad (3)$$

(see Fig. 8). Estimates of the radius valley slope  $d \log R_{p,\text{valley}}/d \log M_\star$  that are based on observational data as found in the literature (Berger et al. 2020; Petigura et al.




Fig. 7: Same as Fig. 6, but the markers are colour-coded against the Roche lobe of each planet normalised to the planetary radius (Top) or to the orbital semi-major axis (Bottom).

2022; Ho & Van Eylen 2023) are mainly the results of works focusing on FGK stars, which lead to steeper slopes (although the accompanying uncertainties are about 40% or higher). The only homogeneous comparison currently available is with the work by L22, who found  $d \log R_{p,\text{valley}}/d \log M_\star = +0.08 \pm 0.12$  (consistent with our estimate), which may again suggest that planets orbiting M dwarfs differ from those orbiting FGK stars in the context of the radius valley.

However, from a theoretical perspective, it is hard to draw firm conclusions about the mechanisms underlying the formation and evolution of exoplanets when studying the radius valley within the  $R_p$ - $M_\star$  space. Rogers et al. (2021) cautioned that the  $R_{p,\text{valley}}$ - $M_\star$  slope shows several degeneracies. They theoretically derived that the expected slope does not only depend on  $M_\star$ , but also on the incident bolometric flux  $S$ , and it can be




Fig. 8: Same as Fig. 6, but this time,  $R_p$  is plotted against stellar mass.

expressed as

$$\left. \frac{d \log R_{p,\text{valley}}}{d \log M_\star} \right|_{\text{th}} \approx \alpha \left( \zeta - \frac{2}{3} \right) + \beta, \quad (4)$$

where  $\alpha \equiv \partial \log R_{p,\text{valley}} / \partial \log S$  and  $\beta \equiv \partial \log R_{p,\text{valley}} / \partial \log M_\star$  are predicted, depending on the scenario at play (either photo-evaporation or core-powered mass loss), while  $\zeta$  is the exponent entering the mass-luminosity relation, that is,  $L_\star \propto M_\star^\zeta$ . Because  $\zeta \gg \alpha$  and  $\zeta \gg |\beta|$  (Rogers et al. 2021), the slope value is mainly controlled by  $\zeta$ , which needs to be properly estimated according to the stellar spectral type. Cuntz & Wang (2018) proposed the following expression for  $\zeta = \zeta(M_\star)$  for low-mass stars:

$$\zeta = -141.7M_\star^4 + 232.4M_\star^3 - 129.1M_\star^2 + 33.29M + 0.215 \quad (5)$$

and averaging out that function over our mass range of interest, we obtained  $\zeta_M = 4.0$ . Plugging in the  $(\alpha, \beta)$  predictions by Rogers et al. (2021) along with  $\zeta_M$  in Eq. (4), we computed  $d \log R_{p,\text{valley}} / d \log M_{\star\text{th}} \approx +0.23$  and  $+0.27$  for the photo-evaporation and core-powered mass-loss models, respectively. The difference with our observationally inferred estimate may suggest that other mechanisms shape the observed properties of planets orbiting M dwarfs (e.g. a significant role of gas-poor formation, for which Lee & Connors (2021) theoretically predicted a  $d \log R_{p,\text{valley}} / d \log M_\star$  down to  $+0.11$ ). However, Wu (2019) first remarked that the specific scaling relation between the planetary core mass and stellar mass further influences Eq. (4), and Rogers et al. (2021) indeed verified that the  $d \log R_{p,\text{valley}} / d \log M_\star$  may be considerably altered when these scalings are accounted for.

### 6.3. Density valley

Finally, as L22 concluded that the demographics of exoplanets can be better visualised by considering the density valley, we repeated the SVM analysis in the  $\hat{\rho}$ - $P$  space, where the normalised density  $\hat{\rho} \equiv \frac{\rho_p}{\rho_{\oplus\text{-like}}}$  and  $\rho_{\oplus\text{-like}}$  is the density that a planet of given mass would have if it had an Earth-like composition. As done by L22, we followed Zeng et al. (2019), who computed that an Earth-like planet of mass  $M_p$  has a radius  $R_{\oplus\text{-like}} = M_p^{1/3.7}$ , where

both the mass and the radius are expressed in Earth units. Therefore, according to Zeng et al. (2019), the density of an Earth-like planet scales as (Earth units)

$$\rho_{\oplus\text{-like}} = M_p^{0.7/3.7}, \quad (6)$$

which is the normalisation factor to derive  $\hat{\rho}$  from  $\rho_p$ . The density valley is shown in Fig. 9 along with the SVM-based best-fit line,

$$\log \hat{\rho}_{\text{valley}} = -0.02_{-0.04}^{+0.12} \log P - 0.313_{-0.076}^{+0.034}. \quad (7)$$

Fig. 9 confirms that the normalised density  $\hat{\rho}$  separates two different populations of exoplanets, as first pointed out by L22. Our quantitative characterisation of the valley yields a slope  $d \log \hat{\rho}_{\text{valley}} / d \log P = -0.02_{-0.04}^{+0.12}$ , which is well consistent with zero, similar to  $d \log \hat{\rho}_{\text{valley}} / d \log P_{\text{L22}} = +0.02 \pm 0.04$  estimated by L22.

The agreement of both our  $d \log R_{p,\text{valley}} / d \log P$  and  $d \log \hat{\rho}_{\text{valley}} / d \log P$  outcomes with the results from L22 may also suggest that the L22 interpretation of planet demographics may be followed. In detail, L22 identified that planets with  $R_p \lesssim 1.6 R_\oplus$  are rocky, planets with  $R_p \gtrsim 2.3 R_\oplus$  are puffy sub-Neptunes, and planets with intermediate radii are water worlds, that is, planets with the same mass content of condensed water and rocks. L22 interpreted the density gap as a division between rocky planets and water worlds, which also agrees with the conclusions by Venturini et al. (2020), who find that the radius gap separates dry from wet planets.

At lower stellar mass, the minimum mass for a planet to undergo type I migration decreases (e.g. Burn et al. 2021). As a result, water worlds are more common around M dwarfs, and their abundance shapes the topology of the radius valley, which is then determined by the favoured inward migration of water worlds rather than by atmospheric loss processes (Venturini et al. in prep.). The migration causes an overlap between rocky planets and water worlds within the mass-radius and  $R_p$ - $P$  space. Hence, the radius valley is partially filled (as also found by L22) and its slope becomes shallower than expected from thermally-driven atmospheric mass-loss mechanisms.

## 7. Conclusions

The M4 V star TOI-732 hosts two transiting planets, namely a close-in USP planet at  $P_b \sim 0.77$  d and an outer one at  $P_c \sim 12.25$  d. They straddle the radius valley and have  $R_b \sim 1.3 R_\oplus$  and  $R_c \sim 2.4 R_\oplus$ . The system has been analysed by C20, N20, and L22, but by collecting 25 CHEOPS LCs and benefiting from a further still unpublished TESS sector, we were able to double the number of space-based observations for a total of  $\sim 140$  transit events observed with both ground- and space-based facilities. Furthermore, in addition to the 127 RV data points already available in the literature, we obtained 38 RV observations with the MAROON-X spectrograph.

We jointly analysed all the available LCs and RV time series using the MCMCI routine by Bonfanti & Gillon (2020), reaching a transit depth precision of 4.4% (resp. 4.0%) and an RV semi-amplitude precision of 6.2% (3.8%) for TOI-732 b (TOI-732 c). Even with respect to the most recent parameters available in the literature, we were able to improve the precision on the transit and RV observables up to a factor  $\sim 2.4$ , with a remarkably positive impact on the mean densities of both planets. We estimated  $\rho_b = 5.8_{-0.8}^{+1.0} \text{ g cm}^{-3}$  and  $\rho_c = 3.24_{-0.43}^{+0.55} \text{ g cm}^{-3}$  (hence  $\sim 15\%$  uncertainty for both), and only  $\sim 20\%$  of the currently known exoplanets around M dwarf are known with a comparable or better precision according to the NASA Exoplanet Archive.




Fig. 9: Normalised density as a function of orbital period. The normalised density is the mean density of the planet divided by the density the planet would have if it had an Earth-like composition (same variable as introduced in L22 to display the density valley). The grey line and its shaded area indicate the density valley with its corresponding error, as in Fig.6.

Based on the internal structure modelling we performed, TOI-732 b probably does not host any gaseous envelope, but it is fully compatible with a rocky composition. Instead, TOI-732 c is compatible with having a volatile layer, with our interior structure model yielding a H-He envelope mass  $M_{\text{gas,c}} = 0.02^{+0.05}_{-0.02} M_{\oplus}$ , which corresponds to a thickness of  $R_{\text{gas,c}} = 0.40^{+0.24}_{-0.27} R_{\oplus}$ . However, based on the Aguihine et al. (2021) models, the mass and radius values of TOI-732 c are also compatible with an Earth-like core surrounded by a steam water layer. From the physical parameters of the planets, we then infer that the inner planet is a super-Earth, while the outer planet is a sub-Neptune. This constitutes a quite common system architecture in the exoplanet field.

We finally built a sample of well-characterised M-dwarf exoplanets (the  $M_{\text{sample}}$ ) with  $R_p < 4 R_{\oplus}$  and whose radii and masses are known to better than 8% and 25%, respectively. After this, we investigated the slopes of the radius valley as a function of the planet orbital periods and of the host stellar mass because theoretical models predict different trends depending on the mechanisms that have underlain planet formation and evolution. Following an SVM approach (e.g. Cortes & Vapnik 1995), we determined a  $d \log R_{p,\text{valley}}/d \log P = -0.065^{+0.024}_{-0.013}$ , differing by  $\sim 2\sigma$  from the  $d \log R_{p,\text{valley}}/d \log P_{\text{H23}} = -0.11 \pm 0.02$  slope derived by Ho & Van Eylen (2023) when targeting FGK stars, which may imply that formation and evolution mechanisms are at play with different weights in FGK and M-dwarf exoplanet systems.

Theoretical predictions would associate a  $d \log R_{p,\text{valley}}/d \log P_{\text{TD}} = -0.10$  with a thermally driven mass-loss scenario (Affolter et al. 2023), while Lopez & Rice (2018) computed an upper limit for the radius valley slope ( $d \log R_{p,\text{valley}}/d \log P_{\text{L18}} = +0.11$ ) derived from gas-empty planet formation models. As our result falls in between, with a negative slope, we may argue that thermally driven mass-loss events can explain the evolution of the majority of M-dwarf exoplanets, but some of the planets in our  $M_{\text{sample}}$  may be compatible with the gas-poor formation scenario. This type of

formation mechanism has recently been invoked to justify the physical properties of some exoplanets hosted by M dwarfs, such as the cases of TOI-1634 b (Cloutier et al. 2021b) or LHS 1903 e (Wilson et al. 2023).

An alternative explanation for the observed radius valley topology instead relies on the abundance of water worlds around low-mass stars. In particular, the favoured inward migration of water worlds suggested by simulations (Venturini et al. 2020; Burn et al. 2021) would cause a partial filling of the radius valley (L22; Venturini et al., in prep.). The radius-valley slope would then become flatter compared to what is theoretically expected if only thermally driven mass-loss mechanisms were at play; this agrees with our  $d \log R_{p,\text{valley}}/d \log P$  estimate. Following L22, we further confirm the presence of a density valley that better separates rocky and water-rich exoplanets. By repeating the SVM analysis in the  $(\hat{\rho}, P)$  plane, we computed a slope of  $d \log \hat{\rho}_{\text{valley}}/d \log P = -0.02^{+0.12}_{-0.04}$ , which agrees well with the  $d \log \hat{\rho}_{\text{valley}}/d \log P_{\text{L22}} = +0.02 \pm 0.04$  value found by L22. Therefore, the interpretation of L22 in terms of planet demographics and formation can be adopted here as well. In summary, when comparing theoretical predictions of  $d \log R_{p,\text{valley}}/d \log P$  values (see Tab. 7) with our findings, our  $d \log R_{p,\text{valley}}/d \log P = -0.065^{+0.024}_{-0.013}$  estimate can be justified by invoking further mechanisms (e.g. gas-poor formation or inward migration) in addition to thermally driven mass-loss phenomena.

*Acknowledgements.* We thank the anonymous referee for the valuable comments, which improve the quality of the manuscript. CHEOPS is an ESA mission in partnership with Switzerland with important contributions to the payload and the ground segment from Austria, Belgium, France, Germany, Hungary, Italy, Portugal, Spain, Sweden, and the United Kingdom. The CHEOPS Consortium would like to gratefully acknowledge the support received by all the agencies, offices, universities, and industries involved. Their flexibility and willingness to explore new approaches were essential to the success of this mission. CHEOPS data analysed in this article will be made available in the CHEOPS mission archive ([https://cheops.unige.ch/archive\\_browser/](https://cheops.unige.ch/archive_browser/)). TWi acknowledges support from the UKSA and the University of Warwick. This work has been carried out within the framework of the NCCR PlanetS supported by the Swiss National Science Foundation under grants 51NF40\_182901 and 51NF40\_205606. ABr was supported by the SNSA. S.G.S. acknowledge support from FCT through FCT contract nr. CEECIND/00826/2018 and POPH/FSE (EC). ML acknowledges support of the Swiss National Science Foundation under grant number PCEFP2\_194576. ASi and CBR acknowledge support from the Swiss Space Office through the ESA PRODEX program. This work was also partially supported by a grant from the Simons Foundation (PI Queloz, grant number 327127). YAI acknowledges support from the Swiss National Science Foundation (SNSF) under grant 200020\_192038. NCSa acknowledges funding by the European Union (ERC, FIERCE, 101052347). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them. V.V.G. is an F.R.S-FNRS Research Associate. RAl, DBa, EPa, and IRi acknowledge financial support from the Agencia Estatal de Investigación of the Ministerio de Ciencia e Innovación MCIN/AEI/10.13039/501100011033 and the ERDF “A way of making Europe” through projects PID2019-107061GB-C61, PID2019-107061GB-C66, PID2021-125627OB-C31, and PID2021-125627OB-C32, from the Centre of Excellence “Severo Ochoa” award to the Instituto de Astrofísica de Canarias (CEX2019-000920-S), from the Centre of Excellence “María de Maeztu” award to the Institut de Ciències de l’Espai (CEX2020-001058-M), and from the Generalitat de Catalunya/CERCA programme. S.C.C.B. acknowledges support from FCT through FCT contracts nr. IF/01312/2014/CP1215/CT0004. LBo, VNa, IPa, GPI, RRA, GSc, and TZi acknowledge support from CHEOPS ASI-INAF agreement n. 2019-29-HH.0. ACC acknowledges support from STFC consolidated grant numbers ST/R000824/1 and ST/V000861/1, and UKSA grant number ST/R003203/1. P.E.C. is funded by the Austrian Science Fund (FWF) Erwin Schroedinger Fellowship, program J4595-N. This project was supported by the CNES. The Belgian participation to CHEOPS has been supported by the Belgian Federal Science Policy Office (BELSPO) in the framework of the PRODEX Program, and by the University of Liège through an ARC grant for Concerted Research Actions financed by the Wallonia-Brussels Federation. L.D. is an F.R.S-FNRS Postdoctoral Researcher. This work was supported by FCT - Fundação para a Ciência e a Tecnologia through national funds and by FEDER through COMPETE2020 through the research

- 1036 grants UIDB/04434/2020, UIDP/04434/2020, 2022.06962.PTDC. O.D.S.D. is  
 1037 supported in the form of work contract (DL 57/2016/CP1364/CT0004) funded  
 1038 by national funds through FCT. B.-O. D. acknowledges support from the Swiss  
 1039 State Secretariat for Education, Research and Innovation (SERI) under contract  
 1040 number MB22.00046. This project has received funding from the European Re-  
 1041 search Council (ERC) under the European Union’s Horizon 2020 research and  
 1042 innovation programme (project FOUR ACES. grant agreement No 724427). It has  
 1043 also been carried out in the frame of the National Centre for Competence in  
 1044 Research PlanetS supported by the Swiss National Science Foundation (SNSF).  
 1045 DE acknowledges financial support from the Swiss National Science Founda-  
 1046 tion for project 200021\_200726. MF and CMP gratefully acknowledge the sup-  
 1047 port of the Swedish National Space Agency (DNR 65/19, 174/18). DG grate-  
 1048 fully acknowledges financial support from the CRT foundation under Grant No.  
 1049 2018.2323 “Gaseous rocky? Unveiling the nature of small worlds”. M.G. is  
 1050 an F.R.S.-FNRS Senior Research Associate. MNG is the ESA CHEOPS Project  
 1051 Scientist and Mission Representative, and as such also responsible for the Guest  
 1052 Observers (GO) Programme. MNG does not relay proprietary information be-  
 1053 tween the GO and Guaranteed Time Observation (GTO) Programmes, and does  
 1054 not decide on the definition and target selection of the GTO Programme. CHe  
 1055 acknowledges support from the European Union H2020-MSCA-ITN-2019 un-  
 1056 der Grant Agreement no. 860470 (CHAMELEON). SH gratefully acknowl-  
 1057 edges CNES funding through the grant 837319. KGI is the ESA CHEOPS  
 1058 Project Scientist and is responsible for the ESA CHEOPS Guest Observers Pro-  
 1059 gramme. She does not participate in, or contribute to, the definition of the Guar-  
 1060 anteed Time Programme of the CHEOPS mission through which observations  
 1061 described in this paper have been taken, nor to any aspect of target selection  
 1062 for the programme. K.W.F.L. was supported by Deutsche Forschungsgemein-  
 1063 schaft grants RA714/14-1 within the DFG Schwerpunkt SPP 1992, Exploring  
 1064 the Diversity of Extrasolar Planets. This work was granted access to the HPC  
 1065 resources of MesoPSL financed by the Région Ile de France and the project  
 1066 Equip@Meso (reference ANR-10-EQPX-29-01) of the programme Investisse-  
 1067 ments d’Avenir supervised by the Agence Nationale pour la Recherche. PM  
 1068 acknowledges support from STFC research grant number ST/M001040/1. GS  
 1069 acknowledges support provided by NASA through the NASA Hubble Fellow-  
 1070 ship grant HST-HF2-51519.001-A awarded by the Space Telescope Science In-  
 1071 stitute, which is operated by the Association of Universities for Research in As-  
 1072 tronomy, Inc., for NASA, under contract NAS5-26555. GyMSz acknowledges  
 1073 the support of the Hungarian National Research, Development and Innovation  
 1074 Office (NKFIH) grant K-125015, a PRODEX Experiment Agreement No.  
 1075 4000137122, the Lendület LP2018-7/2021 grant of the Hungarian Academy of  
 1076 Science and the support of the city of Szombathely. NAW acknowledges UKSA  
 1077 grant ST/R004838/1. R.L. acknowledges funding from University of La Laguna  
 1078 through the Margarita Salas Fellowship from the Spanish Ministry of Universi-  
 1079 ties ref. UNI/551/2021-May 26, and under the EU Next Generation funds. This  
 1080 work has made use of data from the European Space Agency (ESA) mission *Gaia*  
 1081 (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Process-  
 1082 ing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by na-  
 1083 tional institutions, in particular the institutions participating in the *Gaia* Mul-  
 1084 tilateral Agreement. This article is based on observations made with the MuS-  
 1085 CAT2 instrument, developed by ABC, at Telescopio Carlos Sánchez operated on  
 1086 the island of Tenerife by the IAC in the Spanish Observatorio del Teide. This  
 1087 work is partly financed by the Spanish Ministry of Economics and Competitiveness  
 1088 through grants PGC2018-098153-B-C31. This work is partly supported by  
 1089 JSPS KAKENHI Grant Numbers JP17H04574, JP18H01265, and JP18H05439,  
 1090 Grant-in-Aid for JSPS Fellows Grant Number JP20J21872, JST PRESTO Grant  
 1091 Number JPMJPR1775, and a University Research Support Grant from the Na-  
 1092 tional Astronomical Observatory of Japan (NAOJ). This material is based upon  
 1093 work supported by the National Science Foundation Graduate Research Fellow-  
 1094 ship under Grant No. DGE 1746045.
- 1096 **References**
- 1097 Abdurro’uf, Accetta, K., Aerts, C., et al. 2022, *ApJS*, 259, 35  
 1098 Adibekyan, V., Dorn, C., Sousa, S. G., et al. 2021, *Science*, 374, 330  
 1099 Adibekyan, V. Z., Sousa, S. G., Santos, N. C., et al. 2012, *A&A*, 545, A32  
 1100 Affolter, L., Mordasini, C., Oza, A. V., Kubyskhina, D., & Fossati, L. 2023,  
 1101 *A&A*, 676, A119  
 1102 Agol, E., Dorn, C., Grimm, S. L., et al. 2021, *PSJ*, 2, 1  
 1103 Aguichine, A., Mousis, O., Deleuil, M., & Marcq, E. 2021, *ApJ*, 914, 84  
 1104 Alibert, Y. 2017, *A&A*, 606, A69  
 1105 Alibert, Y., Carron, F., Fortier, A., et al. 2013, *A&A*, 558, A109  
 1106 Allard, F. 2014, in *Exploring the Formation and Evolution of Planetary Systems*,  
 1107 ed. M. Booth, B. C. Matthews, & J. R. Graham, Vol. 299, 271–272  
 1108 Almeida-Fernandes, F. & Rocha-Pinto, H. J. 2018, *MNRAS*, 476, 184  
 1109 Almenara, J. M., Bonfils, X., Otegi, J. F., et al. 2022, *A&A*, 665, A91  
 1110 Anglada-Escudé, G. & Butler, R. P. 2012, *ApJS*, 200, 15  
 Antoniadis-Karnavas, A., Sousa, S. G., Delgado-Mena, E., et al. 2020, *A&A*, 1111  
 636, A9 1112  
 Astudillo-Defru, N., Cloutier, R., Wang, S. X., et al. 2020, *A&A*, 636, A58 1113  
 Barkaoui, K., Burdanov, A., Hellier, C., et al. 2019, *AJ*, 157, 43 1114  
 Barkaoui, K., Timmermans, M., Soubkiou, A., et al. 2023, *A&A*, 677, A38 1115  
 Barnes, R. 2017, *Celestial Mechanics and Dynamical Astronomy*, 129, 509 1116  
 Bastian, N., Covey, K. R., & Meyer, M. R. 2010, *ARA&A*, 48, 339 1117  
 Bean, J. L., Seifahrt, A., Hartman, H., et al. 2010, *ApJ*, 713, 410 1118  
 Ben-Hur, A., Horn, D., Siegelmann, H. T., & Vapnik, V. 2002, *J. Mach. Learn.* 1119  
*Res.*, 2, 125–137 1120  
 Benedict, G. F., Henry, T. J., Franz, O. G., et al. 2016, *AJ*, 152, 141 1121  
 Benneke, B., Wong, I., Piaulet, C., et al. 2019, *ApJ*, 887, L14 1122  
 Benz, W., Broeg, C., Fortier, A., et al. 2021, *Experimental Astronomy*, 51, 109 1123  
 Berger, T. A., Huber, D., Gaidos, E., van Saders, J. L., & Weiss, L. M. 2020, *AJ*, 1124  
 160, 108 1125  
 Berger, T. A., Schlieder, J. E., Huber, D., & Barclay, T. 2023, arXiv e-prints, 1126  
 arXiv:2302.00009 1127  
 Blackwell, D. E. & Shallis, M. J. 1977, *MNRAS*, 180, 177 1128  
 Bluhm, P., Luque, R., Espinoza, N., et al. 2020, *A&A*, 639, A132 1129  
 Bonfanti, A., Delrez, L., Hooton, M. J., et al. 2021, *A&A*, 646, A157 1130  
 Bonfanti, A. & Gillon, M. 2020, *A&A*, 635, A6 1131  
 Bonfanti, A., Ortolani, S., & Nascimbeni, V. 2016, *A&A*, 585, A5 1132  
 Bonfanti, A., Ortolani, S., Piotto, G., & Nascimbeni, V. 2015, *A&A*, 575, A18 1133  
 Bonfils, X., Almenara, J. M., Cloutier, R., et al. 2018, *A&A*, 618, A142 1134  
 Bonfils, X., Forveille, T., Delfosse, X., et al. 2005, *A&A*, 443, L15 1135  
 Brandeker, A., Heng, K., Lendl, M., et al. 2022, *A&A*, 659, L4 1136  
 Brown, T. M., Baliber, N., Bianco, F. B., et al. 2013, *PASP*, 125, 1031 1137  
 Burn, R., Schlecker, M., Mordasini, C., et al. 2021, *A&A*, 656, A72 1138  
 Burt, J. A., Dragomir, D., Mollière, P., et al. 2021, *AJ*, 162, 87 1139  
 Butler, R. P., Vogt, S. S., Marcy, G. W., et al. 2004, *ApJ*, 617, 580 1140  
 Cañas, C. I., Kanodia, S., Libby-Roberts, J., et al. 2023, *AJ*, 166, 30 1141  
 Cale, B., Plavchan, P., LeBrun, D., et al. 2019, *AJ*, 158, 170 1142  
 Casagrande, L., Schönrich, R., Asplund, M., et al. 2011, *A&A*, 530, A138 1143  
 Castelli, F. & Kurucz, R. L. 2003, in *IAU Symposium*, Vol. 210, *Modelling of* 1144  
*Stellar Atmospheres*, ed. N. Piskunov, W. W. Weiss, & D. F. Gray, A20 1145  
 Castro-González, A., Demangeon, O. D. S., Lillo-Box, J., et al. 2023, *A&A*, 675,  
 A52 1147  
 Charbonneau, D., Irwin, J., Nutzman, P., & Falco, E. E. 2008, in *American As-* 1148  
*tronomical Society Meeting Abstracts*, Vol. 212, *American Astronomical So-* 1149  
*cety Meeting Abstracts #212*, 44.02 1150  
 Chaturvedi, P., Bluhm, P., Nagel, E., et al. 2022, *A&A*, 666, A155 1151  
 Chen, H. & Rogers, L. A. 2016, *ApJ*, 831, 180 1152  
 Cherubim, C., Cloutier, R., Charbonneau, D., et al. 2023, *AJ*, 165, 167 1153  
 Childs, A. C., Martin, R. G., & Livio, M. 2022, *ApJ*, 937, L41 1154  
 Ciardi, D. R., Fabrycky, D. C., Ford, E. B., et al. 2013, *ApJ*, 763, 41 1155  
 Cloutier, R., Astudillo-Defru, N., Bonfils, X., et al. 2019, *A&A*, 629, A111 1156  
 Cloutier, R., Charbonneau, D., Deming, D., Bonfils, X., & Astudillo-Defru, N. 1157  
 2021a, *AJ*, 162, 174 1158  
 Cloutier, R., Charbonneau, D., Stassun, K. G., et al. 2021b, *AJ*, 162, 79 1159  
 Cloutier, R., Eastman, J. D., Rodriguez, J. E., et al. 2020, *AJ*, 160, 3 1160  
 Cloutier, R. & Menou, K. 2020, *AJ*, 159, 211 1161  
 Cointepas, M., Almenara, J. M., Bonfils, X., et al. 2021, *A&A*, 650, A145 1162  
 Cortes, C. & Vapnik, V. 1995, *Machine Learning*, 20, 273 1163  
 Cosentino, R., Bonanno, G., Bruno, P., et al. 2000, *Mem. Soc. Astron. Italiana*, 1164  
 71, 393 1165  
 Cosentino, R., Lovis, C., Pepe, F., et al. 2012, in *Society of Photo-Optical Instru-* 1166  
*mentation Engineers (SPIE) Conference Series*, Vol. 8446, *Ground-based and* 1167  
*Airborne Instrumentation for Astronomy IV*, ed. I. S. McLean, S. K. Ramsay,  
 & H. Takami, 84461V 1168  
 Crossfield, I. J. M., Malik, M., Hill, M. L., et al. 2022, *ApJ*, 937, L17 1170  
 Cuntz, M. & Wang, Z. 2018, *Research Notes of the American Astronomical* 1171  
*Society*, 2, 19 1172  
 Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, *VizieR Online Data* 1173  
*Catalog*, II/246 1174  
 Delfosse, X., Forveille, T., Mayor, M., et al. 1998, *A&A*, 338, L67 1175  
 Demangeon, O. D. S., Zapatero Osorio, M. R., Alibert, Y., et al. 2021a, *A&A*, 1176  
 653, A41 1177  
 Demangeon, O. D. S., Zapatero Osorio, M. R., Alibert, Y., et al. 2021b, *A&A*, 1178  
 653, A41 1179  
 Diamond-Lowe, H., Kreidberg, L., Harman, C. E., et al. 2022, *AJ*, 164, 172 1180  
 Dorn, C., Venturini, J., Khan, A., et al. 2017, *A&A*, 597, A37 1181  
 Dransfield, G., Timmermans, M., Triaud, A. H. M. J., et al. 2023, *MN-* 1182  
*RAS[arXiv:2305.06206]* 1183  
 Eggleton, P. P. 1983, *ApJ*, 268, 368 1184  
 Emshenhuber, A., Mordasini, C., Burn, R., et al. 2021, *A&A*, 656, A70 1185  
 Erkaev, N. V., Kulikov, Y. N., Lammer, H., et al. 2007, *A&A*, 472, 329 1186  
 Espinoza, N. & Jordán, A. 2015, *MNRAS*, 450, 1879 1187  
 Essack, Z., Shporer, A., Burt, J. A., et al. 2023, *AJ*, 165, 47 1188  
 Foreman-Mackey, D., Agol, E., Ambikasaran, S., & Angus, R. 2017, *AJ*, 154,  
 220 1189  
 1190

- 1191 Fridlund, M., Georgieva, I. Y., Bonfanti, A., & et al. 2023, accepted for publica- 1270  
1192 tion in A&A 1271
- 1193 Fruhwirth-Schnatter, S., Celeux, G., & Robert, C. P. 2018, Handbook of Mixture 1272  
1194 Analysis, 1st edn. (Chapman and Hall/CRC) 1273
- 1195 Fulton, B. J. & Petigura, E. A. 2018, AJ, 156, 264 1274
- 1196 Fulton, B. J., Petigura, E. A., Howard, A. W., et al. 2017, AJ, 154, 109 1275
- 1197 Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2023, A&A, 674, A1 1276
- 1198 Gelman, A. & Rubin, D. B. 1992, Statistical Science, 7, 457 1277
- 1199 Ghachoui, M., Soubkiou, A., Wells, R. D., et al. 2023, A&A, 677, A31 1278
- 1200 Gillon, M., Pont, F., Demory, B. O., et al. 2007, A&A, 472, L13 1279
- 1201 Ginzburg, S., Schlichting, H. E., & Sari, R. 2018, MNRAS, 476, 759 1280
- 1202 Gupta, A. & Schlichting, H. E. 2019, MNRAS, 487, 24 1281
- 1203 Gupta, A. & Schlichting, H. E. 2020, MNRAS, 493, 792 1282
- 1204 Hakim, K., Rivoldini, A., Van Hoolst, T., et al. 2018, Icarus, 313, 61 1283
- 1205 Haldemann, J., Alibert, Y., Mordasini, C., & Benz, W. 2020, A&A, 643, A105 1284
- 1206 Haldemann, J., Dorn, C., Venturini, J., & et al. 2023, A&A in press 1285
- 1207 Hatzes, A. P. 2016, in Astrophysics and Space Science Library, Vol. 428, Meth- 1286  
1208 ods of Detecting Exoplanets: 1st Advanced School on Exoplanetary Science, 1287  
1209 ed. V. Bozza, L. Mancini, & A. Sozzetti, 3 1288
- 1210 Hirano, T., Gaidos, E., Winn, J. N., et al. 2020, ApJ, 890, L27 1289
- 1211 Hirano, T., Livingston, J. H., Fukui, A., et al. 2021, AJ, 162, 161 1290
- 1212 Ho, C. S. K. & Van Eylen, V. 2023, MNRAS, 519, 4056 1291
- 1213 Hoerl, A. E. & Kennard, R. W. 1970, Technometrics, 12, 55 1292
- 1214 Holman, M. J., Winn, J. N., Latham, D. W., et al. 2006, ApJ, 652, 1715 1293
- 1215 Hoyer, S., Guterman, P., Demangeon, O., et al. 2020, A&A, 635, A24 1294
- 1216 Huang, T., Peng, H., & Zhang, K. 2017, Statistica Sinica, 27, 147 1295
- 1217 Incha, E., Vanderburg, A., Jacobs, T., et al. 2023, MNRAS, 523, 474 1296
- 1218 Jehin, E., Gillon, M., Queloz, D., et al. 2011, The Messenger, 145, 2 1297
- 1219 Jenkins, J. M., Twicken, J. D., McCauliff, S., et al. 2016, in Society of Photo- 1298  
1220 Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9913, Soft- 1299  
1221 ware and Cyberinfrastructure for Astronomy IV, 99133E 1300
- 1222 Kaminski, A., Trifonov, T., Caballero, J. A., et al. 2018, A&A, 618, A115 1301
- 1223 Kasting, J. F., Whitmire, D. P., & Reynolds, R. T. 1993, Icarus, 101, 108 1302
- 1224 Kay, C., Opher, M., & Kornbleuth, M. 2016, ApJ, 826, 195 1303
- 1225 Kegerreis, J. A., Eke, V. R., Catling, D. C., et al. 2020, ApJ, 901, L31 1304
- 1226 Kemmer, J., Stock, S., Kossakowski, D., et al. 2020, A&A, 642, A236 1305
- 1227 Kiman, R., Faherty, J. K., Cruz, K. L., et al. 2021, AJ, 161, 277 1306
- 1228 Koppappu, R. K., Ramirez, R., Kasting, J. F., et al. 2013, ApJ, 765, 131 1307
- 1229 Kossakowski, D., Kemmer, J., Bluhm, P., et al. 2021, A&A, 656, A124 1308
- 1230 Kotani, T., Tamura, M., Nishikawa, J., et al. 2018, in Society of Photo-Optical 1309  
1231 Instrumentation Engineers (SPIE) Conference Series, Vol. 10702, Ground- 1310  
1232 based and Airborne Instrumentation for Astronomy VII, ed. C. J. Evans, 1311  
1233 L. Simard, & H. Takami, 1070211 1312
- 1234 Kubyskhina, D. 2023, IAU Symposium, 370, 103 1313
- 1235 Kubyskhina, D. & Fossati, L. 2022, A&A, 668, A178 1314
- 1236 Kubyskhina, D., Fossati, L., Erkaev, N. V., et al. 2018, ApJ, 866, L18 1315
- 1237 Kurucz, R. L. 1993, SYNTHE spectrum synthesis programs and line data (As- 1316  
1238 trophysics Source Code Library) 1317
- 1239 Lam, K. W. F., Csizmadia, S., Astudillo-Defru, N., et al. 2021, Science, 374, 1271 1318
- 1240 Lam, K. W. F., Korth, J., Masuda, K., et al. 2020, AJ, 159, 120 1319
- 1241 Lammer, H., Selsis, F., Ribas, I., et al. 2003, ApJ, 598, L121 1320
- 1242 Lee, E. J., Chiang, E., & Ormel, C. W. 2014, ApJ, 797, 95 1321
- 1243 Lee, E. J. & Connors, N. J. 2021, ApJ, 908, 32 1322
- 1244 Lee, E. J., Karalis, A., & Thorgren, D. P. 2022, ApJ, 941, 186 1323
- 1245 Leleu, A., Alibert, Y., Hara, N. C., et al. 2021, A&A, 649, A26 1324
- 1246 Lillo-Box, J., Figueira, P., Leleu, A., et al. 2020, A&A, 642, A121 1325
- 1247 Lindegren, L., Bastian, U., Biermann, M., et al. 2021, A&A, 649, A4 1326
- 1248 Lobo, A. H., Shields, A. L., Palubski, I. Z., & Wolf, E. 2023, ApJ, 945, 161 1327
- 1249 Lopez, E. D. & Fortney, J. J. 2013, ApJ, 776, 2 1328
- 1250 Lopez, E. D. & Fortney, J. J. 2014, ApJ, 792, 1 1329
- 1251 Lopez, E. D. & Rice, K. 2018, MNRAS, 479, 5303 1330
- 1252 Luger, R. & Barnes, R. 2015, Astrobiology, 15, 119 1331
- 1253 Luger, R., Fulton, B. J., Kunimoto, M., et al. 2022, A&A, 664, A199 1332
- 1254 Luque, R. & Pallé, E. 2022, Science, 377, 1211 1333
- 1255 Luque, R., Pallé, E., Kossakowski, D., et al. 2019, A&A, 628, A39 1334
- 1256 Luque, R., Serrano, L. M., Molaverdikhani, K., et al. 2021, A&A, 645, A41 1335
- 1257 MacDonald, M. G. 2019, MNRAS, 487, 5062 1336
- 1258 Maciel, W. J., Rodrigues, T. S., & Costa, R. D. D. 2011, A&A, 47, 401 1337
- 1259 Maldonado, J., Micela, G., Baratella, M., et al. 2020, A&A, 644, A68 1338
- 1260 Mandel, K. & Agol, E. 2002, ApJ, 580, L171 1339
- 1261 Marboeuf, U., Thiabaud, A., Alibert, Y., Cabral, N., & Benz, W. 2014, A&A, 1340  
1262 570, A36 1341
- 1263 Marcy, G. W., Butler, R. P., Vogt, S. S., Fischer, D., & Lissauer, J. J. 1998, ApJ, 1342  
1264 505, L147 1343
- 1265 Marigo, P., Girardi, L., Bressan, A., et al. 2017, ApJ, 835, 77 1344
- 1266 Martinez, C. F., Cunha, K., Ghezzi, L., & Smith, V. V. 2019, ApJ, 875, 29 1345
- 1267 Matsumura, S., Takeda, G., & Rasio, F. A. 2008, ApJ, 686, L29 1346
- 1268 Mayor, M., Pepe, F., Queloz, D., et al. 2003, The Messenger, 114, 20 1347
- Miguel, Y., Cridland, A., Ormel, C. W., Fortney, J. J., & Ida, S. 2020, MNRAS, 1270  
491, 1998 1271
- Mishra, L., Alibert, Y., Udry, S., & Mordasini, C. 2023, A&A, 670, A68 1272
- Morris, B. M., Delrez, L., Brandeker, A., et al. 2021, A&A, 653, A173 1273
- Narita, N., Fukui, A., Kusakabe, N., et al. 2019, Journal of Astronomical Tele- 1274  
scopes, Instruments, and Systems, 5, 015001 1275
- Nordström, B., Mayor, M., Andersen, J., et al. 2004, A&A, 418, 989 1276
- Nowak, G., Luque, R., Parviainen, H., et al. 2020, A&A, 642, A173 1277
- Ojha, L., Troncone, B., Buffo, J., Journaux, B., & McDonald, G. 2022, Nature 1278  
Communications, 13, 7521 1279
- Oliva, E. 2006, Memorie della Societa Astronomica Italiana Supplementi, 9, 409 1280
- Owen, J. E. & Wu, Y. 2013, ApJ, 775, 105 1281
- Owen, J. E. & Wu, Y. 2017, ApJ, 847, 29 1282
- Parviainen, H., Tingley, B., Deeg, H. J., et al. 2019, A&A, 630, A89 1283
- Pass, E. K., Charbonneau, D., Irwin, J. M., & Winters, J. G. 2022, ApJ, 936, 109 1284
- Pedregosa, F., Varoquaux, G., Gramfort, A., et al. 2011, Journal of Machine 1285  
Learning Research, 12, 2825 1286
- Pepe, F., Cristiani, S., Rebolo, R., et al. 2021, A&A, 645, A96 1287
- Peterson, M. S., Benneke, B., Collins, K., et al. 2023, Nature, 617, 701 1288
- Petigura, E. A., Rogers, J. G., Isaacson, H., et al. 2022, AJ, 163, 179 1289
- Piaulet, C., Benneke, B., Almenara, J. M., et al. 2023, Nature Astronomy, 7, 206 1290
- Pont, F., Zucker, S., & Queloz, D. 2006, MNRAS, 373, 231 1291
- Powers, L. C., Libby-Roberts, J., Lin, A. S. J., et al. 2023, AJ, 166, 44 1292
- Pozuelos, F. J., Timmermans, M., Rackham, B. V., et al. 2023, A&A, 672, A70 1293
- Quirrenbach, A., Amado, P. J., Caballero, J. A., et al. 2014, in Society of Photo- 1294  
Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9147, 1295  
Ground-based and Airborne Instrumentation for Astronomy V, ed. S. K. Ram- 1296  
say, I. S. McLean, & H. Takami, 91471F 1297
- Quirrenbach, A., Amado, P. J., Ribas, I., et al. 2018, in Society of Photo-Optical 1298  
Instrumentation Engineers (SPIE) Conference Series, Vol. 10702, Ground- 1299  
based and Airborne Instrumentation for Astronomy VII, ed. C. J. Evans, 1300  
L. Simard, & H. Takami, 107020W 1301
- Rasmussen, C. E. & Williams, C. K. I. 2005, Gaussian Processes for Machine 1302  
Learning (Adaptive Computation and Machine Learning) (The MIT Press) 1303
- Rayner, J., Tokunaga, A., Jaffe, D., et al. 2022, PASP, 134, 015002 1304
- Rayner, J., Tokunaga, A., Jaffe, D., et al. 2016, in Society of Photo-Optical In- 1305  
strumentation Engineers (SPIE) Conference Series, Vol. 9908, Ground-based 1306  
and Airborne Instrumentation for Astronomy VI, ed. C. J. Evans, L. Simard, 1307  
& H. Takami, 990884 1308
- Reddy, B. E., Lambert, D. L., & Allende Prieto, C. 2006, MNRAS, 367, 1329 1309
- Reiners, A., Basri, G., & Browning, M. 2009, ApJ, 692, 538 1310
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, Journal of Astronomical 1311  
Telescopes, Instruments, and Systems, 1, 014003 1312
- Rogers, J. G., Gupta, A., Owen, J. E., & Schlichting, H. E. 2021, MNRAS, 508, 1313  
5886 1314
- Saar, S. H. & Linsky, J. L. 1985, ApJ, 299, L47 1315
- Salmon, S. J. A. J., Van Grootel, V., Buldgen, G., Dupret, M. A., & Eggenberger, 1316  
P. 2021, A&A, 646, A7 1317
- Schanche, N., Hébrard, G., Collier Cameron, A., et al. 2020, MNRAS, 499, 428 1318
- Schmidt, S. J., Hawley, S. L., West, A. A., et al. 2015, AJ, 149, 158 1319
- Scholz, R. D., Meusinger, H., & JähreiB, H. 2005, A&A, 442, 211 1320
- Schönrich, R., Binney, J., & Dehnen, W. 2010, MNRAS, 403, 1829 1321
- Schwarz, G. 1978, Annals of Statistics, 6, 461 1322
- Schweitzer, A., Passegger, V. M., Cifuentes, C., et al. 2019, A&A, 625, A68 1323
- Scuflaire, R., Théado, S., Montalbán, J., et al. 2008, Ap&SS, 316, 83 1324
- Seifahrt, A., Bean, J. L., Kasper, D., et al. 2022, in Society of Photo-Optical 1325  
Instrumentation Engineers (SPIE) Conference Series, Vol. 12184, Ground- 1326  
based and Airborne Instrumentation for Astronomy IX, ed. C. J. Evans, J. J. 1327  
Bryant, & K. Motohara, 121841G 1328
- Seifahrt, A., Stürmer, J., Bean, J. L., & Schwab, C. 2018, in Society of Photo- 1329  
Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 10702, 1330  
Ground-based and Airborne Instrumentation for Astronomy VII, ed. C. J. 1331  
Evans, L. Simard, & H. Takami, 107026D 1332
- Sergeev, D. E., Lambert, F. H., Mayne, N. J., et al. 2020, ApJ, 894, 84 1333
- Shulyak, D., Reiners, A., Nagel, E., et al. 2019, A&A, 626, A86 1334
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, AJ, 131, 1163 1335
- Sotin, C., Grasset, O., & Mocquet, A. 2007, Icarus, 191, 337 1336
- Soto, M. G., Anglada-Escudé, G., Dreizler, S., et al. 2021, A&A, 649, A144 1337
- Sousa, S. G., Adibekyan, V., Delgado-Mena, E., et al. 2021, A&A, 656, A53 1338
- Stefansson, G., Mahadevan, S., Maney, M., et al. 2020, AJ, 160, 192 1339
- Thiabaud, A., Marboeuf, U., Alibert, Y., et al. 2014, A&A, 562, A27 1340
- Thiabaud, A., Marboeuf, U., Alibert, Y., Leya, I., & Mezger, K. 2015, A&A, 1341  
574, A138 1342
- Tilley, M. A., Segura, A., Meadows, V., Hawley, S., & Davenport, J. 2019, As- 1343  
trobiology, 19, 64 1344
- Tody, D. 1986, in Society of Photo-Optical Instrumentation Engineers (SPIE) 1345  
Conference Series, Vol. 627, Instrumentation in astronomy VI, ed. D. L. 1346  
Crawford, 733 1347

- 1348 Tody, D. 1993, in *Astronomical Society of the Pacific Conference Series*, Vol. 52,  
1349 *Astronomical Data Analysis Software and Systems II*, ed. R. J. Hanisch,  
1350 R. J. V. Brissenden, & J. Barnes, 173
- 1351 Trifonov, T., Caballero, J. A., Morales, J. C., et al. 2021, *Science*, 371, 1038
- 1352 Trifonov, T., Kürster, M., Zechmeister, M., et al. 2018, *A&A*, 609, A117
- 1353 Udry, S., Bonfils, X., Delfosse, X., et al. 2007, *A&A*, 469, L43
- 1354 Van Eylen, V., Agentoft, C., Lundkvist, M. S., et al. 2018, *MNRAS*, 479, 4786
- 1355 Van Eylen, V., Astudillo-Defru, N., Bonfils, X., et al. 2021, *MNRAS*, 507, 2154
- 1356 Venturini, J., Guilera, O. M., Haldemann, J., Ronco, M. P., & Mordasini, C. 2020,  
1357 *A&A*, 643, L1
- 1358 Watson, A. J., Donahue, T. M., & Walker, J. C. G. 1981, *Icarus*, 48, 150
- 1359 Weiss, L. M., Marcy, G. W., Petigura, E. A., et al. 2018, *AJ*, 155, 48
- 1360 West, A. A., Morgan, D. P., Bochanski, J. J., et al. 2011, *AJ*, 141, 97
- 1361 Wielen, R. 1977, *A&A*, 60, 263
- 1362 Wilson, T. G., Simpson, A. M., Collier Cameron, A., & et al. 2023, submitted to  
1363 *Science*
- 1364 Winn, J. N. 2010, in *Exoplanets*, ed. S. Seager (University of Arizona Press,  
1365 Tucson), 55–77
- 1366 Winters, J. G., Cloutier, R., Medina, A. A., et al. 2022, *AJ*, 163, 168
- 1367 Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, 140, 1868
- 1368 Wu, Y. 2019, *ApJ*, 874, 91
- 1369 Wyatt, M. C., Kral, Q., & Sinclair, C. A. 2020, *MNRAS*, 491, 782
- 1370 Zacharias, N., Finch, C. T., Girard, T. M., et al. 2012, *VizieR Online Data Cata-*  
1371 *log*, I/322A
- 1372 Zechmeister, M., Reiners, A., Amado, P. J., et al. 2018, *A&A*, 609, A12
- 1373 Zechmeister, M., Reiners, A., Amado, P. J., et al. 2020, *SERVAL: SpEc-*  
1374 *trum Radial Velocity Analyser*, *Astrophysics Source Code Library*, record  
1375 ascl:2006.011
- 1376 Zeng, L., Jacobsen, S. B., Sasselov, D. D., et al. 2019, *Proceedings of the Na-*  
1377 *tional Academy of Science*, 116, 9723
- 
- 1 Space Research Institute, Austrian Academy of Sciences, 1378  
Schmiedlstrasse 6, A-8042 Graz, Austria 1379  
e-mail: andrea.bonfanti@oeaw.ac.at 1380
- 2 Department of Astronomy & Astrophysics, University of Chicago, 1381  
Chicago, IL 60637, USA 1382
- 3 Department of Physics, University of Warwick, Gibbet Hill Road, 1383  
Coventry CV4 7AL, United Kingdom 1384
- 4 Observatoire astronomique de l’Université de Genève, Chemin Pe- 1385  
gasi 51, 1290 Versoix, Switzerland 1386
- 5 Weltraumforschung und Planetologie, Physikalisches Institut, Uni- 1387  
versity of Bern, Gesellschaftsstrasse 6, 3012 Bern, Switzerland 1388
- 6 Department of Astronomy, Stockholm University, AlbaNova Uni- 1389  
versity Center, 10691 Stockholm, Sweden 1390
- 7 Instituto de Astrofísica e Ciências do Espaço, Universidade do 1391  
Porto, CAUP, Rua das Estrelas, 4150-762 Porto, Portugal 1392
- 8 Center for Space and Habitability, University of Bern, 1393  
Gesellschaftsstrasse 6, 3012 Bern, Switzerland 1394
- 9 ETH Zurich, Department of Physics, Wolfgang-Pauli-Strasse 2, 1395  
CH-8093 Zurich, Switzerland 1396
- 10 Cavendish Laboratory, JJ Thomson Avenue, Cambridge CB3 0HE, 1397  
UK 1398
- 11 Instituto de Astrofísica de Canarias (IAC), E-38205 La Laguna, 1399  
Tenerife, Spain 1400
- 12 Departamento de Astrofísica, Universidad de La Laguna (ULL), E- 1401  
38206 La Laguna, Tenerife, Spain 1402
- 13 Lund Observatory, Dept. of Astronomy and Theoretical Physics, 1403  
Lund University, Box 43, 22100 Lund, Sweden 1404
- 14 Departamento de Física e Astronomia, Faculdade de Ciências, Uni- 1405  
versidade do Porto, Rua do Campo Alegre, 4169-007 Porto, Portugal 1406
- 15 Space sciences, Technologies and Astrophysics Research (STAR) 1407  
Institute, Université de Liège, Allée du 6 Août 19C, 4000 Liège, 1408  
Belgium 1409
- 16 Instituto de Astrofísica de Canarias, Via Lactea s/n, 38200 La La- 1410  
guna, Tenerife, Spain 1411
- 17 Departamento de Astrofísica, Universidad de La Laguna, As- 1412  
trofísico Francisco Sanchez s/n, 38206 La Laguna, Tenerife, Spain 1413
- 18 European Space Agency (ESA), European Space Research and 1414  
Technology Centre (ESTEC), Keplerlaan 1, 2201 AZ Noordwijk, 1415  
The Netherlands 1416
- 19 Admatis, 5. Kandó Kálmán Street, 3534 Miskolc, Hungary 1417
- 20 Depto. de Astrofísica, Centro de Astrobiología (CSIC-INTA), 1418  
ESAC campus, 28692 Villanueva de la Cañada (Madrid), Spain 1419
- 21 INAF, Osservatorio Astronomico di Padova, Vicolo 1420  
dell’Osservatorio 5, 35122 Padova, Italy 1421
- 22 Centre for Exoplanet Science, SUPA School of Physics and Astron- 1422  
omy, University of St Andrews, North Haugh, St Andrews KY16 1423  
9SS, UK 1424
- 23 Institute of Planetary Research, German Aerospace Center (DLR), 1425  
Rutherfordstrasse 2, 12489 Berlin, Germany 1426
- 24 INAF, Osservatorio Astrofisico di Torino, Via Osservatorio, 20, I- 1427  
10025 Pino Torinese To, Italy 1428
- 25 Centre for Mathematical Sciences, Lund University, Box 118, 221 1429  
00 Lund, Sweden 1430
- 26 Aix Marseille Univ, CNRS, CNES, LAM, 38 rue Frédéric Joliot- 1431  
Curie, 13388 Marseille, France 1432
- 27 Astrobiology Research Unit, Université de Liège, Allée du 6 Août 1433  
19C, B-4000 Liège, Belgium 1434
- 28 Centre Vie dans l’Univers, Faculté des sciences, Université de 1435  
Genève, Quai Ernest-Ansermet 30, 1211 Genève 4, Switzerland 1436
- 29 Leiden Observatory, University of Leiden, PO Box 9513, 2300 RA 1437  
Leiden, The Netherlands 1438
- 30 Department of Space, Earth and Environment, Chalmers University 1439  
of Technology, Onsala Space Observatory, 439 92 Onsala, Sweden 1440
- 31 Dipartimento di Fisica, Università degli Studi di Torino, via Pietro 1441  
Giuria 1, I-10125, Torino, Italy 1442
- 32 Department of Astrophysics, University of Vienna, Türkenschanz- 1443  
strasse 17, 1180 Vienna, Austria 1444
- 33 Institute for Theoretical Physics and Computational Physics, Graz 1445  
University of Technology, Petersgasse 16, 8010 Graz, Austria 1446
- 34 Konkoly Observatory, Research Centre for Astronomy and Earth 1447  
Sciences, 1121 Budapest, Konkoly Thege Miklós út 15-17, Hungary 1448
- 35 ELTE Eötvös Loránd University, Institute of Physics, Pázmány 1449  
Péter sétány 1/A, 1117 Budapest, Hungary 1450
- 36 IMCE, UMR8028 CNRS, Observatoire de Paris, PSL Univ., Sor- 1451  
bonne Univ., 77 av. Denfert-Rochereau, 75014 Paris, France 1452
- 37 Institut d’astrophysique de Paris, UMR7095 CNRS, Université 1453  
Pierre & Marie Curie, 98bis Blvd. Arago, 75014 Paris, France 1454
- 38 Astrophysics Group, Keele University, Keele, Staffordshire, 1455  
ST5 5BG, UK 1456


1482 **Appendix A: CHEOPS raw light curves**

Fig. A.1: Raw CHEOPS LCs in chronological order of observations from CH 1 up to CH 6, as presented in Table 3. The main systematic affecting the LCs is due to the highly variable flux pattern correlating with the spacecraft roll angle.




Fig. A.2: Same as Fig. A.1, but for CHEOPS LCs from CH 7 up to CH 12.




Fig. A.3: Same as Fig. A.1, but for CHEOPS LCs from CH 13 up to CH 18.




Fig. A.4: Same as Fig. A.1, but for CHEOPS LCs from CH 19 up to CH 24.





Fig. A.5: Same as Fig. A.1, but for CHEOPS LCs CH 25.

1484 **Appendix B: Ground-based facilities light curves**


1485

Fig. B.1: Phase-folded LCs of TOI-732 b (first column) and TOI-732 c (second column) observed by ground-based facilities in the following filters:  $g'$  (first row),  $r'$  (second row),  $i'$  (third row), and  $z'$  (fourth row).




Fig. B.2: LCs of TOI-732 c taken by ground-based facilities. From *Top to Bottom* going row wise the observation filters are B, V, R, and RG715.




Fig. B.3: LC of TOI-732 b as observed by OAA in the I filter.

1486 **Appendix C: Additional tables and figures**

Table C.1: Quadratic limb darkening (LD) coefficients ( $u_1, u_2$ ) for each photometric filter.

LD	Prior	Posterior
CHEOPS $u_1$	$\mathcal{N}(0.311, 0.054)$	$0.330^{+0.053}_{-0.054}$
CHEOPS $u_2$	$\mathcal{N}(0.383, 0.041)$	$0.387 \pm 0.042$
TESS $u_1$	$\mathcal{N}(0.208, 0.042)$	$0.217 \pm 0.043$
TESS $u_2$	$\mathcal{N}(0.415, 0.030)$	$0.418 \pm 0.032$
$g'$ $u_1$	$\mathcal{N}(0.408, 0.047)$	$0.398^{+0.048}_{-0.047}$
$g'$ $u_2$	$\mathcal{N}(0.386, 0.029)$	$0.382 \pm 0.031$
$r'$ $u_1$	$\mathcal{N}(0.444, 0.078)$	$0.434^{+0.078}_{-0.077}$
$r'$ $u_2$	$\mathcal{N}(0.326, 0.061)$	$0.324^{+0.065}_{-0.066}$
$i'$ $u_1$	$\mathcal{N}(0.310, 0.048)$	$0.320^{+0.051}_{-0.052}$
$i'$ $u_2$	$\mathcal{N}(0.346, 0.043)$	$0.345^{+0.044}_{-0.045}$
$z'$ $u_1$	$\mathcal{N}(0.162, 0.043)$	$0.153^{+0.043}_{-0.044}$
$z'$ $u_2$	$\mathcal{N}(0.439, 0.028)$	$0.437 \pm 0.030$
B $u_1$	$\mathcal{N}(0.399, 0.034)$	$0.400 \pm 0.034$
B $u_2$	$\mathcal{N}(0.401, 0.018)$	$0.401^{+0.019}_{-0.020}$
V $u_1$	$\mathcal{N}(0.400, 0.063)$	$0.393^{+0.066}_{-0.064}$
V $u_2$	$\mathcal{N}(0.386, 0.041)$	$0.385 \pm 0.045$
R $u_1$	$\mathcal{N}(0.413, 0.070)$	$0.412^{+0.072}_{-0.075}$
R $u_2$	$\mathcal{N}(0.325, 0.055)$	$0.322^{+0.059}_{-0.060}$
I $u_1$	$\mathcal{N}(0.270, 0.052)$	$0.271 \pm 0.054$
I $u_2$	$\mathcal{N}(0.379, 0.042)$	$0.377^{+0.045}_{-0.044}$
RG715 $u_1$	$\mathcal{N}(0.193, 0.040)$	$0.193 \pm 0.042$
RG715 $u_2$	$\mathcal{N}(0.427, 0.028)$	$0.426 \pm 0.030$

**Notes.**  $\mathcal{N}(\mu, \sigma)$  denotes a Normal prior with mean  $\mu$  and standard deviation  $\sigma$ .




Fig. C.1: Same as Fig. 6, but with the markers colour-coded against  $R_p T_{\text{eq}}^4$ , which correlates with the core-powered mass loss strength (Gupta & Schlichting 2019).

Table C.2: Polynomial detrending baselines applied to the space-based light curves within the MCMC scheme.

Time series	Planet	Detrending model	Time series	Planet	Detrending model
CH 1	b c	GP(roll) + $t^3 + (xy)^2$	TE 29	b	$dx^1 + (xy)^1$
CH 2	b	GP(roll) + $(xy)^2$	TE 30	b	$dx^1 + (xy)^1$
CH 3	b c	GP(roll) + smear <sup>1</sup> + $(xy)^1$	TE 31	b c	$t^1 + dx^1 + (xy)^1$
CH 4	b	GP(roll) + $t^1 + (xy)^1$	TE 32	b	$t^1 + dx^1$
CH 5	b	GP(roll) + $(xy)^2$	TE 33	b	$t^3 + dx^1$
CH 6	0	GP(roll) + sky <sup>1</sup>	TE 34	b	$t^3 + dx^1 + dy^1$
CH 7	b	GP(roll) + $(xy)^2$	TE 35	b	$dx^1$
CH 8	b	GP(roll) + $t^1 + sky^1 + (xy)^2$	TE 36	b	$dx^1$
CH 9	b c	GP(roll) + $t^1 + (xy)^2$	TE 37	b	$t^1 + dx^1 + (xy)^1$
CH 10	0	GP(roll) + $(xy)^1$	TE 38	b	$t^1 + dx^2$
CH 11	b	GP(roll) + $t^4 + (xy)^2$	TE 39	b	$dx^1$
CH 12	b	GP(roll) + $t^1$	TE 40	b	$dx^1 + (xy)^1$
CH 13	0	GP(roll) + $(xy)^2$	TE 41	b	$dx^1$
CH 14	b	GP(roll) + $t^2 + (xy)^2$	TE 42	b	$t^2 + dx^3 + (xy)^1$
CH 15	0	GP(roll) + $(xy)^2$	TE 43	b	$dx^1$
CH 16	b c	GP(roll) + $(xy)^2$	TE 44	b	$dx^1$
CH 17	b	GP(roll) + $(xy)^2$	TE 45	b	$t^1 + dx^1$
CH 18	b	GP(roll) + $(xy)^2$	TE 46	b	$t^1 + dx^1$
CH 19	b	GP(roll)	TE 47	b	$dx^1$
CH 20	0	GP(roll) + $t^2$	TE 48	b	$dx^1$
CH 21	b	GP(roll)	TE 49	b	$t^2 + dx^1 + (xy)^1$
CH 22	0	GP(roll)	TE 50	b	$dx^1$
CH 23	b	GP(roll)	TE 51	b	$t^2 + dx^1$
CH 24	0	GP(roll) + $t^2$	TE 52	b	$dx^1$
CH 25	0	GP(roll) + $t^2$	TE 53	b	$t^1 + dx^1$
TE 1	b	$dx^1$	TE 54	b	$dx^1$
TE 2	b	$dx^1$	TE 55	b	$dx^1$
TE 3	b	$dx^1$	TE 56	b c	$t^1 + dx^1$
TE 4	b c	$dx^1$	TE 57	b	$dx^1$
TE 5	b	$dx^1$	TE 58	b	$t^1 + dx^1$
TE 6	b	$c$	TE 59	b	$dx^1$
TE 7	b	$dx^1$	TE 60	b	$t^2 + dx^1$
TE 8	b	$dx^1$	TE 61	b	$t^1 + dx^1$
TE 9	b	$dx^1$	TE 62	b	$t^1 + dx^1$
TE 10	b	$c$	TE 63	b	$t^3 + dx^1$
TE 11	b	$c$	TE 64	b	$dx^1$
TE 12	b	$c$	TE 65	b	$t^1 + dx^1$
TE 13	b	$c$	TE 66	b	$dx^1$
TE 14	b	$dx^1$	TE 67	b	$t^2 + dx^1$
TE 15	b	$t^1$	TE 68	b	$t^1 + dx^1$
TE 16	b	$c$	TE 69	b	$dx^1$
TE 17	b c	$dy^1$	TE 70	b c	$t^1 + dx^1$
TE 18	b	$dx^1$	TE 71	b	$dx^1$
TE 19	b	$c$	TE 72	b	$dx^1$
TE 20	b	$c$	TE 73	b	$t^1 + dx^1 + (xy)^1$
TE 21	b	$c$	TE 74	b	$dx^1$
TE 22	b	$c$	TE 75	b	$t^1 + dx^1$
TE 23	b	$c$	TE 76	b	$dx^1$
TE 24	b	$c$	TE 77	b	$t^3 + dx^1$
TE 25	b	$c$	TE 78	b	$dx^1$
TE 26	b	$dx^1$	TE 79	b	$t^1 + dx^1$
TE 27	b	$c$	TE 80	b	$t^1 + dx^1$
TE 28	b	$c$	TE 81	b	$t^2 + dx^1$

**Notes.** CHEOPS LCs further required a GP-based pre-detrending against the roll angle, here denoted with GP(roll). The LC counter refers to the CHEOPS (CH) and TESS (TE) light curves, extracted as detailed in the text in chronological order of observations. In particular, TE LC from 1 to 28, from 29 to 52, and from 53 to 81 are extracted from Sector 9, 35, and 62, respectively. All the ground-based observations reduced as explained in the text only required a normalisation scalar ( $c$ ). See text for further details.

Table C.3: Radial velocity jitter for each RV time series as inferred from the MCMC global analysis, after applying the polynomial detrending as specified in the third column.  $c$  denotes a scalar offset; see text for further details

Instrument	RV jitter [ $\text{m s}^{-1}$ ]	Detrending
HARPS	$1.593^{+0.045}_{-0.026}$	$t^4$
IRD	$0.8391^{+0.0064}_{-0.0063}$	$c$
HARPS-N	$2.151^{+0.043}_{-0.040}$	$t^3$
CARMENES	$2.033^{+0.060}_{-0.057}$	$t^3$
iSHELL	$4.05 \pm 0.30$	$c$
MAROON-X blue Feb 2021	$1.12^{+0.21}_{-0.23}$	$c$
MAROON-X blue Apr 2021	$0.007^{+0.038}_{-0.007}$	$c$
MAROON-X blue May 2021	$0.11^{+0.16}_{-0.11}$	$c$
MAROON-X red Feb 2021	$0.07^{+0.15}_{-0.07}$	$c$
MAROON-X red Apr 2021	$0.06^{+0.20}_{-0.06}$	$c$
MAROON-X red May 2021	$0.616^{+0.052}_{-0.039}$	$d1w^1$

Table C.4: Main planetary parameters of the Msample.

Planet	$P$ [d]	$R_p$ [ $R_\oplus$ ]	$M_p$ [ $M_\oplus$ ]	$\rho_p$ [ $\rho_\oplus$ ]	$R_p$ -location	Reference
TOI-732 b	0.76837931 <sup>+0.00000039</sup> <sub>-0.00000042</sub>	1.325 <sup>+0.057</sup> <sub>-0.058</sub>	2.46 ± 0.19	1.06 <sup>+0.18</sup> <sub>-0.14</sub>	below	This work
TOI-732 c	12.252284 ± 0.000013	2.39 <sup>+0.10</sup> <sub>-0.11</sub>	8.04 <sup>+0.50</sup> <sub>-0.48</sub>	0.59 <sup>+0.10</sup> <sub>-0.08</sub>	above	This work
GJ 1132 b	1.628931 ± 0.000027	1.13 ± 0.056	1.66 ± 0.23	1.15 ± 0.23	below	Bonfils et al. (2018)
GJ 1214 b	1.58040433 ± 0.00000013	2.74 <sup>+0.050</sup> <sub>-0.053</sub>	8.17 ± 0.43	0.396 ± 0.031	above	Cloutier et al. (2021a)
GJ 1252 b	0.51824160 ± 0.00000069	1.180 ± 0.078	1.32 ± 0.28	0.80 ± 0.23	below	Crossfield et al. (2022)
GJ 3090 b	2.853136 <sup>+0.000064</sup> <sub>-0.000038</sub>	2.13 ± 0.11	3.34 ± 0.72	0.346 ± 0.092	above	Almenara et al. (2022)
GJ 3473 b	1.1980035 <sup>+0.0000018</sup> <sub>-0.0000019</sub>	1.264 <sup>+0.050</sup> <sub>-0.049</sub>	1.86 ± 0.30	0.92 ± 0.18	below	Kemmer et al. (2020)
GJ 357 b	3.93072 <sup>+0.00008</sup> <sub>-0.00006</sub>	1.217 <sup>+0.084</sup> <sub>-0.083</sub>	1.84 ± 0.31	1.02 ± 0.27	below	Luque et al. (2019)
GJ 367 b	0.321962 <sup>+0.000010</sup> <sub>-0.000012</sub>	0.718 ± 0.054	0.546 ± 0.078	1.48 ± 0.39	below	Lam et al. (2021)
GJ 486 b	1.467119 <sup>+0.000031</sup> <sub>-0.000030</sub>	1.305 <sup>+0.063</sup> <sub>-0.067</sub>	2.82 <sup>+0.11</sup> <sub>-0.12</sub>	1.27 ± 0.20	below	Trifonov et al. (2021)
HD 260655 b	2.76953 ± 0.000003	1.24 ± 0.023	2.14 ± 0.34	1.12 ± 0.19	below	Luque et al. (2022)
HD 260655 c	5.70588 ± 0.000007	1.533 <sup>+0.051</sup> <sub>-0.046</sub>	3.09 ± 0.48	0.86 ± 0.16	below	Luque et al. (2022)
K2-146 b	2.6698 ± 0.0001	2.25 ± 0.10	5.6 ± 0.7	0.492 ± 0.090	above	Lam et al. (2020)
K2-18 b	32.940045 ± 0.000010	2.61 ± 0.087	8.63 ± 1.35	0.485 ± 0.090	above	Benneke et al. (2019)
K2-25 b	3.48456408 <sup>+0.0000060</sup> <sub>-0.0000050</sub>	3.44 ± 0.12	24.5 <sup>+5.7</sup> <sub>-5.2</sub>	0.60 ± 0.15	above	Stefansson et al. (2020)
K2-3 b	10.05465350 <sup>+0.0000088</sup> <sub>-0.0000091</sub>	2.078 <sup>+0.076</sup> <sub>-0.067</sub>	5.11 <sup>+0.65</sup> <sub>-0.64</sub>	0.569 ± 0.093	above	Diamond-Lowe et al. (2022)
Kepler-138 c	13.78150 <sup>+0.00007</sup> <sub>-0.00009</sub>	1.51 ± 0.04	2.3 <sup>+0.6</sup> <sub>-0.5</sub>	0.67 ± 0.17	below	Piaulet et al. (2023)
L 168-9 b	1.40150 ± 0.00018	1.39 ± 0.09	4.6 ± 0.56	1.71 ± 0.39	below	Astudillo-Defru et al. (2020)
L 98-59 c	3.6904 ± 0.0003	1.35 ± 0.07	2.42 <sup>+0.35</sup> <sub>-0.34</sub>	0.98 ± 0.21	below	Cloutier et al. (2019)
L 98-59 d	7.4507245 <sup>+0.0000081</sup> <sub>-0.0000046</sub>	1.521 <sup>+0.12</sup> <sub>-0.10</sub>	1.94 ± 0.28	0.55 ± 0.14	below	Demangeon et al. (2021b)
LHS 1140 b	24.73694 <sup>+0.000041</sup> <sub>-0.000040</sub>	1.635 ± 0.046	6.38 <sup>+0.46</sup> <sub>-0.44</sub>	1.46 ± 0.16	below	Lillo-Box et al. (2020)
LHS 1140 c	3.77792 ± 0.00003	1.169 <sup>+0.037</sup> <sub>-0.038</sub>	1.76 <sup>+0.17</sup> <sub>-0.16</sub>	1.10 ± 0.15	below	Lillo-Box et al. (2020)
LHS 1478 b	1.9495378 <sup>+0.0000040</sup> <sub>-0.0000041</sub>	1.242 <sup>+0.051</sup> <sub>-0.049</sub>	2.33 ± 0.20	1.22 ± 0.18	below	Soto et al. (2021)
LP 791-18 c	4.9899093 <sup>+0.0000074</sup> <sub>-0.0000072</sub>	2.438 ± 0.096	7.1 ± 0.7	0.490 ± 0.075	above	Peterson et al. (2023)
LTT 1445 A b	5.3587657 <sup>+0.0000043</sup> <sub>-0.0000042</sub>	1.305 <sup>+0.066</sup> <sub>-0.061</sub>	2.87 <sup>+0.26</sup> <sub>-0.25</sub>	1.29 ± 0.22	below	Winters et al. (2022)
LTT 1445 A c	3.1239035 <sup>+0.0000034</sup> <sub>-0.0000036</sub>	1.147 <sup>+0.055</sup> <sub>-0.054</sub>	1.54 <sup>+0.20</sup> <sub>-0.19</sub>	1.02 ± 0.19	below	Winters et al. (2022)
TOI-1075 b	0.6047328 ± 0.0000032	1.791 <sup>+0.12</sup> <sub>-0.08</sub>	9.95 <sup>+1.4</sup> <sub>-1.3</sub>	1.73 ± 0.37	below	Essack et al. (2023)
TOI-1201 b	2.4919863 <sup>+0.0000030</sup> <sub>-0.0000031</sub>	2.415 <sup>+0.091</sup> <sub>-0.090</sub>	6.28 <sup>+0.84</sup> <sub>-0.88</sub>	0.446 ± 0.079	above	Kossakowski et al. (2021)
TOI-1231 b	24.245586 <sup>+0.000064</sup> <sub>-0.000066</sub>	3.65 <sup>+0.16</sup> <sub>-0.15</sub>	15.4 ± 3.3	0.317 ± 0.079	above	Burt et al. (2021)
TOI-1235 b	3.444717 <sup>+0.000040</sup> <sub>-0.000042</sub>	1.694 <sup>+0.080</sup> <sub>-0.077</sub>	5.9 <sup>+0.62</sup> <sub>-0.61</sub>	1.21 ± 0.21	below	Bluhm et al. (2020)
TOI-1634 b	0.9893436 ± 0.0000020	1.749 ± 0.079	10.14 ± 0.95	1.90 ± 0.31	below	Hirano et al. (2021)
TOI-1695 b	3.1342791 <sup>+0.0000071</sup> <sub>-0.0000063</sub>	1.9 <sup>+0.16</sup> <sub>-0.14</sub>	6.36 ± 1.0	0.93 ± 0.26	below	Cherubim et al. (2023)
TOI-244 b	7.397225 <sup>+0.000023</sup> <sub>-0.000026</sub>	1.52 ± 0.12	2.68 ± 0.3	0.76 ± 0.20	below	Castro-González et al. (2023)
TOI-269 b	3.6977104 ± 0.0000037	2.77 ± 0.12	8.8 ± 1.4	0.414 ± 0.085	above	Cointepas et al. (2021)
TOI-270 b	3.3601538 ± 0.0000048	1.206 ± 0.039	1.58 ± 0.26	0.901 ± 0.172	below	Van Eylen et al. (2021)
TOI-270 c	5.6605731 ± 0.0000031	2.355 ± 0.064	6.15 ± 0.37	0.471 ± 0.048	above	Van Eylen et al. (2021)
TOI-270 d	11.379573 ± 0.000013	2.133 ± 0.058	4.78 ± 0.43	0.493 ± 0.060	above	Van Eylen et al. (2021)
TOI-776 b	8.24661 <sup>+0.00005</sup> <sub>-0.00004</sub>	1.85 ± 0.13	4.0 ± 0.9	0.63 ± 0.19	below	Luque et al. (2021)
TRAPPIST-1 b	1.510826 ± 0.000006	1.116 <sup>+0.014</sup> <sub>-0.012</sub>	1.374 ± 0.069	0.989 ± 0.060	below	Agol et al. (2021)
TRAPPIST-1 c	2.421937 ± 0.000018	1.097 <sup>+0.014</sup> <sub>-0.012</sub>	1.308 ± 0.056	0.991 ± 0.055	below	Agol et al. (2021)
TRAPPIST-1 d	4.049219 ± 0.000026	0.788 <sup>+0.011</sup> <sub>-0.010</sub>	0.388 ± 0.012	0.793 ± 0.040	below	Agol et al. (2021)
TRAPPIST-1 e	6.101013 ± 0.000035	0.920 <sup>+0.013</sup> <sub>-0.012</sub>	0.692 ± 0.022	0.889 ± 0.046	below	Agol et al. (2021)
TRAPPIST-1 f	9.20754 ± 0.000032	1.045 <sup>+0.013</sup> <sub>-0.012</sub>	1.039 ± 0.031	0.910 ± 0.042	below	Agol et al. (2021)
TRAPPIST-1 g	12.352446 ± 0.000054	1.129 <sup>+0.015</sup> <sub>-0.013</sub>	1.321 ± 0.038	0.920 ± 0.043	below	Agol et al. (2021)
TRAPPIST-1 h	18.772866 ± 0.000021	0.755 ± 0.014	0.326 ± 0.020	0.757 ± 0.063	below	Agol et al. (2021)

Notes.  $R_p$ -location refers to the location of the exoplanets with respect to the radius valley as derived by the SVM algorithm described in Sec. 6.