



Publication Year	2018
Acceptance in OA	2020-11-06T15:11:22Z
Title	Chemo-dynamical signatures in simulated Milky Way-like galaxies
Authors	SPAGNA, Alessandro, CURIR, Anna, GIAMMARIA, MARCO, LATTANZI, Mario Gilberto, MURANTE, Giuseppe, RE FIORENTIN, Paola
Publisher's version (DOI)	10.1017/S1743921317006019
Handle	http://hdl.handle.net/20.500.12386/28202
Serie	PROCEEDINGS OF THE INTERNATIONAL ASTRONOMICAL UNION
Volume	330

Chemo-dynamical signatures in simulated Milky Way-like galaxies

Alessandro Spagna¹, Anna Curir¹, Marco Giammaria²,
Mario G. Lattanzi¹, Giuseppe Murante³ and Paola Re Fiorentin¹

¹INAF – Osservatorio Astrofisico di Torino
via Osservatorio 20, 10025 Pino Torinese, Italy
email: spagna@oato.inaf.it

²Università degli Studi di Torino, Italy

³INAF – Osservatorio Astronomico di Trieste, Italy

Abstract. We have investigated the chemo-dynamical evolution of a Milky Way-like disk galaxy, AqC4, produced by a cosmological simulation integrating a sub-resolution ISM model. We evidence a global inside-out *and* upside-down disk evolution, that is consistent with a scenario where the “thin disk” stars are formed from the accreted gas close to the galactic plane, while the older “thick disk” stars are originated *in situ* at higher heights. Also, the bar appears the most effective heating mechanism in the inner disk. Finally, no significant metallicity-rotation correlation has been observed, in spite of the presence of a negative [Fe/H] radial gradient.

Keywords. Galaxy: abundance, evolution, structure, kinematics and dynamics

In order to study the chemo-dynamical signatures related to the galactic formation processes, we have analyzed a new Λ CDM cosmological simulation, AqC4, carried out with the GADGET-3 TreePM+SPH code, where star formation, chemical evolution and stellar feedback are described using a sub-grid Multi Phase Particle Integrator (MUPPI) model. The main parameters of this simulation are listed in Table 1, while further details on the model are described by Murante *et al.* (2015).

The morphology of AqC4 at redshift $z = 0$ is well represented by the three main stellar components of a Milky Way-like galaxy (i.e. bulge/spheroid, thin and thick disk). An extensive analysis of the spatial, kinematic, and chemical properties of AqC4 is presented by Giammaria (2017), who confirmed that the 6D (\mathbf{x}, \mathbf{v}) stellar distribution is similar to that observed in the Milky Way. The main difference is the presence of a more massive bulge ($B/T=0.34$), which also causes a faster rotation velocity ($V_\phi \sim 270$ km/s at $R \simeq 5 - 10$ kpc). Moreover, the metallicity distribution of the stellar halo results a few dex higher than that of the Galactic halo.

However, the overall evolution of AqC4 appears quite consistent with similar studies based on independent simulations (e.g. Minchev *et al.* 2012, Bird *et al.* 2013, Martig *et al.* 2014, Ma *et al.* 2017).

Table 1. AqC4 parameters. M_{DM} : mass of DM particles; M_{gas} initial mass of gas particle; ϵ = smoothing parameter; M_{vir} and R_{vir} : DM mass and virial radius at redshift $z=0$; $N_{\text{DM}}, N_{\text{gas}}, N_{\text{star}}$: number of DM particles, gas particles and stellar particles within R_{vir} at $z = 0$; Ω_i : density parameters; H_0 : Hubble constant.

M_{DM} [M_\odot]	M_{gas} [M_\odot]	ϵ [kpc]	M_{vir} [M_\odot]	R_{vir} [kpc]	N_{DM}	N_{gas}	N_{star}	Ω_{m}	Ω_Λ	Ω_{b}	H_0
$2.7 \cdot 10^5$	$5.1 \cdot 10^4$	0.163	$1.49 \cdot 10^{12}$	237	5 518 587	1 348 120	6 919 646	0.25	0.75	0.04	71




Figure 1. Disk $|z|$ distributions with R between 3-5 kpc and 6-8 kpc.

Figure 1 compares the evolution of the vertical distributions at redshift 0, 1, and 2.16, that clearly evidence an *upside-down* disk evolution in the “solar” annulus, $6 \text{ kpc} < R < 8 \text{ kpc}$. Such result supports a formation scenario that firstly generates the ancient “thick disk” stars *in situ*, from an initially turbulent ISM characterized by shorter scale-lengths and higher scale-heights with respect to the younger “thin disk” stars (Haywood *et al.* 2013, Bird *et al.* 2013). Conversely, after about 6 Gyr ($z = 1$), a strong disk thickening is observed in the inner disk, $3 \text{ kpc} < R < 5 \text{ kpc}$, possibly due to the heating induced by the central bar (cfr. Grand *et al.* 2016).

Finally, although a typical negative $[\text{Fe}/\text{H}]$ radial gradient is present in the disk of AqC4, no significant rotation-metallicity relation is present, as observed in our Milky Way (Spagna *et al.* 2010, Lee *et al.* 2011; see also Schönrich & McMillan 2017). This result deserves further investigation as it may be an effect of the specific formation history of this simulation.

References

- Bird, J. C. *et al.* 2013, *ApJ*, 773, 43
 Giammaria, M. 2017, *Tesi laurea magistrale in Fisica - Ind. Astrofisica*, Turin University (Italy)
 Grand, R. J. J., Springel, V., Gómez, F. A. *et al.* 2016, *MNRAS*, 459, 199
 Haywood, M., Di Matteo, P., Lehnert, M. D. *et al.* 2013, *A&A*, 560, A109
 Lee, Y. S., Beers, T. C., An, D. *et al.* 2011, *ApJ*, 738, 187
 Ma, X., Hopkins, P. F., Wetzell, A. R. *et al.* 2017, *MNRAS*, 467, 2430
 Martig, M., Minchev, I., & Flynn, C. 2014, *MNRAS*, 442, 2474
 Minchev, I., Famaey, B., Quillen, A. C. *et al.* 2012, *A&A*, 548, A127
 Murante, G., Monaco, P., Borgani, S., *et al.* 2015, *MNRAS*, 447, 178
 Schönrich, R. & McMillan, P. J. 2017, *MNRAS*, 467, 1154
 Spagna, A., Lattanzi, M. G., Re Fiorentin, P., & Smart, R. L. 2010, *A&A*, 510, L4